

SUPPORTING INFORMATION FOR

Electronic structure and magnetic coupling of pure and Mg-doped

KCuF₃

Fausto Cargnoni,¹ Simone Cenedese,¹ Paolo Ghigna², Mario Italo Trioni¹, Marco Scavini,^{1,3}

1. Consiglio Nazionale delle Ricerche - Istituto di Scienze e Tecnologie Molecolari, Via Golgi 19, 20133 Milano - Italy
2. Dipartimento di Chimica, Università di Pavia, and Unità INSTM di Pavia, Viale Taramelli 13, 27100 Pavia – Italy
3. Dipartimento di Chimica, Università degli Studi di Milano, Via Golgi 19, 20133 Milano - Italy




FIGURE 1SI. Total and projected DOS of pure (left) and Mg doped (right) KCuF_3 . The Fermi level is set to zero. No relevant contribution from K and Mg is present in the energy range of the plots.




FIGURE 2SI. Projected density of states on copper atoms of pure KCuF₃ determined at the B3LYP level of theory for the AF1 phase for the d_{xy} (red line), d_{xz} (blue line) and d_{yz} (green line) orbitals.

Table 1SI. Magnetic coupling parameters between Cu pairs in *ab* planes (J_{ab}) and along the *c* axis (J_c) determined with the BECKE and LYP exchange and correlation functionals, respectively, without any Hartree-Fock exchange contribution.

Material	Transition	J_{ab} (K)	J_c (K)
KCuF ₃	F - AF1		-3079
KCuF ₃	AF2 - AF3		-1373
KCuF ₃	AF1 - AF3	1021	
KCuF ₃	F - AF2	737	

Table 2SI. Magnetic coupling parameters between Cu pairs in *ab* planes (J_{ab}) and along the *c* axis (J_c) determined with the BECKE and LYP exchange and correlation functionals, respectively, with a 40% Hartree-Fock exchange contribution.

Material	Transition	J_{ab} (K)	J_c (K)
KCuF ₃	F - AF1		-150
KCuF ₃	AF2 - AF3		-212
KCuF ₃	AF1 - AF3	10	
KCuF ₃	F - AF2	-39	