

## Research Article

# Quark-Antiquark Effective Potential in Symplectic Quantum Mechanics

Renato Luz <sup>1</sup>, Gustavo Petronilo <sup>1</sup>, Ademir de Santana <sup>1</sup>, Caroline Costa <sup>2</sup>,  
Ronni Amorim <sup>3,4</sup> and Rendisley Paiva <sup>3</sup>

<sup>1</sup>International Center of Physics, Instituto de Física, Universidade de Brasília, 70.910-900 Brasília, DF, Brazil

<sup>2</sup>Instituto de Física Teórica, Universidade Estadual Paulista, Rua Dr. Bento Teobaldo Ferraz, 271-Bloco II, 01140-070 São Paulo, SP, Brazil

<sup>3</sup>International Center of Physics, Universidade de Brasília, Faculdade Gama, 72.444-240 Brasília, DF, Brazil

<sup>4</sup>Canadian Quantum Research Center, 204-3002 32 Ave Vernon, BC, Canada

Correspondence should be addressed to Gustavo Petronilo; [gustavopetronilo@gmail.com](mailto:gustavopetronilo@gmail.com)

Received 29 October 2021; Revised 22 December 2021; Accepted 29 December 2021; Published 7 February 2022

Academic Editor: Ruilin Zhu

Copyright © 2022 Renato Luz et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP<sup>3</sup>.

In this paper, we study within the structure of Symplectic Quantum Mechanics a bidimensional nonrelativistic strong interaction system which represent the bound state of heavy quark-antiquark, where we consider a Cornell potential which consists of Coulomb-type plus linear potentials. First, we solve the Schrödinger equation in the phase space with the linear potential. The solution (ground state) is obtained and analyzed by means of the Wigner function related to Airy function for the  $c\bar{c}$  meson. In the second case, to treat the Schrödinger-like equation in the phase space, a procedure based on the Bohlin transformation is presented and applied to the Cornell potential. In this case, the system is separated into two parts, one analogous to the oscillator and the other we treat using perturbation method. Then, we quantized the Hamiltonian with the aid of stars operators in the phase space representation so that we can determine through the algebraic method the eigenfunctions of the undisturbed Hamiltonian (oscillator solution), and the other part of the Hamiltonian was the perturbation method. The eigenfunctions found (undisturbed plus disturbed) are associated with the Wigner function via Weyl product using the representation theory of Galilei group in the phase space. The Wigner function is analyzed, and the nonclassicality of ground state and first excited state is studied by the nonclassicality indicator or negativity parameter of the Wigner function for this system. In some aspects, we observe that the Wigner function offers an easier way to visualize the nonclassic nature of meson system than the wavefunction does phase space.

## 1. Introduction

With the discovery of the  $J/\Psi$  meson in 1974 [1], the search to describe such heavy quark systems was based on the approach of potential models. The  $J/\Psi$ , which has a mass around 3.5 times that of the proton, is the lowest bound state of a charm and anticharm quark [2]. The  $c\bar{c}$  meson appears to be an excellent device for testing QCD predictions. In quantum chromodynamics (QCD), the computation of meson properties is completely nonperturbative. As a result, direct calculation utilizing lattice QCD (LQCD) techniques is the only general method accessible. However, other strategies are effective for

heavy quarkonia. The light quarks in a meson, the mass of the bound state is much larger than the mass of the quark, move at relativistic speeds. On the other hand, the speed of the charm and bottom quarks in their respective quarkonia is tiny enough that relativistic effects in these states are greatly diminished. The velocity,  $v$ , is believed to be around  $0.3c$  for charmonia. An extension in powers of  $v^2/c^2$  can then be used to approximate the computation. This method is known as nonrelativistic QCD (NRQCD).

The masses of heavy quarkonium states can be calculated using effective potential models. The nonrelativistic motion of the quarks that make up the quarkonium state is

exploited in this technique to imply that they move in a static potential, comparable to nonrelativistic hydrogen atom models. One of the most prominent potential models is the Cornell (or funnel) potential [3]. Specifically, the system is characterized by a linear combination of the Coulomb and linear potentials. The study of these potentials should take into account the two important features of the Quantum Chromodynamics (QCD), namely, asymptotic freedom and quark confinement [4–7]. Another interest is that the Cornell potential can be used to analyze the transition between the confined and unconfined phases of matter [8–10]. This potential is of considerable importance in various branches of physics such as propagation of gravitational waves, particle and nuclear physics [11–15], mathematical modeling of the parton vibrations inside hadronic system [16–20], quantum chromodynamics, and atomic physics [5, 9, 11, 15]. It takes the form:

$$V(q) = \frac{b}{q} + aq, \quad (1)$$

where the first term is responsible for the interaction between quark and its antiquark by gluon exchange at short distances. The second term is responsible for quark confinement at large distances. It is known that the potential  $V(q)$  given in Eq. (1) reproduces two specific features of the strong interaction. The Coulomb-like term displays a property known as asymptotic freedom. This phenomenon tells us that strong interaction coupling constant is a function of momentum transfer. At short distances, the momentum transfer increases in a quark-antiquark collision. In this regard, the coupling constant becomes so small that quarks, and gluons can be considered as approximately free, and their interactions can be treated by a perturbation theory [21–26]. Over large distances, momentum transfer decreases as the strong interaction coupling constant becomes larger. On the other extreme, it features a mechanism called confinement that keeps quarks and antiquarks permanently within hadrons at 1 Fermi separations [22, 25, 27, 28]. Confinement is explained consistently in the flux tube model. The gluon field between a pair of color charges develops a string-like configuration as the distance between quarks and antiquarks is extended (flux tube). As a result, across extended distances, the energy density in the tube between the quarks maintaining the gluon field remains constant [22, 25, 29]. The energy stored in the field is then proportional to the quark separation, giving us the linear term in the potential of Eq. (1). In that way, at relatively larger separation, the potential energy can create new quarks pairs in colorless forms instead of a free quark. The confinement has been a mystery until now since no one has been able to confirm it; although, there has been recent progress using the techniques of lattice QCD [22, 25, 28].

There are several studies with the Cornell potential in literature. For instance, Flores [9, 30] studied the Schrödinger equation with the Cornell potential using variational method and supersymmetric quantum mechanics [9, 10, 30–32]. Bruni et al. [8] calculated the energy configuration for a

quark-antiquark pair from the Nambu-Goto action. It was shown that this configuration energy has the shape of a Cornell potential [8]. Khoka et al. [31, 33] utilized the analytical exact iterative method to solve  $N$ -dimensional Schrödinger equation with an extended Cornell potential [31]. They found the energy and mass spectrum of heavy quarkonia [31]. Omugbe [31] applied the Nikiforov-Uvarov method to obtain the eigensolutions of the radial Schrödinger equation with Cornell potential plus an inversely quadratic potential [11, 31, 34]. Until now, an analysis of such states in quantum phase space via the Wigner function is still lacking. In this work, we investigate confinement term of strong interaction system described by heavy quarks-antiquarks in the phase space.

The first successful formalism of quantum mechanics in phase space was introduced by Wigner in 1932 [35–57]. He was motivated by the problem of finding a way to improve the quantum statistical mechanics [42–47, 50–54, 58–70]. In the Wigner formalism, an operator  $A$  defined in Hilbert space  $\mathcal{H}$ , is associated to a function in a phase space, says  $A \longrightarrow a_W(q, p)$ . The product of two operators,  $AB$ , both defined in  $\mathcal{H}$ , is associated to a Weyl product or star product of their phase space correspondents, i.e. [61–69],

$$a_W \star b_W = a_W(q, p) e^{\frac{i\hbar}{2}\Lambda} b_W(q, p), \quad (2)$$

where  $\Lambda = \overrightarrow{\partial}_p \overleftarrow{\partial}_q - \overleftarrow{\partial}_q \overrightarrow{\partial}_p$ . Such result can be seen as the action of a star operator,  $a_W(q, p) \star$ , in the function  $b_W(q, p)$ . This fact is important for define the star operators  $a_W(q, p) \star$ , which are used to build up representations of symmetry groups in a symplectic manifold [50–52, 64]. This gives rise, for example, to the Klein-Gordon and Dirac equations written in phase space [44, 61, 62, 64]. The connection with Wigner function is derived, providing a physical interpretation for the representation. These symplectic representations provide a way to consider a perturbative approach for Wigner function on the bases of symmetry groups.

The aim of this work is to analyze the behavior of the Wigner function given in terms of Airy function for the ground state of  $c\bar{c}$  meson in the formalism of the symplectic quantum mechanics. For this, we consider only the linear potential that is a model more simple of the Cornell potential. In sequence, we studied the Schrodinger equation in the phase space to obtain the energy eigenvalues and corresponding wave functions by using the Bohlin transformation and perturbation theory for the Cornell potential. Then, we calculate the Wigner function for the heavy  $c\bar{c}$  meson ( $J\psi$ ). Consequently, we calculate the negativity parameter of the Wigner function associated to the ground state and first excited state of  $c\bar{c}$  meson that present nonclassical behavior. The study of the Wigner function for such a system is crucial in order to learn more about the system's chaotic nature.

The paper is organized as follows. In Section 2, we write the Schrödinger equation in phase space, and we perform the relation between phase space amplitude and Wigner function. In Section 3, we solve the Schrödinger equation in phase space for the quark-antiquark potential and calculate the Wigner function. In Section 4, the perturbative

method is presented in order to solve the Schrödinger equation in phase space with the Cornell potential, and the Wigner function is calculated. The conclusion is presented in Section 5.

## 2. Symplectic Quantum Mechanics: Outline and Notation

In this section, we construct a formalism to quantum mechanics in phase space. For this purpose, we introduce an Hilbert space associated to phase space, denoted by  $\mathcal{H}(\Gamma)$ . In this sense, the association of the Hilbert space  $\mathcal{H}$  with the phase space  $\Gamma$  is given by  $\int dpdq \phi^*(q, p)\phi(q, p) < \infty$  where  $\phi(p, q) \in \Gamma$ . Then, we write  $\phi(q, p) = \langle q, p | \phi \rangle$ , with the aid  $\int dpdq |q, p\rangle \langle q, p| = 1$  such that  $\langle \phi |$  is the dual vector of  $|\phi\rangle$ . This symplectic Hilbert space is denoted by  $H(\Gamma)$ . To construct a symplectic representation of quantum mechanics, we define the momentum and position operator

$$\hat{P} = p^* = p - i \frac{\partial}{\partial q}, \quad (3)$$

$$\hat{Q} = p^* = q + i \frac{\partial}{\partial p}, \quad (4)$$

which satisfy the Heisenberg commutation relation  $[\hat{Q}, \hat{P}] = i$ .

Introduced the following operators,

$$\begin{aligned} \hat{K}_i &= m\hat{Q}_i - t\hat{P}_i, \\ \hat{L}_i &= \varepsilon_{ijk}\hat{Q}_j\hat{P}_k, \\ \hat{H} &= \frac{\hat{P}^2}{2m}. \end{aligned} \quad (5)$$

We obtain the following set of commutation rules:

$$\begin{aligned} [\hat{L}_i, \hat{L}_j] &= i\varepsilon_{ijk}\hat{L}_k, \quad [\hat{L}_i, \hat{K}_j] = i\varepsilon_{ijk}\hat{K}_k, \quad [\hat{K}_i, \hat{H}] = i\hat{P}_i, \\ [\hat{L}_i, \hat{P}_j] &= i\varepsilon_{ijk}\hat{P}_k, \quad [\hat{K}_i, \hat{P}_j] = im\delta_{ij}1, \end{aligned} \quad (6)$$

with all the other commutations being zero. This is the Galilei-Lie algebra with a central extension characterized by  $m$ , where  $\hat{P}$ ,  $\hat{K}$ ,  $\hat{L}$ , and  $\hat{H}$  are the generators of translation, Galilean boosts, rotation, and time translation, respectively.

The evolution of the wave function in phase space is

$$\psi(q, p, t) = e^{\hat{H}t}\psi(q, p, 0), \quad (7)$$

derivating with respect to time, and we obtain

$$\partial_t \psi(q, p; t) = \left( \frac{\hat{P}^2}{2m} + V(\hat{Q}) \right) \psi(q, p; t). \quad (8)$$

This is the Schrodinger-like equation in phase space [54].

The association of  $\psi(q, p, t)$  with a function  $f_w$  is given by [54].

$$f_w(q, p, t) = \psi(q, p, t) * \psi^\dagger(q, p, t), \quad (9)$$

which has all the properties of the Wigner function [60]. In the next section, we solve the representation Schrödinger equation in phase space and establish the relation of amplitudes in phase space and the Wigner function for the  $c\bar{c}$  meson system.

## 3. Quark Confinement and Schrödinger Equation in Phase Space

In this section, we analyze the quark confinement in the context of phase space. For this purpose, we consider the second term of Cornell potential given in Eq. (1). Therefore, we find a solution for the Schrödinger equation in phase space with linear potential and construct the associated Wigner function. We write the Schrödinger equation in phase space with the linear potential  $V(q)$  as follows:

$$\frac{p^2}{2m} * \psi(q, p) + \lambda q * \psi(q, p) = E\psi(q, p). \quad (10)$$

And using equations (3) and (4) in (10), we get the Schrodinger equation in the form

$$\frac{1}{2m} \left( p^2 - ip\partial_q - \frac{1}{4}\partial_q^2 \right) \psi + \lambda \left( q + \frac{i}{2}\partial_p \right) \psi = E\psi, \quad (11)$$

where we used  $\hbar = 1$ . By using the transformation  $\omega = (p^2/2m) + \lambda q$ , such that

$$\frac{\partial \psi}{\partial q} = \frac{\partial \psi}{\partial \omega} \frac{\partial \omega}{\partial q} = \lambda \frac{\partial \psi}{\partial \omega}. \quad (12)$$

With this, we obtain

$$\alpha \frac{\partial^2 \psi}{\partial \omega^2} - \omega \psi = -E\psi, \quad (13)$$


where  $\alpha = \lambda^2/8m$ . Then, the Schrodinger equation is reduced to the Airy equation (49)

$$\frac{\partial^2 \psi}{\partial \omega^2} - \kappa \psi = 0, \quad (14)$$

where  $\kappa = \omega - E/\alpha$ . The solution of this equation is

$$\psi(\omega) = C_1 Ai \left[ \alpha^{\frac{2}{3}} \left( \frac{\omega}{\alpha} - \frac{E}{\alpha} \right) \right] + C_2 Bi \left[ \alpha^{\frac{2}{3}} \left( \frac{\omega}{\alpha} - \frac{E}{\alpha} \right) \right], \quad (15)$$

where  $Ai$  and  $Bi$  are homogeneous Airy functions. However,  $Bi$  goes to infinity for  $\omega \rightarrow \infty$ , and this solution is not relevant; so,  $C_2 = 0$  [71]. In this case, the solution of the Eq.

FIGURE 1: Wigner function for ground state of the  $c\bar{c}$  meson.

(14) is reduced to [71]

$$\psi(q, p) = C_1 A_i \left[ \left( \frac{\lambda^2}{8m} \right)^{-1/3} \left( \frac{p^2}{2m} + \lambda q - E \right) \right], \quad (16)$$

where  $\psi(q, p)$  is a real function. As consequence, by virtue of associativity,  $\psi \star \psi \propto \psi$ , we write the solution of Wigner function in terms of the Airy function

$$f_w(q, p) = N A_i \left[ \left( \frac{\lambda^2}{8m} \right)^{-1/3} \left( \frac{p^2}{2m} + \lambda q - E \right) \right], \quad (17)$$

where  $N$  is a normalization constant. By condition  $f_w(q, p) = 0$ , we can determine the energy levels

$$A_i \left[ \left( \frac{\lambda^2}{8m} \right)^{-1/3} \left( \frac{p^2}{2m} + \lambda q - E \right) \right] = 0. \quad (18)$$

So, the energy of system are determined by

$$E = \frac{p^2}{2m} - \alpha^{1/3} r_i, \quad (19)$$

where  $r_i$  is the zero of the Airy function. In our case, we consider only the energy for ground state

$$E = \frac{p^2}{2m} - \left( \frac{\lambda^2}{8m} \right)^{1/3} r_0. \quad (20)$$

In consequence, the normalization factor, see Ref. [71], is determined by

$$\int_0^\infty \int_{-p}^p f_w(q, p) dq dp = 1. \quad (21)$$

Then, we get

$$f_w(q, p) = N A_i \left[ \left( \frac{\lambda^2}{8m} \right)^{-1/3} \left( \frac{p^2}{2m} + \lambda q - E \right) \right], \quad (22)$$

where  $N = \alpha^{-1/3}$  and  $m$  are the reduced mass of the constituent quark and antiquark. In the sequence, we analyze the solution above of the Wigner function for  $c\bar{c}$  meson. The behavior of the Wigner function for the strong interaction of the heavy quarks is shown in (Figures 1 and 2).

For the  $c\bar{c}$  meson, we are taking into account Eq. (22). The  $c\bar{c}$  meson is shown in Figure 1 represented by Wigner function for ground state. The experimental parameter,  $\lambda$  linear confinement parameter,  $m$  reduced mass, and  $q$  maximum relative distance between quark and antiquark are all in units of  $MeV$  from  $c\bar{c}$  meson which is shown in Table 1. Figure 1 is observed for ground state of charmonium meson which the Wigner function provides values negative in the phase space. This behavior is related to the character quantum. Then, it was make cut in graph 3D as illustrated in Figure 2. Figure 2 shows that when  $=0 MeV$ , the Wigner function presents negative values, curve (a). This is because of quantum interference, also visualized in the curves (b) and (c). In particular,  $p = 0 MeV$  corresponds a limit. It is worth noting that when  $p = 0$ , the energy is  $E = 9.7 MeV$ . When varying the kinetic energy to  $= 100 MeV$ , the Wigner function is displaced to the left, curve (b). We can keep increasing the kinetic energy for  $p = 200 MeV$ , which further evidences the distance for the left of the graph of the Wigner function, curve (c). By varying the kinetic energy for  $= 400 MeV$ , it is noticed that the graph of the Wigner function is moved away visibly to the left side, curve (d). Thus, it is observed that there is a limit to the right for the existence of the Wigner function not to be zero. This limit approaches the experimental value that refers maximum relative distance  $q = 4077 MeV^{-1}$  (Table 1). Then, a boundary condition is placed at zero. Consequently, the  $c\bar{c}$  meson only exists




FIGURE 2: The curves (a)–(d) represent cut off graphics of Figure 1 for the ground state  $c\bar{c}$  meson with varying of the kinetic energy.

TABLE 1: The experimental values for linear confinement  $\lambda$ , reduced mass  $m$ , and maximum relative distance  $q$  for  $c\bar{c}$  meson [4, 72].

(a)	(b)	(c)	(d)
$p^2/2m = 0 \text{ MeV}$	$p^2/2m = 100 \text{ MeV}$	$p^2/2m = 200 \text{ MeV}$	$p^2/2m = 400 \text{ MeV}$
$\lambda = 600 \text{ MeV}^2 \text{ (Exp.)}$			
$m = 630 \text{ MeV (Exp.)}$			
$q = 4077 \text{ MeV}^{-1} \text{ (Exp.)}$			

between zero and where Wigner function decay. Note that the analysis in phase space is revealing that for the variation of the kinetic energy there is an upper limit of existence of the charm-anticharm meson (Figure 2). Although it is simple to describe the quark case through a linear potential, the confinement is observed, which is not seen with the configuration space solution.

#### 4. Cornell Potential and Schrödinger Equation in Phase Space

In this section, we analyze the quark-antiquark interaction in phase space context. For this purpose, we consider the Cornell potential given in Eq. (1). In this way, we present a solution for the Schrödinger equation with Cornell potential in phase space and the associated Wigner function. The two-dimensional Hamiltonian for the  $c\bar{c}$  meson is given as

$$H = \frac{p_{1x}^2 + p_{1y}^2}{2m} + \frac{p_{2x}^2 + p_{2y}^2}{2m} + V(r), \quad (23)$$

where  $m$  is reduced mass in units of the MeV. In order to solve the Schrödinger equation for this Hamiltonian, the Bohlin transformation is utilized. Then, the Bohlin map-

ping is defined by [52, 58, 59]

$$x + iy = (q_1^2 - q_2^2) + i(2q_1q_2), \quad (24)$$

$$x = q_1^2 - q_2^2, \quad (25)$$

$$y = 2q_1q_2, \quad (26)$$

by defining

$$P_x + iP_y = \frac{p_1 + ip_2}{2(q_1 + iq_2)}, \quad (27)$$

which leads to

$$P_x = \frac{p_1q_1 + p_2q_2}{2(q_1^2 + q_2^2)}, \quad (28)$$

$$P_y = \frac{p_2q_1 - p_1q_2}{2(q_1^2 + q_2^2)}. \quad (29)$$

Substituting Eqs. (25), (26), (28), and (29) in Eq. (23) leads to the Hamiltonian

$$H = \frac{1}{4} \left[ \frac{(p_1^2 + p_2^2)}{2m(q_1^2 + q_2^2)} \right] + \frac{b}{(q_1^2 + q_2^2)} + a(q_1^2 + q_2^2). \quad (30)$$

Consequently, the hypersurface in phase space defined by  $H = E$  leads to

$$\frac{(p_1^2 + p_2^2)}{2m} - 4E(q_1^2 + q_2^2) + 4a(q_1^2 + q_2^2)^2 = -4b, \quad (31)$$

where  $p_1$  and  $p_2$  are the canonical momenta conjugate. Eq. (31) can be written as

$$\left[ \frac{(p_1^2 + p_2^2)}{2m} - 4E(q_1^2 + q_2^2) + 4a(q_1^2 + q_2^2)^2 + 4b \right] \star \psi(q_1, p_1, q_2, p_2) = 0. \quad (32)$$

Observe that Eq. (32) is obtained from the classical Hamiltonian through of the star product. Therefore, the Bohlin mapping that leads to Eq. (31) is a classical transformation. For this reason, the perturbation theory is used to analyze the Eq. (32). So, the equation in phase space is given by

$$\left[ \widehat{H}_0 + \widehat{H}_1 \right] \star \psi(q_1, p_1, q_2, p_2) = -4b\psi(q_1, p_1, q_2, p_2), \quad (33)$$

where  $\widehat{H}_0 = (p_1^2 + p_2^2)/2 - 4E(q_1^2 + q_2^2)$  and  $\widehat{H}_1 = 4a(q_1^2 + q_2^2)^2$ .

The equation for  $\widehat{H}_0$  has the form

$$\widehat{H}_0 \psi^{(0)}(q_1, p_1, q_2, p_2) = b_{n_1, n_2}^{(0)} \psi^{(0)}(q_1, p_1, q_2, p_2), \quad (34)$$

where  $\psi^{(0)}(q_1, p_1, q_2, p_2)$  corresponds the eigenfunction the unperturbed Hamiltonian.

Defining the operators

$$\begin{aligned} \widehat{a} &= \sqrt{\frac{mW}{2k}} \left( q_1 \star + \frac{ip_1 \star}{mW} \right), \\ \widehat{a}^\dagger &= \sqrt{\frac{mW}{2k}} \left( q_1 \star - \frac{ip_1 \star}{mW} \right), \\ \widehat{b} &= \sqrt{\frac{mW}{2k}} \left( q_2 \star + \frac{ip_2 \star}{mW} \right), \\ \widehat{b}^\dagger &= \sqrt{\frac{mW}{2k}} \left( q_2 \star - \frac{ip_2 \star}{mW} \right), \end{aligned} \quad (35)$$

where  $-4E = mW^2/2$  and the star operators  $q_i \star$ ,  $p_i \star$  are given by

$$\begin{aligned} q_i \star &= q_i + \frac{i}{2} \frac{\partial}{\partial p_i}, \\ p_i \star &= p_i - \frac{i}{2} \frac{\partial}{\partial q_i}. \end{aligned} \quad (36)$$

So, the Hamiltonian is

$$\widehat{H} = \frac{(p_1^2 + p_2^2)}{2} - 4E(q_1^2 + q_2^2) + 4a(q_1^2 + q_2^2)^2. \quad (37)$$

The unperturbed Hamiltonian is defined as

$$\widehat{H}_0 = \frac{(p_1^2 + p_2^2)}{2} - 4E(q_1^2 + q_2^2), \quad (38)$$

and the perturbed part is

$$\widehat{H}_1 = 4a(q_1^2 + q_2^2)^2. \quad (39)$$

The equation that is to be analyzed is given by

$$\widehat{H} \star \psi(q_1, p_1, q_2, p_2) = -4b\psi(q_1, p_1, q_2, p_2). \quad (40)$$

The unperturbed equation is

$$\widehat{H}_0 \psi^{(0)}(q_1, p_1, q_2, p_2) = b_{n_1, n_2}^{(0)} \psi^{(0)}(q_1, p_1, q_2, p_2). \quad (41)$$

The unperturbed part  $\widehat{H}_0$  has solutions given by

$$\psi_{n_1, n_2}^{(0)}(q_1, p_1, q_2, p_2) = \Phi_{n_1}(q_1, p_1) \Gamma_{n_2}(q_2, p_2), \quad (42)$$

where  $\Phi_{n_1}(q_1, p_1)$  and  $\Gamma_{n_2}(q_2, p_2)$  are solutions. The acting operators in the states vectors is

$$\widehat{a} \Phi_{n_1} = \sqrt{n_1} \Phi_{n_1-1}, \quad (43)$$

$$\widehat{a}^\dagger \Phi_{n_1} = \sqrt{n_1 + 1} \Phi_{n_1+1}, \quad (44)$$

$$\widehat{b} \Gamma_{n_2} = \sqrt{n_2} \Gamma_{n_2-1}, \quad (45)$$

$$\widehat{b}^\dagger \Gamma_{n_2} = \sqrt{n_2 + 1} \Gamma_{n_2+1}. \quad (46)$$


Utilizing the relations  $\widehat{a} \Phi_0 = 0$ ,  $\widehat{b} \Gamma_0 = 0$ , the ground state solution is

$$\psi_{0,0}^{(0)}(q_1, p_1, q_2, p_2) = N e^{-(q_1^2 + p_1^2)} L_{n_1}(q_2^2 + p_2^2) e^{-(q_2^2 + p_2^2)} L_{n_2}(q_2^2 + p_2^2), \quad (47)$$

where  $L_{n_1}$  and  $L_{n_2}$  are Laguerre polynomials, and  $N$  is a normalisation constant. The eigenvalue solutions obtained of Eq. (41) are

$$b_{0,0}^{(1)} = \frac{kW}{4} (n_1 + n_2 + 1) + \frac{4ak^2}{mW^2}. \quad (48)$$

The states excited are obtained from Eq. (47) using operators given in Eqs. (43)–(46). So, the solution for the first-



(a) Wigner function for ground state  $n=0$  of the  $c\bar{c}$  meson (b) Wigner function for first excited state  $n=1$  of the  $c\bar{c}$  meson

FIGURE 3: Wigner function for the energy levels  $n=0$  and 1 for the  $c\bar{c}$  meson.

order perturbed is given by

$$\begin{aligned} \psi_{n_1, n_2}^{(1)}(q_1, p_1, q_2, p_2) &= \psi_{n_1, n_2}^{(0)}(q_1, p_1, q_2, p_2) \\ &+ \sum_{\substack{m_1 \neq n_1, \\ m_2 \neq n_2}} \frac{\langle \psi_{m_1, m_2}^{(0)}(q_1, p_1, q_2, p_2) | \widehat{H}_1 | \psi_{n_1, n_2}^{(0)}(q_1, p_1, q_2, p_2) \rangle}{b_{n_1, n_2}^{(0)} - b_{m_1, m_2}^{(0)}} \\ &\times \psi_{m_1, m_2}^{(0)}(q_1, p_1, q_2, p_2). \end{aligned} \quad (49)$$

Note that it is required to solve the following expression before a solution for Eq. (49).

$$I = \left\langle \psi_{m_1, m_2}^{(0)}(q_1, p_1, q_2, p_2) \left| \frac{a}{W^2} \left[ (a^\dagger + a)^2 + (b^\dagger + b)^2 \right]^2 \right| \psi_{n_1, n_2}^{(0)}(q_1, p_1, q_2, p_2) \right\rangle. \quad (50)$$

Using the orthogonality relations,

$$\begin{aligned} \langle \Phi_n^*(q_1, p_2) | \Phi_m(q_1, p_1) \rangle &= \delta_{n,m}, \\ \langle \Gamma_n^*(q_1, p_2) | \Gamma_m(q_1, p_1) \rangle &= \delta_{n,m}. \end{aligned} \quad (51)$$

The ground state of  $c\bar{c}$  meson is

$$\psi_{0,0}^{(1)} = \psi_{0,0}^{(0)} + \frac{a}{2W^3} \left[ -4\sqrt{2}\psi_{2,0}^{(0)} - \psi_{2,2}^{(0)} - \frac{7}{2}\sqrt{2}\psi_{0,2}^{(0)} - \frac{\sqrt{6}}{2}\psi_{0,4}^{(0)} \right]. \quad (52)$$

And for excited states of the  $c\bar{c}$  meson, the wave functions are

$$\begin{aligned} \psi_{1,0}^{(1)} &= \psi_{1,0}^{(0)} + \frac{a}{2W^3} \left[ -\frac{\sqrt{30}}{2}\psi_{5,0}^{(0)} + \frac{-1-11\sqrt{6}}{2}\psi_{3,0}^{(0)} - 3\psi_{3,2}^{(0)} - 5\sqrt{2}\psi_{1,2}^{(0)} - \frac{\sqrt{6}}{2}\psi_{1,4}^{(0)} \right], \\ \psi_{0,1}^{(1)} &= \psi_{0,1}^{(0)} + \frac{a}{2W^3} \left[ -\frac{\sqrt{30}}{2}\psi_{0,5}^{(0)} + \frac{-1-11\sqrt{6}}{2}\psi_{0,3}^{(0)} - 3\psi_{0,3}^{(0)} - 5\sqrt{2}\psi_{2,1}^{(0)} - \frac{\sqrt{6}}{2}\psi_{4,1}^{(0)} \right]. \end{aligned} \quad (53)$$

The Wigner function for the  $c\bar{c}$  meson is given by

$$f_w(q_1, p_1, q_2, p_2) = \psi_{n_1, n_2}^{(1)}(q_1, p_1, q_2, p_2) \hat{a} \psi_{n_1, n_2}^{\dagger(1)}(q_1, p_1, q_2, p_2). \quad (54)$$

To obtain the corrections of the first order of energy that is requiring to solve the matrix,

$$W = \begin{pmatrix} W_{aa} & W_{ab} \\ W_{ba} & W_{bb} \end{pmatrix}, \quad (55)$$

where  $W_{ij} = \langle \psi_i^{(0)} | H_1 | \psi_j^{(0)} \rangle$ . Hence, we get

$$W = \begin{pmatrix} \frac{25ak^2}{m^2W^2} & \frac{2ak^2}{m^2W^2} \\ \frac{17ak^2}{m^2W^2} & \frac{25ak^2}{m^2W^2} \end{pmatrix}. \quad (56)$$

The eigenvalue obtained is  $\lambda_1 = 30, 83(ak^2/m^2W^2)$  and  $\lambda_2 = 19, 17(ak^2/m^2W^2)$ . Then, the corrections of the first order of eigenvalue of Eq. (40) are

$$\begin{aligned} b_{1,0}^{(1)} &= \frac{-kW}{4}(n_1 + n_2 + 1) + 30, 83 \left( \frac{ak^2}{m^2W^2} \right), \\ b_{0,1}^{(1)} &= \frac{-kW}{4}(n_1 + n_2 + 1) + 19, 17 \left( \frac{ak^2}{m^2W^2} \right). \end{aligned} \quad (57)$$

Figures 3(a) and 3(b) show plots of the Wigner function associated to the ground state and first excited state for  $c\bar{c}$  meson. It should be noted that all plots consider the coordinates  $q_2$  and  $p_2$  constant in order to show a three-dimensional figure, while  $q_1 = q$ ,  $p_1 = p$ . In Figure 3(b), one clearly sees the presence of negative values for the Wigner function at the level excited of the  $c\bar{c}$  meson. In Figure 3(a), the Wigner function looks strictly positive to the ground state.

TABLE 2: Negativity parameter  $\eta(\psi)$  for the levels  $n_1, n_2 = 0, 1$ .

$n_1, n_2$	$\eta(\psi)$
0,0	0
0,1	0
1,0	0.230

With the Wigner function determined for the  $c\bar{c}$  meson, we can calculate the negativity parameter for this system, and this parameter is correlated to the nonclassicality of the system [73]. The results of this calculation are shown in (Table 2). As can be seen in (Table 2), the negativity parameter increases when  $n_1$  and  $n_2$  grow. The negativity parameter is a relevant tool in the context of Wigner formalism because one is related to nonclassicality of physical systems as soon as is also related to chaotic behavior. In this way, by examining the negativity parameter, it is possible to investigate elements that the traditional wavefunction formalism does not permit. Then, the study in phase space is crucial in order to understand more about the chaotic nature.

## 5. Concluding Remarks

In this paper, we have studied heavy quarkonium bound states using the Schrödinger-like equation with an approach the Cornell potential in the Symplectic Quantum Mechanics framework. In particular, this sector is characterized by a linear potential which leads to the quark confinement plus one Coulomb-type potential. This is called of Cornell potential that feature experimentally this sector of quantum chromodynamics QCD. Initially, the Wigner function was obtained for the  $c\bar{c}$  meson system. It was observed that analyzing in the phase space allows for simple model to observe the mesons confinement and why it is not seen into configuration space solution. We check out through the analysis in the phase space that due to the variation of kinetic energy, it has a maximum limit of the existence to the meson systems given by graphics obtained. After that, we have obtained the energy eigenvalues and the respective wave functions using perturbation theory. Such that the states, denoted a quasi-amplitude of probability, are associated with Wigner function by the Weyl product. Using the Wigner functions, the negativity parameter for the ground state and first excited state of the system was derived. Finally, we observe through the analysis in the phase space that due to the change of energy levels, the negativity parameter increases Table 2. Which evidenced an increase in the quantum character of the system. It was not seen in the literature recent works.

## Data Availability

Data used to support the findings of this study are included within the article.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

The authors would like to thank CNPq and CAPES of Brazil for their support in this study.

## References

- [1] A. Mirjalili and M. Taki, "Noncommutative correction to the Cornell potential in heavy-quarkonium atoms," *Theoretical and Mathematical Physics*, vol. 186, no. 2, pp. 280–285, 2016.
- [2] T. A. Nahool, A. M. Yasser, M. Anwar, and G. A. Yahya, "Charmonium properties," *East European Journal of Physics*, vol. 3, p. 31, 2020.
- [3] H. S. Chung, J. Lee, and D. Kang, "Cornell potential parameters for S-wave heavy quarkonia," 2008, <https://arxiv.org/abs/0803.3116>.
- [4] H. Mansour and A. Gamal, "Bound state of heavy quarks using a general polynomial potential," *Adv. High Energy Phys.*, vol. 2018, article 7269657, 7 pages, 2018.
- [5] N. R. Soni, B. R. Joshi, R. P. Shah, H. R. Chauhan, and J. N. Pandya, "Spectroscopy using the Cornell potential," *European Physical Journal C: Particles and Fields*, vol. 78, no. 7, 2018.
- [6] E. V. B. Leite, H. Belich, and R. L. L. Vitória, "Effects of the Cornell-type potential on a position-dependent mass system in Kaluza-Klein theory," *Adv. High Energy Phys.*, vol. 2019, article 6740360, pp. 1–7, 2019.
- [7] G. X. A. Petronilo, R. G. G. Amorim, S. C. Ulhoa, A. F. Santos, A. E. Santana, and F. C. Khanna, "Size-effect at finite temperature in a quark-antiquark effective model in phase space," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 36, no. 17, p. 2150121, 2021.
- [8] R. C. L. Bruni, E. F. Capossoli, and H. Boschi-Filho, "Quark-antiquark potential from a deformed AdS/QCD," *Adv. High Energy Phys.*, vol. 2019, article 1901659, 6 pages, 2019.
- [9] A. Vega and J. Flores, "Heavy quarkonium properties from Cornell potential using variational method and supersymmetric quantum mechanics," *Pramana-Journal of Physics*, vol. 87, no. 5, p. 73, 2016.
- [10] P. Czernski, "Bc-mesons in the deconfined phase," *EPJ Web of Conferences*, vol. 81, p. 05008, 2014.
- [11] E. P. Inyang, E. P. Inyang, J. Karniliyus, J. E. Ntibi, and E. S. William, "Diatomic molecules and mass Spectrum of heavy quarkonium system with Kratzer-screened coulomb potential (KSCP) through the solutions of the Schrödinger equation," *European Journal of Applied Physiology*, vol. 3, no. 2, pp. 48–55, 2021.
- [12] F. Tajik, Z. Sharifi, M. Eshghi, M. Hamzavi, M. Bigdeli, and S. M. Ikhdaier, "Eigenspectra and statistical properties of the Klein-Gordon equation with Cornell potential: unequal mixings of scalar and time-like vector potentials," *Physica A*, vol. 535, article 122497, 2019.
- [13] N. Ferkous and A. Bounames, "2D Pauli equation with Hulthén potential in the presence of Aharonov-Bohm effect," *Communications in Theoretical Physics*, vol. 59, no. 6, pp. 679–683, 2013.

- [14] S. M. Ikhdair, "Relativistic bound states of Spinless particle by the Cornell potential model in external fields," *Adv. High Energy Phys.*, vol. 2013, p. 10, 2013.
- [15] F. Ahmed, "Effects of coulomb- and Cornell-types potential on a spin-0 scalar particle under a magnetic field and quantum flux in topological defects space-time," *E. P. L.*, vol. 133, no. 50002, 2021.
- [16] L. A. Trevisan, C. Mirez, and F. M. Andrade, "The exact solution for the Dirac equation with the Cornell potential," *Few-Body Systems*, vol. 55, p. 1055, 2014.
- [17] B. C. Wang, E. C. Brenag, R. G. G. Amorim, V. C. Rispoli, and S. C. Ulhoa, "The noncommutative Coulomb potential," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 36, p. 2150094, 2021.
- [18] H. Mutuk, "Cornell potential: a neural network approach," *Adv. High Energy Phys.*, vol. 2019, article 3105373, 9 pages, 2019.
- [19] S. Koothottil, J. P. Prasanth, and V. M. Bannur, "Cornell potential model for strongly coupled quark gluon plasma," *Proceedings of the DAE Symp. on Nucl. Phys.*, vol. 62, p. , 2017912, 2017.
- [20] A. I. Ahmadov, C. Aydin, and O. Uzun, "Bound state solution of the Schrödinger equation at finite temperature," *Journal of Physics: Conference Series*, vol. 1194, article 012001, 2019.
- [21] M. Robinson, K. Bland, G. Cleaver, and J. Dittmann, "A simple introduction to particle physics part I - foundations and the standard model," 2008, <https://arxiv.org/abs/0810.3328>.
- [22] R. Mann, *An Introduction to Particle Physics and the Standard Model*, CRC press, 2009.
- [23] S. Gasiorowicz, *Elementary Particle Physics*, Wiley, New York, 1966.
- [24] D. Griffiths, *Introduction to Elementary Particles*, John Wiley & Sons, 2008.
- [25] I. J. R. Aitchison and A. J. G. Hey, *Gauge Theories in Particle Physics: A Practical Introduction: From Relativistic Quantum Mechanics to QED*, IOP Publishing, 2012.
- [26] J. J. Sakurai, *Invariance Principles and Elementary Particles*, Princeton University Press, 2015.
- [27] S. Weinberg, *The Discovery of Subatomic Particles*, Cambridge University Press, Revised edition, 2003.
- [28] M. Thomson, *Modern Particle Physics*, Cambridge University Press, 2013.
- [29] H. Mutuk, "S-wave heavy quarkonium spectra: mass, decays, and transitions," *Adv. High Energy Phys.*, vol. 2018, pp. 1–12, 2018.
- [30] E. P. Inyang, E. P. Inyang, E. S. William, and E. E. Ibekwe, "Study on the applicability of Varshni potential to predict the mass-spectra of the quark-antiquark systems in a non-relativistic framework," *Jordan Journal of Physics*, vol. 14, p. 337, 2021.
- [31] E. Omugbe, "Non-relativistic eigensolutions of molecular and heavy quarkonia interacting potentials via the Nikiforov Uvarov method," *Canadian Journal of Physics*, vol. 98, no. 12, pp. 1125–1132, 2020.
- [32] E. P. Inyang, I. O. Akpan, E. P. Inyang, and E. S. William, "Approximate solutions of the Schrödinger equation with Hulthén-Hellmann potentials for a quarkonium system," *Revista Mexicana de Física*, vol. 67, no. 3 May-Jun, p. 482, 2021.
- [33] E. M. Khoka, M. Abu-Shady, and T. A. Abdel-Karim, "Quarkonium masses in the N-dimensional space using the analytical exact iteration method," *International Journal of Theoretical and Applied Mathematics*, vol. 2, p. 86, 2016.
- [34] E. P. Inyang, E. P. Inyang, I. O. Akpan, J. E. Ntibi, and E. S. William, "Analytical solutions of the Schrödinger equation with class of Yukawa potential for a quarkonium system via series expansion method," *European Journal of Applied Physiology*, vol. 2, p. 6, 2020.
- [35] E. Wigner, "On the quantum correction for thermodynamic equilibrium," *Physics Review*, vol. 40, no. 5, pp. 749–759, 1932.
- [36] D. S. Dean, P. Le Doussal, S. N. Majumdar, and G. Schehr, "Wigner function of noninteracting trapped fermions," *Physical Review A*, vol. 97, no. 6, p. 063614, 2018.
- [37] M. Hillery, R. F. O'Connell, M. O. Scully, and E. P. Wigner, "Distribution functions in physics: fundamentals," *Physics Reports*, vol. 106, no. 3, pp. 121–167, 1984.
- [38] W. B. Case, "Wigner functions and Weyl transforms for pedestrians," *American Journal of Physics*, vol. 76, no. 10, pp. 937–946, 2008.
- [39] T. Tilma, M. J. Everitt, J. H. Samson, W. J. Munro, and K. Nemoto, "Wigner functions for arbitrary quantum systems," *Physical Review Letters*, vol. 117, no. 18, p. 180401, 2016.
- [40] D. Chruściński and K. Młodawski, "Wigner function and Schrödinger equation in phase-space representation," *Physical Review A*, vol. 71, no. 5, p. 052104, 2005.
- [41] C. Costa, M. R. Tenser, R. G. G. Amorim, M. C. B. Fernandes, A. E. Santana, and J. D. M. Vianna, "Symplectic field theories: scalar and spinor representations," *Advances in Applied Clifford Algebras*, vol. 28, p. 1, 2018.
- [42] R. G. G. Amorim, M. C. B. Fernandes, F. C. Khanna, A. E. Santana, and J. D. M. Vianna, "Non-commutative geometry and symplectic field theory," *Physics Letters A*, vol. 361, no. 6, pp. 464–471, 2007.
- [43] R. G. G. Amorim, F. C. Khanna, A. P. C. Malbouisson, J. M. C. Malbouisson, and A. E. Santana, "Realization of the noncommutative Seiberg–Witten gauge theory by fields in phase space," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 30, no. 22, p. 1550135, 2015.
- [44] H. Dessano, R. A. S. Paiva, R. G. G. Amorim, S. C. Ulhoa, and A. E. Santana, "Wigner function and non-classicality for oscillator systems," *Brazilian Journal of Physics*, vol. 49, no. 5, pp. 715–725, 2019.
- [45] R. G. G. Amorim, F. C. Khanna, A. E. Santana, and J. D. M. Vianna, "Perturbative symplectic field theory and Wigner function," *Physica A: Statistical Mechanics and its Applications*, vol. 388, no. 18, pp. 3771–3778, 2009.
- [46] F. C. Khanna, A. P. C. Malbouisson, J. M. C. Malbouisson, and A. E. Santana, *Thermal Quantum Field Theory: Algebraic Aspects and Applications*, World Scientific, 2009.
- [47] J. S. da Cruz Filho, R. G. G. Amorim, S. C. Ulhoa, F. C. Khanna, A. E. Santana, and J. D. M. Vianna, "Hénon–Heiles interaction for hydrogen atom in phase space," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 31, no. 10, p. 1650046, 2016.
- [48] G. Torres-Vega and J. H. Frederick, "A quantum mechanical representation in phase space," *The Journal of Chemical Physics*, vol. 98, no. 4, pp. 3103–3120, 1993.

- [49] G. Torres-Vega and J. H. Frederick, "Quantum mechanics in phase space: new approaches to the correspondence principle," *The Journal of Chemical Physics*, vol. 93, no. 12, pp. 8862–8874, 1990.
- [50] R. G. G. Amorim and R. A. S. Paiva, "Quantum physics in phase space: an analysis of simple pendulum," *Advances in Theoretical & Computational Physics*, vol. 1, 2018.
- [51] A. X. Martins, R. A. S. Paiva, G. Petronilo, R. R. Luz, R. G. G. Amorim, and S. C. Ulhoa, "Analytical solution for the gross-Pitaevskii equation in phase space and Wigner function," *Advances in High Energy Physics*, vol. 2020, article 7010957, 6 pages, 2020.
- [52] R. A. S. Paiva, R. G. G. Amorim, S. C. Ulhoa, A. E. Santana, and F. C. Khanna, "Zeeman effect in phase space," *Adv. High Energy Phys.*, vol. 2020, article 4269246, 9 pages, 2020.
- [53] A. X. Martins, T. M. R. Filho, R. G. G. Amorim, R. A. S. Paiva, G. X. A. Petronilo, and S. C. Ulhoa, "Analytical solution for quantum quartic oscillator in phase space," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 35, no. 20, p. 2050100, 2020.
- [54] M. D. Oliveira, M. C. B. Fernandes, F. C. Khanna, A. E. Santana, and J. D. M. Vianna, "Symplectic quantum mechanics," *Annals of Physics*, vol. 312, no. 2, pp. 492–510, 2004.
- [55] T. L. Curtright, D. B. Fairlie, and C. K. Zachos, *A Concise Treatise on Quantum Mechanics in Phase Space*, World Scientific Publishing Company, 2013.
- [56] T. L. Curtright, D. B. Fairlie, and C. K. Zachos, "Features of time-independent Wigner functions," *Physical Review D*, vol. 58, no. 2, p. 025002, 1998.
- [57] L. E. Ballentine, *Quantum Mechanics: A Modern Development*, World Scientific Publishing Company, 1998.
- [58] P. Campos, M. G. R. Martins, M. C. B. Fernandes, and J. D. M. Vianna, "Quantum mechanics on phase space: the hydrogen atom and its Wigner functions," *Annals of Physics*, vol. 390, pp. 60–70, 2018.
- [59] P. Campos, M. G. R. Martins, and J. D. M. Vianna, "Quantum mechanics on phase space and the coulomb potential," *Physics Letters A*, vol. 381, no. 13, pp. 1129–1133, 2017.
- [60] M. B. de Menezes, M. C. B. Fernandes, M. G. R. Martins, A. E. Santana, and J. D. M. Vianna, "Supersymmetric symplectic quantum mechanics," *Annals of Physics*, vol. 389, pp. 111–135, 2018.
- [61] R. G. G. Amorim, S. C. Ulhoa, and E. O. Silva, "On the symplectic Dirac equation," *Brazilian Journal of Physics*, vol. 45, no. 6, pp. 664–672, 2015.
- [62] R. G. G. Amorim, M. C. B. Fernandes, F. C. Khanna, A. E. Santana, and J. D. M. Vianna, "Symplectic Schrödinger equation for many-body systems," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 35, no. 6, p. 2050033, 2020.
- [63] R. G. G. Amorim, J. S. da Cruz Filho, A. F. Santos, and S. C. Ulhoa, "On scalar electromagnetism in phase space," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 33, no. 22, p. 1850131, 2018.
- [64] G. X. A. Petronilo, S. C. Ulhoa, K. V. S. Araújo, R. A. S. Paiva, R. G. G. Amorim, and A. E. Santana, "The Landau problem and non-classicality," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 35, no. 25, p. 2050148, 2020.
- [65] J. D. M. Vianna, M. C. B. Fernandes, and A. E. Santana, "Galilean-covariant Clifford algebras in the phase-space representation," *Foundations of Physics*, vol. 35, no. 1, pp. 109–129, 2005.
- [66] M. C. B. Fernandes, F. C. Khanna, M. G. R. Martins, A. E. Santana, and J. D. M. Vianna, "Non-linear Liouville and Schrödinger equations in phase space," *Physica A*, vol. 389, no. 17, pp. 3409–3419, 2010.
- [67] R. G. G. Amorim, F. C. Khanna, A. P. C. Malbouisson, J. M. C. Malbouisson, and A. E. Santana, "Quantum field theory in phase space," *International Journal of Modern Physics A: Particles and Fields; Gravitation; Cosmology; Nuclear Physics*, vol. 34, no. 8, p. 1950037, 2019.
- [68] R. G. G. Amorim and W. B. Silva, "Quartic potential in phase space," *Physica*, vol. 10, p. 20, 2014.
- [69] M. A. De Gosson, *Symplectic Methods in Harmonic Analysis and in Mathematical Physics*, Springer Science & Business Media, 2011.
- [70] E. E. Perepelkin, B. I. Sadovnikov, N. G. Inozemtseva, and E. V. Burlakov, "Wigner function of a quantum system with polynomial potential," *Journal of Statistical Mechanics*, vol. 2020, no. 5, p. 053105, 2020.
- [71] O. Vallée and M. Soares, *Airy Functions and Applications to Physics*, World Scientific Publishing Company, 2010.
- [72] M. Tanabashi et al., *Review of Particle Physics: Particle Data Groups*, 2018.
- [73] A. Kenfack and K. Zyczkowski, "Quantum semiclass," *Journal of Optics B*, vol. 6, p. 396, 2004.