

Clinical Study

Posterior Transpedicular Dynamic Stabilization versus Total Disc Replacement in the Treatment of Lumbar Painful Degenerative Disc Disease: A Comparison of Clinical Results

Tunc Oktenoglu,¹ Ali Fahir Ozer,² Mehdi Sasani,¹
Yaprak Ataker,³ Cengiz Gomleksiz,⁴ and Irfan Celebi⁵

¹Neurosurgery Department, American Hospital, 34365 Istanbul, Turkey

²Neurosurgery Department, Koc University School of Medicine, 34450 Istanbul, Turkey

³Physical Therapy and Rehabilitation Department, American Hospital, 34365 Istanbul, Turkey

⁴Neurosurgery Department, Mengücek Gazi Training and Research Hospital, School of Medicine, Erzincan University, 2400 Erzincan, Turkey

⁵Radiology Department, Sisli Etfal Hospital, 34360 Istanbul, Turkey

Correspondence should be addressed to Ali Fahir Ozer; alifahirozer@gmail.com

Received 19 October 2012; Accepted 28 November 2012

Academic Editor: Deniz Erbulut

Copyright © 2013 Tunc Oktenoglu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Study Design. Prospective clinical study. *Objective.* This study compares the clinical results of anterior lumbar total disc replacement and posterior transpedicular dynamic stabilization in the treatment of degenerative disc disease. *Summary and Background Data.* Over the last two decades, both techniques have emerged as alternative treatment options to fusion surgery. *Methods.* This study was conducted between 2004 and 2010 with a total of 50 patients (25 in each group). The mean age of the patients in total disc prosthesis group was 37,32 years. The mean age of the patients in posterior dynamic transpedicular stabilization was 43,08. Clinical (VAS and Oswestry) and radiological evaluations (lumbar lordosis and segmental lordosis angles) of the patients were carried out prior to the operation and 3, 12, and 24 months after the operation. We compared the average duration of surgery, blood loss during the surgery and the length of hospital stay of both groups. *Results.* Both techniques offered significant improvements in clinical parameters. There was no significant change in radiologic evaluations after the surgery for both techniques. *Conclusion.* Both dynamic systems provided spine stability. However, the posterior dynamic system had a slight advantage over anterior disc prosthesis because of its convenient application and fewer possible complications.

1. Introduction

Currently, one of the most important causes of chronic low back pain is thought to be a painful disc [1–3]. Some biomechanical and biochemical changes play a role in intervertebral disc degeneration; on the other hand intrinsic, extrinsic, and genetic factors are also important. Compression of the spine, torsional injuries, overload, and congenital anomalies have been shown to contribute to disc degeneration with applying excessive pressure onto intervertebral discs [4–10]. Despite numerous research studies, the etiology and physiopathology of disc degeneration remain unknown [2]. Annular tears

resulting from degeneration of the annulus fibrosis, that contains pain receptors and internal disc ruptures, are the most common cause of pain [11–13]. Today, it is believed that degenerative disc disease (DDD) might cause instability in spine segments, and it is widely accepted that progressive back pain results due to this instability [14–16]. In fact, segmental instability begins when disc height deterioration is initiated by the progression of intervertebral disc degeneration. Instability as a consequence of disc degeneration has been described by Frymoyer [14, 15] as primary segmental instability and by Kirkaldy-Willis and Farfan [2] as the discogenic pain and instability stage in the overall process

of degeneration. Benzel [16] included degenerative disc disease among the chronic instabilities and described the disease as “dysfunctional segmental motion” and “torsional instability.” Fusion is the standard surgical treatment option for painful lumbar degenerative disc disease that is unresponsive to conservative treatment modalities. Nonetheless, the side effects of fusion (pseudarthrosis, adjacent segment disease, and the donor site morbidity) and suboptimal clinical satisfaction rates, which have been reported even in patients with radiologically observed fusion, have led to a search for alternative treatments [17–22].

Numerous dynamic techniques were developed over the last two decades. Recently, these devices were classified as total disc replacement (TDR) and posterior transpedicular dynamic systems (PTDS) [23]. Both PTDS and TDR have been widely used in surgical treatment of degenerative disc diseases of the lumbar spine. Numerous studies showed promising clinical results [24–35]. However, there is no study that compares the TDR and PTDS techniques in the treatment of DDD.

In this prospective study, we evaluated and compared the clinic and radiologic outcome of TDR and PTDS in patients with painful lumbar degenerative disc disease through an extensive literature review.

2. Material and Methods

2.1. Total Disc Replacement Group. We performed TDR on 25 patients (14 females and 11 males). The mean age of the patients was 37.32 (with a range from 25 to 50), and the mean follow-up period was 29.16 months (with a range from 24 to 42 months).

A lumbar total disc replacement (Maverick, Medtronic Sofamor Danek, Memphis, TN, USA) was placed into the intervertebral disc space with open window laparotomy technique [36].

All patients in the TDR group had a lumbar single-level painful disc. 15 patients showed L4-L5 DDD, and 10 patients showed L5-S1 DDD (Figure 1). All of the patients were informed about the surgery, and they signed a written, informed consent form. The inclusion criteria for TDR surgery included a complaint of lower back pain that had duration of at least 12 months and at least six months of conservative treatment without satisfactory results. Other inclusion criteria were that the patients must be less than 50 years old and have no signs of lumbar degenerative spondylolisthesis or osteoarthritis in their facet joints, which was confirmed with computerized tomography (CT) and dynamic plain radiographs. The patients also had to have symptomatic lumbar degenerative disc disease that was visible in magnetic resonance imaging (MRI) as a blackened disc as well as a confirmation of the diagnosis by displaying pain behaviors during discography.

2.2. Posterior Dynamic Transpedicular Stabilization Group. We performed posterior dynamic transpedicular stabilization on 25 patients (13 females and 12 males). The mean age of the patients was 43.08 years (with a range from 24 to 55 years),




FIGURE 1: A 30-year-old woman complained of severe back pain attacks. She had no neurological deficits. (a) T2-weighted MRI scans showed advanced degeneration with Modic changes in the L4-L5 disc. (b) Maverick disc prosthesis was applied.

and the mean follow-up period was 36.48 months (with a range from 24 to 48 months).

Patients in the dynamic posterior stabilization group were operated with the Cosmic (Ulrich GmbH & Co. KG, Ulm, Germany) posterior dynamic transpedicular stabilization system (hinged screw-rigid rod) through transmuscular approach [35].

All cases in the PTDS group had one-level painful disc disease. The operated discs were L4-L5 region (16 cases) and L5-S1 region (9 cases) (Figure 2).

Similar to the patients in the TDR group, the inclusion criteria included a confirmed diagnosis of symptomatic lumbar degenerative disc disease through MRI and positive discography, a complaint of lower back pain that had a duration of at least 12 months, at least 6 months of conservative treatment without satisfactory results, and the absence of apparent instability confirmed with lumbosacral dynamic X-rays.

2.3. Clinical Evaluation. We evaluated and compared the average surgical time, blood loss during the surgery, and the length of the stay in hospital for both groups of patients (Table 1). The visual analog scale (VAS) and the Oswestry Disability Index (ODI) were used for the clinical evaluations and follow-up examinations. Clinical evaluations of the patients were carried out in the data at preoperative period and 3, 12, and 24 months after the surgery (Tables 2 and 3).

2.4. Radiological Evaluation. To diagnose lumbar disc disease, an MRI examination of each patient was performed and a black disc was observed. Pain symptoms were confirmed with the detection of provocative pain through a discography




FIGURE 2: (a) A 50-year-old male complained of severe back pain attacks. (a) T2-weighted MRI scans showed degeneration of the disc at L4-L5 lumbar disc (L5- S1 considered as sacralization). (b) Following posterior transpedicular dynamic stabilization with the Cosmic system.

TABLE 1: Comparison of TDR and PTDS groups.

		TDR ($n = 25$)	PTDS ($n = 25$)	P
Age	Min-Max	25-50	24-55	0.018*
	Mean \pm SD	37.32 \pm 6.62	43.08 \pm 9.65	
Followup (month)	Min-Max	24-42	24-48	0.001**
	Mean \pm SD	29.16 \pm 4.77	36.48 \pm 6.72	
Length of hospital stay (day)	Min-Max	3-5	2-5	0.022*
	Mean \pm SD	3.56 \pm 0.58	3.04 \pm 0.93	
Operation time (minute)	Min-Max	120-260	40-70	0.001**
	Mean \pm SD	181.20 \pm 39.40	52.40 \pm 7.79	
Blood loss (mL.)	Min-Max	300-600	75-175	0.001**
	Mean \pm SD	402.00 \pm 91.83	103.00 \pm 22.03	

Student's t -test, * $P < 0.05$, ** $P < 0.01$.

which was applied to the black disc. Lumbosacral plain and dynamic (hyperflexion and hyperextension) X-rays and CT examinations of the patients were carried out by independent radiology experts in preoperative. Follow-up plain X-Ray studies were obtained 3, 12, and 24 months after the surgery. Control CT study was performed in postoperative 24 months. Loose screws as well as broken screws, instrument migration, subsidence, and spontaneous fusion were evaluated. Additionally lumbar lordosis angle (LL) and segmental lordosis angle (α) data was obtained (Tables 4 and 5) (Figure 3).

2.5. Statistical Analysis. The Number Cruncher Statistical System (NCSS) 2007 and 2008 PASS Statistical Software (Utah, USA) were used for statistical analysis of the data. In addition to descriptive statistical methods (e.g., mean, standard deviation), Student's t -test was used to compare the normally distributed parameters between the two groups. The Mann-Whitney U test was used for the comparison of parameters with nonnormal distribution. Bonferroni test was used to compare the follow-up data with normal distribution,

and paired sample t -test was used for dual comparison. In nonnormal distribution group, the follow-up data compared with Friedman test and Wilcoxon test was used for dual comparison. The significance level was $P < 0.05$.

3. Results

There was statistically significant difference observed between the mean ages and follow-up periods of the groups ($P < 0.05$) (Table 1). The PTDS applied to significantly older patients was compared to TDR group.

There was a statistically significant difference ($P < 0.01$) between the level of blood loss in the two groups. The level of blood loss was significantly higher in the TDR group compared to the PTDS group (Table 1, Figure 4).

The operation time was significantly longer ($P < 0.01$) in the TDR group compared to the posterior dynamic stabilization group (Table 1, Figure 5).

There was significant difference in the length of the hospital stay between the two groups ($P < 0.05$) (Table 1).

TABLE 2: (a) The comparison of VAS scores. (b) The comparison of decrease % in follow-up VAS data.

VAS		TDR (<i>n</i> = 25)	PTDS (<i>n</i> = 25)	^a <i>P</i>
Preop VAS	Min–Max	6–10	6–9	0.219
	Mean ± SD	8.24 ± 1.09[8]	7.96 ± 0.79 [8]	
VAS (3 months)	Min–Max	0–5	1–5	0.588
	Mean ± SD	2.44 ± 1.16 [2] [‡]	2.64 ± 0.91 [2] [‡]	
VAS (12 months)	Min–Max	0–3	0–4	0.087
	Mean ± SD	1.68 ± 0.85 [2] [‡]	1.28 ± 0.94 [1] [‡]	
VAS (24 months)	Min–Max	0–2	0–3	0.240
	Mean ± SD	0.84 ± 0.69 [1] [‡]	0.68 ± 0.85 [1] [‡]	
^b <i>P</i>		0.001**	0.001**	

^a Mann-Whitney *U* test, ^b Friedman test.

[‡] Wilcoxon signed-rank test *P* < 0.001.

** *P* < 0.01.

VAS (decrease %)	TDR (<i>n</i> = 25) Mean ± SD	PTDS (<i>n</i> = 25) Mean ± SD	<i>P</i>
Preop * VAS (3 months)	70.18 ± 14.14	66.18 ± 13.57	0.374
Preop * VAS (12 months)	79.91 ± 10.00	83.71 ± 12.08	0.214
Preop * VAS (24 months)	89.86 ± 8.41	91.30 ± 10.78	0.519

Mann-Whitney *U* test.

TABLE 3: (a) The comparison of ODI scores. (b) The comparison of decrease % in follow-up OSW data.

ODI		TDR (<i>n</i> = 25)	PTDS (<i>n</i> = 25)	^a <i>P</i>
Preop ODI	Min–Max	40–100	46–98	0.847
	Mean ± SD	67.20 ± 20.79	66.16 ± 16.82	
ODI (3 months)	Min–Max	12–56	12–38	0.069
	Mean ± SD	32.32 ± 10.87 [‡]	27.16 ± 8.63 [‡]	
ODI (12 months)	Min–Max	4–34	2–26	0.010*
	Mean ± SD	18.00 ± 7.64 [‡]	12.88 ± 5.75 [‡]	
ODI (24 months)	Min–Max	2–20	2–18	0.408
	Mean ± SD	9.12 ± 4.28 [‡]	8.04 ± 4.53 [‡]	
^b <i>P</i>		0.001**	0.001**	

^a Student's *t*-test, ^b repeated measures test.

[‡] Adjustment for multiple comparisons: Bonferroni *P* < 0.01.

* *P* < 0.05, ** *P* < 0.01.

ODI (decrease %)	TDR (<i>n</i> = 25) Mean ± SD	PTDS (<i>n</i> = 25) Mean ± SD	<i>P</i>
Preop * ODI (3 months)	49.96 ± 16.46	54.91 ± 20.00	0.669
Preop * ODI (12 months)	72.80 ± 9.33	78.22 ± 11.68	0.093
Preop * ODI (24 months)	86.15 ± 6.15	86.33 ± 8.92	0.808

Mann-Whitney *U* test.

Preoperative VAS and ODI levels were not significantly (*P* > 0.05 and *P* > 0.05) different between the groups (Tables 2 and 3).

In both groups the clinical parameters (VAS and ODI) showed significant improvement in all postoperative time

periods when compared to preoperative data (Tables 2 and 3, *P* < 0.01). There were no statistically significant differences observed between the groups for the each follow-up VAS (*P* < 0.05, Table 2). The ODI data showed significant difference only in postop 12 months. The PTDS group had significantly

TABLE 4: The comparison of LL angles.

LL		TDR (n = 25)	PTDS (n = 25)	^a P
Preop LL	Min–Max	25–65	34–72	0.948
	Mean ± SD	49.60 ± 10.46	49.80 ± 11.26	
LL (3 months)	Min–Max	26–65	34–69	0.747
	Mean ± SD	49.52 ± 9.51	48.60 ± 10.52	
LL (12 months)	Min–Max	24–64	30–67	0.764
	Mean ± SD	49.60 ± 10.15	48.72 ± 10.50	
LL (24 months)	Min–Max	22–65	35–65	0.786
	Mean ± SD	49.56 ± 10.38	48.80 ± 9.30	
^b P		0.998	0.890	

^a Student's *t*-test, ^b repeated measures test.
LL: lumbar lordosis.

TABLE 5: The comparison of SL (α) angles.

ALPHA		TDR (n = 25)	PTDS (n = 25)	^a P
Preop ALPHA	Min–Max	4–17	4–30	0.274
	Mean ± SD	10.32 ± 3.06	11.68 ± 5.33	
ALPHA (3 months)	Min–Max	3–19	3–33	0.566
	Mean ± SD	10.40 ± 3.70	11.20 ± 5.84	
ALPHA (12 months)	Min–Max	4–16	2–31	0.392
	Mean ± SD	10.36 ± 2.90	11.52 ± 6.05	
ALPHA (24 months)	Min–Max	5–14	3–30	0.248
	Mean ± SD	10.32 ± 2.28	11.56 ± 4.79	
^b P		0.989	0.858	

^a Student's *t*-test, ^b repeated measures test.
SL: segmental lordosis.




FIGURE 3: Lordosis of the lumbar spine (L1-S1) was measured via the angle between the lines drawn from the lower endplate of L1 and the upper endplate of S1 (LL). Additionally segmental lordosis (α) at the operation level was measured via the angle between the lines drawn from the upper and lower endplates of the vertebrae that form the operation segments.

better outcome in this time period. However this advantage did not persist. There was no significant difference in 24-month scores ($P > 0.05$, Table 3) (Figures 6 and 7).

There were no significant differences observed between the preoperative and postoperative lumbar (LL) and segmental lordosis (alpha) evaluations for both techniques ($P > 0.05$) (Tables 4 and 5).




FIGURE 4: The data of blood loss in surgery for both groups.




FIGURE 5: The data of operation times for both groups.




FIGURE 6: The distribution of VAS scores for both groups.




FIGURE 7: The distribution of ODI scores for both groups.

No surgical morbidity and/or complications observed in the group treated with PTDS. There were two iliac vein injuries that occurred in two patients in the TDR group. These injuries were sutured in the operation with no mortality and residual morbidity.

4. Discussion

Fusion has been widely used as a surgical treatment for painful disc disease. Fusion eliminates the abnormal movements and offers satisfactory outcome. On the other hand, even in patients with 100% fusion achieved with applying 360° fusion method, the satisfaction rate is not necessary optimal and might be low as 30% [19–21, 23, 24]. Donor site problems have also been a significant complication in fusion surgery [20]. Therefore, alternative treatment techniques were developed in an attempt to prevent side effects that are commonly observed after fusion surgery and to improve the patient satisfaction rate. In recent years, dynamic systems that provide spine mobility have been developed to avoid the well-known side effects of fusion technique. Today indications and contraindications of TDR and PTDS are well known [23]. Both techniques can be used for the same indications. A painful black disc can be treated with application of either technique.

TDR was developed over the past ten years as a promising surgery that was preferable over fusion surgery because

the proponents of TDR claimed that the procedure preserves mobility and reduces the risk of adjacent segment disease. After a ten-year effort by Büttner-Janz et al., TDR was announced as a new solution method for painful disc disease [37]. Numerous TDR systems were developed and offered for clinical application [29, 32, 38]. Biomechanical studies showed that the TDR prosthesis stabilizes the spine while providing nearly intact segmental motion [29, 32, 39]. Early clinical results of TDR in the treatment of DDD showed promising outcomes [37, 40–46].

The patients treated with TDR usually have short recuperation times and less postoperative pain compared to fusion procedure. On the other hand, TDR application has several significant limitations including; (a) the patient should be between 30 and 50 years old, (b) there should not be any posterior column disruption, (c) intervertebral disc height should be ≥ 4 mm, and (d) single-level DDD is more appropriate to apply TDR. Beside these limitations TDR is an anterior approach which has its inherent risks such as injury to intra-abdominal organs and vascular structures. Additionally, the lesions in the peritoneal cavity caused by abrasion, ischemia, desiccation, infection, thermal injury, and foreign bodies can result in adhesion formation [47].

TDR is used extensively around the world; however severe complications have been associated with the technique [48, 49]. In this study we observed mild iliac vein injuries during the placement of the lumbar disc prosthesis in the two patients within the TDR group. Other possible disadvantages of TDR technique are as follows: revision surgery is quite difficult, biomechanically the L5-S1 level had no normal segmental motion, and results of two-level TDR use were not considered to be satisfactory according to patients [50]. Putzier et al. [51] concluded that the long-term results of a study by Charité were not satisfactory and they concluded their article yearning to fusion technique.

Guyer et al. [52] published the results of a 5-year study showing that TDR was not superior to fusion. The authors concluded that there was no strong evidence that TDR was superior to fusion, and they suggested that high-quality, randomized controlled trials with relevant control groups and a long-term followup were needed to evaluate the effectiveness and safety of TDR [53].

Posterior dynamic stabilization systems are designed to increase the success of spinal surgery and to eliminate the complications of fusion with rigid instrumentation such as adjacent segment disease (12.2–18.5%) [54] due to the stress-shielding properties (2–3% per year after stabilization) [55], pseudarthrosis (3–55%) [18, 56, 57], device-related osteopenia [58], and loss of motion in fused spinal segments. Besides these side effects of fusion, clinical healing might be suboptimal in cases even with satisfactory radiological results [22, 59]. Therefore, the use of posterior dynamic stabilization in the surgical treatment of DDD may provide greater patient satisfaction, resulting from shorter hospital stays, less recuperation time, and none of the disadvantages related to fusion, which requires more invasive procedures.

Numerous biomechanical studies proved that hinged screw stabilization can stabilize the spine almost as well as the rigid screw stabilization used in fusion surgery in

the treatment of chronic lumbar instability [60, 61]. There are no randomized controlled studies in the literature because PTDS is a new technique. However, there are many retrospective studies that are precursors for future randomized studies. Recently, studies on PTDS have shown very encouraging clinical results and demonstrated that these systems provide stabilization that is similar to the posterior rigid stabilization obtained with fusion surgery [23–28, 30, 31, 34, 35, 60, 62, 63]. There are few studies which concluded that dynamic stabilization is not superior to rigid stabilization [63–66]. Although these results showed that there was no advantage of PTDS over fusion surgery in clinical outcome, on the other hand these studies also showed that PTDS is superior to fusion due to the simplicity of the procedure, low morbidity, and reduced hospitalization time to achieve similar satisfactory outcome as fusion. Similarly numerous studies have shown that posterior dynamic transpedicular stabilization caused less intraoperative blood loss and had a shorter operating time [25, 28, 31, 35]. Furthermore, several studies reported that PTDS slows down intervertebral disc degeneration by removing the load from the degenerative disc tissue and providing better load distribution which is an important advantage of this technique [25, 27, 30].

Based on previous studies, if a disc is in the beginning stage of degeneration and if there is only posterior annulus defect, the disc might repair itself after PTDS. On the other hand, if the disc has advanced degeneration including decreased disc height, significant dehydration, and/or slight bulging, fusion might occur slowly after PTDS. However, in both of these scenarios, the patient would be pain-free. In cases of advanced disc degeneration, the fusion results are satisfactory because fusion occurs easily. If PTDS is applied to advanced disc degeneration cases, the segments might fuse and the results will be the same. Therefore in regard of motion preservation, TDR may be a superior treatment in this group of patients if they have intact facet joints.

Huang et al. [67] reported the advantages and disadvantages of nonfusion technology in spinal surgery. Some of the potential benefits of nonfusion implants were the elimination of possible complications due to bone grafts and pseudarthrosis as well as a reduction in the surgical morbidity and the incidence of adjacent level degeneration. The potential risks of nonfusion implants included mechanical failure, dissolution and migration, subsidence, and same-level degeneration.

Previous studies suggested that lumbar total disc prosthesis would reduce the stress on the adjacent disc, after sagittal balance is restored. Harrop et al. [68] reviewed the literature on lumbar adjacent segment degeneration after fusion and TDR. They concluded that adjacent segment disease had a stronger relationship with fusion than arthroplasty. Stoll et al. [34] reported symptomatic adjacent segment disease in 9% of their posterior dynamic transpedicular stabilization patients after a 38-month follow-up period. Cakir et al. [63] reported their results after performing PTDS with Dynesys and TDR with ProDisc (Synthes-Spine Solutions, New York, NY). They suggested that both dynamic systems were promising alternative options compared to fusion for patients with different pathologies because of reduced morbidity. Cakir et al. [63]

obtained good clinical results with both systems. Both TDR and PTDS result in less adjacent segment disease. Although a reduced incidence of adjacent segment degeneration appears to be the most important advantage of nonfusion systems, this advantage has not been proven.

Considering all of the features of both techniques, PTDS is a less invasive surgery compared to fusion and TDR techniques. Additionally, PTDS has no age limitation and does not require intact posterior spinal column as TDR technique. Finally, anterior lumbar disc prosthesis requires transperitoneal or retroperitoneal intervention and usually requires a multidisciplinary approach (general surgeon, cardiovascular surgeon, and spinal surgeon). Naturally, the complication rate decreases with a conventional surgical approach and increases when complex anatomical structures are involved in the surgery.

5. Conclusion

In this study, we observed that both dynamic techniques TDR and PTDS offered satisfactory outcome in the surgical treatment of lumbar DDD. However, in this limited study, PTDS had several advantages over TDR such as (a) less invasive technique, (b) shorter operation time, (c) less intraoperative bleeding, and (d) lower complication rates. Further prospective, randomized clinical studies with a larger number of patients and with a longer follow-up period are needed to support our findings.

References

- [1] N. Bogduk, "The innervation of the lumbar spine," *Spine*, vol. 8, no. 3, pp. 286–293, 1983.
- [2] W. H. Kirkaldy-Willis and H. F. Farfan, "Instability of the lumbar spine," *Clinical Orthopaedics and Related Research*, vol. 165, pp. 110–123, 1982.
- [3] F. P. Morgan and T. King, "Primary instability of lumbar vertebrae as a common cause of low back pain," *Journal of Bone and Joint Surgery (British Volume)*, vol. 39, no. 1, pp. 6–22, 1957.
- [4] J. MALINSKY, "The ontogenetic development of nerve terminations in the intervertebral discs of man," *Acta Anatomica*, vol. 38, pp. 96–113, 1959.
- [5] M. C. Battié, T. Videman, and E. Parent, "Lumbar disc degeneration: epidemiology and genetic influences," *Spine*, vol. 29, no. 23, pp. 2679–2690, 2004.
- [6] A. Ong, J. Anderson, and J. Roche, "A pilot study of the prevalence of lumbar disc degeneration in elite athletes with lower back pain at the Sydney 2000 Olympic Games," *British Journal of Sports Medicine*, vol. 37, no. 3, pp. 263–266, 2003.
- [7] T. Videman, S. Sarna, M. C. Battie et al., "The long-term effects of physical loading and exercise lifestyles on back-related symptoms, disability, and spinal pathology among men," *Spine*, vol. 20, no. 6, pp. 699–709, 1995.
- [8] C. J. Wallach, L. G. Gilbertson, and J. D. Kang, "Gene therapy applications for intervertebral disc degeneration," *Spine*, vol. 28, no. 15, supplement, pp. S93–S98, 2003.
- [9] H. A. Horner and J. P. G. Urban, "2001 Volvo award winner in basic science studies: effect of nutrient supply on the viability of cells from the nucleus pulposus of the intervertebral disc," *Spine*, vol. 26, no. 23, pp. 2543–2549, 2001.

- [10] L. Ala-Kokko, "Genetic risk factors for lumbar disc disease," *Annals of Medicine*, vol. 34, no. 1, pp. 42–47, 2002.
- [11] M. A. Edgar and J. A. Ghadially, "Innervation of the lumbar spine," *Clinical Orthopaedics and Related Research*, vol. 115, pp. 35–41, 1976.
- [12] M. Grönblad, J. Virri, S. Rönkkö et al., "A controlled biochemical and immunohistochemical study of human synovial-type (group II) phospholipase A2 and inflammatory cells in macroscopically normal, degenerated, and herniated human lumbar disc tissues," *Spine*, vol. 21, no. 22, pp. 2531–2538, 1996.
- [13] H. V. Crock, "Internal disc disruption. A challenge to disc prolapse fifty years on," *Spine*, vol. 11, no. 6, pp. 650–653, 1986.
- [14] J. W. Frymoyer, "Segmental instability," in *The Adult Spine*, J. W. Frymoyer, Ed., pp. 1873–1891, Raven Press, New York, NY, USA, 1991.
- [15] J. W. Frymoyer and D. K. Selby, "Segmental instability: rationale for treatment," *Spine*, vol. 10, no. 3, pp. 280–286, 1985.
- [16] E. C. Benzel, "Stability and instability of the spine," in *Biomechanics of Spine Stabilization*, E. C. Benzel, Ed., pp. 29–43, Thime, New York, NY, USA, 2001.
- [17] S. J. Suratwala, M. R. Pinto, T. J. Gilbert, R. B. Winter, and J. M. Wroblewski, "Functional and radiological outcomes of 360° fusion of three or more motion levels in the lumbar spine for degenerative disc disease," *Spine*, vol. 34, no. 10, pp. E351–E358, 2009.
- [18] A. A. Faundez, J. D. Schwender, Y. Safriel et al., "Clinical and radiological outcome of anterior-posterior fusion versus transforaminal lumbar interbody fusion for symptomatic disc degeneration: a retrospective comparative study of 133 patients," *European Spine Journal*, vol. 18, no. 2, pp. 203–211, 2009.
- [19] B. I. Martin, S. K. Mirza, B. A. Comstock, D. T. Gray, W. Kreuter, and R. A. Deyo, "Reoperation rates following lumbar spine surgery and the influence of spinal fusion procedures," *Spine*, vol. 32, no. 3, pp. 382–387, 2007.
- [20] J. C. Banwart, M. A. Asher, and R. S. Hassanein, "Iliac crest bone graft harvest donor site morbidity: a statistical evaluation," *Spine*, vol. 20, no. 9, pp. 1055–1060, 1995.
- [21] T. R. Lehmann, K. F. Spratt, and J. E. Tozzi, "Long-term follow-up of lower lumbar fusion patients," *Spine*, vol. 12, no. 2, pp. 97–104, 1987.
- [22] P. Fritzell, O. Hägg, and A. Nordwall, "Complications in lumbar fusion surgery for chronic low back pain: comparison of three surgical techniques used in a prospective randomized study. A report from the Swedish lumbar spine study group," *European Spine Journal*, vol. 12, no. 2, pp. 178–189, 2003.
- [23] T. Kaner, M. Sasani, T. Oktenoglu, and A. F. Ozer, "Dynamic stabilization of the spine: a new classification system," *Turkish Neurosurgery*, vol. 20, no. 2, pp. 205–215, 2010.
- [24] T. Kaner, M. Sasani, T. Oktenoglu, M. Cosar, and A. F. Ozer, "Utilizing dynamic rods with dynamic screws in the surgical treatment of chronic instability: a prospective clinical study," *Turkish Neurosurgery*, vol. 19, no. 4, pp. 319–326, 2009.
- [25] T. Kaner, S. Dalbayrak, T. Oktenoglu, M. Sasani, A. L. Aydin, and F. O. Ozer, "Comparison of posterior dynamic and posterior rigid transpedicular stabilization with fusion to treat degenerative spondylolisthesis," *Orthopedics*, vol. 33, no. 5, 2010.
- [26] T. Kaner, M. Sasani, T. Oktenoglu et al., "Minimum two-year follow-up of cases with recurrent disc herniation treated with microdiscectomy and posterior dynamic transpedicular stabilisation," *Open Orthopaedics*, vol. 4, pp. 120–125, 2010.
- [27] T. Kaner, M. Sasani, T. Oktenoglu, M. Cosar, and A. F. Ozer, "Clinical outcomes after posterior dynamic transpedicular stabilization with limited lumbar discectomy: carragee classification system for lumbar disc herniations," *SAS Journal*, vol. 4, no. 3, pp. 92–97, 2010.
- [28] A. F. Ozer, N. R. Crawford, M. Sasani et al., "Dynamic lumbar pedicle screw-rod stabilization: two year follow-up and comparison with fusion," *Open Orthopaedics*, vol. 4, pp. 137–141, 2010.
- [29] M. Sasani, T. Öktenoğlu, K. Tuncay, N. Canbulat, S. Carilli, and F. A. Özer, "Total disc replacement in the treatment of lumbar discogenic pain with disc herniation: a prospective clinical study," *Turkish Neurosurgery*, vol. 19, no. 2, pp. 127–134, 2009.
- [30] M. Putzier, S. V. Schneider, J. F. Funk, S. W. Tohtz, and C. Perka, "The surgical treatment of the lumbar disc prolapse: nucleotomy with additional transpedicular dynamic stabilization versus nucleotomy alone," *Spine*, vol. 30, no. 5, pp. E109–E114, 2005.
- [31] A. Von Strempel, D. Moosmann, C. Stoss et al., "Stabilization of the degenerated lumbar spine in the nonfusion technique with cosmic posterior dynamic system," *The Wall Street Journal*, vol. 1, pp. 40–47, 2006.
- [32] J. C. Le Huec, H. Mathews, Y. Basso et al., "Clinical results of Maverick lumbar total disc replacement: two-year prospective follow-up," *Orthopedic Clinics of North America*, vol. 36, no. 3, pp. 315–322, 2005.
- [33] M. Kanayama, T. Hashimoto, K. Shigenobu et al., "Adjacent-segment morbidity after Graf ligamentoplasty compared with posterolateral lumbar fusion," *Journal of Neurosurgery*, vol. 95, no. 1, pp. 5–10, 2001.
- [34] T. M. Stoll, G. Dubois, and O. Schwarzenbach, "The dynamic neutralization system for the spine: a multi-center study of a novel non-fusion system," *European Spine Journal*, vol. 11, no. 2, supplement, pp. S170–S178, 2002.
- [35] T. Oktenoglu, A. F. Ozer, M. Sasani et al., "Posterior dynamic stabilization in the treatment of lumbar degenerative disc disease: 2-year follow-up," *Minimally Invasive Neurosurgery*, vol. 53, no. 3, pp. 112–116, 2010.
- [36] S. Carilli, T. Oktenoglu, and A. F. Ozer, "Open-window laparotomy during a transperitoneal approach to the lower lumbar vertebrae: new method for reducing complications," *Minimally Invasive Neurosurgery*, vol. 49, no. 4, pp. 227–229, 2006.
- [37] K. Büttner-Janz, K. Schelnack, and Zippel H, "An alternative treatment strategy for lumbar disc damage using the SB Charité modular disc prosthesis," *Zeitschrift für Orthopädie und Ihre Grenzgebiete*, vol. 125, no. 1, pp. 1–6, 1987.
- [38] T. Marnay, *ProDisc Retrospective Clinical Study: 7–11 Year Follow Up*, Spine Solutions, New York, NY, USA, 2002.
- [39] T. J. Errico, "Lumbar disc arthroplasty," *Clinical Orthopaedics and Related Research*, no. 435, pp. 106–117, 2005.
- [40] S. Blumenthal, P. C. McAfee, R. D. Guyer et al., "A prospective, randomized, multicenter food and drug administration investigational device exemptions study of lumbar total disc replacement with the CHARITÉ artificial disc versus lumbar fusion—part I: evaluation of clinical outcomes," *Spine*, vol. 30, no. 14, pp. 1565–1575, 2005.
- [41] J. Zigler, R. Delamarter, J. M. Spivak et al., "Results of the prospective, randomized, multicenter food and drug administration investigational device exemption study of the ProDisc-L total disc replacement versus circumferential fusion for the treatment of 1-level degenerative disc disease," *Spine*, vol. 32, no. 11, pp. 1155–1162, 2007.

- [42] S. H. Hochschuler, D. D. Ohnmeiss, R. D. Guyer, and S. L. Blumenthal, "Artificial disc: preliminary results of a prospective study in the United States," *European Spine Journal*, vol. 11, no. 2, supplement, pp. S106–S110, 2002.
- [43] H. Mayer, K. Wiechert, A. Korge, and I. Qose, "Minimally invasive total disc replacement: surgical technique and preliminary clinical results," *European Spine Journal*, vol. 11, no. 2, supplement, pp. S124–S130, 2002.
- [44] R. Bertagnoli and S. Kumar, "Indications for full prosthetic disc arthroplasty: a correlation of clinical outcome against a variety of indications," *European Spine Journal*, vol. 11, no. 2, supplement, pp. S131–S136, 2002.
- [45] P. C. McAfee, B. Cunningham, G. Holsapple et al., "A prospective, randomized, multicenter food and drug administration investigational device exemption study of lumbar total disc replacement with the CHARITÉtrade; artificial disc versus lumbar fusion—part II: evaluation of radiographic outcomes and correlation of surgical technique accuracy with clinical outcomes," *Spine*, vol. 30, no. 14, pp. 1576–1583, 2005.
- [46] T. David, "Charité artificial disc long term results of one level (10 year)," *Spine*, vol. 32, no. 6, pp. 661–666, 2007.
- [47] F. H. Geisler, S. L. Blumenthal, R. D. Guyer et al., "Neurological complications of lumbar artificial disc replacement and comparison of clinical results with those related to lumbar arthrodesis in the literature: results of a multicenter, prospective, randomized investigational device exemption study of Charité intervertebral disc. Invited submission from the Joint section meeting on disorders of the spine and peripheral nerves, March 2004," *Journal of Neurosurgery: Spine*, vol. 1, no. 2, pp. 143–154, 2004.
- [48] C. Rosen, P. D. Kiester, and T. Q. Lee, "Lumbar disk replacement failures: review of 29 patients and rationale for revision," *Orthopedics*, vol. 32, no. 8, article 562, 2009.
- [49] A. van Ooij, F. Cumhur Oner, and A. J. Verbout, "Complications of artificial disc replacement: a report of 27 patients with the SB Charité disc," *Journal of Spinal Disorders and Techniques*, vol. 16, no. 4, pp. 369–383, 2003.
- [50] C. J. Siepe, H. M. Mayer, K. Wiechert, and A. Korge, "Clinical results of total lumbar disc replacement with ProDisc II: three-year results for different indications," *Spine*, vol. 31, no. 17, pp. 1923–1932, 2006.
- [51] M. Putzier, J. F. Funk, S. V. Schneider et al., "Charité total disc replacement—clinical and radiographical results after an average follow-up of 17 years," *European Spine Journal*, vol. 15, no. 2, pp. 183–195, 2006.
- [52] R. D. Guyer, P. C. McAfee, R. J. Banco et al., "Prospective, randomized, multicenter food and drug administration investigational device exemption study of lumbar total disc replacement with the CHARITÉ artificial disc versus lumbar fusion: five-year follow-up," *Spine Journal*, vol. 9, no. 5, pp. 374–386, 2009.
- [53] K. D. van den Eerenbeemt, R. W. Ostelo, B. J. van Royen, W. C. Peul, and M. W. van Tulder, "Total disc replacement surgery for symptomatic degenerative lumbar disc disease: a systematic review of the literature," *European Spine Journal*, vol. 19, no. 8, pp. 1262–1280, 2010.
- [54] P. Park, H. J. Garton, V. C. Gala, J. T. Hoff, and J. E. McGillicuddy, "Adjacent segment disease after lumbar or lumbosacral fusion: review of the literature," *Spine*, vol. 29, no. 17, pp. 1938–1944, 2004.
- [55] V. K. Goel, T. H. Lim, J. Gwon et al., "Effects of rigidity of an internal fixation device: a comprehensive biomechanical investigation," *Spine*, vol. 16, no. 3, pp. S155–S161, 1991.
- [56] S. M. Juratli, G. M. Franklin, S. K. Mirza, T. M. Wickizer, and D. Fulton-Kehoe, "Lumbar fusion outcomes in Washington State workers' compensation," *Spine*, vol. 31, no. 23, pp. 2715–2723, 2006.
- [57] A. Waguespack, J. Schofferman, P. Slosar, and J. Reynolds, "Etiology of long-term failures of lumbar spine surgery," *Pain Medicine*, vol. 3, no. 1, pp. 18–22, 2002.
- [58] P. C. McAfee, I. D. Farey, C. E. Sutterlin, K. R. Gurr, K. E. Warden, and B. W. Cunningham, "The effect of spinal implant rigidity on vertebral bone density: a canine model," *Spine*, vol. 16, no. 6, supplement, pp. S190–S197, 1991.
- [59] N. Boos and J. K. Webb, "Pedicle screw fixation in spinal disorders: a European view," *European Spine Journal*, vol. 6, no. 1, pp. 2–18, 1997.
- [60] H. Bozkus, M. Şenoğlu, S. Baek et al., "Dynamic lumbar pedicle screw-rod stabilization: in vitro biomechanical comparison with standard rigid pedicle screw-rod stabilization—laboratory investigation," *Journal of Neurosurgery: Spine*, vol. 12, no. 2, pp. 183–189, 2010.
- [61] W. Schmoelz, U. Onder, A. Martin, and A. Von Stempel, "Non-fusion instrumentation of the lumbar spine with a hinged pedicle screw rod system: an in vitro experiment," *European Spine Journal*, vol. 18, no. 10, pp. 1478–1485, 2009.
- [62] V. K. Goel, R. J. Konz, H. T. Chang et al., "Hinged-dynamic posterior device permits greater loads on the graft and similar stability as compared with its equivalent rigid device: a three-dimensional finite element assessment," *Journal of Prosthetics and Orthotics*, vol. 13, no. 1, pp. 17–20, 2001.
- [63] B. Cakir, M. Richter, K. Huch, W. Puhl, and R. Schmidt, "Dynamic stabilization of the lumbar spine," *Orthopedics*, vol. 29, no. 8, pp. 716–722, 2006.
- [64] D. Grob, A. Benini, A. Junge, and A. F. Mannion, "Clinical experience with the dynesys semirigid fixation system for the lumbar spine: surgical and patient-oriented outcome in 50 cases after an average of 2 years," *Spine*, vol. 30, no. 3, pp. 324–331, 2005.
- [65] P. Korovessis, Z. Papazisis, G. Koureas, and E. Lambiris, "Rigid, semirigid versus dynamic instrumentation for degenerative lumbar spinal stenosis: a correlative radiological and clinical analysis of short-term results," *Spine*, vol. 29, no. 7, pp. 735–742, 2004.
- [66] C. C. Würigler-Hauri, A. Kalbarczyk, M. Wiesli, H. Landolt, and J. Fandino, "Dynamic neutralization of the lumbar spine after microsurgical decompression in acquired lumbar spinal stenosis and segmental instability," *Spine*, vol. 33, no. 3, pp. E66–E72, 2008.
- [67] R. C. Huang, F. P. Girardi, M. R. Lim, and F. P. Cammisia, "Advantages and disadvantages of nonfusion technology in spine surgery," *Orthopedic Clinics of North America*, vol. 36, no. 3, pp. 263–269, 2005.
- [68] J. S. Harrop, J. A. Youssef, M. Maltenfort et al., "Lumbar adjacent segment degeneration and disease after arthrodesis and total disc arthroplasty," *Spine*, vol. 33, no. 15, pp. 1701–1707, 2008.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

