Synthesis, Characterization, and Biotoxicity of $\widehat{N N}$ Donor Sulphonamide Imine Silicon(IV) Complexes

Mukta Jain¹ and R. V. Singh²

¹ Department of Chemistry, Seth G. B. Podar College, Nawalgarh, Jhunjhunu, Rajasthan-333042, India ² Department of Chemistry, University of Rajasthan, Jaipur-302004, India

Received 20 April 2004; Revised 30 July 2004; Accepted 8 October 2004

The organosilicon derivatives of 2-[1-(2-furayl)ethyledene]sulphathiazole with organosilicon chlorides have been synthesised and characterized on the basis of analytical, conductance, and spectroscopic techniques. Probable trigonal bipyramidal and octahedral structures for the resulting derivatives have been proposed on the basis of electronic, IR, ¹H, ¹³C NMR, and ²⁹Si NMR spectral studies. In the search for better fungicides, bactericides, nematicides, and insecticides studies were conducted to assess the growth-inhibiting potential of the synthesized complexes against various pathogenic fungal, bacterial strains, root-knot nematode *Meloidogyne incognita*, and insect *Trogoderma granarium*. These studies demonstrate that the concentrations reached levels which are sufficient to inhibit and kill the pathogens, nematode, and insect.

Copyright © 2006 M. Jain and R. V. Singh. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Sulpha drugs are a group of compounds used for eliminating a wide range of infections in human and other animal systems. Many chemotherapeutically important sulpha drugs, like sulphadiazine, sulphathiazole, sulphamerazine, and so forth, possess SO_2NH moiety which is an important toxophoric function [1]. The heterocyclic compounds with both sulphur and nitrogen atoms in the ring system have also been used in the synthesis of biologically active complexes. It is however noteworthy that the biological activity gets enhanced on undergoing complexation with metal ions [2]. Schiff bases and their metal complexes have exhibited biological activity as antibiotics, antiviral, and antitumor agents because of their specific structures. Heteronuclear Schiff base complexes have been found in applications as magnetic materials, catalysts and in the biological engineering field [3–6].

Organosilicon compounds of sulphur-containing ligands have attracted much attention recently due to their biological importance. The sulphur containing ligands are well known for their anticarcinogenic, antibacterial, tuberculostatic, antifungal, insecticidal, and acaricidal activities. It has been reported that the activity of sulphur-containing ligand increases on complexation [7–15]. The interest in organosilicon(IV) compounds is due to their versatile applicability in the pharmaceutical industries. Generally, organosilicon compounds seem to own their antitumour properties to the immuno defensive system of the organism [16–19]. The medical applications and effectiveness of the silatranes in the treatment of wounds and tumours are thought to be related to the role of silicon in the growth of epithelial and connective tissues and hair, where its function is to impart strengths, elasticity, and impermeability to water [20].

The preparation and characterization of one biologically active sulphonamide imine derived from 2-acetylfuran with sulphathiazole and its silicon(IV) complexes form the subject of this paper. The results of these investigations seem to be promising. Based on the coordination sites available in the ligand system, this has been classified as monobasic bidentate ligand (Scheme 1).

EXPERIMENT

Adequate care was taken to keep the organosilicon(IV) complexes, chemicals, and glass apparatus free from moisture; clean and well-dried glass apparatus fitted with quickfit interchangeable standard ground joints was used throughout the experimental work. All the chemicals and solvents used were dried and purified by standard methods.

Physical measurements and analytical methods

Nitrogen and sulfur were estimated by the Kjeldhal's and Messenger's methods, respectively. Silicon was determined



Scheme 1

gravimetrically as SiO₂. Molecular weights were determined by the Rast camphor method (freezing point depression method) using resublimed camphor (MP 178°C).

Conductance measurements

The conductance measurements were carried out in dry dimethylformamide (DMF) at room temperature using a systronics conductivity bridge (model 305) in conjunction with a cell having a cell constant of 1.0.

Electronic spectra

The electronic spectra were recorded on a Perkin Elmer UV visible spectrophotometer in the range 200–600 nm, using dry methanol as the solvent.

IR spectra

Infrared spectra were recorded on a Nicolet Magna FT-IR 550 spectrophotometer in KBr pellets.

Nuclear magnetic resonance measurements

Multinuclear magnetic resonance spectra (¹H, ¹³C, and ²⁹Si) were recorded on an FX 90 Q JEOL spectrometer operating at 90 MHz.

¹H NMR spectra

¹H NMR spectra were recorded in deuterated methanol at 89.55 MHz using tetramethylsilane (TMS) as an internal standard.

¹³C NMR spectra

¹³C NMR spectra were recorded in dry methanol using TMS as the internal standard at 22.49 MHz.

²⁹Si NMR spectra

 29 Si NMR spectra were recorded at 17.75 MHz using deuterated dimethylsulphoxide (DMSO-d₆) as the solvent.

Preparation of the ligand

The sulphonamide imine was prepared by the condensation of 2-acetylfuran with sulphathiazole in equimolar ratio in absolute alcohol. The contents were refluxed for 3–4 hours and the solid which separated out was filtered off, recrystallized from the same solvent (ethanol), and dried in vacuo. The physical properties and microanalysis of this sulphonamide imine are recorded in Table 1.

Synthesis of the organosilicon(IV) complexes

For the synthesis of the complexes, first the sodium salt of the ligand was prepared by dissolving sodium metal (0.04-0.07 g) in 30 mL of methanol. Now to the weighed amount of organosilicon chlorides in 1 : 1 (0.38-0.51 g) or 1 : 2 molar ratios (0.11-0.29) in 20 mL methanol, the above prepared sodium salt of the ligand was added. The solution was refluxed for a period of 15–17 hours. The white precipitate of sodium chloride, formed during the course of the reaction, was removed by filtration and the filtrate was dried under reduced pressure. The resulting product was repeatedly washed with a mixture of methanol and *n*-hexane (1 : 1 v/v) and then finally dried for 3–4 hours. The purity was further checked by TLC using silica gel G. The details of these reactions and the analyses of the resulting products are recorded in Table 1.

RESULTS AND DISCUSSION

The 1 : 1 and/or 1 : 2 molar reactions of Me₂SiCl₂, Ph₂SiCl₂, and Ph₃SiCl with sulphonamide imine have led to the formation of Me₂SiCl(2-Ac-F-St), Me₂Si(2-Ac-F-St)₂, Ph₂SiCl(2-Ac-F-St), Ph₂Si(2-Ac-F-St)₂, and Ph₃Si(2-Ac-F-St) types of complexes. The reactions have been carried out in perfectly dry methanolic medium and proceed smoothly with the precipitation of NaCl. These reactions can be represented by the general equations in Scheme 2 showing the formations of the sodium salt and the complexes.

The resulting coloured solids are soluble in most of the common organic solvents. These have been found to be monomeric as evidenced by their molecular weight determinations. The low values of molar conductivity (10– $27 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$) of the resulting silicon complexes in anhydrous DMF show them to be nonelectrolytes in nature.

UV spectra

The electronic spectra of the sulphonamide imine and its 1 : 1 and 1 : 2 organosilicon(IV) complexes have been recorded. The spectrum of the ligand shows a broad band at 370 nm which can be assigned to the n- π * transitions of the azomethine group. This band shows a blue shift in the silicon complexes appearing at 351, 353, 359, and 355, 362 nm for 1 : 1 and 1 : 2 derivatives, respectively, due to the polarisation within the >C=N chromophore caused due to formation of covalent silicon–nitrogen bond. The bands at 255 and 285 nm are due to π - π * transitions, within the benzene ring and (>C=N) band of the azomethine group,

						Elemental analysis (%)							
Compound	Re	actant	(g)	Colour	Yield	MP	С	Η	Ν	S	Si	Cl	Mol Wt
Compound	M^*	LH^*	Na*	and state	(%)	(°C)	Found						
							(Calcd)						
(2-Ac-F-StH)				Light vellow	73	124_130	51.62	3.51	11.84	18.19			325
				Light yenow	75	124-130	(51.86)	(3.77)	(12.09)	(18.45)			(347.39)
MesSiCl(2-Ac-F-St)	0.38	1.02	0.07	Dark brown	74	71_73	46.18	3.88	9.26	14.19	6.07	8.00	412
$\operatorname{Me}_2\operatorname{SICI}(2-\operatorname{AC-I})$	0.50 1.02	1.02	0.07	solid	/1	/1 /5	(46.40)	(4.12)	(9.55)	(14.57)	(6.38)	(8.05)	(439.99)
$Me_{Si}(2-Ac_{F}-St)$	0.11	0.61	0.04	Light brown	76	109_111	50.99	3.67	10.77	16.70	3.48		738
Wic201(2-7:c-1-0t)2	0.11	0.01	0.01	solid	70	109–111	(51.18)	(4.02)	(11.19)	(17.07)	(3.74)		(750.92)
$Pb_{SiC}(2 \land c \in St)$	0.48	0.66	0.04	Dark brown	74	140 151	57.15	3.58	7.09	11.00	4.71	5.92	542
$r_{12}31C1(2-AC-1-5t)$	0.40	0.00	0.04	solid	74	149-131	(57.48)	(3.93)	(7.44)	(11.36)	(4.97)	(6.28)	(564.13)
Dh C:(2 A = E Ct)	0.20	0.01	0.05	Dark brown	77	165 167	57.31	3.74	9.42	14.19	3.00		858
$PII_2 SI(2-AC-F-SI)_2$	0.29	0.81	0.05	solid	//	155-157	(57.64)	(3.91)	(9.60)	(14.65)	(3.20)	_	(875.05)
$Ph_Si(2 - Ac_F_St)$	0.51	0.60	0.04	Duorum oolid	81	90_92	65.02	4.12	6.68	10.19	4.22	_	588
1 11301(2-11-01)	0.51	0.00	0.04	brown sond	01	90-92	(65.42)	(4.49)	(6.93)	(10.58)	(4.63)		(605.78)

TABLE 1: Analysis and physical properties of the ligand and its silicon complexes.

*M = silicon compound, LH = ligand, and Na = sodium metal.

respectively. The K band $\pi - \pi^*$ showed a red shift due to the overlap of the central metal d-orbital with the p-orbital of the donor atom which causes an increase in conjugation and the B-bands undergo a hypsochromic shift in the complexes [21], see Table 2.

IR spectra

The assignments of characteristic IR frequencies for the resulting complexes may be discussed as follows.

The IR spectra of these derivatives do not show any band in the region 3400-3150 cm⁻¹ which could be assigned to ν NH. This clearly indicates the deprotonation of the ligand as a result of complexation with the silicon atom. A sharp band at 1628 cm⁻¹ due to $\nu(>C=N)$ frequency of the free azomethine group in the ligand shifts to the lower frequency (ca 15 cm⁻¹) in the silicon complexes and indicating thereby the coordination of the azomethine nitrogen to the silicon atom. A shift of this frequency to the higher and lower wave number side as well as the "no change" has also been reported in the literature [16].

In dimethylsilicon(IV) complexes, a band at ca 1420 cm⁻¹ has been ascribed to the asymmetric deformation vibrations of (CH₃-Si) group, whereas the band at ca 1270 cm⁻¹ has been ascribed to the symmetric deformation mode of (CH₃-Si) group. New bands are observed in the spectra of the complexes at ca 570–582 cm⁻¹ due to the ν (Si \leftarrow N) vibrations. These remain absent in the spectrum of the ligand. A band due to ν (Si-Cl) at ca 423 and 439 cm⁻¹ is observed in 1 : 1 diorganosilicon(IV) derivatives. It has been reported [16] that the cis form of such complexes gives rise to two ν (Si \leftarrow N) bands, whereas in the transform only one IR active ν (Si \leftarrow N) band is observed. The presence of

only one $\nu(\text{Si} \leftarrow \text{N})$ band in the present case suggests that the complexes exist in the transform, see Table 3

¹H NMR spectra

The proton magnetic resonance spectral data of sulphonamide imine and its corresponding silicon complexes have been recorded in DMSO-d₆. The chemical shift values relative to the TMS peak are listed in Table 4.

The broad signal due to the -NH proton in the ligand disappears in the case of silicon complexes showing the coordination of silicon to nitrogen after the deprotonation of the functional group. The azomethine proton signal due to methyl proton $(H_3 - \overset{\downarrow}{C} = N)$ appears at δ 2.10 ppm in the ligand. The downfield shift of this position in the spectra of the complexes substantiates the coordination of azomethine nitrogen to the silicon atom. The additional signal in the region δ (1.01 and 1.13 ppm) in Me₂SiCl(2-Ac-F-St) and Me₂Si(2-

Ac-F-St)₂ types of complexes are due to Me₂Si group. The ligand shows a complex pattern in the region δ 8.10– 6.92 ppm for the aromatic protons and this is observed in the region δ 8.78–6.95 ppm in the spectra of the organosilicon(IV) complexes. This shifting also supports the coordination through the nitrogen atom.

¹³C NMR spectra

The conclusions drawn from the UV, IR, and ¹H NMR spectra are concurrent with the ¹³C NMR spectral data regarding the confirmation of the proposed structure. ¹³C NMR spectra of the ligand and its silicon complexes were also recorded in dry DMSO. The shifting of the signals due to carbon attached to the azomethine nitrogen in the spectra of



SCHEME 2: General equations showing the formations of the sodium salt and the complexes (R = Me and Ph).

the complexes further supports the involvement of this group in complexation [15]. Data are recorded in Table 5.

²⁹Si NMR spectra

The ²⁹Si NMR spectra of Me₂SiCl(2-Ac-F-St), Ph₂SiCl(2-Ac-F-St), and Ph₃Si(2-Ac-F-St) give sharp signals at δ -91 to δ -98 ppm and the spectra of Me₂Si(2-Ac-F-St)₂ and Ph₂Si(2-Ac-F-St)₂ give sharp signals at δ -128 to δ -110 ppm, which clearly indicates the penta- and hexa-coordinated environment, respectively, around the silicon atom. Though, the exact geometries of these complexes can be suggested on the basis of X-ray crystal structure; inspite of our best efforts we could not develop a suitable crystal for the X-ray studies. Hence, X-ray data could not be included in the present paper.

TABLE 2: UV spectral data of the ligand and its silicon complexes.

Ligand/complay	n - π^* (nm)	π - π^* (nm)	π - π^* (nm)
Ligand/complex	>C=N	C ₆ H ₅ ring	>C=N
(2-Ac-F-StH)	370	255	285
Me ₂ SiCl(2-Ac-F-St)	359	273	281
$Me_2Si(2-Ac-F-St)_2$	362	276	278
Ph ₂ SiCl(2-Ac-F-St)	351	280	275
$Ph_2Si(2-Ac-F-St)_2$	355	285	271
Ph ₃ Si(2-Ac-F-St)	353	290	268

Thus, on the basis of the above spectral features, as well as the analytical data, the penta-coordinated trigonal bipyramidal and hexa-coordinated octahedral geometries shown in Figure 1 have been suggested for the organosilicon(IV) complexes.

Compound/ligand	$\nu(\rm NH)$	ν (C=N)	$\nu(Si \leftarrow N)$	ν (Si-Cl)
(2-Ac-F-StH)	3400-3150 (m)*	1628 (vs)*	_	_
Me ₂ SiCl(2-Ac-F-St)		1622	577 w*	423 m
Me ₂ Si(2-Ac-F-St) ₂		1625	582 w	_
Ph ₂ SiCl(2-Ac-F-St)		1619	574 w	439 m
Ph ₂ Si(2-Ac-F-St) ₂		1613	576 w	_
Ph ₃ Si(2-Ac-F-St)		1616	570 w	_

TABLE 3: IR spectral data (cm⁻¹) of the ligand and its silicon complexes.

* m = medium, vs = very strong, and w = weak.

TABLE 4: ¹H NMR spectral data (δ , ppm) of the ligand and its silicon complexes.

Ligand/complex	Si-CH ₃	CH ₃	NH	Aromatic proton	²⁹ Si NMR
(2-Ac-F-StH)	_	2.10 (3H, s*)	10.54 (br*, 1H)	8.10-6.92 (m)*	
Me ₂ SiCl(2-Ac-F-St)	1.01 (1s, 6H)	2.25 (3H, s)		8.36-7.20 (m)	-98 (ppm)
Me ₂ Si(2-Ac-F-St) ₂	1.13 (s, 6H)	2.17 (6H, s)		8.784-7.00 (m)	-128 (ppm)
Ph ₂ SiCl(2-Ac-F-St)	_	2.22 (3H, s)	_	8.48-6.95 (m)	-94 (ppm)
Ph ₂ Si(2-Ac-F-St) ₂	_	2.15 (6H, s)	_	8.56-7.30 (m)	-110 (ppm)
Ph ₃ Si(2-Ac-F-St)		2.19 (3H, s)	_	8.51-7.14 (m)	-91 (ppm)

* m = multiplet, br = broad, and s = singlet.

BIOLOGICAL ASPECTS

Fungicidal, bactericidal, nematicidal, and insecticidal activities of the sulphonamide imine and its respective organosilicon(IV) complexes against pathogenic fungi, bacteria, rootknot nematode, and insect are recorded in Tables 6–12.

Antifungal screening

Like plant cells, fungi also possess cell walls but they cannot perform photosynthesis, moulds spoil food, damage potato, and crop plants (corn and wheat). They also cause rotting of clothes, shoes, and wooden materials. Some fungi cause diseases like athlete's foot and ring worm.

Method

The antifungal activities were evaluated against *Macrophomina phaseolina*, *Aspergillus niger*, *Fusarium oxysporum*, and *Alternaria alternata* by agar plate technique [22]. The compounds were dissolved in 25, 50, and 100 ppm concentrations in methanol and then mixed with the medium. The linear growth of the fungus was obtained by measuring the diameter of the colony after 96 hours. The inhibition percentage was calculated as 100 $(D_{fc} - D_{ft})/D_{fc}$, where D_{fc} and D_{ft} are the diameters of the fungus colony in the control and the test plates, respectively.

Antibacterial screening

Of all the microorganisms, bacteria are the most abundant. They generally reproduce quite fast, such as *P cepacicola* which reproduces itself every 9.5 minutes. However, some bacteria are very slow growing, such as those that cause tuberculosis and leprosy. This makes early diagnosis of these diseases rather difficult. The most common bacteria used for scientific research is *E coli*. Its normal living place is the lower human intestine (COLON).

Method

Bactericidal activities were evaluated by the paper disc plate method [23]. The nutrient agar medium (peptone, beef extract, NaCl, and agar-agar) and 5 mm diameter paper discs (Whatman No. 1) were used. The compounds were dissolved in methanol in 500 and 1000 ppm concentrations. The filter paper discs were soaked in different solutions of the compounds, dried, and then placed in the petri plates previously seeded with the test organisms (*P cepacicola, E coli, K aerogenous,* and *S aureus*). The plates were incubated for 24–30 hours at $28 \pm 2^{\circ}$ C and the inhibition zone around each disc was measured.

Observations

The free ligand and its respective metal chelates were screened against selected fungi and bacteria to assess their potential as antimicrobial agents. The results are quite promising. The antimicrobial data reveal that the complexes are superior than the free ligands. The enhanced activity of the silicon chelates may be ascribed to the increased lipophilic nature of these complexes arising due to the chelation [24]. The toxicity increased as the concentration was increased. Further, the results of bioactivity were compared with the conventional fungicide, *Bavistin*, and the conventional bactericide, *Streptomycin*, taken as standards in either case.

Lizzan d/agamentary	Anomethine Center	S: CU	C ₁	C ₂	C ₃	C_4	C ₉	C
Ligand/complex	Azometime C-atom	3I-CH3	C_5	C_6	C ₇	C_8	C ₁₀	C_{11}
(2 - Ac - E - StH)	155.01		146.02	138.99	120.98	143.94	152.60	151.80
(2-76-1-5011)	155.71		128.01	122.46	124.01	125.98	150.00	151.00
$M_{0} \in Cl(2, \Lambda_{0} \in \Omega_{0})$	144.76	12.09	148.91	139.21	121.12	142.92	149.20	150.70
$Me_2SiCI(2-Ac-F-St)$	144.70	15.98	125.96	120.24	123.67	119.76	150.95	150.70
	140 51	15.01	147.69	140.96	121.32	143.01	149.45	140.05
$Me_2 SI(2-AC-F-SI)_2$	148.51		124.21	120.96	122.21	119.10	151.00	149.03
$\mathbf{D}_{\mathbf{L}}^{\mathbf{L}}$ $\mathbf{C}_{\mathbf{L}}^{\mathbf{L}}$ \mathbf{L} \mathbf{L} \mathbf{L} \mathbf{L} \mathbf{L}	152.46	_	146.36	137.01	120.76	142.10	151.20	140.70
$Pn_2SICI(2-AC-F-St)$	155.46		126.01	121.78	119.98	122.46	150.15	149./0
	146.56		145.16	138.06	127.92	143.21	147.20	150.10
$Ph_2Si(2-Ac-F-St)_2$	146.76	—	128.96	121.02	120.21	123.74	149.80	150.12
	15100		144.05	133.42	120.81	143.40	148.78	
Ph ₃ Si(2-Ac-F-St)	154.90	—	127.01	121.98	123.32	124.86	150.55	149.40

TABLE 5: ¹³C NMR spectral data (δ , ppm) of the ligand and its silicon complexes.





Where, R = Me / Ph and X = Ph/Cl

FIGURE 1: (a) The penta-coordinated trigonal bipyramidal and (b) hexa-coordinated octahedral geometries R = Me/Ph and X = Ph/Cl.

In fungicide activity, most of the organosilicon(IV) complexes were able to inhibit and kill the pathogens at 50 ppm concentration, whilst 100 ppm concentration proved invariably fatal. None of the fungi was able to withstand this concentration. In bactericidal activity, the complexes exhibited remarkable potential in inhibiting the growth of pathogens. Many of the complexes were found to be even more toxic than the standard. Thus, it can be postulated that further intensive studies of these complexes in this direction as well as in agriculture could lead to the interesting results.

Nematicidal activity

Development of the concept of pest management and their implementation have led to a greater appreciation of the need for a wide range of tactics for nematode control. The objective of nematode control is to improve growth and yield of plants, which can be achieved through a reduction of the nematode population in soil or in plants, or through a reduction of their damage. Chemical method can be used to control nematodes [25]. *M incognita* produce galls on the roots of many host plants and responsible for 44.87 percent of yield loss in brinjal [26].

Method

First of all we applied different concentrations (25, 50, and 100) in ppm of complexes and ligand on root-knot nematode *M* incognita spp. in a step-by-step [27] procedure. For experiment, egg masses were separated from heavily infected brinjal roots and washed under running water. After cutting the roots, one percent of sodium hypochlorite solution was added, shaked, and then sieved through 150 and 400 sieves. Then the eggs of nematode were counted and replicated three times. At this experiment, temperature range was $30 \pm 2^{\circ}$ C.

Observations

Maximum hatching was recorded in control. All the metal complexes are more toxic than the ligand and all bimolar complexes are more active than unimolar organosilicon derivatives. Dimethylsilicon(IV) complexes are less hazardous than diphenylsilicon(IV) complexes. The activity increases with increasing the concentration of the solutions.

Mode of action [15]

Much smaller amounts of the nonfumigant and fumigant nematicides are needed in plant protection against nematode because the indirect hematostatic effects of non-fumigant nematicides resulting from impairment of neuromuscular activity, interfere with movement, feeding, invasion, development, reproduction, fecundity, and hatching of nematodes

Ligand/complex	Aspergillus niger			Macrophomina phaseolina			Fusar	Fusarium oxysporum			Alternaria alternata		
Ligand/complex	25	50	100	25	50	100	25	50	100	25	50	100	
$(2 \Delta c_E StH)$	34	53	61	35	50	68	39	56	65	43	60	66	
(2-/10-1-5011)	(50.72)	(38.37)	(37.75)	(51.38)	(39.02)	(29.16)	(44.28)	(38.46)	(35.00)	(39.44)	(30.23)	(34.00)	
M_{0} SiCl(2 Ac E St)	37	56	72	38	52	71	42	59	71	45	62	68	
Wic25iCi(2-/ic-1-5t)	(46.37)	(34.88)	(36.11)	(47.22)	(36.58)	(26.04)	(40.00)	(35.16)	(29.00)	(36.62)	(27.91)	(32)	
$M_{\alpha} S(2 \Lambda_{\alpha} E S_{4})$	42	63	78	41	57	74	46	65	74	47	65	72	
Wic201(2-AC-1-0t)2	(39.13)	(26.74)	(20.40)	(43.05)	(30.48)	(22.44)	(35.28)	(28.57)	(26.00)	(33.80)	(24.42)	(28.00)	
Ph. SiCl(2 $\triangle c$ E St)	38	57	76	40	53	72	43	61	73	46	63	70	
1 H ₂ 5ICI(2-AC-1-5t)	(44.92)	(33.72)	(22.44)	(44.44)	(35.36)	(26.53)	(38.57)	(32.96)	(27.00)	(35.21)	(26.74)	(30.00)	
$Ph_{Si}(2 \land c \in St)$	44	66	82	47	61	80	48	67	78	49	66	76	
(36.23	(36.23)	(23.25)	(16.32)	(34.72)	(25.60)	(18.36)	(31.42)	(26.37)	(22.00)	(30.99)	(23.26)	(24.00)	
$Ph_{Si}(2 A_{C} F_{St})$	40	60	80	42	54	73	45	63	75	47	64	71	
r11301(2-AC-F-SL)	(42.02)	(30.23)	(18.36)	(41.66)	(34.14)	(25.51)	(35.71)	(30.76)	(25.00)	(33.80)	(25.58)	(29.00)	
Bavistin	69	86	98	72	82	96	70	91	100	71	86	100	

TABLE 6: Fungicidal screening data of the ligand and its silicon complexes inhibition percentage after 96 hours and SD values (25, 50, and 100 are concentrations in ppm).

TABLE 7: Bactericidal screening data of the ligand and its silicon complexes diameter inhibition zone (mm) after 24 hours (500 and 1000 are concentrations in ppm).

Ligand/complex	Esherichia coli (–)		Klebsiella aerogenous (–)		Pseudom	onas cepacicola (–)	Staphylococcus aureus (+)	
Ligand/complex	500	1000	500	1000	500	1000	500	1000
(2-Ac-F-StH)	6	6	6	11	10	12	9	13
$Me_2SiCl(2\text{-}Ac\text{-}F\text{-}St)$	8	12	9	15	12	14	12	14
$Me_2Si(2-Ac-F-St)_2$	10	16	11	17	15	17	16	18
Ph ₂ SiCl(2-Ac-F-St)	10	14	10	16	14	16	15	16
Ph ₂ Si(2-Ac-F-St) ₂	13	18	14	19	17	19	18	19
Ph ₃ Si(2-Ac-F-St)	11	16	12	17	15	17	16	17
Streptomycin	1	2	3	5	2	5	15	17

TABLE 8: Nematicidal screening data of the ligand and its silicon complexes (25, 50, and 100 are concentrations in ppm).

Ligand/complex	(% of hatching <i>M incognita</i>)						
Ligand/complex	25	50	100				
(2-Ac-F-StH)	22.5	19.0	15.0				
$Me_2SiCl(2-Ac-F-St)$	20.2	16.5	No hatching				
$Me_2Si(2-Ac-F-St)_2$	18.5	14.9	No hatching				
Ph ₂ SiCl(2-Ac-F-St)	19.6	16.4	No hatching				
$Ph_2Si(2-Ac-F-St)_2$	15.9	11.9	No hatching				
Ph ₃ Si(2-Ac-F-St)	18.6	14.0	No hatching				

which are considered more important than their direct killing action.

Insecticidal activity

Many insects cause injury to economic plants by feeding on them externally: by chewing their leaves or other part: In order to raise more food, man has devised methods to alter normal population growth of many insect pests by reducing their chance for survival. To control the insect pests, the man since long has been employing various strategies which include mechanical, physical, chemical, and biological methods [28].

Methods

Ovicidal

To determine the efficacy of complexes as ovicide, eggs were treated by contact method. By spreading 1 mL of complex solutions on petri dishes (5.0 cm diameter), a thin film of 100 and 200 concentrations were prepared. The solvent was allowed to evaporate 200 eggs for 0–24 hours and kept in contact with the insecticidal film through out their incubation period. A control with each experiment was also run in which the eggs were kept in 1 mL of solvent. By Abott's formula [29], percentage of egg of mortality and percentage of corrected egg of mortality were calculated.

% corrected mortality =
$$\frac{K_T - K_C}{100 - K_C} \times 100$$
, (1)

where $K_T = \%$ kill in treated, $K_C = \%$ kill in control.

Ligand/complex	Dose level	Average no. of eggs hatched	Average no. of eggs unhatched	% eggs hatching	% eggs unhatched	% corrected mortality
$(2 \land c \in StH)$	100	15	5	75	25	21.05
(2-76-1-5111)	200	11	9	55	45	42.10
$M_{0} S(C)(2 \wedge c E S(t))$	100	13	7	65	35	31.57
Me231CI(2-AC-F-St)	200	9	11	45	55	52.63
	100	9	11	45	55	52.63
$Me_2SI(2-AC-F-SI)_2$	200	7	13	35	65	63.15
Dh $C(C)(2 \wedge a \in C_{4})$	100	11	9	55	45	42.10
FII25ICI(2-AC-F-5t)	200	7	13	35	65	63.15
$\mathbf{D}_{\mathbf{L}} \in (\mathbf{C}, \mathbf{A}, \mathbf{C}, \mathbf{C})$	100	8	12	40	60	57.89
$PII_2 SI(2-AC-F-SI)_2$	200	5	15	25	75	73.68
Dh C:(2 A a E Ct)	100	10	10	50	50	47.36
$Pn_3 S1(2-AC-F-St)$	200	7	13	35	65	63.15
Control		19	1	95	5	_

TABLE 9: Ovicidal screening data of the ligand and its silicon complexes (100 and 200 are concentrations in ppm).

TABLE 10: Larvicidal screening data of the ligand and its silicon complexes (100 and 200 are concentrations in ppm).

Ligand/complex	Doca laval	Average no. of	Average no. of	% pupal	% larval	% corrected
Ligand/complex	Dose level	pupal formed	dead larvae	formation	mortality	mortality
$(2 \Lambda_c E StH)$	100	16	4	80	20	15.78
(2-710-1-5011)	200	13	7	65	35	31.57
M_{2} SiCl(2 Ac E St)	100	13	7	65	35	31.57
Me251CI(2-AC-F-5t)	200	10	10	50	50	47.36
M_{2} Si(2 Ac E St)	100	9	11	45	55	52.63
$Me_2 SI(2-Ac-1-St)_2$	200	7	13	35	65	63.15
Dh $S(C)(2 \wedge c \in St)$	100	11	9	55	45	42.10
FII23ICI(2-AC-F-3t)	200	8	12	40	60	57.89
Dh $Si(2 \land c \in St)$	100	7	13	35	65	63.15
$F_{112}31(2-AC-F-3t)_2$	200	5	15	25	75	73.68
Dh C:(2 A = E Ct)	100	11	9	55	45	42.10
r11301(2-AC-F-St)	200	6	14	30	70	68.42
Control	_	19	1	95	5	_

Larvicidal

By feeding method larvicidal efficacy of the synthesized chemicals was assessed. The last instar larvae were separated from subculture and kept in vials containing 5 g of topically treated wheat grains with 1 mL of chemicals. Until the pupal formation, larvae were allowed to continue their development on this diet, replicated thrice, each dose. The food was treated with solvent only in control. By Abott's formula, larval of mortality and percentage of corrected of mortality were calculated.

Pupicidal

From the subculture, the last larval instars were stored out and were kept in separate container. Pupal of known age (0-12 hours) were taken out and were dipped in the desired concentration (100 and 200) of the chemicals along with a control of three replicates that were set for each dose and total emergence and pupal of mortality were recorded after 96 hours. By Abott's formula, percentage of pupal of mortality and percentage of pupal corrected of mortality were calculated.

Adulticidal

By contact method the adulticidal action was assessed. With 1 mL of respective doses, 5 g of wheat grains were treated. The solvent was allowed to evaporate completely. Along with a control, experiment was replicated thrice. Newly emerged adults were taken from the subculture and were released in the plastic vials containing treated food. After 48-hour observations were taken and by Abott's formula, percentage of corrected of mortality was calculated.

Mode of action [30]

Some insecticides are physical poisons causing asphyxiation, some are protoplasmic poisons, a few are respiratory poisons,

	1	8 8	1	`		11 /
Ligand/complex	Dose level	Average no. of adults emerged	Average no. of pupal mortality	% emerged adult	% pupal mortality	% corrected mortality
$(2 \land c \in StH)$	100	15	5	75	25	21.05
(2-AC-1-5011)	200	14	6	70	30	26.31
M_0 , SiCl(2, Ac E St	100	14	6	70	30	26.31
Me251CI(2-AC-1-50)	200	11	9	55	45	42.10
$M_{2} C(2 A_{2} E Ct)$	100	10	10	50	50	47.36
$NIC_2 SI(2-AC-1-St)_2$	200	6	14	30	70	68.42
Dh SiC $(2 \land c \in St)$	100	12	8	60	40	38.84
$F_{112}SICI(2-AC-1-St)$	200	8	12	40	60	57.89
Dh Si(2 Ac E St)	100	9	11	45	55	52.63
$FII_2SI(2-AC-F-St)_2$	200	5	15	25	75	73.68
Dh Si(2 Ac E St)	100	11	9	55	45	42.10
FII33I(2-AC-F-3t)	200	7	13	35	65	63.15
Control	_	19	1	95	5	

TABLE 11: Pupicidal screening data of the ligand and its silicon complexes (100 and 200 are concentrations in ppm).

TABLE 12: Adulticidal screening data of the ligand and its silicon complexes (100 and 200 are concentrations in ppm).

Ligand/complex	Dose level	Average no. of adults in each vial	Average mortality after 48 hours	% adult mortality	% corrected mortality
$(2 \Delta c - E StH)$	100	20	4	20	15.78
(2-76-1-5011)	200	20	6	30	26.31
M_{e} -SiCl(2-Ac-E-St)	100	20	7	35	31.57
Wic20101(2-71C-1-51)	200	20	11	55	52.63
Me $Si(2 - Ac - E - St)$	100	20	11	55	52.63
$MC_2 OI(2 - MC - 1 - Ot)_2$	200	20	14	70	68.42
$Dh_{SiCl}(2 \land c \in St)$	100	20	9	45	42.10
FII25ICI(2-AC-1-5t)	200	20	12	60	57.10
$Dh_{Si}(2 \land c \in St)$	100	20	12	60	57.89
$F H_2 SI(2 - AC - F - SC)_2$	200	20	15	75	57.89
$Dh_{Si}(2 \land c \in St)$	100	20	10	50	73.68
r11301(2-AC-F-St)	200	20	4	70	47.36
Control		20	1	5	68.42

but the majority of them are nerve poisons. The action of insecticides upsets the normal behaviour and actions of the target organisms.

Ovicidal, larvicidal, pupicidal, and adulticidal results are shown in Tables 9, 10, 11, and 12. The data indicated the same observations as were observed in nematicidal activity.

ACKNOWLEDGMENTS

The author Dr Mukta Jain is highly grateful to Honourable Shri Kanti Kumar R Podar and Shri Moti Chand Maloo for their cooperation, constant encouragement, moral support, and blessings during this research work. The authors are also thankful to UGC, New Delhi, India, for the financial assistance vide Grant no. F-12-18(SR-I)/2004.

REFERENCES

- Singh H, Srivastava VK, Shukla SN, Srivastava MK, Upadhyay MK. Synthesis, characterization and fungitoxicity of Mn(II), Fe(II), Co(II), Ni(II) and Zn(II) complexes of N-phenyl-5phenyl-1,3,4-oxydiazol-2-sulphonamide and 5-phenyl-1,3,4oxydiazol-2-iminosulphonamide. *Indian Journal of Chemistry*. 1994;33A:350–351.
- [2] Chaturvedi KK, Singh RV, Tandon JP. The reactions of Boron and Aluminium Isopropoxides with the imines derived from sulphonamides. *Journal für Praktische Chemie*. 1985;327(1):144–150.
- [3] Zhang H-Y, Lei J, Chen Y-Y, Wu Q-A, Zhang Y-S, Gao L-H. Synthesis of the N,N'-bis(ferrocenylmethylene)-1,2phenylenediamine Schiff base and six rare earth metal complexes. Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry. 2001;31(6):973–981.

- [4] Hong-Jian S, Xiao-Yan L, Xue-Gui C, De-Xin L. Studies on complexes of transition metals with Schiff bases II. Studies on the non-isothermal kinetics of thermal decomposition of a complexes of Nickel(II) with N-Salicyclidencglycinc. *Hua Xue Xue Bao. Acta Chimica Sinica*. 1992;13:1168–1170. Chem. Abstr., 1993. 118, 261640X.
- [5] Abele E. Activation of silicon bonds by fluoride ion in the organic synthesis in the new millenium: A Review. *Main Group Metal Chemistry*. 2005;28(2):45–69.
- [6] Ji-Tao W, Sheng-Hua G, Xiang-Min L, Feng-Quan L. Syntheses and characterization of zirconum(IV), tin(IV) and titanium(IV) complexes with acylhydrazones and thioacylhydrazones. *Hua Xue Xue Bao. Acta Chimica Sinica*. 1993;14(5):645– 650. Chem. Abstr., 1993, 119, 240289v.
- [7] Singh K. Synthesis and spectral studies of methylsilicon(IV) chloride complexes with biologically active heterocyclic benzothiazolines. *Indian Journal of Chemistry*. 1991;30A:283–285.
- [8] Belwal S, Singh RV. Bioactive versatile azomethine complexes of organotin(IV) and organosilicon(IV). Applied Organometallic Chemistry. 1998;12(1):39–46.
- [9] Phor A, Chaudhary A, Jain M, Swaroop R, Singh RV. Antimicrobial and antifertility activities of newly synthesized triorganosilicon(IV) complexes. *Main Group Metal Chemistry*. 2001;24:439–444.
- [10] Singh RV, Jain M, Nagpal P. Nitrogen-sulphur and nitrogenoxygen donor ferrocene containing ligands and their diorganosilicon(IV) complexes. *Main Group Metal Chemistry.* 2002; 25(10):615–619.
- [11] Jain M, Singh RV. Pesticidal and nematicidal properties of organosilicon(IV) derivatives of nitrogen-nitrogen donor sulphonamide imine. *International Journal for Chemical Science*. 2003;1(1):17–24.
- [12] Jain M, Kumar D, Singh RV. Toxicological aspects of the bioactive versatile sulphonamide-imine complexes of organosilicon(IV). *Main Group Metal Chemistry*. 2003;26(2):99–109.
- [13] Jain M, Gaur S, Diwedi SC, Joshi SC, Bansal A, Singh RV. Nematicidal, insecticidal, antifertility, antifungal and antibacterial activities of salicylanilide sulphathiazole and its manganese, silicon and tin complexes. *Phosphorus, Sulfur, and Silicon.* 2004;179:1517–1537.
- [14] Jain M, Singh RV. Spectral and antimicrobial studies of organosilicon(IV) complexes of a bidentate Schiff base having nitrogen-nitrogen donor system. *Main Group Metal Chemistry*. 2003;26(4):237.
- [15] Jain M, Gaur S, Singh VP, Singh RV. Organosilicon(IV) and organotin(IV) complexes as biocides and nematicides: synthetic, spectroscopic and biological studies of N^N donor sulfonamide imine and its chelates. *Applied Organometallic Chemistry*. 2004;18(2):73–82.
- [16] Saxena C, Singh RV. Diorganosilicon(IV) complexes of fluoroimines: synthetic, spectroscopic and biological aspects. *Phosphorus, Sulfur, and Silicon.* 1994;97:17–26.
- [17] Belwal S, Tanjea H, Dandia A, Singh RV. Novel bioactive thio- and semicarbazide ligands and their organo-silicon(IV) and organotin(IV) complexes. *Phosphorus, Sulfur, and Silicon.* 1997;127:49–58.
- [18] Belwal S, Saini RK, Singh RV. Synthesis and characterization of bioactive organotin(IV) and organosilicon(IV) complexes. *Indian Journal of Chemistry*. 1998;37A:245–248.
- [19] Singh MS, Bhagwat VW, Raju MD, Tiwari SK. Synthesis and characterization of hypervalent organosilicon compounds. *Indian Journal of Chemistry*. 1999;38A:716–719.

- [20] Saxena C, Singh RV. Organosilicon(IV) complexes of deprotomated form of semicarbazenes. Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry. 1992;22(8):1061–1072.
- [21] Singh RV, Joshi SC, Gajraj A, Nagpal P. Studies of biologically potent organotin(IV) and organosilicon(IV) complexes of a sulfur donor ligand derived from 1-acetylferrocene. *Applied Organometallic Chemistry*. 2002;16(12):713–720.
- [22] Fahmi N, Singh RV. Complexes of oxovanadium(V) with benzothiazolines. *Transition Metal Chemistry*. 1994;19:453–456.
- [23] Fahmi N, Singh RV. Synthetic, structural and biological studies of oxovanadium(V) complexes of azomethines. *Indian Journal of Chemistry*. 1998;37A:1126–1129.
- [24] Fahmi N, Jadon SCS, Singh RV. Biochemical studies on biologically active thio-ligands benzothiazolines and their manganese(II) complexes. *Phosphorus, Sulfur, and Silicon.* 1993; 81:133.
- [25] Trivedi PC, ed. Nematode Diseases in Plants. 1st ed. New Delhi, India: CBS; 1998.
- [26] Jain M, Nehra S, Trivedi PC, Singh RV. Biologically potent heterocyclic sulphonamide imines and their manganese(II) complexes. *Heterocyclic Communications*. 2003;9(1):95–98.
- [27] Jain M, Nehra S, Trivedi PC, Singh RV. Nematicidal, fungicidal and bactericidal activities of manganese(II) complexes with heterocyclic sulphonamide imines. *Metal-Based Drugs*. 2002;9:53–59.
- [28] Dwivedi SC, Mathur M. Evaluation of plant extracts as repellent against IV inster larvae of diamond blackmoth, Plutella xylostella (Lepidoptera plutellidae). J. Exotoxicol. Environ. Monit. 1999;9(3):163–166.
- [29] Dwivedi SC, Bajaj M. Efficacy of Acacia nilotica leaf extract as pupicidal against Trogoderma granarium (everts). *Indian J. Exp.* 2000;3:153–155.
- [30] Srivastava RP, Saxena RC. A Textbook of Insect Toxicology. New Delhi, India: Himanshu; 1989.



International Journal of Medicinal Chemistry



Organic Chemistry International





International Journal of Analytical Chemistry



Advances in Physical Chemistry



Research International

Catalysts



