

Research Article

Synthesis and Characterization of a Linear $[\text{Mn}_3(\text{O}_2\text{CMe})_4(\text{py})_8]^{2+}$ Complex

Eleni E. Moushi,¹ Christos Kizas,¹ Vassilios Nastopoulos,² and Anastasios J. Tasiopoulos¹

¹ Department of Chemistry, University of Cyprus, 1678 Nicosia, Cyprus

² Department of Chemistry, University of Patras, 26500 Patras, Greece

Correspondence should be addressed to Anastasios J. Tasiopoulos, atasio@ucy.ac.cy

Received 16 March 2010; Accepted 25 March 2010

Academic Editor: Spyros Perlepes

Copyright © 2010 Eleni E. Moushi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Two new compounds that consist of the linear trinuclear manganese(II) cation $[\text{Mn}_3(\text{O}_2\text{CMe})_4(\text{py})_8]^{2+}$ cocrystallizing with different counteranions (I_3^- , [1]; ClO_4^- , [2]) are reported. Complex 1 was prepared from the reaction of $[\text{Mn}(\text{O}_2\text{CMe})_2] \cdot 4\text{H}_2\text{O}$ with I_2 in $\text{MeCO}_2\text{H}/\text{py}$, whereas complex 2 was isolated from the reaction of $[\text{Mn}_3\text{O}(\text{O}_2\text{CMe})_6(\text{py})_3] \cdot \text{py}$ with $[\text{Mn}(\text{ClO}_4)_2] \cdot 6\text{H}_2\text{O}$ in MeCN/py . The crystal structures of both compounds were determined by single crystal X-ray crystallography. Magnetic susceptibility studies that were performed in microcrystalline powder of 1 in the 2–300 K range revealed the presence of antiferromagnetic exchange interactions that resulted in an $S = 5/2$ ground spin state.

1. Introduction

Oligonuclear Mn carboxylate clusters have attracted significant interest since they have been located in the active site of metalloenzymes [1] and also often have interesting and sometimes novel magnetic properties [2]. Undoubtedly, the most well-known oligonuclear cluster that appears in biological systems is the tetranuclear Mn complex that is present in the active site of photosystem II and is responsible for the light driven oxidation of water to molecular dioxygen [3–7]. Other Mn compounds observed in the active sites of metalloenzymes involve mononuclear (e.g., in Mn-superoxide dismutases) [8] and dinuclear (e.g., in Mn-catalases) complexes [8, 9]. In all those compounds, the ligation of the Mn ions is provided mainly by O- and N-donor atoms from the various aminoacid residues present in the metalloproteins. In order to prepare functional and structural models of the Mn compounds that are present in metalloenzymes, efforts have been centered on the synthesis and study of manganese carboxylate complexes with various chelating N-donor ligands, such as 2,2'-bipyridine (bpy) [10–13], 1,10-phenanthroline (phen) [10, 14–16], and 2-(2-pyridyl)benzimidazole [17]. As a result a

plethora of dinuclear, trinuclear and tetranuclear manganese compounds containing carboxylate groups or/and nitrogen-donor ligands have been prepared and characterized [4, 7, 10–30]. Such complexes are of significant interest not only as potential functional and structural models of the metal clusters present in Mn-containing metalloenzymes but also as precursors for the isolation of new model compounds. In particular, trinuclear Mn compounds have attracted significant attention since they appear as discrete metal clusters with various topologies including linear [10–26], triangular [27], V-shaped [28], and so forth, clusters and also as building blocks in multidimensional coordination polymers [29]. Linear trinuclear manganese (II) clusters with various molecular formulas such as $[\text{Mn}_3(\text{O}_2\text{CR})_6(\text{L})_2]$ [10–20] and $[\text{Mn}_3(\text{O}_2\text{CR})_4(\text{L}')_2]$ [21–23] have been prepared with several types of carboxylates, bidentate (L), and tridentate or tetradentate (L') chelates and also terminal ligands.

Herein, we report the synthesis, structural characterization, and magnetic properties of a new linear manganese(II) cation, $[\text{Mn}_3(\text{O}_2\text{CMe})_4(\text{py})_8]^{2+}$ which cocrystallizes with two different counteranions (I_3^- , [1] and ClO_4^- , [2]). The cation of 1 and 2 represents the first linear trinuclear Mn^{II} unit that contains only carboxylate and pyridine ligands and

TABLE 1: Crystallographic data for complexes **1** and **2**.

	1	2
Formula ^(a)	C ₄₈ H ₅₂ Mn ₃ N ₈ O ₈ I ₆	C ₄₈ H ₅₂ Mn ₃ N ₈ O ₁₆ Cl ₂
M_w	1795.20	1232.70
Crystal System	Monoclinic	Monoclinic
Space group	P2 ₁ /n	P2 ₁ /n
$a/\text{Å}$	15.2694(7)	21.7552(5)
$b/\text{Å}$	13.8883(4)	11.0081(2)
$c/\text{Å}$	15.2919(6)	23.4535(4)
$\beta/^\circ$	109.041(5)	107.917(2)
$V/\text{Å}^3$	3065.5(2)	5344.3(2)
Z	2	4
T/K	100(2)	100(2)
$\lambda^{(b)}$, Å	0.71073	0.71073
D_c , g/cm ⁻³	1.945	1.532
μ (Mo K α)/mm ⁻¹	3.682	0.874
Refl. collected/unique (R_{int})	23443/7336 (0.0455)	34188/9376 (0.0733)
Obs. refl. [$I > 2\sigma(I)$].	5655	4469
$R1\%^{(c)}$	0.0285	0.0382
$wR2\%^{(d)}$	0.0640	0.0636
Goodness of fit on F^2	0.952	0.736
Largest diff. peak/hole/e ⁻ /Å ⁻³	1.172/-1.043	0.366/-0.298

^(a)Including counteranions. ^(b)Graphite monochromator. ^(c) $R1 = \sum|F_o| - |F_c|/\sum|F_o|$. ^(d) $wR2 = [\sum[w(F_o^2 - F_c^2)^2]/\sum[wF_o^2]^2]^{1/2}$, $w = 1/[\sigma^2(F_o^2) + (m \cdot p)^2 + n \cdot p]$, $p = [\max(F_o^2, 0) + 2F_c^2]/3$, and m and n are constants.

a rare example of a linear Mn^{II}₃ cluster that is stabilized with carboxylate and terminal ligands without containing any polydentate chelates [30].

2. Experimental

2.1. Materials. All manipulations were performed under aerobic conditions using materials (reagent grade) and solvents as received; water was distilled in-house. [Mn₃O(O₂CMe)₆(py)₃] \cdot py was prepared as described elsewhere [27]. *Warning: Although we encountered no problems, appropriate care should be taken in the use of the potentially explosives perchlorate anion.*

2.2. Syntheses of Compounds

2.2.1. [Mn₃(O₂CMe)₄(py)₈](I₃)₂ [1]. Solid I₂ (2.07 g, 8.16 mmol) was added to the yellowish solution of [Mn(O₂CMe)₂] \cdot 4H₂O (2.00 g, 8.16 mmol) in MeCOOH/py (10/20 mL). The resulting red-brown solution was left under magnetic stirring for ~45 minutes, filtered off and the filtrate was left undisturbed at room temperature. After a few weeks, dark brown crystals of **1** suitable for X-ray crystallography were formed. The crystals were collected by filtration, washed with MeCOOH/py (5/10 mL) and dried in vacuum. The yield was ~60% based on total Mn content. A sample for crystallography was maintained in contact with the mother liquor to prevent the loss of interstitial solvent. *Anal. Calc.* for C₄₈H₅₂Mn₃N₈O₈I₆ [1]: C, 32.11; H, 2.92; N, 6.24. Found: C 31.89; H 2.79; N 6.10%. IR data (KBr pellet,

cm⁻¹): $\tilde{\nu} = 3435$ (m), 3059 (m), 1599 (s), 1580 (s), 1564 (s, br), 1483 (m), 1441 (s, br), 1350 (m), 1215 (m), 1151 (m), 1067 (m), 1038 (m), 1005 (m), 752 (m), 700 (s), 683 (m), 650 (m), 629 (m).

2.2.2. [Mn₃(O₂CMe)₄(py)₈](ClO₄)₂ [2]

Method A. To a solution of [Mn₃O(O₂CMe)₆(py)₃] \cdot py (0.294 g, 0.345 mmol) in MeCN/py (10/2 mL) was added Mn(ClO₄)₂ \cdot 6H₂O (0.125 g, 0.345 mmol) and pdH₂ (0.10 mL, 0.105 g, 1.38 mmol) and the mixture was left under magnetic stirring for ~30 minutes. The resulting dark red-brown slurry was filtered off and the dark red-brown filtrate was left undisturbed at room temperature. After few weeks yellow crystals appeared, suitable for X-ray structural determination. The crystals were isolated by filtration, washed with a copious amount of MeCN/py, and dried in vacuum; yield, ~20% based on total ClO₄⁻ content. A sample for crystallography was maintained in contact with the mother liquor to prevent the loss of interstitial solvent. *Anal. Calc.* for C₄₈H₅₂Mn₃N₈O₁₆Cl₂ [2]: C, 46.77; H, 4.25; N, 9.09. Found: C 46.63; H 4.09; N 8.95%.

Method B. Method A was repeated in a mixture of MeCN/py (10/4 mL) without using H₂pd. The yield was ~9% based on total ClO₄⁻ content.

2.3. X-Ray Crystallography. Data were collected on an Oxford-Diffraction Xcalibur diffractometer, equipped with a CCD area detector and a graphite monochromator utilizing




FIGURE 1: A partially labeled plot of the cation of **1**. Color code: Mn, purple; O, red; N, green; C, grey. H atoms are omitted for clarity.

Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$). Suitable crystals were attached to glass fibers using paratone-N oil and transferred to a goniostat where they were cooled for data collection. Unit cell dimensions were determined and refined by using 12271 ($3.07 \leq \theta \leq 30.27^\circ$) and 5746 ($3.06 \leq \theta \leq 30.29^\circ$) reflections for **1** and **2**, respectively. Empirical absorption corrections (multiscan based on symmetry-related measurements) were applied using CrysAlis RED software [31]. The structures were solved by direct methods using SIR92 [32], and refined on F^2 using full-matrix least squares with SHELXL97 [33]. Software packages used: CrysAlis CCD [31] for data collection, CrysAlis RED [31] for cell refinement and data reduction, WINGX for geometric calculations [34], and DIAMOND [35] and MERCURY [36] for molecular graphics. The non-H atoms were treated anisotropically, whereas the hydrogen atoms were placed in calculated, ideal positions and refined as riding on their respective carbon atoms. Unit cell data and structure refinement details are listed in Table 1.


2.4. Physical Measurements. Elemental analyses were performed by the in-house facilities of the Chemistry Department, University of Cyprus. IR spectra were recorded on KBr pellets in the $4000\text{--}400 \text{ cm}^{-1}$ range using a Shimadzu Prestige-21 spectrometer. Variable-temperature DC magnetic susceptibility data down to 1.80 K were collected on a Quantum Design MPMS-XL SQUID magnetometer equipped with a 70 kG (7 T) DC magnet. Diamagnetic corrections were applied to the observed paramagnetic susceptibilities using Pascal's constants. Samples were embedded in solid eicosane, unless otherwise stated, to prevent torquing.

3. Results and Discussions

3.1. Syntheses. Both complexes were prepared serendipitously during our investigations on two different synthetic methods. The first one involved the use of iodine as an oxidizing

agent in various reactions of $[\text{Mn}(\text{O}_2\text{CMe})_2] \cdot 4\text{H}_2\text{O}$, while the second one included the employment of 1,3-propanediol (pdH_2) in reactions with $[\text{Mn}_3\text{O}(\text{O}_2\text{CMe})_6(\text{py})_3] \cdot \text{py}$.

One of the most successful strategies to polynuclear Mn clusters has been the oxidation of a Mn^{2+} starting material with the use of various oxidizing agents, often in the presence of a chelating ligand. Several oxidants have been employed for this purpose such as MnO_4^- , Ce^{IV} , peroxides, bromate, and iodine to form high-oxidation state Mn species [4, 7, 28]. Although the use of iodine as oxidant in Mn cluster chemistry has been reported in the past [4, 7], the oxidation of Mn^{2+} salts from iodine under various conditions is a rather unexplored synthetic method. Compound **1** was prepared during our investigations on reactions of $[\text{Mn}(\text{O}_2\text{CMe})_2] \cdot 4\text{H}_2\text{O}$ with iodine in $\text{MeCOOH}/\text{pyridine}$. A large amount of MeCOOH was used in order to avoid the formation of various Mn oxides/hydroxides that precipitate at basic conditions. Thus, the reaction of $[\text{Mn}(\text{O}_2\text{CMe})_2] \cdot 4\text{H}_2\text{O}$ with solid I_2 in a 1 : 1 ratio in MeCOOH/py (10/20 mL) resulted in the formation of dark brown crystals of **1** in $\sim 60\%$ yield. The formation of **1** is summarized in (1):



Despite the presence of an oxidant (I_2) in the reaction mixture, the final product (compound **1**) contains only Mn^{2+} ions. We believe that species that contain Mn ions in higher oxidation states are also formed but are quite soluble and thus do not precipitate from the reaction solution.

Another synthetic method to new polynuclear Mn clusters employed recently by our group involves the use of aliphatic diols such as pdH_2 in Mn cluster chemistry. These studies have resulted in a number of new polynuclear clusters and coordination polymers with coordinated

TABLE 2: Selected interatomic distances (Å) and angles for complex 1.

Bond Distances (Å)			
Mn1···Mn2			3.799(2)
Mn1–O1			2.154(2)
Mn1–O3			2.196(2)
Mn1–N1			2.247(2)
Mn2–O2			2.093(2)
Mn2–O3			2.234(2)
Mn2–N3			2.235(2)
Mn2–O4			2.276(2)
Mn2–N2			2.288(2)
Mn2–N4			2.295(2)
Bond Angles (°)			
O1–Mn1–O1	180.0	O2–Mn2–O3	103.82(7)
O1–Mn1–O3	90.90(7)	O2–Mn2–N3	108.98(7)
O1'–Mn1–O3	89.10(7)	O3–Mn2–N3	146.90(7)
O3–Mn1–O3'	180.0	N3–Mn2–O4	89.50(7)
O1–Mn1–N1	90.57(7)	O2–Mn2–N2	90.28(8)
O1'–Mn1–N1	89.43(7)	O3–Mn2–N2	96.50(7)
O3–Mn1–N1	87.20(7)	N3–Mn2–N2	87.73(8)
O3'–Mn1–N1	92.80(7)	O4–Mn2–N2	86.28(7)
O1–Mn1–N1	90.57(7)	O3–Mn2–N4	89.30(7)
N1–Mn1–N1'	180.0(2)	N2–Mn2–N4	173.85(8)
		Mn1–O3–Mn2	118.07(7)

TABLE 3: Bond valence sum (BVS)^(a,b) calculations for complexes 1 and 2.

	Complex 1			Complex 2		
	Mn ^{II}	Mn ^{III}	Mn ^{IV}	Mn ^{II}	Mn ^{III}	Mn ^{IV}
Mn1	<u>2.00</u>	1.87	1.90	<u>1.92</u>	1.81	1.81
Mn2	<u>1.90</u>	1.79	1.80	<u>2.05</u>	1.91	1.95
				<u>1.91</u>	1.80	1.81

^(a)The underlined value is the one closest to the charge for which it was calculated. ^(b)The oxidation state is the nearest whole number to the underlined value.

pdH₂ ligands [37–40]. Many of these compounds were isolated from reactions that were involving the use of [Mn₃O(O₂CMe)₆(py)₃]·py as a starting material [37, 38]. These studies, apart from compounds that contain coordinated pdH₂ ligands, have also resulted in complexes that do not include the diol in their asymmetric unit, with 2 being one of the members of this family. Thus, compound 2 was initially prepared from the reaction of [Mn₃O(O₂CMe)₆(py)₃]·py with Mn(ClO₄)₂·6H₂O in the presence of pdH₂ in a 1:1:4 ratio in MeCN/py (10/2 mL) in 20% yield. When the identity of 2 was established and known that it contained neither coordinated nor lattice pdH₂/pd²⁻ ligands, the reaction resulted in the formation of 2 was repeated without including pdH₂ in the reaction mixture. This reaction gave a few crystals of 2. Various modifications were applied in this reaction in order to optimize its yield. Finally, the larger yield (achieved when no pdH₂ was included in the reaction mixture) was ~9% and obtained when an extra amount of pyridine (4 more mL) was

added to the reaction solution. The exact role of pdH₂ in the assembly of 2 and how its use results in larger reaction yield still remain unidentified.

3.2. Description of the Structures. The molecular structure of complex 1 is presented in Figure 1 and selected interatomic distances and angles for 1 are listed in Table 2. Bond valence sum (BVS) calculations for the metal ions of 1 and 2 are given in Table 3. The crystal structures of 1 and 2 present a striking similarity with the main difference between them being their counter-ions and thus only that of 1 will be described here.

Compound 1 crystallizes in the monoclinic P2₁/n space group and comprises the [Mn₃(O₂CMe)₄(py)₈]²⁺ cation and two I₃⁻ counteranions. The cation of 1 (Figure 1) consists of a linear array of three Mn^{II} ions coordinated by four acetate groups and eight terminal pyridine molecules. The oxidation states of the Mn ions were determined by BVS calculations (Table 3), charge considerations, and inspection




FIGURE 2: A representation of the packing of complex **1**. Mn, purple; O, red; C, grey. H atoms are omitted for clarity.

of metric parameters. The central metal ion of the trinuclear unit (Mn1), which is located on a crystallographic inversion center, is ligated by four oxygen atoms from four different acetate ligands and two molecules of pyridine adopting a distorted octahedral coordination geometry. All four acetate ligands bridge two Mn ions with two of them operating in the common *syn-syn- η^1 : η^1 : μ_2* fashion, whereas the other two function in the less common monoatomically bridging *η^2 : η^1 : μ_2* mode. The above mentioned carboxylate bridging modes have also been observed in several other linear trinuclear manganese (II) complexes [10–23]. However, in most linear Mn^{II}_3 complexes each pair of Mn^{II} ions is held together by at least three bridging ligands, whereas in **1** the neighboring Mn ions are connected through two bridging ligands only. One exception in this situation is the compound $[\text{Mn}_3(\text{O}_2\text{CMe})_6(\text{H}_2\text{O})(\text{phen})_2]$ where one pair of Mn ions is linked through two acetate ligands, whereas the second one is held together by three bridging MeCOO^- ligands [16]. The consequence of the presence of less bridging ligands in **1** is the larger $\text{Mn}\cdots\text{Mn}$ separation (3.799 (2) Å) compared to the values observed in other linear trinuclear Mn^{II} complexes which are within the range of 3.2–3.7 Å [10–23]. The observed separation of 3.799 Å is slightly smaller than that (3.868 (4) Å) between the Mn ions bridged by two acetate ligands in $[\text{Mn}_3(\text{O}_2\text{CMe})_6(\text{H}_2\text{O})(\text{phen})_2]$. However, the $\text{Mn}\cdots\text{Mn}$ distance in the other pair of Mn ions of the latter is significantly shorter (3.489 Å) and thus the average $\text{Mn}\cdots\text{Mn}$ separation falls within the range observed for the other linear trinuclear Mn^{II} complexes.

The distorted octahedral coordination environment around each terminal metal ion (Mn2) is completed by three pyridine molecules. The $\text{Mn}_2\text{N}_3\text{O}_3$ octahedron is significantly distorted, with the main distortion arising from the acute O3–Mn2–O4 angle (58.24 (7)°). The $\text{Mn}_1\text{N}_2\text{O}_4$ octahedron is almost perfect. All Mn–N and Mn–O bond lengths of the two crystallographically independent manganese ions are within the expected range for octahedral high-spin Mn^{II} complexes.




FIGURE 3: Plot of $\chi_M T$ versus T for **1**. The solid line is the fit of the experimental data; see the text for the fit parameters.

A close examination of the packing of **1** revealed that the trinuclear molecules are nearly perpendicular to each other (Figure 2) and there are no significant hydrogen bonding interactions between neighboring units of **1**.

3.3. Magnetic Properties. Solid-state dc magnetic susceptibility studies were performed on a powdered crystalline sample of **1** in a 0.1 T field and in the 5.0–300 K temperature range. The obtained data are plotted as $\chi_M T$ versus T in Figure 3.

The $\chi_M T$ product at 300 K for **1** is $12.98 \text{ cm}^3 \text{ mol}^{-1} \text{ K}$, slightly smaller than the value expected for three Mn^{II} ($S = 5/2$) noninteracting ions ($13.125 \text{ cm}^3 \text{ mol}^{-1} \text{ K}$, $g = 2$) indicating the existence of antiferromagnetic exchange interactions. This is corroborated by the continuous decrease of $\chi_M T$ upon cooling down to $10.63 \text{ cm}^3 \text{ mol}^{-1} \text{ K}$ at ~ 50 K. Below that temperature, the decrease is more abrupt, with $\chi_M T$ reaching a value of $4.49 \text{ cm}^3 \text{ mol}^{-1} \text{ K}$ at 5 K. The 5 K $\chi_M T$ value is very close to the spin-only ($g = 2$) value of $4.375 \text{ cm}^3 \text{ mol}^{-1} \text{ K}$ for a spin ground state $S = 5/2$. These results are indicative of antiferromagnetic exchange interactions between the Mn ions of **1** that lead to a spin ground state of $S = 5/2$.

The magnetic susceptibility was simulated taking into account only one isotropic intracluster magnetic interaction, J , between Mn1 and Mn2 centers since the exchange interaction between the terminal Mn ions of **1** and also of most of the known linear Mn^{II}_3 complexes is negligible ($J' = 0$) [10, 11, 15, 16] because of the large $\text{Mn}\cdots\text{Mn}$ separation (for **1** $\text{Mn}2\cdots\text{Mn}2' = 7.598(1) \text{ Å}$). Application of the van Vleck equation [41] to the Kambe's vector coupling scheme [42] allows the determination of a theoretical χ_M versus T expression for **1** from the following Hamiltonian:

$$H = -2J(\hat{S}_1\hat{S}_2 + \hat{S}_1\hat{S}_{2'}), \quad (2)$$

using the numbering scheme of Figure 1, where $S_1 = S_2 = S_{2'} = 5/2$. This expression was used to fit the experimental data giving $J = -1.50 \text{ K}$ and $g = 2.00$ (solid line, Figure 3).

A temperature-independent paramagnetism (TIP) term was held constant at $600 \times 10^{-6} \text{ cm}^3 \text{ mol}^{-1} \text{ K}$.

The obtained J value is smaller than values reported in the literature for other linear Mn^{II}_3 clusters with three bridging ligands per manganese pair which in most cases range from ~ -2.5 to $\sim -7 \text{ K}$ [18]. This behaviour could be rationalized on the basis of the existence of only two bridging ligands per manganese pair and larger $\text{Mn} \cdots \text{Mn}$ separations in **1** as was discussed in detail above (description of the structures). There are, however, examples of linear Mn^{II}_3 clusters with J values comparable to that of **1**, such as $[\text{Mn}_3(\text{L}^1)_2(\mu\text{-O}_2\text{CMe})_4] \cdot 2\text{Et}_2\text{O}$ ($\text{HL}^1 = (1\text{-hydroxy-4-nitrobenzyl})((2\text{-pyridyl)methyl})((1\text{-methylimidazol-2-yl)methyl)amine$) ($J = -1.7 \text{ K}$) [21].

4. Conclusions

A new linear trinuclear manganese(II) complex $[\text{Mn}_3(\text{O}_2\text{CMe})_4(\text{py})_8]^{2+}$ cocrystallizing with I_3^- [1] and ClO_4^- [2] has been synthesized serendipitously. Compound **1** was prepared in an attempt to oxidize $[\text{Mn}(\text{O}_2\text{CMe})_2] \cdot 4\text{H}_2\text{O}$ with I_2 in MeCOOH/py , whereas compound **2** was initially isolated during our investigations on reactions of $[\text{Mn}_3\text{O}(\text{O}_2\text{CMe})_6(\text{py})_3] \cdot \text{py}$ with $\text{Mn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ in the presence of pdH_2 in MeCN/py and was resynthesized in lower yield without adding pdH_2 in the reaction mixture. Although several linear trinuclear Mn^{II} complexes have been prepared and studied, the cation of **1** and **2** has several novel structural features including: (i) different type of ligation since **1** and **2** are the first examples of linear trinuclear Mn clusters with only acetate and pyridine ligands and (ii) different number of bridging ligands between each pair of Mn^{II} ions, since **1** and **2** are rare examples of linear Mn^{II}_3 clusters with only two bridging ligands linking each pair of Mn^{II} ions. Variable temperature dc magnetic susceptibility studies revealed the existence of antiferromagnetic interactions between the Mn ions of **1** resulting in an $S_T = 5/2$ spin ground state.

Acknowledgments

The authors thank the Cyprus Research Promotion Foundation (Grant: TEXNO/0506/06), for financial support of this research. This paper is dedicated to Professor Nick Hadjiliadis for his retirement.

References

- [1] V. L. Pecoraro, *Manganese Redox Enzymes*, VCH Publishers, New York, NY, USA, 1992.
- [2] R. Bagai and G. Christou, "The Drosophila of single-molecule magnetism: $[\text{Mn}_{12}\text{O}_{12}(\text{O}_2\text{CR})_{16}(\text{H}_2\text{O})_4]$," *Chemical Society Reviews*, vol. 38, no. 4, pp. 1011–1026, 2009.
- [3] J. Barber, "Photosynthetic energy conversion: natural and artificial," *Chemical Society Reviews*, vol. 38, no. 1, pp. 185–196, 2009.
- [4] S. Mukhopadhyay, S. K. Mandal, S. Bhaduri, and W. H. Armstrong, "Manganese clusters with relevance to photosystem II," *Chemical Reviews*, vol. 104, no. 9, pp. 3981–4026, 2004.
- [5] A. Mishra, W. Wernsdorfer, K. A. Abboud, and G. Christou, "The first high oxidation state manganese-calcium cluster: relevance to the water oxidizing complex of photosynthesis," *Chemical Communications*, no. 1, pp. 54–56, 2005.
- [6] W. Ruttinger and G. C. Dismukes, "Synthetic water-oxidation catalysts for artificial photosynthetic water oxidation," *Chemical Reviews*, vol. 97, no. 1, pp. 1–24, 1997.
- [7] R. Manchanda, G. W. Brudvig, and R. H. Crabtree, "High-valent oxomanganese clusters: structural and mechanistic work relevant to the oxygen-evolving center in photosystem II," *Coordination Chemistry Reviews*, vol. 144, pp. 1–38, 1995.
- [8] V. L. Pecoraro, M. J. Baldwin, and A. Gelasco, "Interaction of manganese with dioxygen and its reduced derivatives," *Chemical Reviews*, vol. 94, no. 3, pp. 807–826, 1994.
- [9] A. J. Wu, J. E. Penner-Hahn, and V. L. Pecoraro, "Structural, spectroscopic, and reactivity models for the manganese catalases," *Chemical Reviews*, vol. 104, no. 2, pp. 903–938, 2004.
- [10] S. G. Baca, Y. Sevryugina, R. Clérac, I. Malaestean, N. Gerbeleu, and M. A. Petrukhina, "Linear trinuclear manganese(II) complexes: crystal structures and magnetic properties," *Inorganic Chemistry Communications*, vol. 8, no. 5, pp. 474–478, 2005.
- [11] S. Menage, S. E. Vitols, P. Bergerat, et al., "Structure of the linear trinuclear complex $\text{Mn}_3^{\text{II}}(\text{CH}_3\text{CO}_2)_6(\text{bpy})_2$. Determination of the J electron-exchange parameter through magnetic susceptibility and high-field magnetization measurements," *Inorganic Chemistry*, vol. 30, no. 12, pp. 2666–2671, 1991.
- [12] G. Christou, "Manganese carboxylate chemistry and Its biological relevance," *Accounts of Chemical Research*, vol. 22, no. 9, pp. 328–335, 1989.
- [13] G. Fernandez, M. Corbella, J. Mahia, and M. A. Maestro, "Polynuclear Mn^{II} complexes with chloroacetate bridge—syntheses, structure, and magnetic properties," *European Journal of Inorganic Chemistry*, no. 9, pp. 2502–2510, 2002.
- [14] K. Tsuneyoshi, H. Kobayashi, and H. Miyamae, "Structure of hexa- μ -acetato- $1\text{-K}^3\text{O} : 2\text{K}^3\text{O}' : 2\text{K}^3\text{O} : 3\text{K}^3\text{O}'$ -bis(1,10-phenanthroline)-1- $\text{K}^2\text{N}^1, \text{N}^{10} ; 2\text{K}^2\text{N}^1, \text{N}^{10}$ -trimanganese(II)," *Acta Crystallographica Section C-Crystal Structure Communication*, vol. 49, pp. 233–236, 1993.
- [15] R. L. Rardin, P. Poganiuch, A. Bino, et al., "Synthesis and characterization of trinuclear iron(II) and manganese(II) carboxylate complexes: structural trends in low valent iron and manganese carboxylates," *Journal of the American Chemical Society*, vol. 114, no. 13, pp. 5240–5249, 1992.
- [16] M. Wang, C.-B. Ma, H.-S. Wang, C.-N. Chen, and Q.-T. Liu, "Synthesis and characterization of two linear trinuclear complexes $[\text{M}_3(\text{MeCOO})_6(\text{H}_2\text{O})(\text{phen})_2] \cdot \text{H}_2\text{O}$ ($\text{M} = \text{Co}, \text{Mn}$)," *Journal of Molecular Structure*, vol. 873, no. 1–3, pp. 94–100, 2008.
- [17] V. Tangoulis, D. A. Malamatar, K. Soulti, et al., "Manganese(II/II/II) and manganese(III/II/III) trinuclear compounds. Structure and solid and solution behavior," *Inorganic Chemistry*, vol. 35, no. 17, pp. 4974–4983, 1996.
- [18] V. Gómez, D. A. Malamatar, K. Soulti, et al., "Versatility in the coordination modes of n -chlorobenzoate ligands: synthesis, structure and magnetic properties of three types of polynuclear Mn^{II} Compounds," *European Journal of Inorganic Chemistry*, no. 29–30, pp. 4471–4482, 2009.
- [19] C. C. Stoumpos, I. A. Gass, C. J. Milios, et al., "The use of di-2-pyridyl ketone in manganese(II) benzoate chemistry: two novel Mn_3^{II} linkage isomers containing the ketone form of the ligand and a neutral Mn_4^{II} cubane containing the ligand in its gem-diolate(-1) form," *Inorganic Chemistry Communications*, vol. 11, no. 2, pp. 196–202, 2008.

- [20] C. J. Milios, T. C. Stamatatos, P. Kyritsis, et al., "Phenyl 2-pyridyl ketone and its oxime in manganese carboxylate chemistry: synthesis, characterisation, X-ray studies and magnetic properties of mononuclear, trinuclear and octanuclear complexes," *European Journal of Inorganic Chemistry*, no. 14, pp. 2885–2901, 2004.
- [21] M. Kloskowski, D. Pursche, R.-D. Hoffmann, et al., "Novel trinuclear $Mn^{II}/Mn^{II}/Mn^{II}$ complexes—crystal structures and catalytic properties," *Zeitschrift für Anorganische und Allgemeine Chemie*, vol. 633, no. 1, pp. 106–112, 2007.
- [22] H. Asada, K. Hayashi, S. Negoro, M. Fujiwara, and T. Matsushita, "Preparation and structures of trinuclear manganese(II) complexes with *N*-2-pyridylmethylidene-2-hydroxy-5-substituted-phenylamine," *Inorganic Chemistry Communications*, vol. 6, no. 2, pp. 193–196, 2003.
- [23] D. Pursche, M. U. Triller, N. Reddig, A. Rompel, and B. Krebs, "Synthesis and characterization of $[Mn_3(ppi)_2(\mu-OAc)_4(H_2O)_2] \cdot 2MeOH$ —unusual structural properties of a trinuclear oxygen-rich manganese complex," *Zeitschrift für Anorganische und Allgemeine Chemie*, vol. 629, no. 1, pp. 24–28, 2003.
- [24] M. Hirotsu, M. Kojima, W. Mori, and Y. Yoshikawa, "Steric control of polynuclear manganese complexes by the use of tripodal tetradentate ligands," *Bulletin of the Chemical Society of Japan*, vol. 71, no. 12, pp. 2873–2884, 1998.
- [25] D. P. Kessissoglou, M. L. Kirk, M. S. Lah, et al., "Structural and magnetic characterization of trinuclear, mixed-valence manganese acetates," *Inorganic Chemistry*, vol. 31, no. 26, pp. 5424–5432, 1992.
- [26] Y.-G. Li, L. Lecren, W. Wernsdorfer, and R. Clérac, "Antiferromagnetic order in a supramolecular assembly of manganese trimers based on imidazole and Schiff-base ligands," *Inorganic Chemistry Communications*, vol. 7, no. 12, pp. 1281–1284, 2004.
- [27] J. B. Vincent, H.-R. Chang, K. Folting, J. C. Huffman, G. Christou, and D. N. Hendrickson, "Preparation and physical properties of trinuclear oxo-centered manganese complexes of the general formulation $[Mn_3O(O_2CR)_6L_3]^{0,+}$ ($R = Me$ or Ph ; $L =$ a neutral donor group) and the crystal structures of $[Mn_3O(O_2CMe)_6(pyr)_3](pyr)$ and $[Mn_3O(O_2CPh)_6(pyr)_2(H_2O)] \cdot 0.5MeCN$," *Journal of the American Chemical Society*, vol. 109, no. 19, pp. 5703–5711, 1987.
- [28] S. Bhaduri, M. Pink, and G. Christou, "Towards a synthetic model of the photosynthetic water oxidizing complex: $[Mn_3O_4(O_2CMe)_4(bpy)_2]$ containing the $[Mn_3^{IV}(\mu-O_4)]^{4+}$ core," *Chemical Communications*, no. 20, pp. 2352–2353, 2002.
- [29] S. Zartilas, E. E. Moushi, V. Nastopoulos, A. K. Boudalis, and A. J. Tasiopoulos, "Two new coordination polymers containing the triangular $[Mn_3O(O_2CR)_6]^{0/+}$ units," *Inorganica Chimica Acta*, vol. 361, no. 14–15, pp. 4100–4106, 2008.
- [30] R. A. Reynolds III, W. R. Dunham, and D. Coucouvanis, "Kinetic lability, structural diversity, and oxidation reactions of new oligomeric, anionic carboxylate-pyridine complexes," *Inorganic Chemistry*, vol. 37, no. 6, pp. 1232–1241, 1998.
- [31] Oxford diffraction, *CrysAlis CCD and CrysAlis RED, Version 1.171.32.15*, Oxford Diffraction Ltd, Oxford, UK, 2008.
- [32] A. Altomare, G. Cascarano, C. Giacovazzo, et al., "SIR92—a program for automatic solution of crystal structures by direct methods," *Journal of Applied Crystallography*, vol. 27, no. 6, p. 435, 1994.
- [33] G. M. Sheldrick, *SHELXL97—A Program for The Refinement of Crystal Structure*, University of Göttingen, Göttingen, Germany, 1997.
- [34] L. J. Farrugia, "WinGX suite for single crystal small molecule crystallography," *Journal of Applied Crystallography*, vol. 32, no. 4, pp. 837–838, 1999.
- [35] K. Brandenburg, *DIAMOND. Version 3.1d.*, Crystal Impact GbR, Bonn, Germany, 2006.
- [36] C. F. Macrae, P. R. Edgington, P. McCabe, et al., "Mercury: visualization and analysis of crystal structures," *Journal of Applied Crystallography*, vol. 39, no. 3, pp. 453–457, 2006.
- [37] E. E. Moushi, T. C. Stamatatos, W. Wernsdorfer, V. Nastopoulos, G. Christou, and A. J. Tasiopoulos, "A family of 3D coordination polymers composed of Mn_{19} magnetic units," *Angewandte Chemie International Edition*, vol. 45, no. 46, pp. 7722–7725, 2006.
- [38] E. E. Moushi, C. Lampropoulos, W. Wernsdorfer, V. Nastopoulos, G. Christou, and A. J. Tasiopoulos, "A large $[Mn_{10}Na]_4$ loops," *Inorganic Chemistry*, vol. 46, no. 10, pp. 3795–3797, 2007.
- [39] E. E. Moushi, T. C. Stamatatos, W. Wernsdorfer, V. Nastopoulos, G. Christou, and A. J. Tasiopoulos, "A Mn_{17} octahedron with a giant ground-state spin: occurrence in discrete form and as multidimensional coordination polymers," *Inorganic Chemistry*, vol. 48, no. 12, pp. 5049–5051, 2009.
- [40] E. E. Moushi, T. C. Stamatatos, V. Nastopoulos, G. Christou, and A. J. Tasiopoulos, "A new family of octanuclear Mn complexes with a rod-like topology," *Polyhedron*, vol. 28, no. 15, pp. 3203–3208, 2009.
- [41] J. H. Van Vleck, *The Theory of Electric and Magnetic Susceptibility*, Oxford University Press, Oxford, UK, 1932.
- [42] K. Kambe, "On the paramagnetic susceptibilities of some polynuclear complex salts," *Journal of the Physical Society of Japan*, vol. 5, pp. 48–51, 1950.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

