

Research Article

Molecular Characterization of the Ghrelin and Ghrelin Receptor Genes and Effects on Fat Deposition in Chicken and Duck

Q. Nie,¹ M. Fang,^{1,2} L. Xie,¹ X. Peng,¹ H. Xu,¹ C. Luo,¹ D. Zhang,¹ and X. Zhang¹

¹Department of Animal Genetics, Breeding and Reproduction, College of Animal Science, South China Agricultural University, Guangzhou 510642, China

²Department of Laboratory Animal Science, Medical College of Jinan University, Guangzhou, Guangdong 510632, China

Correspondence should be addressed to X. Zhang, xqzhang@scau.edu.cn

Received 4 August 2009; Accepted 19 October 2009

Recommended by M. Ilyas Kamboh

Ghrelin (GHRL) and its receptor (GHSR) are involved in various bioactivities. In this study, the complete cDNA and 5' flanking region of the duck *GHRL* (*dGHRL*) gene and a 3717 bp fragment of the duck *GHSR* (*dGHSR*) gene were obtained. A total of 19, 8, 43, and 48 SNPs identified in 2751, 1358, 3671, and 3567 bp of the chicken *GHRL* (*cGHRL*), chicken *GHSR* (*cGHSR*), *dGHRL*, and *dGHSR* genes, respectively. Both *cGHRL* and *dGHRL* were expressed predominantly in the proventriculus, whereas the highest mRNA levels of *cGHSR* and *dGHSR* were detected in the breast muscle and pituitary. Association analysis showed that C-2047G, A-2355C, and A-2220C of the *cGHRL* gene were significantly associated with abdominal fat weight (AFW; $P = .01$), crude protein content of leg muscle (CPCLM; $P = .02$), and CPCLM ($P = .0009$), respectively. C-1459T of the *cGHSR* gene was also significantly associated with CPCLM ($P = .0004$). C-729T of *dGHRL* and A3427T of *dGHSR* were both significantly associated with subcutaneous fat thickness (SFT; $P = .04$). It was indicated by this study that the *GHRL* and *GHSR* genes were related to fat deposition in both chicken and duck.

Copyright © 2009 Q. Nie et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Small synthetic molecules termed Growth Hormone Secretagogues (GHSs) act on the pituitary gland and the hypothalamus to regulate growth hormone (GH) release. The Growth Hormone Secretagogue Receptor (GHSR or GHS-R), a G protein-coupled receptor, has been cloned from human and pig and is mainly expressed in the pituitary, hypothalamus, and hippocampus [1]. In 1999, a novel 28-amino acid gut-brain peptide named Ghrelin (GHRL) was first isolated from the mammalian stomach [2]. Similar to the synthetic molecules of GHSs, GHRL proved to be an endogenous ligand for GHSR and involved in GH secretion, food intake, and energy homeostasis [2, 3]. Even though the mature GHRL varies in structure among mammals, birds, and fish, the *n*-octanoylation of the Ser-3 residue is highly conserved and essential for biological activity [2, 4]. It is now well known that GHRL interacts with GHSR to mediate a variety of bioactivities through the neuroendocrine pathway.

The *GHRL* and *GHSR* genes are reportedly related to obesity in human and mouse. Body mass index and body fat

were shown to be negatively correlated with plasma GHRL levels [5–8]. Two polymorphisms, Arg51Gln and Leu72Met, in the human *GHRL* gene were identified in obese subjects, and Leu72Met was significantly associated with body mass index and fat mass in some human populations [9, 10]. In addition to these two nonsynonymous SNPs, two other variations of Gln90Leu and a frameshift mutation (2 bp deletion at codon 34) were also reported in the human *GHRL* gene, and the 90Leu allele frequency was significantly higher in extremely obese children and adolescents (0.063) than in normal-weight students [11]. A novel SNP, T3056C in intron 2 of the *GHRL* gene, was significantly associated with the acylation of GHRL and some fatness traits, including high body mass index, body weight, fat mass, and skinfold thickness [12]. Among 11 SNPs identified in the 5' flanking region of the *GHRL* gene, SNP-501A > C was significantly associated with body mass index in human [13]. In mice, the peripheral administration of GHRL was found to regulate body weight, adiposity, and UCP mRNA expression [14]. It was also interesting that central administration of GHRL enhanced rat fat ingestion [15].

Until now, the research progress on the *GHRL* and *GHSR* genes was achieved in chicken. The chicken *GHRL* (*cGHRL*) gene cDNA was first cloned by Kaiya et al. [16], and it encoded a 116-amino acid GHRL precursor and a 26-amino acid mature GHRL. The complete *cGHRL* gene comprised five exons and four introns. The first exon did not encode any amino acids, a characteristic that was conserved in mammals [17, 18]. The *cGHSR* gene was composed of two exons separated by an intron and encoded a normal *GHSR* with seven transmembrane domains (TM1 to TM7); however, a variant *GHSR* without TM6 due to a 48-bp deletion was found [19]. A total of 19 and 37 SNPs were identified in the *cGHRL* and *cGHSR* genes, respectively [20], and an 8-bp indel in the 5' untranslated region (UTR) and some SNPs of the *cGHRL* gene were related to chicken growth and muscle fiber traits [21–23]. A 6-bp indel of the *cGHSR* gene was significantly linked with fat traits [21]. The duck *GHRL* (*dGHRL*) gene cDNA was also cloned, and the encoded dGHRL was highly similar to cGHRL [24]. Nevertheless, no further studies on the *dGHRL* and duck *GHSR* (*dGHSR*) genes have been reported.

Fat deposition is significant in both chicken and duck. Most fat accumulates as abdominal fat in chicken but as subcutaneous fat in duck; these characteristics make the chicken and duck ideal for use in this study. RACE, genome walking, real time PCR, and association analysis were performed to characterize the *GHRL* and *GHSR* genes and reveal their effects on fat deposition in chicken and duck.

2. Materials and Methods

2.1. Animals and Sample Preparation. A total of 411 birds (219 males and 193 females) from 8 chicken populations (POP1 to POP8) and 139 birds from 9 duck populations (POP9 to POP17) were used in this study (Table 1). These samples consisted of six chicken breeds, including Xinghua (XH), White Recessive Rock (WRR), Taihe Silkies (TS), Leghorn (LH), Lingnan Yellow (LY), Huiyang Beard (HB), and an F2 full-sib hybrid population (XH&WRR) described by Fang et al. [23], as well as four duck breeds, Sanshui White (SW), Peking (PK), Partridge (PT), and Lake (LK). Either blood or live tissues were collected from these birds at 9 weeks of age for the purpose of sequence cloning, variation, mRNA quantitative expression, cell culture, and marker trait association analyses (Table 1).

The SW commercial population (POP14), including 100 ducks (52 males and 48 females), were slaughtered at 90 days old, and body weights at hatch (g) and 90 days old (g), carcass weight (g), fat thickness under skin (mm), fat width (mm), eviscerated weight (g), abdominal fat pad weight (g), and abdominal fat pad ratio (%) were recorded. All birds were raised in separate pens and fed with commercial corn-soybean-based diets that met NRC requirements. All chickens were collected from Guangdong Wens Foodstuff Corporation Ltd. (Guangdong, China), and all ducks were provided by Guangdong Sanshui Lianke Duck Farm (Guangdong, China).

Genomic DNA was extracted from EDTA-anticoagulated blood. Total RNA was extracted from the isolated tissues using an improved Trizol isolation method (Invitrogen, California, USA) following the manufacturer's instructions. One microgram of total RNA was used to obtain cDNA by reverse transcription with use of the Rever Tra Ace Transcriptase Kit (Toyobo, Japan) and random hexamers.

2.2. Primers. Thirty-three primer pairs (PM1 to PM33) were designed and consisted of PM1 to PM6 (6) for the *cGHRL* gene, PM7 to PM10 (4) for the *cGHSR* gene, PM11 for the chicken β -actin gene, PM12 to PM24 (13) for the *dGHRL* gene, PM25 to PM32 (8) for the *dGHSR* gene, and PM33 for the duck β -actin gene (Table 2). PM12 and PM26 were both genomic primers, each of which consisting of three reverse primers and a random forward primer supplied by the Genomic Walking Kit. PM14 was a primer pair for 3' RACE, and the remaining 30 primer pairs were gene-specific primers. The GENETOOL software (<http://www.biologysoft.com/>) was used to design the primers based on reported sequences of the chicken *GHRL* (GenBank AB075215), *GHSR* (GenBank AB095994), and β -actin genes (GenBank NM_205518), as well as duck *GHRL* (GenBank AY338466), *GHSR* (GenBank EU005225), β -actin (GenBank EF667345), and goose *GHRL* (GenBank AY338465). All primers were commercially synthesized by Shanghai Biosune Biotechnology Co. Ltd. (Shanghai, China).

2.3. PCR, PCR-RFLP, and Sequencing. PCR was performed in 25 μ L reactions containing 50 ng chicken genomic DNA, 1 \times PCR buffer, 12.5 pmol primers, 100 μ M of each dNTP, 1.5 mM MgCl₂, and 1.0 U *Taq* DNA polymerase (Sangon Biological Engineering Technology Company, Shanghai, China). PCR was run in a Mastercycler gradient (M. J. Research Co. Ltd., USA) with the following procedure: 3 minutes at 94°C, followed by 35 cycles of 30 seconds at 94°C, 45 seconds at the annealing temperature (58–62°C) and 1 minute at 72°C, and a final extension of 5 minutes at 72°C. PCR products were checked for size and quality on 1% agarose gels.

The PCR product of PM5 was sequenced for genotyping of the A-2220C, A-2246G, A-2355C, C-2399del, C-2030T, and C-2047G polymorphisms in the *cGHRL* gene. The PCR product of the PM7 reaction was digested with *Tat I* with the PCR-RFLP method for genotyping C-1896A in the *cGHSR* gene. The PCR product of the PM8 reaction was digested by *Hin1 II* with the PCR-RFLP method for genotyping the C-1459T SNP in the *cGHSR* gene. The PCR product of the PM9 reaction was digested with *Tail I* with the PCR-RFLP method for genotyping the A-1022C SNP in the *cGHSR* gene. The PCR product of the PM20 reaction was digested with *Csp6 I* with the PCR-RFLP method for genotyping the C-729T SNP in the *dGHRL* gene. The PCR product of the PM22 reaction was digested with *Tail I* with the PCR-RFLP method for genotyping the T985C SNP in the *dGHRL* gene. The PCR product of the PM30 reaction was digested with *Csp6 I* with the PCR-RFLP method for genotyping

TABLE 1: Chicken and duck samples used in this study.

No.	Species	Population ⁽¹⁾	Samples	Blood/live tissues	Purpose
POP1	Chicken	XH	6 (3 ♂; 3 ♀)		
POP2	Chicken	WRR	6 (3 ♂; 3 ♀)	Blood for DNA	<i>GHRL</i> and <i>GHSR</i> gene variation
POP3	Chicken	TS	6 (3 ♂; 3 ♀)		
POP4	Chicken	LH	6 (3 ♂; 3 ♀)		
POP5	Chicken	XH&WRR	373 (200 ♂; 173 ♀)		
POP6	Chicken	LY	6 (3 ♂; 3 ♀)	18 tissues ⁽²⁾	Real time PCR
POP7	Chicken	HY	6 (3 ♂; 3 ♀)		
POP8	Chicken	LY	2 ♀	Subcutaneous and abdominal adipose tissues	Cell culture
POP9	Duck	SW	1 ♂	Blood for DNA and proventriculus tissue	<i>GHRL</i> gene sequence cloning
POP10	Duck	SW	6 (3 ♂; 3 ♀)	Blood for DNA	<i>GHRL</i> and <i>GHSR</i> gene variation
POP11	Duck	PK	6 (3 ♂; 3 ♀)		
POP12	Duck	PT	6 (3 ♂; 3 ♀)		
POP13	Duck	LK	6 (3 ♂; 3 ♀)		
POP14	Duck	SW	100 (52 ♂; 48 ♀)	Blood for DNA	Association analysis
POP15	Duck	SW	6 (3 ♂; 3 ♀)	18 tissues	Real time PCR
POP16	Duck	PT	6 (3 ♂; 3 ♀)		
POP17	Duck	SW	2 ♀	Subcutaneous and abdominal adipose tissues	Cell culture

⁽¹⁾XH = Xinghua chicken, WRR = White Recessive Rock, TS = Taihe Silkies, LH = Leghorn, XH&WRR = an F2 population crossed by XH and WRR, LY = Lingnan Yellow, HY = Huiyang Beard, SW = Sanshui White duck, PK = Peking duck, PT = Partridge duck, and LK = Lake duck.

⁽²⁾ The 18 tissues used were pituitary, cerebrum, lung, abdominal fat, liver, testis, proventriculus, ovary, subcutaneous fat, spleen, kidney, oviduct, leg muscle, uropygial gland, hypothalamus, glandularis, small intestine, cerebellum, heart and breast muscle.

the T404C SNP in the *dGHSR* gene. The PCR product of the PM31 reaction was digested with *Csp6 I* with the PCR-RFLP method for genotyping the A3427G SNP in the *dGHSR* gene. Digestion products were detected by agarose gel (1.5%) electrophoresis, and genotypes were determined by specific profiles (Supplementary Figure 1 in Supplementary Material available on-line at doi: 10.1155/2009/567120).

2.4. RT-PCR and 3' RACE. Total RNA was extracted from the proventriculus from an SW duck by the TRIzol reagent (Invitrogen, Carlsbad, CA) following the manufacturer's instructions. A proventriculus cDNA library was prepared using the SMART RACE cDNA amplification kit (Clontech Laboratories, Mountain View, CA). Primer-directed RT-PCR and 3' rapid amplification of cDNA ends (RACE) were used to generate duck *GHRL* cDNAs, amplified using primers P1 to P3 (Table 2). A cDNA library was prepared using the SMART RACE cDNA amplification kit (Clontech Laboratories, Mountain View, CA). The 3' ends of the cDNAs were amplified with a specific forward primer and reverse random primers from the SMART RACE cDNA amplification kit following the manufacturer's instructions.

2.5. Genome Walking. To obtain the 5' flanking region of the duck *GHRL* and *GHSR* genes, genome walking was performed using the Genome Walking Kit (TakaRa, Japan). The duck genomic DNA was purified using the Genomic DNA Purification Kit (U-gene, China) and was then used for genome walking. The genome walking procedure included three reactions: the first PCR reaction used the outer adaptor

primer AP1 from the kit and an outer gene-specific primer PM12R1 (or PM26R1); the second reaction used primers AP2 and PM26R2; the third reaction used primers AP3 and PM26R3. Each reaction was performed step-by-step following the kit instructions. PCR products of the third reaction were purified using the Gel Extraction Kit (U-gene, China), cloned into the pGEM-T easy vector (Promega, USA), and sequenced by Shanghai Biosune Biotechnology Co. Ltd. (Shanghai, China).

2.6. Quantitative Real Time PCR. The mRNA levels of *GHRL* and *GHSR* in 18 tissues of chickens and ducks were determined by quantitative real time PCR with SYBR green dye using β -actin as an internal positive control (IPC). The 15 μ L PCR mixture contained 1 μ L of cDNA, 7.5 μ M Super Mix (Toyobo, Japan), 10 μ M of each primer (PM6, PM10, PM11, PM24, PM32, and PM33), and 6.0 μ L DEPC H₂O. Real time PCR was run on an ABI 7500 (Applied Biosystems, Foster City, CA, USA) using the following program: initial denaturation at 95°C for 4 minutes, then 40 cycles of 94°C for 30 seconds, 63°C for 30 seconds, and 72°C for 45 seconds. Each sample was repeated three times and the average mRNA level was obtained for further analysis. Two random PCR products of PM6, PM10, PM11, PM24, PM32, and PM33 were sequenced by Shanghai Biosune Biotechnology Co. Ltd. (Shanghai, China) to confirm that specific fragments of the *cGHRL*, *cGHSR*, chicken β -actin, *dGHRL*, *dGHSR*, and duck β -actin genes were obtained. Quantitative values were obtained from the Ct values, which were the inverse ratio relative to the starting PCR product. The relative quantification was obtained by $2^{-\Delta\text{Ct}}$ in which

TABLE 2: Descriptions of the 33 primers used in this study.

Gene	Primer	Forward/reverse	Sequence (5' to 3')	Annealing temp (°C)	Purpose
<i>cGHRL</i>	PM1	PM1F	tgctgaaggaccgaaaacaaa	56	SNP identification
		PM1R	gccttgacagatgccttagtg		
	PM2	PM2F	aagaaagctggtaactgcactag	55	SNP identification
		PM2R	ggtagggctggtaggagta		
	PM3	PM3F	tgcttctgctactcttttcat	63	SNP identification
		PM3R	ggccaaggaggagtgtct		
PM4	PM4F	cggcagacactcctcct	57	SNP identification	
	PM4R	gccatcctccaactgtgtat			
PM5	PM5F	tatgcttctgctactcttt	58	Genotyping by sequencing	
	PM5R	tggaagcgcactataacc			
PM6	PM6F	catacagcaaaaaggatac	63	Real time PCR	
	PM6R	tgtggtgtccttcagct			
<i>cGHSR</i>	PM7	PM7F	gtgggtcagggcatcaaactc	58	SNP identification and PCR-RFLP
		PM7R	tcgaggctgctgaattttatg		
	PM8	PM8F	ggctcttcttttggttgtct	60	SNP identification and PCR-RFLP
		PM8R	tcgccctctctgattcacct		
	PM9	PM9F	tgatgccactggctctgagaat	58	Genotyping by PCR-RFLP
		PM9R	tatccagctgccatgtaaat		
PM10	PM10F	tggcgtcgagcatgagaat	63	Real time PCR	
	PM10R	ccacgactagcatcttcacag			
Chicken β -actin	PM11	PM11F	ccccaaagccaacagagaga	63	Real time PCR
		PM11R	ggtggtgaagctgtagcctctc		
<i>dGHRL</i>	PM12	PM12F	Genomic Walking Kit	According to the kit	Genomic Walking
		PM12R1	ttccatcagttctagacgagt		
		PM12R2	tgttgtccactactctgatact		
	PM13	PM12R3	gcccctgctttatctgtatc	56	RT-PCR
		PM13F	ccgctggtagaagaaac		
	PM14	PM13R	aagctacacaccacgtgaat	54	3' RACE
		PM14F	gaggcaagctgaaggacaaccaca		
	PM15	PM14R	3' RACE Kit	56	RT-PCR
		PM15F	gctggtttcccgtgaattc		
	PM16	PM15R	tgttgtccactactctgatact	54	Intron 1 amplification
		PM16F	gctggtttcccgtgaattc		
	PM17	PM16R	tgttgtccactactctgatact	56	Intron 2 amplification
		PM17F	tggtttggctggctctagt		
	PM18	PM7R	PM12R3	57	Intron 3 amplification
		PM18F	aaagcaggggcagaagat		
	PM19	PM18R	PM12R2	54	Intron 4 amplification
		PM19F	agggtcctggccaaaaat		
	PM20	PM19R	PM12R1	56	SNP identification and PCR-RFLP
		PM20F	cgcattgtagccttcacac		
	PM21	PM20R	agccgatgggttagcagagag	56	SNP identification
		PM21F	tcggctccatcagttgcagttat		
	PM22	PM21R	cggcagtgatattttgctgttg	56	SNP identification and PCR-RFLP
		PM22F	tcccacgacagagtagtttgag		
	PM23	PM22R	gcctcccctgctcctaa	54	SNP identification
PM23F		atagcagcattttagaagtga			
PM24	PM23R	ttgcctcagcttttcaact	63	Real time PCR	
	PM24F	cgtgtaattcctctctgctaa			
	PM24R	cgatgtattttgctgtgttt			

TABLE 2: Continued.

Gene	Primer	Forward/reverse	Sequence (5' to 3')	Annealing temp (°C)	Purpose
<i>dGHSR</i>	PM25	PM25F	ccctgagcaccaacgagt	56	Intron 1 amplification
		PM25R	ggcaaccagcagagtatga		
		PM26F	Genomic Walking Kit		
	PM26	PM26R1	cggaccgatgttcttctcttt	According to the kit	Genomic Walking
		PM26R2	gacgggcaggaaaaagaagac		
		PM26R3	ggcccagaggatgaggatga		
	PM27	PM27F	tggcgttctccgacctgctcat	57	SNP identification
		PM27R	acacagacctcagaaacac		
	PM28	PM28F	cgggtggcttaataacaacag	56	SNP identification
		PM28R	tgcccttttctcagtttctc		
	PM29	PM29F	cggaccataattaacacctgaa	56	SNP identification
		PM29R	caggtagaaggagcaaaaggaca		
	PM30	PM30F	tggcgttctccgacctgctcat	56	Genotyping by PCR-RFLP
		PM30R	acacagacctcagaaacac		
	PM31	PM31F	gcaggtgtttagatatggct	56	Genotyping by PCR-RFLP
PM31R		cgtgaaaaggcaaccagcagag			
PM32	PM32F	ccctgagcaccaacgagt	63	Real time PCR	
	PM32R	ccaccacaactaacattttcaca			
Duck β -actin	PM33	PM33F PM33R	acgccaacacgggtgctg gggtccggattcatcatactc	63	Real time PCR

$\Delta Ct = Ct_{\text{target gene}} - Ct_{\text{IPC}}$. The data are expressed as the mean \pm S.E. Statistics were analyzed by SAS Students *t*-test that measures analysis of variance with a significance level of 0.05.

2.7. Statistics Methods

2.7.1. Sequence BLAST, Prediction of Transcription Factors, and Haplotypes. The homology of GHRL among human, mouse, chicken, turkey, emu, goose, and duck was studied. Identity percentages were obtained from amino acid BLAST with the DNASTAR software (<http://www.dnastar.com/>). The obtained sequences of GHRL and GHSR were aligned with the Mega 3.1 software (<http://www.megasoftware.net/>). The 798-bp 5' flanking region of the *dGHRL* gene obtained in this study was used to predict the potential binding sites for transcription factors by an online service (http://www.fruitfly.org/cgi-bin/seq_tools/promoter.pl). In the chicken F2 population (XH&WRR) and the duck commercial population (SW), haplotypes were inferred based on the genotype data with the PHASE 2.1 software (<http://en.wikipedia.org/wiki/Phase>) [25].

2.7.2. Nucleotide Diversity (θ). To evaluate the nucleotide diversity, the normalized numbers of SNPs (θ) were calculated by dividing K , the number of observed nucleotide changes, by the total sequence length in base pairs (L) and correcting for the sample size (n) as described in [26], using the following formula:

$$\theta = \frac{K}{\sum_{i=1}^{n-1} i^{-1} L}. \quad (1)$$

2.7.3. Marker-Trait Association Analysis. The effects of 13 SNPs identified in this study on fatty traits were estimated by marker-trait association analysis. These SNPs included six in the *cGHRL* gene (C-2030T, C-2047G, A-2220C, A-2246G, A-2355C, and C-2399del), three in the *cGHSR* gene (C-1896A, C-1459T, and A-1022C), two in the *dGHRL* gene (C-729T and T985C), and two in the *dGHSR* gene (T404C and A3427G). Marker-trait association analysis was performed using the SAS GLM procedure (*SAS Institute, 1996*) and the genetic effects were analyzed by a mixed procedure according to the following model:

$$Y = \mu + G + D + H + S + e, \quad (2)$$

where Y is an observation on the trait, μ is the overall population mean, G is the fixed effect of the genotype, D is the random effect of dam, H is the fixed effect of hatch, S is the fixed effect of sex (male or female), and e is the residual random error.

3. Results

3.1. Cloning the Full-Length cDNA of the *dGHRL* Gene. The full-length cDNA of the *dGHRL* gene (835 bp) was amplified using primers PM13 to PM15. It comprised 115 bp of the 5' untranslated region (5' UTR), the 351 bp open reading frame (ORF) encoding a 116-amino acid (aa) peptide, and 369 bp of the 3' UTR (Figure 1) (GenBank EF613551). The SW duck GHRL precursor (116 aa) included a signal peptide (23 aa), the mature GHRL (26 aa), and a long C-terminal peptide (67 aa). Compared with the other reported mallard *dGHRL* precursor sequence (NCBI accession number: AY338466), there were some different amino acids (W16R, G21A, A53T,

```

gctggttttcccggtgaattcctctctgctaacccatcggtccatcagttgcagttattacaatttagttacagaagaa
atcactttttgacacaccaacctaatgaagaaaatcATGTTTCTCAGAGGTACTCTGCTGGGAATTCCTTTTCAGCAIC
      M F L R G T L L G I L L F S I
CTCCGGACAGAAACTGCTTTGGCTGGCTCTAGTTTTTTAAGCCCTGAATTTAAAAAATACAGCAACAAAATGATCCAACA
L R T E T A L A G S S F L S P E F K K I Q Q Q N D P T
AAAAACAACAGCAAAAATACATCGCCGAGGCACAGAGGGCTTCTGGGATGCAGATAAAGCAGGCAGAGAAGATGGCAACGAC
K T T A K I H R R G T E G F W D A D K A G T E D G N D
AGTATTGAACTCAGTTTCATGTTCCCTTTGAAATGGTGCAAGATAACAGAAGAGGAATATCAAGAGTATGGACAACACG
S I E L K F H V P F E I G V K I T E E E Y Q E Y G Q T
CTGGA AAAAGATGCTACAGGACATTCTCAAAGACAATGCTAAAGAAACTCCAGTGA AAAAGCTGAggcaagctgaaggacaa
L E K M L Q D I L K D N A K E T P V K S *
ccacagcacttgatgagctgatttaagtttactttaaaatgcaattattactgatttcacagctgtaagaagaaggaa
gtacatgcacctgtgtagtttgcctgacaaaacagacatcatcacagatggacttttaagtaaaaatacaagatctacaac
caaaaatagtacaggcaatgttcattgcaggtggatgtgaggcttatttcctcttttccttagtaactactgtgatta
ttctaatgacatgcttgaatctatagaactcgtctagaactgatggaacatcttcttacgcttctgtacacatct
ataatagdaaaataagctaatgctgaaactgt
    
```

FIGURE 1: cDNA and deduced amino acid sequence of the *dGHRL* gene. The arrowheads indicate the locations of introns. Polyadenylation signal is boxed. The asterisk (*) represents the putative stop codon (TGA). The nucleotide sequence is deposited in GenBank (EF613551).

SW duck	MFLRGTLGILLFSILRTETALAGSSFLSPEFKKIQQQNDPTKTTAKIHRRGTEGFWD-A
Mallard	MFLRGTLGILLFSILRTETALAGSSFLSPEFKKIQQQNDPTKTTAKIHRRGTEGFWD-A
Goose	MFLRGTLGILLFSILWTETALAGSSFLSPEFKKIQQQNDPAKATAKIHRRGTEGFWD-T
Turkey	MFLRLALLGILLLSILGTETAQAGSSFLSPAYKNIQQQKDRKPTARLHRRGTEGFWD-I
Emu	MFLRGALLVILLFVLTETTLAAGSSFLSPDYKKIQQRKDPKPTTKLHRRGVEGFSD-T
Mouse	MLSSGTICSLILLMLWMDMAMAGSSFLSPEHQKAAQRKESKKPPAKLQPRALQEWLHFE
Human	MLSSGTICSLILLMLWMDMAMAGSSFLSPEHQKAAQRKESKKPPAKLQPRALQEWLHFE
	*: :: :***:* : : ***** :: *::: * *
SW duck	DKAGTEDGNDLSIELKFHVPFIEIGVKITEEEYQYEQTLKMLQDILKDNAKETPVKS
Mallard	DKAGTEDGNDLSIELKFHVPFIEIGVKITEEEYQYEQTLKMLQDILKDNAKETPVKS
Goose	DKTGAEDDNNVELKFNVPFIEIGVKITEEEYQYEQTLKMLQDILEENAKETPVKN
Turkey	DETAGEDDNNNSVDIKFNVPFIEIGVKITEREYQYEQALEKMLQDIFEENAKETQTKD
Emu	DEAWAEDDNNNSIEIKFNVPFIEIGVKITEEQYQYEQMLEKVLGDILEENTKETRMKN
Mouse	DRGQAEETEELEIRFNAPFDVGIKLSGAQYQQHGRALGKFLQDILWEEVKEAPADK
Human	DRGQAEETEELEIRFNAPFDVGIKLSGAQYQQHGRALGKFLQDILWEEVKEAPADK
	*. *: : :***:* : : ***** : * * * * * :

FIGURE 2: Alignment of GHRL sequences among 7 species. The mature peptides were boxed.

T59A, A64T, and G71S). The putative SW *dGHRL* precursor showed identities of 88.8%, 72.4%, 69.8%, 71.6%, 37.9%, and 37.1% with its counterparts of goose, chicken, turkey, emu, mouse, and human, respectively (Figure 2). As far as the mature peptides are concerned, however, *dGHRL* had much higher homology with the other six species (92.3%, 65.4%, 65.4%, 69.2%, 53.8%, and 50.0% with goose, chicken, turkey, emu, mouse, and human, resp.) (Figure 2).

3.2. Genomic Organization of the *dGHRL* and *dGHSR* Genes. The sequence ensemble generated with the PM12 to PM19 products gave rise to a 3670 bp genomic fragment of the *dGHRL* gene, which included 798 bp of the 5' flanking region, 883 bp of five exons (134, 137, 114, 109, and 389 bp for exon 1 to exon 5) and 1989 bp of four introns (168, 469, 458, and 894 bp for intron 1 to intron 4). Like other species, the four introns of the *dGHRL* gene all followed the "GT-AG" rule, and moreover, exon 1 did not encode any amino acids (Supplementary Figure 2). The obtained *dGHRL* genomic sequences were submitted to the NCBI database (GenBank EF613552).

The 3717 bp sequence of the *dGHSR* gene was deduced by genomic walking (PM26), PCR amplification (PM25) and sequence ensemble with reported sequences (Genbank

EU005225). It comprised two incomplete exons (591 and 148 bp for exon 1 and exon 2) and one intron (2978 bp) that followed the "GT-AG" intron rule (Supplementary Figure 2). The obtained *dGHSR* gene sequences were also submitted to the NCBI database (GenBank FJ194548).

3.3. Online Prediction of Transcription Factor Binding Sites in the *dGHRL* Gene. Online prediction of transcription factor binding sites in the 5' flanking region of the *dGHRL* gene showed that a typical TATA box was present in 23 bp upstream of the transcription start site, and potential binding sites for some transcription factors were found, including one AML-1a site (runt-factor AML-1), two cap 1 sites (cap signal for transcription initiation), two CdxA 1 sites, three GATA-1 sites (GATA-binding factor 1), as well as one Oct-1 site (octamer factor 1) (Figure 3).

3.4. SNPs and Nucleotide Diversity. A total of 19 SNPs (10 transitions, 8 transversions, and 1 indel) and 1 polyA polymorphism were identified in 2751 bp of the 5' UTR region of the *cGHRL* gene, while 8 SNPs (5 transitions and 3 transversions) were identified in 1358 bp of the 5' UTR of the *cGHSR* gene (Table 3). Forty-three SNPs (27 transitions, 15 transversions, and 1 triallelic SNP), 1 SSR, and 1 9-bp

gattctcgagtagtggtgttctcctctagaagaaccttatttctgaagctatctataatagctaaataacatcctagtttc	80
tccaacaaaccacaatttaattacctccctatataccaaatcagcatgcaatgttctgttcttctgttataaacgcgatg	160
gtagccttccacacacacacacaagaagtgtctaacagttttacattcagacattgctttggaatcacgcagcaagaagtgt	240
gtgttctttcatttttcatttctatttaagatcttaactgctcaactgaaaaggctgctttacattccaggatcttacc	320
cttctccatttttcagcagtgcaatctgatcaatgtaacattttgttttaagtaaatggaagcaatgttctcaggta	400
tcattctcagttccagttctttcagcaaaagacaaacaaaatttgataaaacagaaccagttacacatttcagataacaa	480
taaaaatgctttgaggtcatttcttttgaaaaatcacattacagccatttaatttttaaaaagaacaatgttaacgcca	560
tgggaagaaacaaaacaaaacctctcactccagtcctacaccatttctattataaataaaaacaaacgcagcaaacgg	640
atgtatgcagtttgacaccttctgttgattttgtttatcttacaaagctgaccgagtttccaaaggaatgtgagtagacatata	720
ggctgcatttccagaaatgtcatggcacagaacaaatattctctctataaaacgcttcggaaagcagcagccttgg	798

FIGURE 3: Online prediction of transcription factor binding sites and the TATA box in the 5' flanking region of the *dGHRL* gene.

indel were identified in 3671 bp of the *dGHRL* gene, while 48 SNPs (35 transitions and 13 transversions) and 5 indels were found in 3567 bp of the *dGHSR* gene (Table 3). Eleven SNPs were located in the coding regions of the *dGHRL* gene, but only four SNPs were nonsynonymous and led to amino acid changes in the C-terminal extension peptide. For the *dGHSR* gene, however, no nonsynonymous SNPs were found within five SNPs in the coding regions (Table 3). A triallelic SNP (GA/C at nt 1774) was observed in the coding region of the *dGHRL* gene, which was related to three different amino acids (A/P/T at aa 59) (Table 3).

The average bps per SNP were 145, 170, 86, and 75 for the *cGHRL*, *cGHSR*, *dGHRL*, and *dGHSR* genes, respectively. The nucleotide diversities (θ) corrected for sample size were 1.83×10^{-3} , 1.56×10^{-3} , 3.10×10^{-3} , and 3.56×10^{-3} for the *cGHRL*, *cGHSR*, *dGHRL*, and *dGHSR* genes, respectively.

3.5. Tissue-Specific Expression of the *cGHRL*, *cGHSR*, *dGHRL*, and *dGHSR* Genes. The highest mRNA level of the *cGHRL* gene was found in the proventriculus and then the pituitary, cerebrum, abdominal fat, subcutaneous fat, hypothalamus, small intestine, heart, and breast muscle. The highest mRNA level of the *dGHRL* gene was also found in the proventriculus and then the abdominal fat, testicles, subcutaneous fat, uropygial gland, and breast muscle (Figure 4; Supplementary Table 1).

The highest mRNA level of the *cGHSR* gene was detected in the breast muscle and then the subcutaneous fat, leg muscle, abdominal fat, heart, spleen, liver, uropygial gland, cerebrum, proventriculus, pituitary, and testicles. The highest mRNA level of the *dGHSR* gene was detected in the pituitary and then the subcutaneous fat, hypothalamus, small intestine, testis, cerebellum, and cerebrum, whereas little mRNA was found in the other tissues (Figure 4; Supplementary Table 1).

When the HB and LY chicken breeds were compared, significant differences in the *GHRL* mRNA levels were found in the proventriculus, subcutaneous fat, lung, and kidney ($P < .05$), whereas significant differences in the *GHSR* mRNA levels were found in the hypothalamus and breast muscle ($P < .05$). When the PT duck and SW duck breeds were compared, significant differences in the *GHRL* mRNA

levels were found in the proventriculus and subcutaneous fat ($P < .05$), whereas no significant differences in the *GHSR* mRNA levels were found in any tissues ($P > .05$) (Figure 4; Supplementary Table 1).

3.6. Associations of SNPs with Fatty Traits in Chicken and Duck. For the *cGHRL* gene, C-2047G and A-2355C were found to be significantly associated with abdominal fat weight (AFW; $P = .01 < .05$) and crude protein content of leg muscle (CPCLM; $P = .02 < .05$), and A-2220C was significantly associated with CPCLM ($P = .0009 < .01$). In addition, haplotypes across six SNPs (C-2030T, C-2047G, A-2220C, A-2246G, A-2355C, and C-2399del) were significantly associated with AFW ($P = .04 < .05$), abdominal fat ratio (AFR; $P = .04 < .05$), and CPCLM ($P < .0001$). As far as the three genotypes of each locus were concerned, CC of C-2047G, CC of A-2220C, and AA of A-2355C had the highest values of association with AFW and CPCLM, indicating that the advantageous alleles for fat deposition were C-2047, A-2220, and A-2355 (Table 4). For the *cGHSR* gene, C-1459T was shown to be significantly associated with CPCLM ($P = .0004 < .01$). Individuals with the CC genotype had the highest CPCLM association, indicating that C rather than T was the dominant allele for CPCLM (Table 4).

In duck, C-729T and haplotypes of C-729T and T985C of the *dGHRL* gene were both significantly associated with subcutaneous fat thickness (SFT; $P = .04$ and $.05$, resp.). Individuals with the CT genotype of C-729T had the highest SFT. For the *dGHSR* gene, A3427G and haplotypes of T404C and A3427G were both significantly associated with SFT ($P = .04$ and $.05$, resp.). Individuals with the TT genotype of C3427T had the highest SFT, indicating that T rather than C was the dominant allele for fat deposition (Table 4).

4. Discussion

4.1. Molecular Characterization of the *GHRL* and *GHSR* Genes. The genomic organization of the *GHRL* and *GHSR* genes was conserved in chicken and duck (Supplementary Figure 2). The cDNAs of the *cGHRL* and *cGHSR* genes were reported by Kaiya et al. [16] and Tanaka et al. [19],

TABLE 3: SNPs identified in the *cGHRL*, *cGHSR*, *dGHRL*, and *dGHSR* genes.

Gene	No.	SNP ¹	Region	Note ^(2,3,4)	No.	SNP	Region	Note ^(2,3,4)
<i>cGHRL</i>	1	C-495T	5' flanking	19369777 ⁽²⁾ <i>Dde I</i> ⁽³⁾	11	A-2220C	5' flanking	19371502
	2	C-517T	5' flanking	19369799	12	A-2246G	5' flanking	19371528 <i>Tas I</i>
	3	C-873G	5' flanking	19370155	13	A-2355C	5' flanking	19371637
	4	T-1038C	5' flanking	19370320	14	C-2399del	5' flanking	19371681
	5	C-1253T	5' flanking	19370535 <i>Hin1 II</i>	15	G-2861T	5' flanking	19372143
	6	T-1562G	5' flanking	19370844	16	G-3264C	5' flanking	19372546
	7	A-1589G	5' flanking	19370871	17	C-3290T	5' flanking	19372572 <i>Tai I</i>
	8	G-1950A	5' flanking	19371232	18	G-3321T	5' flanking	19372603
	9	C-2030T	5' flanking	19371312	19	T-3324C	5' flanking	19372606
	10	C-2046G	5' flanking	19371329				
<i>cGHSR</i>	1	A-1907G	5' flanking	18787994	5	T-1288C	5' flanking	18789055
	2	C-1896A	5' flanking	18788005 <i>Tat I</i>	6	A-1022C	5' flanking	18789321 <i>Tail I</i>
	3	C-1608T	5' flanking	18788293	7	T-892C	5' flanking	18789762 <i>Bsa II</i>
	4	C-1459T	5' flanking	18788442 <i>Hin1 II</i>	8	G-834C	5' flanking	18789820 <i>Hin6 I</i>
<i>dGHRL</i>	1	G221A	5' flanking		23	T1128C	Exon 2	
	2	T256C	5' flanking		24	T1145C	Exon 2	G5 Syn ⁽⁴⁾ , <i>Csp6 I</i>
	3	T263C	5' flanking		25	C1179G	Exon 2	R17G, <i>Msp I</i>
	4	G273T	5' flanking		26	A1244T	Intron 2	
	5	A276C	5' flanking		27	T1278C	Intron 2	
	6	A303G	5' flanking		28	G1390A	Intron 2	
	7	A342C	5' flanking		29	A1756G	Exon 3	T53A, <i>Hin 6 I</i>
	8	T401C	5' flanking	<i>Csp6 I</i>	30	G1761A	Exon 3	E54 Syn
	9	C405T	5' flanking		31	G1774A/C	Exon 3	A59P/T
	10	A532C	5' flanking		32	A1789G	Exon 3	T64A
	11	A628G	5' flanking		33	C1803T	Exon 3	N68 Syn
	12	C638T	5' flanking		34	A1807G	Exon 3	G70S
	13	A645G	5' flanking		35	A1943C	Intron 3	<i>HpyCH4IV</i>
	14	C686G	5' flanking		36	C2070T	Intron 3	
	15	C691A	5' flanking		37	T2115C	Intron 3	<i>TailI</i>
	16	A736T	5' flanking		38	C2179T	Intron 3	
	17	C778A	5' flanking		39	A2324G	Exon 4	E89 Syn
	18	C1028A	Intron 1		40	G2345A	Exon 4	T96 Syn
	19	A1070G	Intron 1		41	A2351G	Exon 4	E98 Syn
	20	A1108G	Exon 2		42	G2409C	Intron 4	
	21	A1118G	Exon 2		43	T2509C	Intron 4	
	22	A1117G	Exon 2					
<i>dGHSR</i>	1	C320T	Exon 1	A Syn, <i>Msp I</i> ⁽⁴⁾	25	C2815A	Intron	
	2	T359C	Exon 1	N Syn	26	C2843T	Intron	
	3	G398C	Exon 1	T Syn, <i>Bsa II</i>	27	C2861T	Intron	
	4	T404C	Exon 1	Y Syn, <i>Csp6 I</i>	28	G2863C	Intron	
	5	G548A	Exon 1	T Syn	29	C2883T	Intron	
	6	C779T	Intron		30	T2906C	Intron	
	7	T1364A	Intron		31	C2939A	Intron	
	8	C1437T	Intron		32	T2958A	Intron	
	9	T1456C	Intron		33	G2993A	Intron	
	10	T1469C	Intron		34	G3001A	Intron	

TABLE 3: Continued.

Gene	No.	SNP ¹	Region	Note ^(2,3,4)	No.	SNP	Region	Note ^(2,3,4)
<i>dGHSR</i>	11	C1478T	Intron		35	A3304G	Intron	
	12	A2393T	Intron		36	A3037G	Intron	
	13	C2430T	Intron		37	T3050G	Intron	
	14	A2431G	Intron		38	G3080A	Intron	
	15	T2578C	Intron		39	T3208A	Intron	<i>Taa I</i>
	16	T2591A	Intron	<i>Tail I</i>	40	C3329T	Intron	<i>Taa I</i>
	17	T2619A	Intron		41	T3340A	Intron	
	18	C2657T	Intron		42	G3362A	Intron	
	19	C2681T	Intron		43	G3363A	Intron	
	20	A2682G	Intron		44	T3380C	Intron	
	21	G2746A	Intron		45	C3424T	Intron	<i>Csp6 I</i>
	22	T2792C	Intron		46	A3427G	Intron	
	23	G2794T	Intron		47	A3451G	Intron	
	24	G2803A	Intron		48	C3543T	Intron	

⁽¹⁾The first nucleotide of start codon was marked as +1, and the next upstream nucleotide was -1. SNP position was determined based on reported sequences of GenBank EF613552 (*dGHRL* gene) and FJ194548 (*dGHSR* gene), respectively. ⁽²⁾SNP position was determined according to the released chicken genomic sequence (<http://genome.ucsc.edu/>). ⁽³⁾Restriction enzyme was used for PCR-RFLP. ⁽⁴⁾SNP caused a codon and amino acid change.




FIGURE 4: Relative mRNA expression levels of the *GHRL* and *GHSR* genes in adult chicken and duck tissues. The vertical axis indicates the $2^{-\Delta C_t}$ value (mean \pm S.E.M., $n = 6$). Pituitary, Cerebrum, Lung, Abdominal fat, Liver, Testis, Gizzard, Subcutaneous fat, Spleen, Kidney, Leg muscle, Uropygial gland, Hypothalamus, Small intestine, Cerebellum, Heart, Breast muscle, and Proventriculus are considered.

respectively. The genome sequences of the *cGHRL* and *cGHSR* genes were also previously known. The reported *cGHRL* and *cGHSR* genes were 2706 and 4121 bp long, respectively, and comprised five and two exons each [17, 19, 27]. From the cloning and sequencing performed in this study, the *dGHRL* cDNA (GenBank EF613551) had a 351-bp

ORF that encodes a 116-aa precursor. The predicted mature *dGHRL* was a 26-aa peptide, and only two amino acids were different from *cGHRL*. The first seven amino acids, including a serine residue at position 3 (site of *n*-octanoylation), of the mature *dGHRL* are identical to those of chicken, turkey, goose, and emu (Figure 2) [17, 24].

TABLE 4: Effects of SNPs on fatty traits in chicken and duck.

Gene	SNP	Traits	P-value	Genotype and traits		
<i>cGHRL</i>	C-2047G	AFW	.01	29.01±1.089 ^a (C/C)	25.57±3.46 ^b (C/G)	23.30±5.676 ^b (G/G)
	A-2220C	CPCLM	.0009	20.71±0.1892 ^a (A/A)	20.03±0.319 ^a (A/C)	22.47±0.7554 ^B (C/C)
	A-2355C	CPCLM	.0203	21.83±0.4864 ^a (A/A)	20.75±0.228 ^b (A/C)	20.58±0.1889 ^b (C/C)
<i>cGHSR</i>	C-1456T	CPCLM	.0004	67.77±4.598 ^a (C/C)	66.88±1.097 ^a (C/T)	63.40±0.7162 ^B (T/T)
<i>dGHRL</i>	C-729T	SFT	.0408	1.52±0.22 ^a (T/T)	2.44±0.32 ^b (C/T)	2.02±0.48 ^{ab} (C/C)
<i>dGHSR</i>	C-3427T	SFT	.0407	0.80±0.77 ^a (C/C)	1.79±0.30 ^b (C/T)	1.93±0.23 ^b (T/T)

Note that only significant associations are shown in this table. AFW = abdominal fat weight; CPCLM = crude protein content of leg muscle; SFT = subcutaneous fat thickness. Values within a row without a common superscript letter differ and values with superscript letters differ significantly ($P < .05$ and $P < .01$ each). Genotypes of a and b mean they differed significantly ($P < .05$) and genotypes of a and B mean they differed great significantly ($P < .01$).

The obtained *dGHRL* gene sequence (GenBank EF613552) also comprised five exons and four introns, and the first exon did not encode any amino acids. This characteristic was conserved in chicken, turkey, human, and rodent [16–18, 28, 29]. The obtained 3717 bp sequence of the *dGHSR* gene (GenBank FJ194548) consisted of two exons and one intron and encoded a 347-aa mature receptor protein. All introns of the *cGHRL*, *cGHSR*, *dGHRL*, and *dGHSR* genes followed the GT-AG rule, which is conserved in eukaryotic genes.

Online prediction indicated that a typical TATA box and a number of transcriptional factor binding sites (AML-1a, cap 1, CdxA 1, GATA-1, and Oct-1) were present in the 5' flanking region of the *dGHRL* gene. These transcriptional factor binding sites were also found in the *GHRL* gene promoter regions of human and chicken [29, 30], and a "TATATAA" sequence was reported to be located at the *cGHRL* gene promoter [18]. GATA regulatory motifs were first found in the promoters of globulin and other erythroid-specific genes [31]. AML-1 is also a key regulator of hematopoiesis and plays an important role in development of all hematopoietic lineages [32]. It is possible that *dGHRL* gene expression in hematopoietic cells would be regulated by GATA-1 and AML-1a. The transcription factor cap-1 is a key factor of the cap-dependent translation initiation mechanism. Oct-1 is one of the transcription factors with a POU homeodomain and is expressed in most tissues including the brain [33]. Moreover, Oct-1 plays very important roles in the transcriptional regulation of gonadotropin-releasing hormone and aldolase C gene expression in the brain [34, 35]. CdxA is one of the vertebrate caudal proteins that play important roles in establishment of the body plan during early development [36]. These transcription factor binding sites in the putative promoter region might affect *dGHRL* gene expression.

The *GHRL* and *GHSR* genes were found to be more variable in poultry than in mammals. As indicated by this study, the SNP densities were 145, 170, 86, and 75 bp per SNP for the *cGHRL* (2751 bp/19 SNP), *cGHSR* (1358/8), *dGHRL* (3671/43), and *dGHSR* (3567/48) genes, respectively (Table 3). In humans, a total of 11 SNPs were found in 1657 bp of *GHRL* gene with an SNP density of 1 SNP/151 bp [13]. The reported density for the human *GHSR* gene (5 SNP of 1101 bp) was 220 bp per SNP [37]. Based on the dbSNP data (<http://www.ncbi.nlm.nih.gov>), a total of 8 and

32 SNPs were reported in 2922 and 6494 bp of the mouse *GHRL* and *GHSR* genes, with estimated densities of 365 and 203 bp per SNP, respectively. It was indicated that the higher variations of the *GHRL* and *GHSR* genes were found in chicken and duck compared with those of human and mouse. Moreover, as far as these two species were concerned, higher SNP densities were found in duck compared to chicken, as the nucleotide diversity of the *dGHRL* (3.10×10^{-3}) and *dGHSR* (3.56×10^{-3}) genes were much higher than those of *cGHRL* (1.83×10^{-3}) and *cGHSR* (1.56×10^{-3}). Our recent study on the chicken and duck *THRSP* α genes also showed higher variations in the duck genome (data not shown).

4.2. Effects of the *GHRL* and *GHSR* Genes on Fat Deposition. As indicated in this study, the *GHRL* and *GHSR* genes were associated with fatness traits in chicken and duck, which is consistent with some previous studies in humans. Many studies showed associations of the *GHRL* gene with adiposity [10–12, 38] and Type 2 diabetes [39] in humans. Additionally, some polymorphisms in the *GHSR* gene were found to be associated with obesity, bulimia nervosa, and pharmacological abnormalities in humans [40–43]. In chickens, the association of *cGHRL* polymorphisms with growth traits has been reported [22, 23]. Moreover, significant association of the *cGHSR* gene with several fatness traits was reported by a recent study [21]. In this study, the *GHRL* and *GHSR* genes were found to be associated with several fatness traits like AFW, CPCLM, and SFT in both chicken and duck.

Analysis of mRNA levels also showed the effects of the *GHRL* and *GHSR* genes on fat deposition in poultry. Even though the *GHRL* gene was predominantly expressed in the proventriculus, high mRNA levels of the *GHRL* and *GHSR* genes were detected in subcutaneous fat and abdominal fat in both chicken and duck (Figure 4). This suggests that both *GHRL* and *GHSR* genes may be involved in fat mobilization in poultry. Moreover, much higher *GHRL* mRNA levels were found in LY (2.07 ± 0.06) rather than HB (0.07 ± 0.05) chickens; higher levels were found in SW (0.78 ± 0.01) compared to those in PT (0.13 ± 0.01) ducks (Supplementary Table 1). SW ducks and LY chickens are two fast-growing commercial breeds with higher fat contents compared with those of the other two native breeds of PT ducks and HB chickens. The higher *GHRL* mRNA

in higher-fat breeds suggested that GHRL was probably advantageous for fat deposition in poultry.

5. Conclusion

From these investigations, the variations and expression patterns of the *cGHRL*, *cGHSR*, *dGHRL*, and *dGHSR* genes were characterized, and some SNPs of these genes were related to fatty traits.

Acknowledgments

This work was funded by projects under the Major State Basic Research Development (973) Program of China (2006CB102107) and the National Natural Science Foundation of China (30600429). Q. Nie and M. Fang contributed this work equally.

References

- [1] A. D. Howard, S. D. Feighner, D. F. Cully, et al., "A receptor in pituitary and hypothalamus that functions in growth hormone release," *Science*, vol. 273, no. 5277, pp. 974–977, 1996.
- [2] M. Kojima, H. Hosoda, Y. Date, M. Nakazato, H. Matsuo, and K. Kangawa, "Ghrelin is a growth-hormone-releasing acylated peptide from stomach," *Nature*, vol. 402, no. 6762, pp. 656–660, 1999.
- [3] Y. Date, M. Kojima, H. Hosoda, et al., "Ghrelin, a novel growth hormone-releasing acylated peptide, is synthesized in a distinct endocrine cell type in the gastrointestinal tracts of rats and humans," *Endocrinology*, vol. 141, no. 11, pp. 4255–4261, 2000.
- [4] M. Matsumoto, H. Hosoda, Y. Kitajima, et al., "Structure-activity relationship of ghrelin: pharmacological study of ghrelin peptides," *Biochemical and Biophysical Research Communications*, vol. 287, no. 1, pp. 142–146, 2001.
- [5] V. Monti, J. J. Carlson, S. C. Hunt, and T. D. Adams, "Relationship of ghrelin and leptin hormones with body mass index and waist circumference in a random sample of adults," *Journal of the American Dietetic Association*, vol. 106, no. 6, pp. 822–828, 2006.
- [6] C. Stylianou, A. Galli-Tsinopoulou, D. Farmakiotis, et al., "Ghrelin and leptin levels in obese adolescents. Relationship with body fat and insulin resistance," *Hormones*, vol. 6, no. 4, pp. 295–303, 2007.
- [7] J. H. N. Lindeman, H. Pijl, F. M. H. van Dielen, E. G. W. M. Lentjes, C. van Leuven, and T. Kooistra, "Ghrelin and the hypostatotropism of obesity," *Obesity Research*, vol. 10, no. 11, pp. 1161–1166, 2002.
- [8] C. M. Foster, A. Barkan, J. Z. Kasa-Vubu, and C. Jaffe, "Ghrelin concentrations reflect body mass index rather than feeding status in obese girls," *Pediatric Research*, vol. 62, no. 6, pp. 731–734, 2007.
- [9] O. Ukkola, E. Ravussin, P. Jacobson, et al., "Role of ghrelin polymorphisms in obesity based on three different studies," *Obesity Research*, vol. 10, no. 8, pp. 782–791, 2002.
- [10] O. Ukkola, E. Ravussin, P. Jacobson, et al., "Mutations in the preproghrelin/ghrelin gene associated with obesity in humans," *The Journal of Clinical Endocrinology & Metabolism*, vol. 86, no. 8, pp. 3996–3999, 2001.
- [11] A. Hinney, A. Hoch, F. Geller, et al., "Ghrelin gene: identification of missense variants and a frameshift mutation in extremely obese children and adolescents and healthy normal weight students," *The Journal of Clinical Endocrinology & Metabolism*, vol. 87, no. 6, pp. 2716–2719, 2002.
- [12] R. Ando, S.-I. Kawakami, T. Bungo, et al., "Feeding responses to several neuropeptide Y receptor agonists in the neonatal chick," *European Journal of Pharmacology*, vol. 427, no. 1, pp. 53–59, 2001.
- [13] J. Vartiainen, Y. A. Kesäniemi, and O. Ukkola, "Sequencing analysis of ghrelin gene 5' flanking region: relations between the sequence variants, fasting plasma total ghrelin concentrations, and body mass index," *Metabolism*, vol. 55, no. 10, pp. 1420–1425, 2006.
- [14] T. Tsubone, T. Masaki, I. Katsuragi, K. Tanaka, T. Kakuma, and H. Yoshimatsu, "Leptin downregulates ghrelin levels in streptozotocin-induced diabetic mice," *American Journal of Physiology*, vol. 289, no. 6, pp. R1703–R1706, 2005.
- [15] T. Shimbara, M. S. Mondal, T. Kawagoe, et al., "Central administration of ghrelin preferentially enhances fat ingestion," *Neuroscience Letters*, vol. 369, no. 1, pp. 75–79, 2004.
- [16] H. Kaiya, S. van der Geyten, M. Kojima, et al., "Chicken ghrelin: purification, cDNA cloning, and biological activity," *Endocrinology*, vol. 143, no. 9, pp. 3454–3463, 2002.
- [17] Q. Nie, H. Zeng, M. Lei, et al., "Genomic organisation of the chicken ghrelin gene and its single nucleotide polymorphisms detected by denaturing high-performance liquid chromatography," *British Poultry Science*, vol. 45, no. 5, pp. 611–618, 2004.
- [18] M. P. Richards, S. M. Poch, and J. P. McMurtry, "Characterization of turkey and chicken ghrelin genes, and regulation of ghrelin and ghrelin receptor mRNA levels in broiler chickens," *General and Comparative Endocrinology*, vol. 145, no. 3, pp. 298–310, 2006.
- [19] M. Tanaka, T. Miyazaki, I. Yamamoto, et al., "Molecular characterization of chicken growth hormone secretagogue receptor gene," *General and Comparative Endocrinology*, vol. 134, no. 2, pp. 198–202, 2003.
- [20] Q. Nie, M. Lei, J. Ouyang, H. Zeng, G. Yang, and X. Zhang, "Identification and characterization of single nucleotide polymorphisms in 12 chicken growth-correlated genes by denaturing high performance liquid chromatography," *Genetics Selection Evolution*, vol. 37, no. 3, pp. 339–360, 2005.
- [21] M. Lei, C. Luo, X. Peng, et al., "Polymorphism of growth-correlated genes associated with fatness and muscle fiber traits in chickens," *Poultry Science*, vol. 86, no. 5, pp. 835–842, 2007.
- [22] C. C. Li, K. Li, J. Li, et al., "Polymorphism of ghrelin gene in twelve Chinese indigenous chicken breeds and its relationship with chicken growth traits," *Asian-Australasian Journal of Animal Sciences*, vol. 19, no. 2, pp. 153–159, 2006.
- [23] M. Fang, Q. Nie, C. Luo, D. Zhang, and X. Zhang, "An 8 bp indel in exon 1 of ghrelin gene associated with chicken growth," *Domestic Animal Endocrinology*, vol. 32, no. 3, pp. 216–225, 2007.
- [24] J. Yuan, J. Zhou, X. Hu, and N. Li, "Molecular cloning and comparison of avian preproghrelin genes," *Biochemical Genetics*, vol. 45, no. 3–4, pp. 185–194, 2007.
- [25] SAS Institute, *SAS® User's Guide. Statistics. Version 6.12*, SAS Institute, Cary, NC, USA, 1996.
- [26] M. Cargill, D. Altshuler, J. Ireland, et al., "Characterization of single-nucleotide polymorphisms in coding regions of human genes," *Nature Genetics*, vol. 22, no. 3, pp. 231–238, 1999.
- [27] S. M. E. Geelissen, I. M. E. Beck, V. M. Darras, E. R. Kühn, and S. van der Geyten, "Distribution and regulation of chicken

- growth hormone secretagogue receptor isoforms," *General and Comparative Endocrinology*, vol. 134, no. 2, pp. 167–174, 2003.
- [28] N. Nakai, M. Kaneko, N. Nakao, et al., "Identification of promoter region of ghrelin gene in human medullary thyroid carcinoma cell line," *Life Sciences*, vol. 75, no. 18, pp. 2193–2201, 2004.
- [29] M. Tanaka, Y. Hayashida, T. Iguchi, N. Nakao, N. Nakai, and K. Nakashima, "Organization of the mouse ghrelin gene and promoter: occurrence of a short noncoding first exon," *Endocrinology*, vol. 142, no. 8, pp. 3697–3700, 2001.
- [30] N. Kanamoto, T. Akamizu, T. Tagami, et al., "Genomic structure and characterization of the 5' -flanking region of the human ghrelin gene," *Endocrinology*, vol. 145, no. 9, pp. 4144–4153, 2004.
- [31] T. Evans, M. Reitman, and G. Felsenfeld, "An erythrocyte-specific DNA-binding factor recognizes a regulatory sequence common to all chicken globin genes," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 85, no. 16, pp. 5976–5980, 1988.
- [32] E. Ogawa, M. Maruyama, H. Kagoshima, et al., "PEBP2/PEA2 represents a family of transcription factors homologous to the products of the *Drosophila* runt gene and the human AML1 gene," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 90, no. 14, pp. 6859–6863, 1993.
- [33] M. A. M. Groenen, R. J. M. Dijkhof, J. J. van der Poel, R. van Diggelen, and E. Verstege, "Multiple octamer binding sites in the promoter region of the bovine α 2-casein gene," *Nucleic Acids Research*, vol. 20, no. 16, pp. 4311–4318, 1992.
- [34] S. A. Eraly, S. B. Nelson, K. M. Huang, and P. L. Mellon, "Oct-1 binds promoter elements required for transcription of the GnRH gene," *Molecular Endocrinology*, vol. 12, no. 4, pp. 469–481, 1998.
- [35] H. Skala, A. Porteu, M. Thomas, et al., "Upstream elements involved in vivo in activation of the brain-specific rat aldolase C gene: role of binding sites for POU and winged helix proteins," *The Journal of Biological Chemistry*, vol. 273, no. 48, pp. 31806–31814, 1999.
- [36] Y. Margalit, S. Yarus, E. Shapira, Y. Gruenbaum, and A. Fainsod, "Isolation and characterization of target sequences of the chicken CdxA homeobox gene," *Nucleic Acids Research*, vol. 21, no. 21, pp. 4915–4922, 1993.
- [37] J. Vartiainen, S. M. Pöykkö, T. Räisänen, Y. A. Kesäniemi, and O. Ukkola, "Sequencing analysis of the ghrelin receptor (growth hormone secretagogue receptor type 1a) gene," *European Journal of Endocrinology*, vol. 150, no. 4, pp. 457–463, 2004.
- [38] E. Miraglia del Giudice, N. Santoro, G. Cirillo, et al., "Molecular screening of the ghrelin gene in Italian obese children: the Leu72Met variant is associated with an earlier onset of obesity," *International Journal of Obesity*, vol. 28, no. 3, pp. 447–450, 2004.
- [39] S. M. Pöykkö, O. Ukkola, H. Kauma, M. J. Savolainen, and Y. A. Kesäniemi, "Ghrelin Arg51Gln mutation is a risk factor for Type 2 diabetes and hypertension in a random sample of middle-aged subjects," *Diabetologia*, vol. 46, no. 4, pp. 455–458, 2003.
- [40] H.-J. Wang, F. Geller, A. Dempfle, et al., "Ghrelin receptor gene: identification of several sequence variants in extremely obese children and adolescents, healthy normal-weight and underweight students, and children with short normal stature," *The Journal of Clinical Endocrinology & Metabolism*, vol. 89, no. 1, pp. 157–162, 2004.
- [41] A. Baessler, M. J. Hasinoff, M. Fischer, et al., "Genetic linkage and association of the growth hormone secretagogue receptor (ghrelin receptor) gene in human obesity," *Diabetes*, vol. 54, no. 1, pp. 259–267, 2005.
- [42] K. Miyasaka, H. Hosoya, A. Sekime, et al., "Association of ghrelin receptor gene polymorphism with bulimia nervosa in a Japanese population," *Journal of Neural Transmission*, vol. 113, no. 9, pp. 1279–1285, 2006.
- [43] G. Liu, J.-P. Fortin, M. Beinborn, and A. S. Kopin, "Four missense mutations in the ghrelin receptor result in distinct pharmacological abnormalities," *The Journal of Pharmacology and Experimental Therapeutics*, vol. 322, no. 3, pp. 1036–1043, 2007.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

