Research Article **The Safety of Cruciferous Plants in Humans: A Systematic Review**

Ori Scott,¹ Elaine Galicia-Connolly,² Denise Adams,² Soleil Surette,² Sunita Vohra,^{2,3} and Jerome Y. Yager¹

¹ Pediatric Neurosciences, Department of Pediatrics, Stollery Children's Hospital, University of Alberta, Room 3-469, Edmonton Clinic Health Academy (ECHA), 11405-87 Avenue, Edmonton, AB, Canada T6G 1C9

² CARE Program, Department of Pediatrics, University of Alberta, Edmonton Clinic Health Academy (ECHA), 11405-87 Avenue,

Edmonton, AB, Canada T6G 1C9

³ School of Public Health, University of Alberta, Edmonton Clinic Health Academy (ECHA), 11405-87 Avenue, Edmonton, AB, Canada T6G 1C9

Correspondence should be addressed to Jerome Y. Yager, jyager@ualberta.ca

Received 27 July 2011; Accepted 11 October 2011

Academic Editor: Ikhlas A. Khan

Copyright © 2012 Ori Scott et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Some cruciferous plants may serve as preventive treatments for several medical conditions; our objective was to systematically investigate their safety in humans. Four electronic databases were searched, and, of 10,831 references identified, 50 were included. Data were extracted by two independent reviewers, whereafter the association between interventions and adverse events was assessed. Adverse events in 53 subjects were identified through clinical trials; of these, altered drug metabolism was rated as certainly/ likely caused by cruciferous plants. Adverse events in 1247 subjects were identified through observational studies, of which none received high causality ratings. Adverse events in 35 subjects were identified through case reports, of which allergies and warfarin resistance were rated as certainly/likely caused by cruciferous plants. We conclude that cruciferous plants are safe in humans, with the exception of allergies. Individuals treated with warfarin should consult their physician. Further investigation of uses of cruciferous plants in preventative medicine is warranted.

1. Introduction

In the last decade, a rapid rise in the demand for natural health products has become evident. This prominent trend stems from increasing awareness of the potential for significant adverse effects caused by pharmacologic interventions, along with a growing interest in preventive medicine strategies [1].

A group of plants that has been shown to possess strong anti-inflammatory and antioxidative abilities is the cruciferous plants of the *Brassica* genus. Brassica is a genus of plants in the Brassicaceae family. Vegetables of the family Brassicaceae (also called Cruciferae) are generally referred to as cruciferous vegetables. These vegetables are widely cultivated, with many genera, species, and cultivars being raised for food production. The most common Brassica vegetables eaten by people are in a single species (*B. oleracea*), including kale, collard greens, cabbage, Brussels sprouts, kohlrabi, broccoli, and cauliflower. Numerous other species in the genus are also edible such as mustard (*B. nigra*), Chinese cabbage (*B. rapa*), and oilseed rape (*B. napus;* sometimes referred to as "rapeseed oil" or "canola") [2].

Some members of this group have received wide acclaim as potential natural preventers or attenuators of several health conditions such as coronary artery disease, gastritis, and cancer [3–8]. Recent studies reveal that these therapeutic effects may extend to the fetus as well: an international casecontrol study showed that maternal consumption of cruciferous vegetables during pregnancy decreases the risk of childhood anaplastic astrocarcinomas [9]; a study in rats found that offspring of spontaneously hypertensive rats that consumed broccoli sprouts during gestation had lower blood pressure and reduced markers of oxidative stress and inflammation, compared to controls [10].

The impetus for this systematic review stems from work done in our laboratory that has shown the potential benefit of supplementing the diet of pregnant dams with broccoli sprouts in preventing perinatal brain injury. In this regard,

PICOS	Eligibility criteria	Data extracted
Patient	All human subjects were included	Age, concurrent medical conditions and treatments, reason for intervention (where relevant)
Intervention	All types of exposure (oral, topical, or respiratory) to cruciferous plants, their derivatives, or their constituents	Plant or substance exposed to, route of exposure, duration of exposure, dose (if available)
Comparators	Reports with or without a comparator group. Reports without control groups were included in order to include all potential adverse events	Numbers in the intervention and comparator groups (when relevant)
Outcome	Reports of presence or absence of adverse events	Presence or absence of adverse events, description of adverse event, acute management of adverse event (for case reports), outcome (when available), and causality
Study design	All study designs which were relevant to the assessment of safety were included. Studies in all languages were included and translated when necessary	Type of study design and setting

TABLE 1: Eligibility criteria and data extracted¹.

¹PICOS: patient, intervention, comparators, outcome, study design.

we have established a model of chronic placental insufficiency that results in fetal growth restriction and damage to the white matter of the brain, reminiscent of periventricular leukomalacia, the anatomic hallmark of cerebral palsy. Supplementing the maternal diet during the last trimester of pregnancy and first weeks of infant nutrition prevented the white matter injury and accompanying behavioural deficits in the offspring. These findings suggest, for the first time, a safe and efficacious approach to the prevention of developmental disability and cerebral palsy [11]. In order to provide the background in which to determine whether this therapy would be acceptable for humans, a thorough determination of the safety of the cruciferous species of plants in humans is necessary.

Researchers now believe that the key to the therapeutic abilities of cruciferous plants is their high content of phase-2 enzyme inducers such as sulforaphane, which induce the transcription of genes found under the control of the antioxidant response element (ARE). This causes an upregulation of the endogenous antioxidant glutathione, crucial to the cell's ability to withstand oxidative stress [12–14].

Several reviews evaluated the safety and health benefits of some components of cruciferous plants, such as indole-3-carbinol and isothiocyanates, and found them safe for humans [15, 16]. However, to our knowledge, a systematic review evaluating the safety of cruciferous plants, namely, the degree of certainty that they would not cause any adverse effects, has not yet been published.

The purpose of this paper is to systematically collect and synthesize all published reports of human adverse events associated with exposure to cruciferous plants. This approach of analyzing safety of a multi-ingredient natural product is especially important when considering consumption of the whole plant as a therapeutic means, as opposed to treatment with an isolated component or metabolite of a plant.

2. Methods

This systematic paper was undertaken in line with the relevant criteria of the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) statement [17]. The following methods used in the systematic review, including identification, screening, eligibility criteria, inclusion, and data extraction, were agreed upon by the authors *a priori*.

References were identified through comprehensive search strategies, which were developed in conjunction with a research librarian. An electronic search of the following four databases was performed: Medline, Embase, Pascal, and IPA (International Pharmaceutical Abstracts), from inception to July 2010, irrespective of language. The following search terms were used: isothiocyanate, Brassica, cruciferous, glucoraphanin, sulforaphane, broccoli, kale, cabbage, cauliflower, collard green, Brussels sprout. MESH headings and keywords were searched, and truncation was used as needed. A list of the search strategies is available as supplemental material available online at doi: 10.1155/2012/503241.

The research question and eligibility criteria were developed by using PICOS (patient, intervention, comparators, outcome, study design). The inclusion criteria were any studies reporting original data, addressing safety of cruciferous plants, their derivatives, or their constituents, to humans. Details of the eligibility criteria are reported in Table 1.

The references were imported into a bibliographic database (RefWorks). Titles and abstracts of identified studies were independently screened by two reviewers. Full texts for potentially relevant studies were obtained and reviewed for inclusion based on predetermined criteria. Disagreement was resolved by consensus, and if necessary a third party was consulted (JYY).

Data was extracted by two independent reviewers, using standardized forms. Disagreement was resolved by consensus. Data relating to the patient or group, the intervention, the comparator group (where relevant), the outcomes measured (adverse events or lack thereof), and the study design were extracted as detailed in Table 1.

The degree of association between the intervention and the adverse event was independently assessed by the two reviewers using the Causality Algorithm used by Health Canada and the WHO Collaborating Centre for International Drug Monitoring (Health Canada 2009). Categories for assessment were certainly, likely, possibly, unlikely, conditional/unclassified, and Unassessible/Unclassifiable. Disagreement between reviewers was resolved by discussion. Guidelines used to assess causality are available as supplemental material.

The studies were finally categorized into 3 groups: (1) trials, (2) observational studies, and (3) case reports. Studies were classified as controlled clinical trials when involving one or more test treatments, at least one control treatment, specified outcome measures for evaluating the studied intervention, and when the method for assigning patients to the test treatment was not a mathematical randomization technique. Randomized controlled trials, on the other hand, were defined as studies in which randomization using mathematical techniques, such as the use of a random numbers table, is employed to assign patients to test or control treatments. Studies were classified as single-case experimental design trials when the subjects served as his/her own control, rather than using another individual or a group.

We did not perform a meta-analysis as the inclusion criteria for subjects, interventions, clinical outcomes, and study designs were very heterogeneous; conducting a meta-analysis using these data would not have been appropriate.

3. Results

3.1. Study Characteristics. Searches resulted in a total of 10,831 references. After screening of titles and abstracts, the full texts of 367 articles were obtained. Of these, 50 articles met all inclusion criteria. Of the total included publications, 45 were published in English, 2 were published in German, and one each was published in Italian, French, and Spanish.

Articles were excluded for the following reasons: not human subjects (210), not evaluating adverse effects or lack thereof (21), intervention or substance exposure does not include cruciferous plants (2), not presenting original data (47). Thirty-seven articles discussing toxic oil syndrome (37) were excluded post hoc for reasons presented below.

Of the 50 included studies, 13 were trials (6 single-case experimental design trials, 4 controlled clinical trials, and 3 randomized control trials (RCTs)), 13 were observational studies (7 cohort studies, 3 case-control studies, 2 cross-sectional studies, and 1 qualitative survey), and 24 were case reports (Figure 1).

3.2. Trials. Thirteen trials were included, reporting adverse events in 53/496 patients (10.7%) [18–30]. Three of them were randomized controlled trials (RCTs), 4 were controlled clinical trials, and 6 were single-case experimental design (SCED) trials. Six of the trials were classified as safety trials, meaning that their stated primary aim was to assess safety of

interventions [18–23], whereas 7 were nonsafety trials which evaluated efficacy of interventions but also reported adverse events [24–30] (Table 2).

The safety trials [18–23] included 1 RCT, 1 controlled clinical trial, and 4 SCED trials. The only safety RCT was a double-blinded trial in 12 healthy subjects, ages 28–57, of whom 9 consumed daily doses of broccoli sprout extract combined with myrosinase (an enzyme found in all Brassica which converts glucosinolates to active isothiocyanates) over a period of 7 days. Extracts contained 25 μ mol glucosinolate, 100 μ mol glucosinolate, or 25 μ mol isothiocyanate. No adverse events were reported [18].

The second safety trial was a double-blind controlled clinical trial in 76 subjects, ages 3–39. Thirty-eight subjects had a reported mustard allergy, and 38 age- and gender-matched controls suffered from dust-mite allergy but had no mustard allergy. Both groups underwent food challenges with mustard and a variety of other allergens. Of the 38 patients, 14 had a positive mustard challenge, of which 12 had oral allergy syndrome (pruritus, mild angioedema of lips, tongue, and throat), 1 had more severe angioedema with bronchial asthma, and 1 had systemic anaphylaxis. It is important to mention that of the 38 patients, 35 were atopic (including rhinitis, and/or bronchial asthma, atopic dermatitis) [19].

The last four safety trials were all single-case experimental design trials (SCED). The first was a trial in 10 healthy subjects, ages 21-30, which strived to determine the effect of cruciferous vegetables on phenacetin and antipyrine metabolism. Subjects started by consuming a control diet for 13 days, after which they were placed on a diet containing 300 g/day of Brussels sprouts and 200 g/day of cabbage over a period of 7 days, and finally returned to the control diet for the last 10 days. In parallel, every morning the subjects were given either 900 mg of phenacetin or 1.8 mg/kg of antipyrine. The mean plasma concentrations of either drug were measured at different intervals after administration. Researchers found that the cruciferous diet enhanced the phenacetin metabolism in the gastrointestinal tract and/or during its first pass in the liver, increasing its rate of elimination. No changes in antipyrine metabolism were noted [20]. It is worth noting that the US Food and Drug Administration ordered the withdrawal of drugs containing phenacetin in November 1983, owing to its carcinogenic and kidney-damaging properties (Federal Register of October 5, 1983 (48 FR 45466)).

The second SCED trial included 10 healthy subjects, ages 23–35, and aimed to determine the effect of cruciferous vegetables on acetaminophen and oxazepam metabolism. Subjects started by consuming a control diet for 10 days, after which they were put on a diet containing 300 g/day of Brussels sprouts and 200 g/day of cabbage over a period of 10 days, and finally returned to the control diet for the last 10 days. In parallel, every morning the subjects were given either 45 mg of oxazepam or 1500 mg of acetaminophen. The mean plasma concentrations of either drug were measured at different intervals after administration, and a 24 h urinary recovery of conjugates was performed. Researchers found that the cruciferous diet enhanced the acetaminophen metabolism and glucuronide conjugation; however, they mention



FIGURE 1: PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) flow diagram of study identification, inclusion, and exclusion.

that the decreased levels of cysteine conjugate in urine actually indicate reduced toxicity. No changes in oxazepam metabolism were noted [21].

The third safety SCED trial included 10 healthy subjects, ages 22–40; the goal of this study was to determine the effect of a diet rich in Brussels sprouts on warfarin metabolism. Subjects consumed their customary diets for 3 days and were then instructed to consume 400 g/day of Brussels sprouts for the next 18 days. Warfarin administration (20 mg) was provided twice on days 1 and 18. Mean plasma concentrations of warfarin were measured at different time periods, as well as prothrombin activity, and clearance rates were calculated. Researchers showed that the diet had resulted in accelerated warfarin metabolism and decreased anticoagulation [22].

The last safety SCED trial included 6 subjects, ages 20–38, with a history of recurrent allergic reactions to consumption

of raw cabbage. This study aimed to determine the sensitivity profile of such patients to other cruciferous vegetables (cauliflower, mustard, and broccoli), as well as to cooked versus raw cabbage. Skin prick tests were performed for a variety of allergens; later, food challenges were performed with cooked cabbage, and levels of cabbage-specific IgE were measured in patients' sera. All patients were found to be atopic (5 with allergic rhinitis); all had positive skin prick tests to mugwort (Artemisia vulgaris) pollen and other aeroallergens (for instance, pollen of plants from the genera Parietaria and Olea), and 5 had positive skin prick tests to hazelnuts, walnuts, and peanuts. Mustard sensitivity (cross-reactivity with cabbage) was detected in all patients. Patients demonstrated no allergic reaction to cooked cabbage. All patients were found to have cabbage-specific IgE antibodies in their sera [23].

		2			;		1					
	St	tudy details		Patie	ent details		Inte	ervention)	Dutcome	
ference	Study design ¹	Setting	No. of patients/ comparators ²	Age range	Concurrent conditions/ treatments	Plant/ substance	Route ³	Duration ⁴	Dose	Adverse events	Outcome ⁵	Causality ⁶
fety trial: apiro al. 2006 8]	s RCT	Hospital	9/3	28–57 yrs	Healthy	Broccoli sprout extract	Oral	7 days	3 doses a day, 25 μmol glucosinolate 100 μmol glucosinolate, or 25 μmol isothiocyanate	No adverse events	NA	NA
gueroa al. 2005 9]	CCT	Hospital	38/38	3–39 yrs	Reported mustard allergy. Atopy and allergies to a variety of other allergens	Mustard and a variety of other allergens	Oral	Acute exposure	Oral: increasing doses of mustard (80–6480 mg)	14 showed positive mustard challenge (12 had oral allergy syndrome, 1 had more severe angioedema with bronchial asthma, and 1 had anaphylaxis)	Resolved after symptomatic treatment	Possibly
ıntuck al. 1979 0]	SCED	Multisite hospital and homes	10	21–32 yrs	Healthy	Cabbage and Brussels sprouts	Oral	7 days	200 g/d cabbage and 300 g/d Brusselss sprouts	All showed accelerated phenacetin metabolism	Resolved after diet was discontinued	Likely
intuck al. 1984 1]	SCED	Multisite hospital and homes	10	23–35 yrs	Healthy	Cabbage and Brusselss sprouts	Oral	10 days	200 g/d cabbage and 300 g/d Brusselss sprouts	All showed accelerated metabolism, enhanced glucuronide conjugation	Resolved after diet was discontinued	Likely
vesen al. 1988 2]	SCED	Multi-site hospital and homes	10	22-40 yrs	Healthy	Brusselss sprouts, lightly steamed	Oral	2 weeks	400 g/d	All showed accelerated warfarin metabolism	Resolved after diet was discontinued	Likely

TABLE 2: Trials of cruciferous plants, their derivatives, or their constituents in humans.

		Causality ⁶	Possibly	Possibly	NA	NA	NA	NA
	Jutcome	Outcome ⁵	NA	Full resolution: in adult after returning to original dose. In pediatric cases spontaneously	NA	NA	VA	AN
	Õ	Adverse events	All showed positive skin prick tests and IgE tests to raw cabbage; positive skin prick tests to a variety of allergens	Imbalance and tremor in one adult who was given twice the original dose by the researchers and unsteadiness with nausea in two pediatric patient at who by mistake took higher doses	No adverse events	No adverse events	No adverse events	No adverse events
		Dose	NA	200 mg twice a day for adults; for kids based on weight	Not specified	20 g/d	Not specified	200 mg twice a day for adults; pediatric dosage was determined by weight
-	ervention	Duration ⁴	Acute exposure	9–24 months	12 days	1 year	4 months	10–86 months
unuea.	Int	Route ³	Oral and topical allergy tests	Oral	Oral	Oral	Oral	Oral
IABLE 2: CON		Plant/ substance	Cabbage (raw versus cooked) and a variety of other allergens	Indole-3- carbinol supplement	Broccoli sprout infusion	Mustard oil	Cauliflower leaves powder, in biscuits, or shakarpara	Indole-3- carbinol supplement
	Patient details	Concurrent conditions/ treatments	Recurrent allergic reactions after consuming cabbage. Atopy	Recurrent respiratory papillomatosis	Healthy	Acute myocardial infarction	Nutritional deficits: low serum hemoglobin and retinol	Recurrent respiratory papillomatosis
- 6		Age range	20–38 yrs	2.5–61 yrs	25–65 yrs	Mean: 48 yrs	10–12 yrs	5-71 yrs
		No. of patients/ comparators ²	Q	18	100/100	120/118	33/33	33
-	tudy details	Setting	Hospital	Hospital	Community	Multisite hospital and homes	Community	Multi-site hospital and homes
Ċ	N	Study design ¹	SCED	CCT	RCT	RCT	CCT	CCT
		Reference	Vovolis et al. 2009 [23]	Nonsafety trials Rosen et al. 1998 [24]	Kensler et al. 2005 [25]	Singh et al. 1997 [26]	Jood et al. 2001 [27]	Rosen and Bryson 2004 [28]

TABLE 2: Continued.

Study details Study Setting design ¹ Setting CCT Single site SCED Single site
Study Study (Study design ¹ Sett design ¹ Sett CCT Sing SCED Sing

²Comparator numbers appear only for controlled trials.

³Route of exposure. ⁴Duration of exposure. ⁵Degree of resolution in cases were adverse events were reported; NA, not applicable. ⁶The degree of association between the intervention and the adverse event, as rated by reviewers; NA, not applicable.

The nonsafety trials [24–30] included 2 RCTs, 4 controlled clinical trials, and 1 single-case experimental design trial, of which only one identified adverse events.

The trial which reported the adverse events was a controlled clinical trial in 18 subjects, ages 2.5-61, who suffered from recurrent respiratory papillomatosis and underwent complete surgical removal. All subjects received oral indole-3-carbinol supplementation for a period of 9-24 months in order to examine whether the supplements reduced the occurrence of the papillomas. Doses were 200 mg twice a day for adults, and individually calculated for pediatric patients based on body weight. Adverse events were reported in 3 patients: 1 adult, who was receiving a higher dose (400 mg twice a day) for 10 days according to the researchers' recommendation, suffered from imbalance and tremor and 2 girls, 2.5 and 12 years old, who took an overdose by mistake, suffered from unsteadiness and nausea. The symptoms in the adult resolved after returning to original dose and resolved spontaneously in the girls within a few hours to a day. According to the authors, the supplement did not cause any other side effects or complications (including acceleration of the disease) and was generally well tolerated [24]. It is unclear from this paper which other medications these specific individuals were taking, and whether they had any other acute health conditions which may have contributed to the adverse events described.

All other non-safety trials reported no adverse events with oral consumption of broccoli sprout infusion/homogenate, mustard oil, cauliflower leaves powder or indole-3carbinol supplements, or with topical application of broccoli sprouts extract [25–30]. These studies included both adult and pediatric populations, whose concurrent medical conditions included none (healthy), nutritional deficits (haemoglobin and retinol), recurrent respiratory papillomatosis, and acute myocardial infarction (Table 2).

3.3. Observational Studies. Thirteen observational studies were included, reporting adverse events related to cruciferous plants in 1,247 patients [31–43]. Seven of them were cohort studies, 3 were case-control studies, 2 were cross-sectional studies, and 1 was a qualitative survey (Table 3).

Three cohort studies, one case-control study, and one cross-sectional survey examined prevalence of allergic reactions to oilseed rape [31–35]. In the first cohort, 7/1478 subjects who were naturally exposed to oilseed rape had either a positive skin test, a positive radio-allergosorbent test, or nasal sensitivity in response to oilseed rape exposure, whereas the proportion of such allergic reactions was 14/37 in those who were occupationally exposed to oilseed rape [31].

In the second cohort, only 147/4468 subjects with suspected oilseed allergy showed positive skin prick test to oilseed rape; most of the subjects in this study were allergic to a variety of other antigens [32].

In the third cohort, 12/22 village residents reported increased allergic symptoms during a year when oilseed rape surrounded the village, compared with a year when another crop surrounded the village. However, authors mention that the symptoms reported did not correlate with oilseed rape pollen levels measured. They were therefore not sure what the true cause could be [33].

In a case-control study, 37 people complained of seasonal allergic symptoms and bronchial reactivity in response to an unknown allergen, of whom 23 were tested. However, only 2 were found to be allergic to oilseed rape and only 10 (including those 2) were found to be atopic. The authors concluded that the symptoms could not be attributed to oilseed rape in most of these cases of seasonal allergy [34].

In the cross-sectional survey, 683/869 of village residents who were exposed to oilseed rape complained of seasonal cough, wheeze, and headaches. However, the authors concluded that the proportion of people who suffered from such symptoms was not much higher in subjects living in close proximity to oilseed rape in comparison to control subjects who do not; this suggests that the seasonal symptoms in rural areas cannot be attributed to oilseed rape allergy alone [35].

Yet another observational study dealing with allergic reactions was a cohort in 259 individuals with suspected contact allergy to foods containing allyl-isothiocyanate. Of 259 subjects who underwent allergy skin tests, 43 had a questionable reaction, of whom 15 had irritation and 3 had follicular reaction. Only two showed a true positive reaction, but one was lost to followup [36].

Three studies suggested a possible connection between cruciferous vegetables and cancer. The first was a cohort study in 64,327 women in Japan, ages 40-79, which examined the possible connection between dietary habits and risk of ovarian cancer death, based on food-frequency questionnaires. Whereas no adverse events were reported with consumption of cabbage or green leafy vegetables, a positive association was established between moderate-high consumption of Chinese cabbage and ovarian cancer. The authors suggest that this might be attributed to the fact, that, in many cases, Chinese cabbage is eaten pickled, as pickled food was proven to increase the risk of cancer [37]. The researchers did not adjust for total energy consumption, or for comalignancies such as breast/endometrial cancer; the first is important as several studies found an association between total energy consumption and increased risk for various types of cancer [38, 39]. The latter is crucial, as many women with a history of such comalignancies are at a higher risk of developing ovarian cancer than the general population [40, 41].

The second suggested a connection between cruciferous vegetables and cancer in a case-control study from Kuwait in which 313 thyroid cancer patients were paired with age- and gender-matched controls; the study's aim was to examine the relationship between different sociodemographic, medical or dietary factors and thyroid cancer, based on questionnaires. Whereas no adverse events were associated with broccoli consumption, a nonstatistically significant positive association was established between moderate-high consumption of cauliflower or cabbage and thyroid cancer (P = 0.08 and P = 0.16, resp.). The authors conclude that no clear association between consumption of cruciferous vegetables and thyroid cancer could be established [42].

The last study bringing up a possible association between cruciferous vegetable consumption and cancer is a case-control study which paired 246 thyroid cancer patients from

	Causality ⁸	ed Possibly	Possibly	ent Possibly	st Possibly
utcome	Outcome ⁷	Not specifi	NA	Improveme when not exposed	Spontaneou resolution
C	Adverse events ⁶	In naturally exposed, 4/1478 showed positive skin test and 3/1478 showed positive RAST and nasal sensitivity. In occupationally exposed, 9/37 showed positive skin test and 5/37 showed positive RAST and nasal sensitivity	147 showed positive skin prick test to oilseed rape	Allergy (sneezing, coughing, eye irritation) in 10	In 43patients: ?+ reaction, of whom 15 had irritation and 3 had follicular reaction. Two showed a true + reaction but one lost to follow up
	Dose ⁵	A NA	Ϋ́Α	NA	NA
ntion	Duration ⁴	Acute to chronic	1 year	Seasonal	Acute
Intervei	Route ³	Respiratory and topical allergy tests	Respiratory	Respiratory	Topical allergy test
	Plant/ substance	Oilseed rape	Oilseed rape	Oilseed rape	Allyl isothiocyanate 0.1 % in petrolatum
tient details	Concurrent conditions/ treatments	Atopy in some subjects	Suspected inhalant allergy to oilseed rape. Multiple allergies to other pollen allergens	Not given	Suspected contact allergy to foods containing allyl isorhiocvanate
Pa	Age range	Adults	Not given	Adults	Not given
tails	No. of patients/ comparators ²	1515 (1478: natural exposure, 37: occupational exposure)	4468	, 22	259
Study de	Setting	Multi-site	Single site	Community	Hospital
	Study design ¹	Cohort	Cohort	Cohort	Cohort
	Reference	Fell et al. 1992 [31]	Hemmer et al. 1997 [32]	Parrat et al. 1995 [33]	Lerbaek et al. 2004 [36]

		Causality ⁸	Possibly	NA	NA	Possibly (for 2 who are allergic) Unlikely (for the rest)
	utcome	Outcome ⁷	Death	NA	NA	Not mentioned
	0	Adverse events ⁶	No adverse events with consumption of cabbage or green leafy vegetables. Of 100 women who reported moderate-high consumption of Chinese cabbage, 46 had ovarian cancer	No adverse events	No adverse events	Of the 23 cases tested, only 2 were found to be truly allergic to oilseed rape, and only 10 (including these 2) were atopic
		Dose ⁵	Varies: 0 times a week to almost every day	5– 7 mg/kg/day	NA	NA
	ntion	Duration ⁴	Chronic (exact time frame given)	7 days	chronic (reported intake in last 12 months)	Seasonal
Continued.	Interve	Route ³	Oral	Oral	Oral	Respiratory
TABLE 3: (Plant/ substance	Cabbage, Chinese cabbage, green leafy vegetables, and other foods	Indole-3- carbinol	Cruciferous vegetables	Oilseed rape
	tient details	Concurrent conditions/ treatments	Not given	No history of recent or chronic illness, drug use, or recent changes in weight	Male smokers with no history of cancer, not using vitamins A, E or beta-carotene in excess	Seasonal allergic symptoms and bronchial reactivity
	Pat	Age range	40– 79 yrs	22- 48 yrs	50– 69 yrs	17– 54 yrs
	ails	No. of patients/ comparators ²	64327 women	12	27111	37/24
	Study det	Setting	Community	Community	Community	Hospital
		Study design ¹	Cohort	Cohort	Cohort	CCS
		Reference	Sakauchi et al. 2007 [37]	Michnovicz and Bradlow 1997 [44]	Michaud et al. 2002 [45]	Soutar et al. 1995 [34]

10

Study deta	Study deta	ta	ils	Pat	tient details	IABLE 3:	Lonunuea. Interven	ıtion		0	utcome	
idy Setting No. of Age Concurrent ign ¹ Setting patients/ range conditions/ comparators ² range treatments	No. of Age Concurrent Setting patients/ range conditions/ comparators ² range treatments	No. of Age Concurrent patients/ range conditions/ comparators ² range treatments	Age Concurrent range conditions/ treatments	Concurrent conditions/ treatments		Plant/ substance	Route ³	Duration ⁴	Dose ⁵	Adverse events ⁶	Outcome ⁷	Causality ⁸
Cases: thyro cancer. Eith cases or controls: asthma, diabetes mellitus, gal diabetes mellitus, gal diabetes mellitus, gal diabetes control 313/313 5- disease, tension, lupous, polyposis co skin allergy, skin allergy, skin allergy, skin allergy	Cases: thyro cancer. Eith cases or controls: asthma, diabetes mellitus, gal mellitus, gal bladder ancer 313/313 5- disease, bladder tension, hyper- tension, hyper- skin allergy, skin allergy, skin allergy, skin disease (not specified)	Cases: thyro cancer. Eitho cases or controls: asthma, diabetes mellitus, gal bladder bladder bladder bladder tension, hyper- tension, lupous, polyposis co skin allergy, skin allergy, skin disease (not specified)	Cases: thyro cancer. Eitho cases or controls: asthma, diabetes mellitus, gal bladder 70 yrs hyper- tension, lupous, polyposis co skin allergy, skin allergy, skin disease (not specified)	Cases: thyro cancer. Eitho cases or controls: asthma, diabetes mellitus, gal bladder disease, hyper- tension, lupous, polyposis co skin allergy, skin disease (not specified)	li, l	Cabbage, cauliflower, broccoli	Oral	Chronic (exact frime given)	Vāries: 0 to 7 days a week	No adverse events with consumption of broccoli. 63/101 people with high cabbage consumption and 55/91 people with high cauliflower consumption had thyroid cancer; however, <i>P</i> trends were not statistically significant (0.08 and 0.16, resp.)	Not mentioned	Possibly
.S Community 246/440 18- Not given 75 yrs	Community 246/440 18- Not given	r 246/440 18- Not given	18– 75 yrs Not given	Not given		White and red cabbages, cauliflower, broccoli, Brusselss sprouts	Oral	Chronic (exact time frame given)	Varies: <2 to >6 times a week	56/110 people who at anytime lived in areas in Sweden where goiter and iodine deficiency were endemic until the 1960's and who reported moderate-high consumption of cruciferous vegetables had thyroid cancer	Not mentioned	Possibly
Both villages: S Community 869/867 14– 448 smokers, 50 yrs 325 ex-smokers	Both villages: Community 869/867 14– 448 smokers, 50 yrs 325 ex-smokers	Both villages: 869/867 14– 448 smokers, 50 yrs 325 ex-smokers	Both villages: 14– 448 smokers, 50 yrs 325 ex-smokers	Both villages: 448 smokers, 325 ex-smokers	1	Oilseed rape	Respiratory	Chronic: months	NA	683 of 869 who were exposed had seasonal cough, wheeze, and headaches	Not mentioned	Possibly

						TABLE 3:	Continued.					
		Study de	etails	Pat	tient details		Interve	ntion		0	utcome	
Reference	Study design ¹	Setting	No. of patients/ comparators ²	Age range	Concurrent conditions/ treatments	Plant/ substance	Route ³	Duration ⁴	Dose ⁵	Adverse events ⁶	Outcome ⁷	Causality ⁸
Sato et al. 2004 [46]	CSS	Communit	y 438	39– 60 yrs	No history of gastric cancer or gastric ulcer	Broccoli	Oral	Chronic (exact time frame given)	Varies: never to few times a week	46/186 people who consumed broccoli once or more a week had changes in enzymes which might indicate chronic atrophic gastritis	Ongoing	Possibly
Lust et al. 1996 [47]	S	Communit	y 273	<4 mo	Not given	Various foods, including cruciferous vegetables (specifically cabbage, cauliflower, broccoli)	Oral (through breast milk)	Unclear (mothers were asked whether they ate different items the previous week)	Not given	63/273 exhibited colic symptoms (abdominal pain, irritability, intense crying)	Not mentioned	Possibly
¹ CCS, case-c ² Comparaton ³ Route of exj ⁴ Duration of ⁵ NA, not app ⁶ RAST, radio ⁶ RAST, radio ⁷ Degree of re ⁸ The degree of	nntrol stur numbers oosure. exposure. licable, in allergosorl solution ii f associati	ly; CSS, cross- appear only fo cases where dc cases were ad on between th	sectional study; QS, r case-control and, se cannot be quant verse events were ru e intervention and	, qualitati cross-sect ified. pported; h	ve survey. ional surveys. VA, not applicable. se event, as rated bj	y reviewers; NA, nc	ot applicable.					

12

Sweden and Norway with 440 age- and gender-matched controls. The goal of the study was to evaluate the relationship between certain dietary habits and the risk of follicular and papillary thyroid carcinomas. With regard to cruciferous vegetables, researchers found that for people who, at any time, resided in areas of Sweden where goiter and iodine deficiency were endemic until the 1960's, there was an increased risk for thyroid cancer with consumption of cruciferous vegetables. These findings were not reproduced anywhere else in Sweden or Norway. Authors suggest that this dual effect might be related to interactions between cruciferous vegetables and other food components such as iodine [43]. They also expressed surprise, as their findings stood in contradiction to those of many research groups that found a protective effect of cruciferous vegetables against thyroid cancer [48–51].

The next observational study included in our paper was a cross-sectional study which aimed to determine the association between broccoli consumption and chronic atrophic gastritis in 438 men, ages 39–60 years. The authors found that consumption of broccoli once or more weekly increased the risk for chronic atrophic gastritis, based on serological tests for pepsinogen I and II. However, the authors acknowledge some major drawbacks to their work. First, *H. Pylori* infection, which increases the risk for chronic atrophic gastritis, was not measured. Second, serologic criteria were used for diagnosis, meaning that the results were prone to measurement errors. Moreover, it is unclear whether the changes in enzyme profiles were accompanied by any clinical manifestations [46].

The last observational study reporting adverse events was a qualitative survey which examined the relationship between consumption of certain foods by breast-feeding mothers, and the appearance of colic symptoms in their babies (4 months old or younger). The survey conductors found a positive association between consumption of cruciferous vegetables (cabbage, cauliflower, and broccoli) and colic symptoms: abdominal pain, irritability, and intense crying [47]. This study exhibited several major limitations: first, all determinants of colic symptoms, as defined by the authors (intense crying, irritability, and abdominal pain) can be subjected to subjective interpretation; the diagnosis of infantile colic, as established over 50 years ago, requires a healthy baby to exhibit "periods of intense, unexplained fussing/crying lasting more than 3 hours a day, more than 3 days a week for more than 3 weeks" [52]. In the study, however, no such objective temporal requirements were defined; any presentation of the symptoms, regardless of duration, was considered to constitute colic. Furthermore, since the questionnaire was administered over the course of one week only, instead of the 3 weeks which are required for diagnosis of colic, it is doubtful whether such diagnosis can be made at all. Moreover, since food intake and colic symptoms were reported only for a week, it is hard to establish proper temporal relationships between the two.

Two cohort studies included in our paper identified no adverse events with exposure to cruciferous plants or their constituents; the first study included 12 subjects who ingested 5–7 mg/kg/day of indole-3-carbinol [44]. The second study, which followed 27,111 male smokers over a period of

12 months, attempted to discover whether intakes of fruits, vegetables, carotenoids, or vitamins A, E, or C could be associated with risk of bladder cancer in these people. The study determined that no association between bladder cancer risk and chronic consumption of cruciferous vegetables could be established [45].

3.4. Case Reports. Twenty-four case reports were included in the review, reporting adverse events in 35 individuals. The patients ranged from 17 to 70 years of age, whose concurrent medical conditions included allergies, both to cruciferous plants and to a variety of other allergens, asthma, bronchitis, atopy, cardiovascular diseases, and different skin conditions (eczema, pruritus, eryhtema, dermatitis, dryness and scaling, and blisters) (Table 4). The adverse events occurred after topical, respiratory, or oral exposure to different cruciferous plants, their constituents, or their derivatives. Twenty-one case reports reported allergic or hypersensitivity reactions in 31 individuals, including allergic contact dermatitis/contact urticaria, contact hypersensitivity, aggravation of eczema, cutaneous lesions similar to pityriasis rosea, asthma, rhinoconjunctivitis, aggravation of cough and chest pain, local swelling and itching, and anaphylaxis. The suspected agents triggering these reactions were turnip seeds, cabbage, broccoli, oilseed rape (flour or pollen), cauliflower, mustard (a variety of preparation forms), isothiocyanates in paint, and Diplotaxis erucoides (Brassica erucoides) pollen. It is important to note that 18 of the individuals (approximately 50% of total) had previous risk factors such as atopy, allergies to allergens unrelated to cruciferous plants (for instance, grasses, dust mite, nuts), different skin conditions, or asthma. Furthermore, 12 of those people had been exposed to the allergens occupationally, for chronic periods of time, presumably leading to the adverse events [53–73].

Two case reports documented warfarin resistance in 3 patients with cardiovascular diseases. The first patient had a prosthetic aortic valve and a history of myocardial infarctions with prolonged prothrombin time [74]. The second had pulmonary embolism, and the third suffered from an unspecified cardiovascular disease [75]. The adverse event in the first patient was attributed to excessive consumption of lettuce and greens (turnip, mustard greens, broccoli); the patient was a 35 years old woman who intended to lose weight by consuming only the above-mentioned foods, which led to a consumption of $6000 \,\mu\text{g}/\text{day}$ of vitamin K (60 times higher than the recommended consumption). After 5 weeks of dieting, she felt substernal chest pain and was referred to the hospital, where a myocardial infarction was diagnosed. She was treated with heparin and nitroglycerine. The second and third cases were attributed to chronic consumption of up to 450 g/day of broccoli; one of them required treatment with Coumadin. All three cases ended up in full resolution of adverse events and a recommendation to restrict consumption of vitamin K-rich foods [74, 75].

The last case report reported a suspected toxic irritative dermatitis (nonallergic) in an individual who applied a home-made mustard wrap in order to relieve the symptoms of bronchitis; authors believe the wrap may have contained some toxic compounds and concluded that the reaction was

		Comments	Skin test also revealed allergy to mustard, cauliflower, and broccoli. Reexposure to coleslaw via oral ingestion after 2 weeks triggered same symptoms	Occupational exposure	Bird fancier. Patient was also found to be allergic to avian antigens
		Causality ⁵	n Certainly	Likely	Possibly
		Outcome ⁴	Full resolution	Not mentioned	Not mentioned
	Outcome	Acute management	Subcutaneous epinephrine, discharged on oral antihistamines and steroids		Not mentioned
		Adverse events	Anaphylaxis: pain and swelling in mouth and throat with difficult breathing. Swelling of lip, tongue, soft palate. No diffuse urticaria, wheezing or hypotension	Patient 1: rhinoconjunctivitis Patient 2: rhinoconjunctivitis and asthma. Patients were tested and found to be allergic to Diplotaxis erucoides pollen	Rhinitis and asthma. Patient was tested and found to be allergic to turnip seeds
		Dose ³	Not specified	NA	NA
•	tervention	Duration ²	2 acute ingestions	y 2 months every year	Chronic dd (exact y not mentioned)
nt details Int	IJ	Route ¹	Oral	l Respirato	Topical ar respirator
		Plant/ substance	Cabbage (in coleslaw)	Diplotaxis erucoides (wal rocket) pollen	Turnip seeds
	ent details	Concurrent conditions/ treatments	In the past allergic rhinitis to numerous inhalant allergens: grasses, milds, ragweed, and dust mite. Immunotherapy to grass and dust allergens stopped a year prior to reported events	Not given	Ex-smoker, with a personal history of house dust mite allergic rhinitis and asthma, and a family history of atopy
ſ	Pati	Age	21	29 and 37	38
	Reference		Blaiss et al. 1987 [53]	Brito et al. 2001 [54]	Compes et al. 2007 [55]

TABLE 4: Case reports of adverse events associated with human exposure to cruciferous plants, their derivatives or their constituents.

Journal of Biomedicine and Biotechnology

					TA	BLE 4: Continued.					
Reference	Pat	tient details		In	tervention			Outcome			
	Age	Concurrent conditions/ treatments	Plant/ substance	Route ¹	Duration ²	Dose ³	Adverse events	Acute management	Outcome ⁴	Causality ⁵	Comments
Caldan 1981 [56]	28	Eczema and hay fever. In the past contact with other vegetables caused itching	Cabbage	Topical	Not reported (unclear whether chronic eczema is related to cabbage exposure)	NA	Contact urticaria. Patch test revealed allergy to "green leaves" (cabbage and Brusselss sprouts)	Not mentioned	Not mentioned	Possibly	Paper lacks many details regarding both urticaria and eczema
Chakrabarti et al. 2003 [57]	20	3-year history of severe eczema and recurrent blisters on palms	Broccoli	Topical	Chronic (exact duration not mentioned)	NA	Allergic contact dermatitis (worsening of eczema)	Subcutaneous epinephrine, discharged on oral antihistamines and steroids	Partial resolution: eczema improved upon avoidance of topical contact with broccoli and other vegetables	Possibly	Patient refused patch tests to other vegetables, but it is possible that she is allergic not only to broccoli
Dannaker et al. 1987 [58]	38	2 year history of hand dermatitis; 6 months history of dryness and scaling at angles of mouth (Unclear whether her hand dermatitis is related to mustard exposure)	Mustard in salad dressings	Topical	Chronic, fo the past 3 years	r NA	Acute episode of allergic contact dermatitis	Not mentioned	Not mentioned	Possibly	Occupational exposure

Journal of Biomedicine and Biotechnology

		Comments	Occupational exposure	No details of what other components the dish included
		Causality ⁵	Possibly	Possibly
		Outcome ⁴	Not mentioned	Full resolution
	Outcome	Acute management	Not mentioned	Required emergency room consult
		Adverse events	Episodes of asthma and rhinoconjunctivitis Allergy tests showed that skin prick tests were only mildly positive for oilseed rape flour, and the peak respiratory flow only slightly decreased during exposure	Oropharyngeal itching, facial and hand swelling, severe bronchospasm
BLE 4: Continued.		Dose ³	NA	Not specified
TA	tervention	Duration ²	Chronic Chronic (exact y duration not mentioned	Acute ingestion
	In	Route ¹	Respirato	Oral
		Plant/ substance	Oilseed rape flour	Cauliflower
	tient details	Concurrent conditions/ treatments	Not given	Allergies to some fruit and nuts, seasonal allergic rhinoconjunctivitis
	Pa	Age	48	70
	Reference		Di Giacomo et al. 1998 [59]	Hernandez et al. 2005 [60]

. C

	Comments	None	Patient was also found to be allergic to other foods
	Causality ⁵	Likely	Likely
	Outcome ⁴	Full resolution	Full resolution
Outcome	Acute management	Not mentioned	IV adrenaline, antihistamine, and corticosteroids
	Adverse events	Patient 1: episodes of pruritus, swelling of tongue, dysphonia, facial edema, and progressive upper respiratory difficulty. On one occasion, he experienced hypotension, and on another urticaria and palpebral edema. Patient 2: episodes of pruritus, swelling of lips and tongue, and edema. Patient 3: dysphonia, dysphonia	Anaphylaxis induced by exercise after ingestion of cabbage or mustard
	Dose ³	Not specified	AN
itervention	Duration ²	Acute ingestions	Acute ingestions
II	Route ¹	Oral	Oral
	Plant/ substance	Mustard sauce	Cabbage and mustard
ent details	Concurrent conditions/ treatments	Patient 1: history of IgE-dependent rhinitis since childhood; urticaria angioedema episodes related with shellfish. Patient 2: history of IgE-dependent asthma and rhinitis and urticaria to latex. Patient 3: history of IgE-dependent rhinitis	For the past 11 years: episodes of exercise-induced anaphylaxis after eating, once or twice a year
Pati	Age	43, 17, and 19	40
Reference		Jorro et al. 1995 [61]	Lingelbach et al. 2003 [62]

TABLE 4: Continued.

		Comments	Apart from ingestion, patient was also occupa- tionally exposed to rapeseed
		Causality ⁵	Possibly
		Outcome ⁴	Partial resolution: eczema improved upon avoidance of mustard and rapeseed
	Outcome	Acute management	Not mentioned
		Adverse events	Vesicular episodes (worsening of eczema). Positive skin prick test to crushed seeds of rapeseed
BLE 4: Continued.		Dose ³	Not specified
TAF	rvention	Duration ²	Acute ingestions
	Inte	Route ¹	Oral (however, allergy testing was topical)
		Plant/ substance	Mustard and rapeseed
	ient details	Concurrent conditions/ treatments	Vesicular hand eczema for the past 9 years, sometimes worsened for no clear reason, and also after every mustard ingestion
	Pat	Age	40
	Reference		Meding 1985 [63]

Ć ÷ Ľ

	Comments	Patient 1 had been using mustard oil for years and only lately did she start experiencing adverse event. Patient 2 did not report any exposure to mustard	Occupational exposure
	Causality ⁵	Possibly (patient 1), unlikely (patient 2)	Possibly
	Outcome ⁴	Improved with treatment, but authors do not mention what happens when not exposed	Full resolution
Outcome	Acute management	Not mentioned (it is mentioned was given, but no further details are provided)	Required emergency room consult
	Adverse events	Patient 1: itching and erythmatous papular lesions on forearms, arms, neck, forehead, ear lobules, and sides of face for the past 2: itching and erythmatous papules on forearms, forehead, cheeks, ear lobules, neck, and dorsum of feet for the past 1.5 years. Upon allergy testing, patient 1 found to be allergic to mustard, jowar flour, and wheat flour. Patient 2 found to be allergic to mustard khal and maize	Generalized urticaria, facial and oropharyngeal angioedema. Upon allergy testing (with cabbage or cauliflower): severe rhinoconjunctivitis and an early asthmatic reaction
	Dose ³	Ϋ́Z	Not specified
ervention	Duration ²	Chronic (exact duration not mentioned)	Acute ingestion
Int	Route ¹	Topical	Oral
	Plant/ substance	Patient 1: mustard; Patient 2: mustard khal	Cabbage
nt details	Concurrent conditions/ treatments	Patient 1 had suffered from dermatitis ever since she was 15, which worsened when handling cattle food. Patient 2: itching and erythematous papules for the past 1.5 years, for which she was treated	Allergic rhinoconjunctivitis to pollen. For past 7 years: episodes of ocular and nasal itching, sneezing, watery nose, tearing, dry cough, chest tightness, and dyspnea after inhaling cauliflower or cabbage vapors
Patić	Age	35 and 47	41
Reference		Pasricha et al. 1985 [64]	Quirce et al. 2005 [65]

TABLE 4: Continued.

		Comments	Occupational exposure. No testing done for food allergies in either patients	Occupational exposure	Some mustard allergy tests were positive but rubbing test was negative. Cross- reactivity with other Brassica
		Causality ⁵	Likely	Possibly	Possibly
		Outcome ⁴	Full resolution. In patient 2: recurrence when eating mustard/radish	Patient 1: full resolution after avoiding broccoli. Patient 2: no resolution even when avoids allergens	Not mentioned
	Outcome	Acute management	Patient 1: not mentioned; Patient 2: hospitalized, treated w/theophylline	Not mentioned	Topical and oral antihistamines
		Adverse events	Patient 1: face flushing, asthma attack. Patient 2: asthma attack	Patient 1: acute episodes of contact dermatitis after handling broccoli; Patient 2: acute episodes of contact dermatitis after handling broccoli or cauliflower	At 4 years of age-topical exposure caused eye lacrimation and cauterization, and dyspnea. Upon ingestion at age of 19: angioedema and bronchospasm
BLE 4: Continued.		Dose ³	Not specified	Υ N	Not specified
TAI	ervention	Duration ²	Acute ingestions	Chronic (exact duration not mentioned)	varies (acute to chronic)
	Int	Route ¹	Oral	Topical	Oral and topical
		Plant/ substance	Patient 1: mustard and horseradish. Patient 2: mustard, radish, turnip	Broccoli or cauliflower	Mustard (sauce and oil)
	ent details	Concurrent conditions/ treatments	Patient 1: asthma triggered by isocyanate paint. Patient 2: ex-smoker, asthma triggered by isocyanate paint	Patient 1: for past 7 years: pruritus, erythema, vesicles, fissures, and peeling in both hands, as well as facial angioedema, within 6–8 h after handling broccoli. Patient 2: asthma; in past 4 years, papules and vesicles in both hands and eyelids 24–36 hours after exposure to cauliflower or broccoli	Not given
	Pati	Age	42 and 34	36 and 54	19
	Reference		Rosenberg and Gervais 1986 [66]	Sanchez- Guerrero and Escudero 1998 [67]	Schulze and Wollina 2003 [68]

20

		Comments	Occupational exposure; allergy testing not performed	None
		Causality ⁵	Possibly	Possibly
		Outcome ⁴	Not mentioned	Full resolution
	Outcome	Acute management	Not mentioned	Not mentioned
		Adverse events	Aggravation of cough and chest pain	Patient 1–3: urticaria, facial edema, rhinocon- junctivitis/rhinitis. Patient 4: facial edema, bronchospasm, rhinitis. Patient 5: urticaria, facial edema, and bronchospasm. All found to be allergid to mustard
ste 4: conunuea.		Dose ³	NA	Not specified
IAB	rvention	Duration ²	Chronic	Acute (ingestion) to seasonal (pollen inhalation)
	Inte	Route ¹	Respiratory	Oral and respiratory
		Plant/ substance	Oilseed rape dust	Mustard sauce or mustard pollen
	ant details	Concurrent conditions/ treatments	Smoker; cough and chest pain	Patient 1: house dust allergy. Patient 2: seasonal rhinitis. Patient 3: peach allergy; seasonal rhinitis; familial atopy. Patient 4: seasonal rhinoconjuctivitis and bronchial asthma, family pollen allergy.
	Patié	Age	43	34, 31, 25, 52, and 33
	Reference		Suh et al. 1998 [69]	Valero et al. 1995 [70]

Journal of Biomedicine and Biotechnology

TABLE 4: Continued.

		Comments	Occupational exposure. Aggravation of eczema also occurs after contact with onions, tulip bulbs, rubber gloves, and pesticides, but to a lesser extent	None	None	Vitamin K consumption was 60 times more than the norm $(360 \mu g)$
		Causality ⁵	Possibly	Possibly	Certainly	Likely
		Outcome ⁴	Not mentioned	Full resolution	Full resolution followed by recurrence on re-exposure	Full resolution
	Outcome	Acute management	Not mentioned	Not mentioned	Corticosteroids and antihistamines	Referred to hospital, where she was treated with nitroglycerin and heparin
		Adverse events	Aggravation of eczema. Upon allergy testing was found allergic to both cauliflower and other Brassica: Brusselss sprouts and red cabbage	Acute episodes of urticaria and angioneurotic edema of face and neck. Upon allergy testing was found to be allergic to mustard	Cutaneous lesions similar to pityriasis rosea (appeared after first application)	Diet-induced warfarin resistance which led to substernal chest pain and myocardial infarction
		Dose ³	VV	Not specified	NA	Not specified. Authors do mention, however, that vitamin K intake was 6000 µg per day
	tervention	Duration ²	Chronic (exact duration not mentioned)	Acute ingestions	Several acute applications	5 weeks
,	п	Route ¹	Topical	Oral	Topical	Oral
		Plant/ substance	Cauliflower	g Mustard (in i mustard sauce or mayonnaise)	Mustard oil	Lettuce and greens (e.g., turnip, mustard greens, broccoli)
:	tient details	Concurrent conditions/ treatments	Eczema in past 10 years	As a child, severe atopic dermatitis, rhinitis and swelling of throat in reaction to fish or egg. Currently, rhinitis when exposed to cats or dogs	Healthy	Prosthetic aortic valve, history of myocardial infarctions with prolonged prothrombin time. Treated with warfarin and dipyridamole
1	Pai	Age	43	25	25	4 35
	Reference		van Ketel et al. 1975 [71]	Widstrom and Johansson 1986 [72]	Zawar 2005 [73]	Walker 198 [74]

TABLE 4: Continued.

22

6			-	TAB	3LE 4: Continued.					
ge P	atient details Concurrent conditions/	Plant/	ln Route ¹	ıtervention Duration ²	Dose ³	Adverse events	Outcome Acute	Outcome ⁴	Causality ⁵	Comments
	treatments Patient 1: pulmonary	substatice					management			
Vot çiven	embolism; Patient 2: cardiovascular disease (not specified which). Treated with heparin and Coumadin	Broccoli	Oral	Chronic (exact duration not mentioned)	Up to 450 g/day	Warfarin resistance	Coumadin anticoagula- tion in patient 2	Full resolution	Likely	None
	(=warfarin)									Authors
15	Bronchitis	Home-made mustard wrap, containing ground mustard seeds and water	Topical	Acute application (20 minutes)	Not specified	Toxic irritative dermatitis	Topical corticosteroids	Full resolution	Possibly	labeled case as toxic irritative dermatitis and NOT an allergic reaction and warn against use of
										home-made medications
ure.										

²Duration of exposure. ³NA, not applicable, in cases where dose cannot be quantified. ⁴Degree of resolution in cases here adverse events were reported; NA, not applicable. ⁵The degree of association between the intervention and the adverse event, as rated by reviewers; NA, not applicable.

not an allergic one. Hence, they advised against the use of home-made wraps. The patient was treated with corticosteroids with full resolution. This case was very brief and lacked a great deal of information regarding previous medical history and whether other medications had been used to treat the patient's bronchitis. Moreover, no test was conducted to verify the cause of the adverse event [76].

3.5. Excluded Papers. Of the 317 papers excluded, 280 did not fulfill the criteria for our systematic review (i.e., they did not evaluate adverse effects or lack thereof, intervention or substance exposure did not include cruciferous plants, or did not present original data). Another 37 articles were excluded since they dealt with toxic oil syndrome. In order to promote transparency in our decision-making process, we shall explain the rationale behind excluding this latter group of papers.

Toxic oil syndrome is the name given to a disease outbreak in Spain in 1981. Its first appearance was as an acute lung disease, which was followed by a range of other chronic symptoms affecting the lungs, the liver, the kidneys, the skin, the joints, the central nervous system, and the immune system [77-83]. The cause for the disease was traced to the consumption of cheap, refined, rapeseed oil that had been intended for industrial use rather than for human consumption. It was sold as "olive oil" and was therefore used for cooking. According to the World Health Organization, toxic compounds derived during the refinement process, used to remove the aniline and to denature oils intended for industrial use, were responsible for causing the disease. Hence, toxic oil syndrome has come to be considered a chemical incident, rather than a side effect associated with consumption of rapeseed oil, since consumption of non-industrial rapeseed oil does not cause the disease. In fact, the specific refinement process of the oil that caused the disease was so unique that experimental studies performed in a variety of laboratory animals have hitherto failed to reproduce the symptoms of human toxic oil syndrome [84].

4. Discussion

Research conducted in the last decade supports the notion that some cruciferous plants and their derivatives may serve to prevent or attenuate several medical conditions. To further this field of investigation, we conducted this systematic paper to determine the safety parameters surrounding the use of cruciferous plants. In this regard, our review identified adverse events in 1335 individuals, out of a total of 101,198 individuals who were included in all studies. Of these, 1292 adverse events were ranked as only possibly or unlikely to be caused by cruciferous plants. Only 43 were determined to have been certainly or likely caused by exposure to cruciferous plants. Adverse events, which could certainly or likely be attributed to members of the Brassica genus, included allergic reactions (including anaphylaxis), changes in metabolism of acetaminophen, phenacetin and warfarin, and warfarin resistance. Most adverse events reported in our paper were allergic or hypersensitivity reactions; however, their significance for the general healthy population is hard to infer, due to

confounding medical histories or low causality ratings. Adverse events from both allergy trials and case reports were mostly in individuals who suffered from atopy, allergies to cruciferous plants and/or other allergens (for instance, grasses, ragweed, dust mite, pollen, dogs, cats and different fruit and nuts), asthma, or a variety of skin conditions such as eczema, pruritus, erythema, dermatitis, dryness and scaling, or blisters. A possible conclusion from these case reports, however, is that occupational exposure to Brassica plants may predispose to develop an allergy to these plants. With regards to observational studies which investigated allergies to cruciferous plants or their derivatives, all concluded that the cases could not be attributed to the suspected culprits.

A second category of adverse events included changes in metabolism of phenacetin, acetaminophen, and warfarin. While phenacetin has been banned for use by the FDA, acetaminophen and warfarin continue to be widely used. As such, it was important to understand if these changes in metabolism pose any risk to those who use these drugs. As for acetaminophen, the authors explain that the decreased levels of the cysteine conjugate of acetaminophen found in subjects' urine actually indicate decreased toxicity; thus, cruciferous plants may not pose any risk to individuals who consume acetaminophen. With regards to warfarin, however, the finding that cruciferous vegetables alter its metabolism might bear greater clinical relevance. It has been suggested that foods high in vitamin K, such as cruciferous vegetables, may interact with warfarin and its anticoagulant activity. In all case reports describing warfarin resistance, the patients had consumed very large amounts of cruciferous vegetables (up to 450 g/day or the equivalent of 6000 μ g vitamin K/day), amounts not warranted for consumption by the general population. However, it is possible that lower doses would have some effects on anticoagulation as well. Similarly large amounts of cruciferous vegetables (400 g/day) were given to subjects in the SCED trial which examined changes in warfarin resistance. As many foods interact with warfarin, among which are mango, avocado, fish oil, soy milk, and foods high in vitamin K [84], it is imperative for physicians to discuss the consumption of such foods (including cruciferous vegetables) with their patients when starting treatment with anticoagulants.

Reports of other types of adverse events identified in our systematic review, including cancer, chronic atrophic gastritis, infantile colic, and toxicity-related events, could only be ranked as possibly or unlikely to be caused by exposure to cruciferous plants, as there was not enough evidence to establish a stronger causal relationship between the suspected culprits and the adverse events. Further research employing better methodology should be undertaken in order to establish whether positive association exists between consumption of Brassica vegetables and any of the above-mentioned adverse events.

Finally, a number of studies did not find any adverse events associated with consumption of cruciferous plants or their derivatives in subjects with a variety of health conditions, including history of myocardial infarction, nutritional deficits, or recurrent respiratory papillomatosis, or in healthy individuals. In summary, to date, adverse events presumably related to cruciferous plants have been reported in 1335 individuals, identified through 50 studies which included a total of 101,198 individuals. However, the number of those which were certainly/likely caused by Brassica plants (including allergic reactions, changes in drug metabolism, and warfarin resistance) was much lower: only 43 of all reported adverse events received such high causality rating. For comparison, the literature shows that the pooled prevalence of adverse drug reactions ranges from 4.2 to 6% [85]. This frequency, if applied to cruciferous plants, would translate into 4250– 6072 reports of adverse events out of the included 101,198 individuals, a value much higher than what was found in our current study.

When analysing our findings, one must bear in mind that the total number of individuals included in all studies comprises both those who were definitely exposed to Brassica plants and those who were presumptively exposed as the degree of exposure could not be ascertained in all cases. This might cause the frequency of occurrence of adverse events to appear lower than the actual value. On the other hand, our study also included case reports of individuals, all of whom were exposed, and reported adverse events. In this regard, the frequency of adverse events tends to be an overestimation, since individuals who are exposed, but do not experience an adverse event, are likely not to report. Hence, the inclusion of these case report, in which essentially all individuals exposed report adverse events, may serve to balance this underestimation. We advise the reader to treat the broad pooled frequency presented in the previous paragraph (1335/101,198) as merely gross estimations of the actual frequency, as conducting a meta-analysis per se was deemed inappropriate given the heterogeneity of study parameters (subjects, interventions, clinical outcomes, and study designs).

We conclude that cruciferous plants are generally safe for human consumption and use. However, individuals with known allergies/hypersensitivities to a certain member of the *Brassica* genus, or those taking warfarin, should consult with their physician before consuming such vegetables. In the future, if cruciferous derivatives are to be investigated as potential therapeutic agents, we recommend that adverse events be monitored. As our findings reflect positively on the safety of Brassica plants in humans and evidence from the current experimental literature of their benefit in certain disease states, we would encourage further exploration of their potential use in the clinical setting. The use of *brassica* or similar food products may have the potential to provide a safer alternative for the treatment of disease in comparison to current pharmaceutical interventions.

Conflict of Interests

The authors declare no conflict of interests.

Author Contribution

JYY initiated the research question; OS, EGC, DA, SS, SV, and JYY designed research; OS, EGC, DA, and SS conducted

research; DA and SS provided essential materials; OS and EGC, analyzed data; OS wrote the paper; OS, EGC and JYY had primary responsibility for final content. SV provided guidance on study design and interpretation, including paper revisions. All authors read and approved the final paper.

Acknowledgments

This work was supported by grants from the Alva Foundation, the Heart and Stroke Foundation of Alberta, NWT and Nunavut, and NeuroDevNet, a Canadian National Centres of Excellence Grant. The authors would like to thank Ms. Miriam Schiffgen for translation of papers from German to English.

References

- E. Goldman, "Practical strategies for implementing integrative medicine in a primary care setting," *Journal of Medical Practice Management*, vol. 24, no. 2, pp. 97–101, 2008.
- [2] S. Prakash and K. Hinata, *Taxonomy, Cytogenetics, and Origin of Crop Brassicas: A Review*, Swedish Natural Science Research Council, Stockholm, Sweden, 1980.
- [3] S. Y. Kim, S. Yoon, S. M. Kwon, K. S. Park, and Y. C. Lee-Kim, "Kale Juice improves coronary artery disease risk factors in hypercholesterolemic men," *Biomedical and Environmental Sciences*, vol. 21, no. 2, pp. 91–97, 2008.
- [4] A. Yanaka, J. W. Fahey, A. Fukumoto et al., "Dietary sulforaphane-rich broccoli sprouts reduce colonization and attenuate gastritis in Helicobacter pylori-infected mice and humans," *Cancer Prevention Research*, vol. 2, no. 4, pp. 353–360, 2009.
- [5] P. Brennan, C. C. Hsu, N. Moullan et al., "Effect of cruciferous vegetables on lung cancer in patients stratified by genetic status: a mendelian randomisation approach," *The Lancet*, vol. 366, no. 9496, pp. 1558–1560, 2005.
- [6] P. H. Chyou, A. M. Y. Nomura, J. H. Hankin, and G. M. Stemmermann, "A case-cohort study of diet and stomach cancer," *Cancer Research*, vol. 50, no. 23, pp. 7501–7504, 1990.
- [7] M. Hara, T. Hanaoka, M. Kobayashi et al., "Cruciferous vegetables, mushrooms, and gastrointestinal cancer risks in a multicenter, hospital-based case-control study in Japan," *Nutrition and Cancer*, vol. 46, no. 2, pp. 138–147, 2003.
- [8] L. I. Wang, E. L. Giovannucci, D. Hunter, D. Neuberg, L. Su, and D. C. Christiani, "Dietary intake of Cruciferous vegetables, Glutathione S-transferase (GST) polymorphisms and lung cancer risk in a Caucasian population," *Cancer Causes* and Control, vol. 15, no. 10, pp. 977–985, 2004.
- [9] J. M. Pogoda, S. Preston-Martin, G. Howe et al., "An international case-control study of maternal diet during pregnancy and childhood brain tumor risk: a histology-specific analysis by food group," *Annals of Epidemiology*, vol. 19, no. 3, pp. 148– 160, 2009.
- [10] M. H. Noyan-Ashraf, L. Wu, R. Wang, and B. H. J. Juurlink, "Dietary approaches to positively influence fetal determinants of adult health," *The FASEB Journal*, vol. 20, no. 2, pp. 371– 373, 2006.
- [11] J. Y. Yager, C. M. Jahraus, and B. H. J. Juurlink, "Dietary phase 2 enzyme inducers are neuroprotective to the immature brain following hypoxia-ischemia," *Ped Research*, vol. 53, no. 4, p. 24A, 2003.

- [12] B. H. J. Juurlink, "Therapeutic potential of dietary phase 2 enzyme inducers in ameliorating diseases that have an underlying inflammatory component," *Canadian Journal of Physiology and Pharmacology*, vol. 79, no. 3, pp. 266–282, 2001.
- [13] J. W. Fahey and P. Talalay, "Antioxidant functions of sulforaphane: a potent inducer of phase II detoxication enzymes," *Food and Chemical Toxicology*, vol. 37, no. 9-10, pp. 973–979, 1999.
- [14] J. W. Fahey, Y. Zhang, and P. Talalay, "Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 19, pp. 10367–10372, 1997.
- [15] D. M. Minich and J. S. Bland, "A review of the clinical efficacy and safety of cruciferous vegetable phytochemicals," *Nutrition Reviews*, vol. 65, no. 6, pp. 259–267, 2007.
- [16] S. Watanabe, X. G. Zhuo, and M. Kimira, "Food safety and epidemiology: new database of functional food factors," *Bio-Factors*, vol. 22, no. 1-4, pp. 213–219, 2004.
- [17] A. Liberati, D. G. Altman, J. Tetzlaff et al., "The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration," *Annals of Internal Medicine*, vol. 151, no. 4, pp. W-65–W-94, 2009.
- [18] T. A. Shapiro, J. W. Fahey, A. T. Dinkova-Kostova et al., "Safety, tolerance, and metabolism of broccoli sprout glucosinolates and isothiocyanates: a clinical phase I study," *Nutrition and Cancer*, vol. 55, no. 1, pp. 53–62, 2006.
- [19] J. Figueroa, C. Blanco, A. G. Dumpiérrez et al., "Mustard allergy confirmed by double-blind placebo-controlled food challenges: clinical features and cross-reactivity with mugwort pollen and plant-derived foods," *Allergy*, vol. 60, no. 1, pp. 48– 55, 2005.
- [20] E. J. Pantuck, C. B. Pantuck, and W. A. Garland, "Stimulatory effect of brussels sprouts and cabbage on human drug metabolism," *Clinical Pharmacology and Therapeutics*, vol. 25, no. 1, pp. 88–95, 1979.
- [21] E. J. Pantuck, C. B. Pantuck, and K. E. Anderson, "Effect of brussels sprouts and cabbage on drug conjugation," *Clinical Pharmacology and Therapeutics*, vol. 35, no. 2, pp. 161–169, 1984.
- [22] L. Ovesen, S. Lyduch, and M. L. Idorn, "The effect of a diet rich in brussels sprouts on warfarin pharmacokinetics," *European Journal of Clinical Pharmacology*, vol. 34, no. 5, pp. 521–523, 1988.
- [23] V. Vovolis, G. Poulios, and N. Koutsostathis, "IgE-mediated allergy to raw cabbage but not to cooked," *Allergy*, vol. 64, no. 6, pp. 964–965, 2009.
- [24] C. A. Rosen, G. E. Woodson, J. W. Thompson, A. P. Hengesteg, and H. L. Bradlow, "Preliminary results of the use of indole-3carbinol for recurrent respiratory papillomatosis," *Otolaryngology—Head and Neck Surgery*, vol. 118, no. 6, pp. 810–815, 1998.
- [25] T. W. Kensler, J. G. Chen, P. A. Egner et al., "Effects of glucosinolate-rich broccoli sprouts on urinary levels of aflatoxin-DNA adducts and phenanthrene tetraols in a randomized clinical trial in He Zuo township, Qidong, People's Republic of China," *Cancer Epidemiology Biomarkers and Prevention*, vol. 14, no. 11, pp. 2605–2613, 2005.
- [26] R. B. Singh, M. A. Niaz, J. P. Sharma, R. Kumar, V. Rastogi, and M. Moshiri, "Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute

myocardial infarction: the Indian experiment of infarct survival—4," *Cardiovascular Drugs and Therapy*, vol. 11, no. 3, pp. 485–491, 1997.

- [27] S. Jood, M. Gupta, S. K. Yadav, and N. Khetarpaul, "Effect of supplementation on haemoglobin and serum retinol levels and nutritional status of school children of Northern India," *Nutrition and Health*, vol. 15, no. 2, pp. 97–111, 2001.
- [28] C. A. Rosen and P. C. Bryson, "Indole-3-carbinol for recurrent respiratory papillomatosis: long-term results," *Journal of Voice*, vol. 18, no. 2, pp. 248–253, 2004.
- [29] M. A. Riedl, A. Saxon, and D. Diaz-Sanchez, "Oral sulforaphane increases phase II antioxidant enzymes in the human upper airway," *Clinical Immunology*, vol. 130, no. 3, pp. 244– 251, 2009.
- [30] A. T. Dinkova-Kostova, J. W. Fahey, K. L. Wade et al., "Induction of the phase 2 response in mouse and human skin by sulforaphane-containing broccoli sprout extracts," *Cancer Epidemiology Biomarkers and Prevention*, vol. 16, no. 4, pp. 847– 851, 2007.
- [31] P. J. Fell, S. Soulsby, M. M. Blight, and J. Brostoff, "Oilseed rape—a new allergen?" *Clinical and Experimental Allergy*, vol. 22, no. 4, pp. 501–505, 1992.
- [32] W. Hemmer, M. Focke, F. Wantke, S. Jäger, M. Götz, and R. Jarisch, "Oilseed rape pollen is a potentially relevant allergen," *Clinical and Experimental Allergy*, vol. 27, no. 2, pp. 156–161, 1997.
- [33] D. Parratt, W. H. Macfarlane Smith, G. Thomson, L. A. Cameron, and R. D. Butcher, "Evidence that oilseed rape (Brassica napus ssp. oleifera) causes respiratory illness in rural dwellers," *Scottish Medical Journal*, vol. 40, no. 3, pp. 74–76, 1995.
- [34] A. Soutar, C. Harker, A. Seaton, and G. Packe, "Oilseed rape and bronchial reactivity," *Occupational and Environmental Medicine*, vol. 52, no. 9, pp. 575–580, 1995.
- [35] A. Soutar, C. Harker, A. Seaton, M. Brooke, and I. Marr, "Oilseed rape and seasonal symptoms: epidemiological and environmental studies," *Thorax*, vol. 49, no. 4, pp. 352–356, 1994.
- [36] A. Lerbæk, S. C. Rastogi, and T. Menné, "Allergic contact dermatitis from allyl isothiocyanate in a Danish cohort of 259 selected patients," *Contact Dermatitis*, vol. 51, no. 2, pp. 79–83, 2004.
- [37] F. Sakauchi, M. M. H. Khan, M. Mori et al., "Dietary habits and risk of ovarian cancer death in a large-scale cohort study (JACC study) in Japan," *Nutrition and Cancer*, vol. 57, no. 2, pp. 138–145, 2007.
- [38] L. Y. Sue, C. Schairer, X. Ma et al., "Energy intake and risk of postmenopausal breast cancer: an expanded analysis in the prostate, lung, colorectal, and ovarian cancer screening trial (PLCO) cohort," *Cancer Epidemiology Biomarkers and Prevention*, vol. 18, no. 11, pp. 2842–2850, 2009.
- [39] C. P. J. Caygill, A. Charlett, and M. J. Hill, "Relationship between the intake of high-fibre foods and energy and the risk of cancer of the large bowel and breast," *European Journal of Cancer Prevention*, vol. 7, no. 2, pp. S11–S17, 1998.
- [40] A. Latif, H. J. McBurney, S. A. Roberts et al., "Breast cancer susceptibility variants alter risk in familial ovarian cancer," *Familial Cancer*, vol. 9, no. 4, pp. 503–506, 2010.
- [41] V. Bissonauth, B. Shatenstein, E. Fafard et al., "Risk of breast cancer among French-Canadian women, noncarriers of more frequent BRCA1/2 mutations and consumption of total energy, coffee, and alcohol," *The Breast Journal*, vol. 15, supplement 1, pp. S63–S71, 2009.

- [42] A. Memon, A. Varghese, and A. Suresh, "Benign thyroid disease and dietary factors in thyroid cancer: a case-control study in Kuwait," *British Journal of Cancer*, vol. 86, no. 11, pp. 1745– 1750, 2002.
- [43] M. R. Galanti, L. Hansson, R. Bergström et al., "Diet and the risk of papillary and follicular thyroid carcinoma: a population-based case-control study in Sweden and Norway," *Cancer Causes and Control*, vol. 8, no. 2, pp. 205–214, 1997.
- [44] J. J. Michnovicz and H. L. Bradlow, "Altered estrogen metabolism and excretion in humans following consumption of indole-3-carbinol," *Nutrition and Cancer*, vol. 16, no. 1, pp. 59–66, 1991.
- [45] D. S. Michaud, P. Pietinen, P. R. Taylor, M. Virtanen, J. Virtamo, and D. Albanes, "Intakes of fruits and vegetables, carotenoids and vitamins A, E, C in relation to the risk of bladder cancer in the ATBC cohort study," *British Journal of Cancer*, vol. 87, no. 9, pp. 960–965, 2002.
- [46] K. Sato, N. Kawakami, T. Ohtsu et al., "Broccoli consumption and chronic atrophic gastritis among Japanese males: an epidemiological investigation," *Acta Medica Okayama*, vol. 58, no. 3, pp. 127–133, 2004.
- [47] K. D. Lust, J. E. Brown, and W. Thomas, "Maternal intake of cruciterous vegetables and other foods and colic symptoms in exclusively breast-fed infants," *Journal of the American Dietetic Association*, vol. 96, no. 1, pp. 46–48, 1996.
- [48] E. Ron, R. A. Kleinerman, and J. D. Boice Jr., "A populationbased case-control study of thyroid cancer," *Journal of the National Cancer Institute*, vol. 79, no. 1, pp. 1–12, 1987.
- [49] L. N. Kolonel, J. H. Hankin, L. R. Wilkens, F. H. Fukunaga, and M. W. Hinds, "An epidemiologic study of thyroid cancer in Hawaii," *Cancer Causes and Control*, vol. 1, no. 3, pp. 223– 234, 1990.
- [50] S. Franceschi, A. Fassina, R. Talamini et al., "Risk factors for thyroid cancer in Northern Italy," *International Journal of Epidemiology*, vol. 18, no. 3, pp. 578–584, 1989.
- [51] G. Wingren, T. Hatschek, and O. Axelson, "Determinants of papillary cancer of the thyroid," *American Journal of Epidemi*ology, vol. 138, no. 7, pp. 482–491, 1993.
- [52] M. A. Wessel, J. C. Cobb, E. B. Jackson, G. S. Harris Jr., and A. C. Detwiler, "Paroxysmal fussing in infancy, sometimes called colic," *Pediatrics*, vol. 14, no. 5, pp. 421–435, 1954.
- [53] M. S. Blaiss, M. L. McCants, and S. B. Lehrer, "Anaphylaxis to cabbage: detection of allergens," *Annals of Allergy*, vol. 58, no. 4, pp. 248–250, 1987.
- [54] F. F. Brito, P. Mur, B. Bartolomé et al., "Rhinoconjunctivitis and occupational asthma caused by Diplotaxis erucoides (wall rocket)," *Journal of Allergy and Clinical Immunology*, vol. 108, no. 1, pp. 125–127, 2001.
- [55] E. Compés, O. Palomares, M. Fernández-Nieto, C. Escudero, and J. Cuesta-Herranz, "Allergy to turnip seeds in a bird fancier," *Allergy*, vol. 62, no. 12, pp. 1472–1473, 2007.
- [56] C. D. Caldan, "Contact urticaria from cabbage (brassica)," Contact Dermatitis, vol. 7, no. 5, p. 279, 1981.
- [57] A. Chakrabarti, L. Prais, and I. S. Foulds, "Allergic contact dermatitis to broccoli," *British Journal of Dermatology*, vol. 148, no. 1, pp. 172–173, 2003.
- [58] C. J. Dannaker and I. R. White, "Cutaneous allergy to mustard in salad maker," *Contact Dermatitis*, vol. 16, no. 4, pp. 212–214, 1987.
- [59] G. R. Di Giacomo, P. Boschetto, P. Maestrelli, and G. Moro, "Asthma and rhino-conjunctivitis form exposure to rape flour: a clinical case report," *Medicina del Lavoro*, vol. 89, no. 3, pp. 226–231, 1998 (Italian).

- [60] E. Hernández, S. Quirce, M. Villalba, J. Cuesta, and J. Sastre, "Anaphylaxis caused by cauliflower," *Journal of Investigational Allergology and Clinical Immunology*, vol. 15, no. 2, pp. 158–159, 2005.
- [61] G. Jorro, C. Morales, J. V. Braso, and A. Pelaez, "Mustard allergy: three cases of systemic reaction to ingestion of mustard sauce," *Journal of Investigational Allergology and Clinical Immunology*, vol. 5, no. 1, pp. 54–56, 1995.
- [62] A. Lingelbach, J. Rakoski, and J. Ring, "Exercise-induced anaphylaxis to cabbage and mustard," *Allergy and Clinical Immunology International*, vol. 15, no. 4, pp. 181–183, 2003.
- [63] B. Meding, "Immediate hypersensitivity to mustard and rape," *Contact Dermatitis*, vol. 13, no. 2, pp. 121–122, 1985.
- [64] J. S. Pasricha, R. Gupta, and S. K. Gupta, "Contact hypersensitivity to mustard khal and mustard oil," *Indian Journal of Dermatology, Venereology and Leprology*, vol. 51, no. 2, pp. 108–110, 1985.
- [65] S. Quirce, M. F. Madero, M. Fernández-Nieto, A. Jiménez, and J. Sastre, "Occupational asthma due to the inhalation of cauliflower and cabbage vapors," *Allergy*, vol. 60, no. 7, pp. 969–970, 2005.
- [66] N. Rosenberg and P. Gervais, "Occupational asthma and food allergy: an association of underestimated frequency," *Presse Medicale*, vol. 15, no. 34, pp. 1712–1714, 1986 (French).
- [67] I. M. Sánchez-Guerrero and A. I. Escudero, "Occupational contact to broccoli," *Allergy*, vol. 53, no. 6, pp. 621–621, 1998.
- [68] I. S. Schulze and U. Wollina, "Mustard allergy," Kosmetische Medizin, vol. 24, no. 2, pp. 63–65, 2003 (German).
- [69] C. H. Suh, H. S. Park, D. H. Nahm, and H. Y. Kim, "Oilseed rape allergy presented as occupational asthma in the grain industry," *Clinical and Experimental Allergy*, vol. 28, no. 9, pp. 1159–1163, 1998.
- [70] A. L. Valero, P. Amat, M. Bescos, M. Lluch, E. Serra, and A. Malet, "Mustard seed allergy: report of five cases," *Revista Espanola de Alergologia e Inmunologia Clinica*, vol. 10, no. 4, pp. 193–198, 1995 (Spanish).
- [71] W. G. van Ketel, "A cauliflower allergy," *Contact Dermatitis*, vol. 1, no. 5, pp. 324–325, 1975.
- [72] L. Widström and S. G. O. Johansson, "IgE-mediated anaphylaxis to mustard," *Acta Dermato-Venereologica*, vol. 66, no. 1, pp. 70–71, 1986.
- [73] V. Zawar, "Pityriasis rosea-like eruptions due to mustard oil application," *Indian Journal of Dermatology, Venereology and Leprology*, vol. 71, no. 4, pp. 282–284, 2005.
- [74] F. B. Walker 4th, "Myocardial infarction after diet-induced warfarin resistance," *Archives of Internal Medicine*, vol. 144, no. 10, pp. 2089–2090, 1984.
- [75] S. J. Kempin, "Warfarin resistance caused by broccoli," *The New England Journal of Medicine*, vol. 308, no. 20, pp. 1229–1230, 1983.
- [76] J. Geier, "Mustard wrap-dermatitis," *Dermatosen in Beruf und Umwelt*, vol. 39, no. 1, pp. 17–18, 1991 (German).
- [77] V. Gutierrez-Millet, J. Navas-Palacios, J. Gomez-Reino, and J. L. Fernandez-Epifanio, "Renal involvement in toxic oil syndrome," *The Lancet*, vol. 1, no. 8281, p. 1120, 1982.
- [78] A. Alonso-Ruiz, A. C. Zea-Mendoza, and J. M. Salazar-Vallinas, "Toxic oil syndrome: a syndrome with features overlapping those of various forms of scleroderma," *Seminars in Arthritis and Rheumatism*, vol. 15, no. 3, pp. 200–212, 1986.
- [79] R. G. Phelps and R. Fleischmajer, "Clinical, pathologic, and immunopathologic manifestations of the toxic oil syndrome. Analysis of fourteen cases," *Journal of the American Academy of Dermatology*, vol. 18, no. 2, pp. 313–324, 1988.

- [80] M. Rodriguez, E. Noguera, and R. S. Del Villar, "Toxic synovitis from denatured rapeseed oil," *Arthritis and Rheumatism*, vol. 25, no. 12, pp. 1477–1480, 1982.
- [81] J. R. Ricoy, A. Cabello, J. Rodriguez, and I. Tellez, "Neuropathological studies on the toxic syndrome related to adulterated rapeseed oil in Spain," *Brain*, vol. 106, no. 4, pp. 817–835, 1983.
- [82] R. Velicia, C. Sanz, and F. Martinez-Barredo, "Hepatic disease in the Spanish toxic oil syndrome. A thirty months follow-up," *Journal of Hepatology*, vol. 3, no. 1, pp. 59–65, 1986.
- [83] World Health Organization, *Toxic Oil Syndrome: Ten Years of Progress*, WHO Regional Office for Europe, Copenhagen, Denmark, 2004.
- [84] A. M. Holbrook, J. A. Pereira, R. Labiris et al., "Systematic overview of warfarin and its drug and food interactions," *Archives of Internal Medicine*, vol. 165, no. 10, pp. 1095–1106, 2005.
- [85] N. Muehlberger, S. Schneeweiss, and J. Hasford, "Adverse drug reaction monitoring cost and benefit considerations—part I: frequency of adverse drug reactions causing hospital admissions," *Pharmacoepidemiology and Drug Safety*, vol. 6, no. 3, pp. S71–S77, 1997.



Journal of Tropical Medicine

Journal of Toxins

KU





The Scientific World Journal

 $(\mathbf{0})$

Hindawi

Submit your manuscripts at http://www.hindawi.com



Autoimmune Diseases





Anesthesiology Research and Practice





Advances in Pharmacological

Emergency Medicine International

BioMed

Research International



Pain Research and Treatment



Journal of Pharmaceutics





International Journal of Medicinal Chemistry



Journal of Addiction





Journal of Drug Delivery

