

## Research Article

# Immunoinformatic Docking Approach for the Analysis of KIR3DL1/HLA-B Interaction

### Alba Grifoni,<sup>1</sup> Carla Montesano,<sup>1</sup> Atanas Patronov,<sup>2</sup> Vittorio Colizzi,<sup>1</sup> and Massimo Amicosante<sup>2,3</sup>

<sup>1</sup> Department of Biology, University of Rome "Tor Vergata", Via della Ricerca Scientifica 1, 00133 Rome, Italy

<sup>2</sup> ProxAgen Ltd., blv. Sitnyakovo 35, 1505 Sofia, Bulgaria

<sup>3</sup> Department of Biomedicine and Prevention, University of Rome "Tor Vergata", Via Montpellier 1, 00133 Rome, Italy

Correspondence should be addressed to Massimo Amicosante; massimo.amicosante@uniroma2.it

Received 30 April 2013; Accepted 12 July 2013

Academic Editor: Rita Casadio

Copyright © 2013 Alba Grifoni et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

KIR3DL1 is among the most interesting receptors studied, within the killer immunoglobulin receptor (KIR) family. Human leukocyte antigen (HLA) class I Bw4 epitope inhibits strongly Natural Killer (NK) cell's activity through interaction with KIR3DL1 receptor, while Bw6 generally does not. This interaction has been indicated to play an important role in the immune control of different viral infectious diseases. However, the structural interaction between the KIR3DL1 receptor and different HLA-B alleles has been scarcely studied. To understand the complexity of KIR3DL1-HLA-B interaction, HLA-B alleles carrying Bw4/Bw6 epitope and KIR3DL1\*001 allele in presence of different peptides has been evaluated by using a structural immunoinformatic approach. Different energy minimization force fields (ff) have been tested and NOVA ff enables the successful prediction of ligand-receptor interaction. HLA-B alleles carrying Bw4 epitope present the highest capability of interaction with KIR3DL1\*001 compared to the HLA-B alleles presenting Bw6. The presence of the epitope Bw4 determines a conformational change which leads to a stronger interaction between nonpolymorphic arginine at position 79 of HLA-B and KIR3DL1\*001 136–142 loop. The data shed new light on the modalities of KIR3DL1 interaction with HLA-B alleles essential for the modulation of NK immune-mediated response.

#### 1. Background

Killer immunoglobulin receptor (KIR)3DL1 is among the most interesting receptors studied within the KIR family. It interacts with polymorphisms located on human leukocyte antigen (HLA)-A and HLA-B molecules that have been widely associated with the control of several infections and autoimmune disorders [1–6]. Interestingly, not only the HLA-B molecule *per se* is involved in the inhibition of natural killer (NK) cell cytotoxic activity, but a critical role is also played by the peptide loaded in the HLA-B-binding pocket [7]. In particular, the relative positions 7 and 8 of the peptide bound to HLA class I are critical for the recognition since they directly interact with amino acids from HLA-B at positions 77, 80, 81, 82, and 83 which form Bw4/Bw6 epitope [7, 8]. Bw4 epitope (Asn 77, Ile 80, Ala 81, Leu 82, and Arg 83) has been associated with a strong inhibition of NK cell's activity

through KIR3DL1 receptor interaction, while Bw6 epitope (Ser 77, Asn 80, Leu 81, Arg 82, and Gly 83) does not interact with KIR3DL1 [3, 9, 10], though a weak bound with one of the alleles has been described [11].

The KIRs/HLA-B complex crystal structure has been resolved and deposited in the Protein Data Bank [12]. In particular, KIR3DL1 D1 domain is capable to interact both with the HLA bond peptide and the HLA class I molecule alpha-1 domain including the Bw4/Bw6 epitope [12]. This notion supports the observed differences in NK mediated cytotoxic effects due to peptides with different amino acids residues in the relative positions P7 and P8 [2, 8, 13, 14].

The basis for this effect might be due to different, not mutually exclusive mechanisms: (i) direct steric effect, (ii) lack of complementarities between the peptide and KIR3DL1\*001, and (iii) the induced conformational change in the Bw4 epitope [12]. To assess the complexity of KIR3DL1-HLA-B interaction and its role in the immune regulation we studied the formation of complexes between different HLA-B alleles carrying Bw4/Bw6 and KIR3DL1\*001 allele in presence of different peptides using a structural immunoinformatic approach.

#### 2. Material and Methods

Seven HLA-B alleles have been evaluated according to carrying Bw4/Bw6 epitope supertype [15] and allele frequency in different human populations [16] (Table 1). Quality of the models based on X-ray crystallography experiments or homology modeling in SWISS model workspace [17] has been evaluated with PROSESS server [18]. In order to compare different HLA-B alleles, peptides with the similar EC<sub>50</sub> (200 nM) were selected as references from IEDB database [19]. In the analysis of HLA-B alleles belonging to the same supertype, the same reference bound peptide has been used in all evaluated HLA-B molecules, even if presenting a different EC<sub>50</sub> for the tested alleles.

Homology modeling has been applied using HLA-B57\_ KIR3DL1 complex as a template (PDB ID: 3VH8). A MOTIF alignment [20] of different HLA-B structures with HLA-B57 followed by HLA-B replacement has been performed. The different HLA-bound peptides have been generated through single-point amino acid substitution.

All models were subjected to energy minimization by using Yet Another Scientific Artificial Reality Application, ("YASARA" http://www.yasara.com/) and the implemented AMBER and NOVA force fields (ff). The same minimization protocol was used with both force fields. This is represented by a combination of steepest descent performed at the beginning and followed up by a simulated annealing. The calculation is performed until the convergence criteria are met or the maximum number of steps is exceeded.

To study differences between AMBER and NOVA ff and to prevent bias due to a low number of data points, HLA-B alleles analyzed have at least 100 data points for both binders ( $EC_{50} < 500 \text{ nm}$ ) and nonbinders ( $EC_{50} > 20000 \text{ nm}$ ) peptides in the IEDB database (HLA-B\*35:01: binders = 162, non-binders = 108; HLA-B\*51:01: binders = 142, nonbinders = 220; HLA-B\*57:01: binders = 144, non-binders = 205; HLA-B\*58:01: binders = 154, non-binders = 516) [19].

HLA-B\_KIR3DL1 complexes were therefore analyzed using NOVA ff for minimization followed by AMBER ff for energy calculation as previously described [21].

The binding free energy ( $\Delta G$ ) has been calculated as the sum of the interactive and solvation energies. KIR3DL1 binding free energy has been calculated as the difference between KIR3DL1-HLA-B-peptide complex and the sum of HLA-peptide complex and KIR3DL1 alone as previously reported [22].

The difference in the KIR3DL1 binding free energy ( $\Delta\Delta G$ ) in presence of different HLA bound peptides has been calculated as differences between the  $\Delta G$  of the KIR3DL1/ HLA-peptide-(substituted) complex and the KIR3DL1/HLApeptide-(reference) complex as previously reported [23]. The more negative is the value the stronger is the predicted KIR3DL1 interaction.

Statistical analysis has been performed using the paired nonparametric Wilcoxon test, on GraphPad Prism 5.0 (San Diego, CA, USA). Graphics are represented as whiskers of 10– 90 percentiles of the distribution obtained for the 180 singleamino acid substitutions in the bound peptide to HLA-B allele.

#### 3. Results and Discussion

3.1. Immunoinformatic Approach. To select the best strategy for evaluation of the HLA-peptide interaction with KIR3DL1 receptor, different energy minimization ff have been tested in parallel. Among them, AMBER and NOVA ff are the most used in bioinformatics approach [24, 25]. To compare the applicability on our system of the two ff, amino acid substitution in the HLA bound peptide has been performed according to a list of binding (EC<sub>50</sub> < 500 nM) and nonbinding peptides (EC<sub>50</sub> > 20000 nM) from IEDB. NOVA ff demonstrated better performance compared to AMBER for distinguishing the binding and nonbinding peptides (HLA-B NOVA ff: mean value binders  $-129.6 \pm 198$  kj/mol, mean value nonbinders  $-399.6 \pm 181.8$  kj/mol, P = 0.0286; HLA-B AMBER ff: mean value binders  $-637.6 \pm 154.9$  kj/mol, mean value nonbinders  $-929.4 \pm 423.8$ , P = 0.1714).

The use of AMBER allowed a greater distortion than NOVA in the cocrystallization structure of HLA-B\*57:01-KIR3DL1 when used for energy minimization (AMBER minimized structure RMSD = 0.725; NOVA minimized structure RMSD = 0.573). This is in agreement with previous observation of other molecular systems comparing the two ff [24]. Therefore, we considered NOVA ff for the further study, since it was capable of predicting successfully the ligand-receptor interaction due to the lower molecular distortion caused on the entire complex.

3.2. Role of HLA Bw4/Bw6 Epitope in KIR3DL1\*001 Interaction. To evaluate the contribution of Bw4 and Bw6 epitopes together with the peptide substitution in KIR3DL1/HLA interaction, different HLA-B alleles carrying Bw4 or Bw6 epitope have been analyzed (Table 1).

The presence of epitope Bw4 determines a higher strength of interaction with KIR3DL1\*001 allele respect to epitope Bw6 (P < 0.0001, Figure 1(a)). This is in agreement with previous experimental and association studies showing a strongest interaction of Bw4 epitope with KIR3DL1 respect to Bw6 [3, 9, 10].

The distribution of the  $\Delta\Delta G$  for all the substituted peptides for each HLA-B allele has been further analyzed. HLA-B\*07:02 has the weakest interaction with KIR3DL1\*001 followed by HLA-B\*35:01 as previously reported in experimental data obtained through flow cytometry and cytotoxic activity assay [26–28]. On the contrary, HLA-B\*27:05 has the strongest interaction with KIR3DL1\*001 followed by HLA-B\*57:01 and HLA-B\*58:01 (Figure 1(b)). The KIR3DL1 interaction capability of these three HLA-B alleles has been

TABLE 1: List of HLA-B alleles selected for immune-informatics approach. Alleles were selected depending on Bw4/Bw6 epitope, supertype [15], allele frequency in different population (the observed range of frequency among all the human populations is indicated) [16], and presence of crystallization structure in PDB databank. For each HLA-B supertype, using IEDB database a binding peptide with a similar  $EC_{50}$  value ( $\approx 200 \text{ nM}$ ) was selected.

HLA	Epitope	Supertype	Allele frequency	Peptide EC <sub>50</sub> $\approx 200  \text{nM}$	PDB code
B*07:02	Bw6	B7	0-0.1	MPVGGQSSF	
B*14:02	Bw6	B27	0-0.07	MPAYIRNTL	
B*35:01	Bw6	B7	0-0.1	MPVGGQSSF	3LKN
B*27:05	Bw4	B27	0-0.2	MPAYIRNTL	3BP4
B*51:01	Bw4	B7	0-0.08	MPVGGQSSF	
B*57:01	Bw4	B58	0-0.06	YTAVVPLVY	3VH8
B*58:01	Bw4	B58	0-0.1	YTAVVPLVY	



FIGURE 1: Contribution of epitope Bw4/Bw6 in KIR3DL1 interaction. Here is shown the strength of interaction between HLA-B alleles carrying epitope Bw4 and Bw6 as the difference in the estimated binding free energy ( $\Delta\Delta G$ ) between the substituted peptide and the reference. The higher is the value the stronger is the interaction. (a) Overall interaction between HLA-B alleles grouped based on carrying epitope Bw4 or Bw6. (b) Interaction of each HLA-B allele with KIR3DL1\*001: the comparison between HLA-B alleles belonging to the same supertype has been performed. Each HLA-B distribution is composed by 180 points corresponding to each complex with a single-peptide amino acid substitution and is reported as 10–90 percentile whiskers. All the statistical analysis has been performed using paired, nonparametric, Wilcoxon test.

independently observed and confirmed in previous literature data mostly through cytotoxic assay [1, 10, 11, 26, 29, 30].

Worth-noting, among HLA-B alleles carrying Bw6 epitope, HLA-B\*14:02 has the highest interaction with KIR3DL1\*001. However, it presents a weaker interaction compared to the Bw4 allele HLA-B\*27:05, the other allele belonging to the B27 supertype (P < 0.0001; Figure 1(b)).

On the contrary, among HLA-B alleles carrying Bw4 epitope, HLA-B\*51:01 has the weakest interaction with KIR3DL1\*001. However, the interaction is stronger respect to Bw6 alleles HLA-B\*07:02 and HLA-B\*35:01, all belonging to the B7 supertype (B\*07:02 versus B\*51:01, P < 0.0001; B\*35:01 versus B\*51:01, P = 0.0023; Figure 1(b)).

Finally, no difference in KIR3DL1\*001 interaction has been observed between HLA-B\*57:01 and HLA-B\*58:01



FIGURE 2: HLA-B and KIR3DL1\*001 interaction binding site. Here is shown the interaction of KIR3DL1\*001 with HLA-B and the most significant differences in the contact surface between Bw4 and Bw6 epitope ((a) and (b) resp.). In green KIR3DL1, in grey HLA-B and in orange the peptide are shown. For each amino acid residue in the site of interaction Van Deer Walls surface was calculated and amino acid residues responsible for the contact surface are named and shown with the parent color as ball sticks.

(P = 0.1022; Figure 1(b)) belonging to B58 supertype both carrying Bw4 epitope.

3.3. Role of Arg79 in KIR3DL1/HLA-B Interaction. The contact residues in the HLA-B and KIR3DL1\*001 interaction have been investigated in the different HLA-B alleles carrying Bw4 and Bw6 epitope. The main difference has been observed in amino acid residues 136–142 of D1 domain of KIR3DL1\*001 interacting with the nonpolymorphic arginine 79 of the HLA-B (Figure 2). In presence of Bw4 epitope (Figure 2(a)), Arg79 is directed towards the 136–142 loop, while in presence of Bw6 epitope Arg79 changes the conformation (Figure 2(b)). This represents the most critical structural aspect at the basis of the Bw4/Bw6 different interaction capability of KIR3DL1 with HLA-B molecule.

#### 4. Conclusions

Here we confirm for the first time via structural immunoinformatic approach, the highest interaction of HLA-B alleles carrying Bw4 epitope with KIR3DL1\*001 compared to epitope Bw6. The presence of epitope Bw4 is determining a conformational change which strengthens the interaction between the nonpolymorphic HLA-B Arg79 residue and KIR3DL1\*001 136–142 loop. In conclusion, the study of KIR3DL1-HLA-B aims to explain the mechanisms of immune modulation of NK cells immune response and may provide the basis for the design of more specific immunotherapeutic approaches in infectious diseases and immune disorders.

#### **Conflict of Interests**

The authors declare that they have no commercial or other conflict of interests within the presentation of the data reported in the paper.

#### Acknowledgments

This study has been supported in part by the University of Rome "Tor Vergata." Alba Grifoni is a recipient of a PhD fellowship in "Immunology and applied biotechnology" (cycle XXVI) of the University of Rome "Tor Vergata."

#### References

- N. T. Young, N. A. Rust, M. J. Dallman, V. Cerundolo, P. J. Morris, and K. I. Welsh, "Independent contributions of HLA epitopes and killer inhibitory receptor expression to the functional alloreactive specificity of natural killer cells," *Human Immunology*, vol. 59, no. 11, pp. 700–712, 1998.
- [2] E. O. Long and S. Rajagopalan, "HLA class I recognition by killer cell Ig-like receptors," *Seminars in Immunology*, vol. 12, no. 2, pp. 101–108, 2000.
- [3] P. Parham, P. J. Norman, L. Abi-Rached, and L. A. Guethlein, "Variable NK cell receptors exemplified by human KIR3DL1/S1," *Journal of Immunology*, vol. 187, no. 1, pp. 11–19, 2011.
- [4] M. P. Martin, Y. Qi, X. Gao et al., "Innate partnership of HLA-B and KIR3DL1 subtypes against HIV-1," *Nature Genetics*, vol. 39, no. 6, pp. 733–740, 2007.
- [5] P. H. Salim, M. Jobim, L. F. Jobim, and R. M. Xavier, "Autoimmune rheumatic diseases and their association with killer immunoglobulin-like receptor genes," *Revista Brasileira de Reumatologia*, vol. 51, no. 4, pp. 351–364, 2011.
- [6] M. Carrington and G. Alter, "Innate immune control of HIV," *Cold Spring Harbor Perspectives in Medicine*, vol. 2, Article ID a007070, 2012.
- [7] M. Peruzzi, N. Wagtmann, and E. O. Long, "A p70 killer cell inhibitory receptor specific for several HLA-B allotypes discriminates among peptides bound to HLA-B\* 2705," *Journal* of *Experimental Medicine*, vol. 184, no. 4, pp. 1585–1590, 1996.
- [8] M. Peruzzi, K. C. Parker, E. O. Long, and M. S. Malnati, "Peptide sequence requirements for the recognition of HLA-B\* 2705 by specific natural killer cells," *Journal of Immunology*, vol. 157, no. 8, pp. 3350–3356, 1996.
- [9] J. E. Gumperz, N. M. Valiante, P. Parham, L. L. Lanier, and D. Tyan, "Heterogeneous phenotypes of expression of the NKB1

natural killer cell class I receptor among individuals of different human histocompatibility leukocyte antigens types appear genetically regulated, but not linked to major histocompatibility complex haplotype," *Journal of Experimental Medicine*, vol. 183, no. 4, pp. 1817–1827, 1996.

- [10] V. Litwin, J. Gumperz, P. Parham, J. H. Phillips, and L. L. Lanier, "NKB1: a natural killer cell receptor involved in the recognition of polymorphic HLA-B molecules," *Journal of Experimental Medicine*, vol. 180, no. 2, pp. 537–543, 1994.
- [11] L. Fu, B. Hazes, and D. N. Burshtyn, "The first Ig domain of KIR3DL1 contacts MHC class I at a secondary site," *Journal of Immunology*, vol. 187, no. 4, pp. 1816–1825, 2011.
- [12] J. P. Vivian, R. C. Duncan, R. Berry et al., "Killer cell immunoglobulin-like receptor 3DL1-mediated recognition of human leukocyte antigen B," *Nature*, vol. 479, no. 7373, pp. 401–405, 2011.
- [13] L. Fadda, G. M. O'Connor, S. Kumar et al., "Common HIV-1 peptide variants mediate differential binding of KIR3DL1 to HLA-Bw4 molecules," *Journal of Virology*, vol. 85, no. 12, pp. 5970–5974, 2011.
- [14] G. B. E. Stewart-Jones, K. di Gleria, S. Kollnberger, A. J. McMichael, E. Y. Jones, and P. Bowness, "Crystal structures and KIR3DL1 recognition of three immunodominant viral peptides complexed to HLA-B\*2705," *European Journal of Immunology*, vol. 35, no. 2, pp. 341–351, 2005.
- [15] J. Sidney, B. Peters, N. Frahm, C. Brander, and A. Sette, "HLA class I supertypes: a revised and updated classification," *BMC Immunology*, vol. 9, article 1, 2008.
- [16] F. F. Gonzalez-Galarza, S. Christmas, D. Middleton, and A. R. Jones, "Allele frequency net: a database and online repository for immune gene frequencies in worldwide populations," *Nucleic Acids Research*, vol. 39, no. 1, pp. D913–D919, 2011.
- [17] K. Arnold, L. Bordoli, J. Kopp, and T. Schwede, "The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling," *Bioinformatics*, vol. 22, no. 2, pp. 195–201, 2006.
- [18] M. Berjanskii, Y. Liang, J. Zhou et al., "PROSESS: a protein structure evaluation suite and server," *Nucleic Acids Research*, vol. 38, no. 2, pp. W633–W640, 2010.
- [19] R. Vita, L. Zarebski, J. A. Greenbaum et al., "The immune epitope database 2.0," *Nucleic Acids Research*, vol. 38, no. 1, pp. D854–D862, 2010.
- [20] E. Krieger, K. Joo, J. Lee et al., "Improving physical realism, stereochemistry, and side-chain accuracy in homology modeling: four approaches that performed well in CASP8," *Proteins*, vol. 77, no. 9, pp. 114–122, 2009.
- [21] A. Grifoni, C. Montesano, P. Palma, A. Salerno, V. Colizzi, and M. Amicosante, "Role of HLA-B alpha-3 domain amino acid position 194 in HIV disease progression," *Molecular Immunol*ogy, vol. 53, article 410, 2012.
- [22] A. P. Graves, D. M. Shivakumar, S. E. Boyce, M. P. Jacobson, D. A. Case, and B. K. Shoichet, "Rescoring docking hit lists for model cavity sites: predictions and experimental testing," *Journal of Molecular Biology*, vol. 377, no. 3, pp. 914–934, 2008.
- [23] T. A. Binkowski, S. R. Marino, and A. Joachimiak, "Predicting HLA class I non-permissive amino acid residues substitutions," *PLoS ONE*, vol. 7, Article ID e41710, 2012.
- [24] E. Krieger, G. Koraimann, and G. Vriend, "Increasing the precision of comparative models with YASARA NOVA—a selfparameterizing force field," *Proteins*, vol. 47, no. 3, pp. 393–402, 2002.

- [25] Y. Duan, C. Wu, S. Chowdhury et al., "A point-charge force field for molecular mechanics simulations of proteins based on condensed-phase quantum mechanical calculations," *Journal of Computational Chemistry*, vol. 24, no. 16, pp. 1999–2012, 2003.
- [26] Z. B. Kurago, C. T. Lutz, K. D. Smith, and M. Colonna, "NK cell natural cytotoxicity and IFN-γ production are not always coordinately regulated: engagement of DX9 KIR+ NK cells by HLA-B7 variants and target cells," *Journal of Immunology*, vol. 160, no. 4, pp. 1573–1580, 1998.
- [27] S. Rojo, N. Wagtmann, and E. O. Long, "Binding of a soluble p70 killer cell inhibitory receptor to HLA-B(\*)5101: requirement for all three p70 immunoglobulin domains," *European Journal of Immunology*, vol. 27, no. 2, pp. 568–571, 1997.
- [28] H. Thananchai, G. Gillespie, M. P. Martin et al., "Cutting edge: allele-specific and peptide-dependent interactions between KIR3DL1 and HLA-A and HLA-B," *Journal of Immunology*, vol. 178, no. 1, pp. 33–37, 2007.
- [29] W. H. Carr, M. J. Pando, and P. Parham, "KIR3DL1 polymorphisms that affect NK cell inhibition by HLA-Bw4 ligand," *Journal of Immunology*, vol. 175, no. 8, pp. 5222–5229, 2005.
- [30] G. M. O'Connor, K. J. Guinan, R. T. Cunningham, D. Middleton, P. Parham, and C. M. Gardiner, "Functional polymorphism of the KIR3DL1/S1 receptor on human NK cells," *Journal of Immunology*, vol. 178, no. 1, pp. 235–241, 2007.



BioMed Research International









International Journal of Genomics











The Scientific World Journal



Genetics Research International



Anatomy Research International



International Journal of Microbiology



Biochemistry Research International



Journal of Marine Biology







International Journal of Evolutionary Biology



Molecular Biology International