

Review Article Some In Vitro/In Vivo Chemically-Induced Experimental Models of Liver Oxidative Stress in Rats

Rumyana Simeonova, Magdalena Kondeva-Burdina, Vessela Vitcheva, and Mitka Mitcheva

Laboratory of Drug Metabolism and Drug Toxicity, Department of Pharmacology, Pharmacotherapy and Toxicology, Faculty of Pharmacy, Medical University 2 Dunav Street, 1000 Sofia, Bulgaria

Correspondence should be addressed to Rumyana Simeonova; rsimeonova@yahoo.com

Received 27 April 2013; Accepted 24 October 2013; Published 16 January 2014

Academic Editor: Afaf K. El-Ansary

Copyright © 2014 Rumyana Simeonova et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oxidative stress is critically involved in a variety of diseases. Reactive oxygen species (ROS) are highly toxic molecules that are generated during the body's metabolic reactions and can react with and damage some cellular molecules such as lipids, proteins, or DNA. Liver is an important target of the oxidative stress because of its exposure to various prooxidant toxic compounds as well as of its metabolic function and ability to transform some xenobiotics to reactive toxic metabolites (as ROS). To investigate the processes of liver injuries and especially liver oxidative damages there are many experimental models, some of which we discuss further.

1. Introduction

Oxidative stress is an imbalance between the production and scavenging of reactive oxygen and nitrogen species (ROS and RNS) and free radicals that can induce lipid peroxidation, DNA fragmentation, and protein oxidation [1]. These damages result in the loss of membrane integrity, structural and functional changes in proteins, and gene mutations [2]. Normally, the affected cells are trying to neutralise reactive molecules by deploying their antioxidative defense that include reduced glutathione (GSH), alpha-tocopherol, ascorbic acid, antioxidant enzymes catalase (CAT), superoxide dismutase (SOD), glutathione peroxidase (GPx), glutathione reductase (GR), and glutathione-S-transferase (GST).

Oxidative stress is critically involved in a variety of diseases. ROS are highly dangerous molecules that are generated during the body's metabolic reactions and can react with and damage some cellular molecules such as lipids, proteins, or DNA.

Liver plays a pivotal role in the regulation of various physiological processes in the body such as carbohydrate metabolism and storage, fat metabolism, bile acid synthesis, and so forth besides being the most important organ involved in the detoxification of various drugs as well as xenobiotics in our body [3]. It is highly susceptible to damage by xenobiotics owing to its continuous exposure to these toxicants via the portal blood circulation [4]. Various chemicals, like carbon tetrachloride (CCl₄), tert-butyl hydroperoxide (t-BHP), alcohol, paracetamol, galactosamine (GalN), and others, can cause potential damage to the liver cells leading to progressive dysfunction. Most of the hepatotoxic chemicals cause damage to the hepatocytes by inducing lipid peroxidation [5, 6]. Thus, the disorders associated with liver are numerous and varied.

One of the most important liver toxicity mechanisms might be a consequence of cell damage by ROS and RNS. Kupffer cells release reactive oxygen species (ROS), cytokines, and chemokines, which induce neutrophil extravasation and activation. Also the liver expresses many cytochrome P450 isoforms, including ethanol-induced CYP2E1. CYP2E1 generates ROS, activates many toxicologically important substrates, and may be the central pathway by which some substances cause oxidative stress (ethanol, carbon tetrachloride, etc.) [7].

In this review we summarize some commonly used toxic models employed in the study of hepatotoxicity and hepatoprotection. A number of models of hepatic disorders support the notion that ROS have a causal role in liver injuries. Experimental liver injuries are induced by specific toxic compounds, because the formation of ROS is stimulated by a number of xenobiotics.

2. Carbon Tetrachloride (CCl₄)

Carbon tetrachloride (CCl₄) is the most widely used model to develop oxidative stress and liver toxicity in rats. Hepatic injury through carbon tetrachloride induced lipid peroxidation is well known and has been extensively used in the experimental models to understand the cellular mechanisms behind oxidative damage and further to evaluate the therapeutic potential of drugs and dietary antioxidants [8].

CCl₄ is activated by cytochrome CYP2E1, CYP2B1, or CYP2B2, and possibly CYP3A, to form the trichloromethyl radical, CCl3^{*} [9]. This radical can bind to cellular molecules (nucleic acid, protein, lipid), impairing crucial cellular processes such as lipid metabolism, with the potential outcome of fatty degeneration (steatosis) [10]. This radical can also react with oxygen to form the trichloromethylperoxy radical CCl_3OO^* , a highly reactive species. CCl_3OO^* initiates the chain reaction of lipid peroxidation, which attacks and destroys polyunsaturated fatty acids [9]. Among the degradation products of fatty acids are reactive aldehydes, malondialdehyde (MDA), and 4-hydroxynonenal, which bind easily to functional groups of proteins and inhibit important enzyme activities. Disturbed cellular processes are most likely due to increased levels of these thiobarbituric acid reactive species (TBARS) [11], lactate dehydrogenase (LDH) leakage as a result of membrane breakdown and concomitant increase in membrane permeability [12], loss of cell protection, witnessed by GSH depletion and as a result of all these changescell death.

In our laboratory we use some *in vitro* and *in vivo* hepatotoxicity models based on CCl_4 -induced liver damage in Wistar rats and in spontaneously hypertensive rats (SHR). *In vitro* experiments are carried out in primary isolated rat hepatocytes [13] or liver microsomes [14]. Cell incubation with CCl_4 (86 μ mol L⁻¹) leads to a significant decrease in cell viability, increased LDH leakage, decreased levels of cellular GSH, and elevation in MDA quantity. Enzyme-induced LPO is started with 20 mM CCl_4 in the presence of 1 mM NADPH [14]. For *in vivo* experiments Wistar rats are challenged with a single dose (2 mL/kg) of 20% of CCl_4 in olive oil [15]. These *in vitro/in vivo* CCl_4 -induced liver injury models are useful for investigations on hepatoprotective and antioxidant properties of some plant-derived biologically active compounds [13–17].

We found that ROS, produced by CCl_4 , decrease the activities not only of antioxidant enzymes such as catalase (CAT), superoxide dismutase (SOD), glutathione peroxidase (GPx), glutathione reductase (GR), and glutathione-S-transferase (GST) [18], but also the activities of some drug metabolizing enzymes such as CYP2E1 and CYP3A, involved in their production [15].

3. Tert-Butyl Hydroperoxide (t-BHP)

The cellular system of energy supply localized in mitochondria is another target of many hepatotoxic substances causing oxidative stress and is one of the most important mechanisms through which hepatotoxic factors induced apoptotic and necrotic processes [19].

Tert-butyl hydroperoxide caused necrosis through inducing mitochondrial reactive oxygen formation [20]. As a prooxidant, t-BHP was widely used and many effects on cell metabolism have been described, for example, changes in calcium homeostasis [21], increase of lipid peroxidation, or decrease of mitochondrial membrane potential [22, 23]. Two mechanisms for t-BHP action were proposed: depletion of cellular stores of GSH and oxidation of functionally important SH groups on mitochondrial enzymes [24] and/or changes of mitochondrial membrane integrity induced by peroxidation of membrane lipids [22, 23]. The metabolism of t-BHP to free radicals undergoes through several steps. In microsomal suspension, in the absence of NADPH, it has been shown to undergo one-electron oxidation to a peroxyl radical (1), whereas in the presence of NADPH it has been shown to undergo one-electron reduction to an alkoxyl radical (2). In isolated mitochondria and intact cells, the t-BHP has been shown to undergo β -scission to the methyl radical (3). All these radicals cause lipid peroxidation process [25, 26]:

$$(CH_3)_3COOH \longrightarrow (CH_3)_3COO^{\bullet} + e^- + H^+$$
 (1)

$$(CH_3)_3COOH + e^- \longrightarrow (CH_3)_3CO^{\bullet} + OH^-$$
 (2)

$$(CH_3)_3 CO^{\bullet} \longrightarrow (CH_3)_2 CO + {}^{\bullet}CH_3$$
 (3)

Experiments on isolated hepatocytes are thus a useful model system for evaluation of the toxic effect of various prooxidants which act directly on mitochondrial enzymes. In our experiments using freshly isolated rat hepatocytes we found that t-BHP (75 μ mol L⁻¹) decreases cell viability [27, 28]. It causes leakage of lactate dehydrogenase (LDH) and formation of malondialdehyde in hepatocytes. Furthermore, t-BHP causes the depletion of cellular GSH levels. These data correlate with the results obtained by many authors [23–25].

Enhanced formation of ROS has been suggested to play a role in some liver disease processes, including alcoholinduced liver injury [29–31], paracetamol-induced liver failure [32, 33], and many others. Many other drugs as isoniazide, amiodarone, and valproic acid as well as widely used and abused substances as nicotine and cocaine damage liver cells by producing toxic ROS. Because of their widespread consumption, they are also used as experimental models of liver injuries.

4. Ethanol

Acute and chronic ethanol treatments increase the production of ROS, lower cellular antioxidant levels, and enhance oxidative stress in many tissues, especially the liver. Ethanolinduced oxidative stress plays a major role in the mechanisms by which ethanol produces liver injury [34].

The liver expresses many cytochrome P450 isoforms, including ethanol-induced CYP2E1. CYP2E1 generates ROS, activates many toxicologically important substrates, and may

be the central pathway by which ethanol causes oxidative stress [7].

CYP2E1 metabolizes and activates many toxicologically important substrates, including ethanol, carbon tetrachloride, acetaminophen, and N-nitrosodimethylamine, to more toxic products [35, 36]. Induction of CYP2E1 by ethanol is a central pathway by which ethanol generates oxidative stress. In our intragastric model of ethanol feeding (3 g/kg, 14 days) of spontaneously hypertensive rats (SHR) a prominent induction of CYP2E1 occurs along with significant alcohol liver injury [37]. Lipid peroxidation also occurs, and ethanol-induced liver pathology correlates with CYP2E1 levels and elevated lipid peroxidation [38]. Chronic ethanol consumption is associated with reduced liver GSH and alphatocopherol level and with reduced superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx) activity [39]. Our results, concerning normotensive rats (Wistar-Kyoto), are in accordance with these data, whereas alcohol intake in SHR increases significantly SOD and CAT activities by approximately 50% [37]. We suggested that the differences in antioxidant status and the effect of ethanol between the strains might be due to the oxidative stress state in the hypertensive pathological model. Additionally we found that multiple ethanol treatment resulted in less pronounced effect on the assessed parameters (MDA, GSH, nNOS) in the female SHR, compared to male SHR [38]. These results might be due to a protective effect of estrogens on the oxidative stress and to a stimulation of the antioxidant defense systems, in liver.

5. Paracetamol

Paracetamol (PCM) is primarily metabolized by sulfation and glucuronidation, but with an increasing dose rate; these pathways become saturated and a greater proportion of the drug is available for oxidation by the microsomal cytochrome P-450 system [40]. N-Acetyl-P-benzoquinone Imine (NAPQI) is the product of this pathway which is thought to be responsible for the subsequent hepatic damage [41]. Nacetyl-P-benzoquinone imine (NAPQI) is a highly reactive electrophile and is detoxified in liver by either reduction to the parent compound, acetaminophen, or conjugation at the metaposition with glutathione, in which both reactions consume GSH [42].

Glutathione (GSH) plays an important role in protecting cells from electrophilic compounds and free radicals such as reactive oxygen species generated during cellular metabolism. Reduced glutathione can act as a reductant, reducing hydrogen peroxide and lipid hydroperoxides directly to H_2O , a reaction catalyzed by GSH-Px [43]. Depletion of intracellular GSH, under conditions of continuous intracellular oxidative stress, leads to oxidation and damage of lipids, proteins, and DNA by the reactive oxygen species [44, 45].

The importance of glutathione in PCM toxicity is further emphasized by the large body of evidence which indicates that interventions which increase GSH content can dramatically reduce PCM and NAPQI-induced hepatic injury [46, 47]. Our previous studies have shown that GSH content in animal livers decreases after PCM overdose (2 g/kg, i.p. single dose) [15] and have proved that biologically active compounds derived from plants are useful for treatment of PCM-induced liver disorders, because of a stimulation of GSH synthesis.

Oxidative stress is also considered to be involved in the induction of hepatotoxicity by PCM. Oxidation of PCM by CYPs may generate ROS. Hydrogen peroxide and superoxide are produced during metabolic activation of PCM in the mixed function oxidase system [47].

6. Nicotine

During smoking, nicotine is rapidly absorbed into the circulatory system where more than 80% is metabolized in the liver [48]. Liver is an important organ and is responsible for biotransformation of drugs and other toxins to remove them from the body. Nicotine from heavy smoking increases the risk of developing some dangerous liver disorders by one of the main mechanisms being the oxidative stress. Increased production of free radicals or decreased function of the defense system play an important role in nicotine toxicity [49]. Also maternal nicotine exposure induces oxidative stress and causes histopathological changes in the lung and liver of lactating offspring [50]. Nicotine induces oxidative stress both in vivo and in vitro that causes a peroxidant/antioxidant imbalance in blood cells, blood plasma, and other tissues [51]. Some authors [50, 52] reported that nicotine induces oxidative stress and depleted antioxidant defense mechanisms through reduction of glutathione peroxidase in liver, lung, and kidney. Oxidative stress generates free radicals that attack the membrane lipids resulting in the formation of malondialdehyde (MDA), which causes peroxidative tissue damage [53]. Animal's studies have shown significantly higher liver and serum levels of MDA, conjugated dienes, hydroperoxides, and free fatty acids in rats intoxicated by nicotine [54, 55].

Nicotine is not recognized as a common experimental model for liver injuries, but because of its well-established prooxidant mechanisms of hepatotoxicity, and widespread consumption is used from many authors [51, 53, 55, 56] for investigations of antioxidant and protective properties of natural compounds.

In our previous experiments [57, 58] enhanced level of tissue lipid peroxides in nicotine treated rats (1 mg/kg i.p; 6,5 mg/kg p.o.) has been shown to be accompanied by a significant decrease in the levels of GSH, glutathione peroxidase (GPx), superoxide dismutase (SOD), and catalase (CAT) and increased glutathione reductase (GR) activity in Wistar rat liver.

7. D-Galactosamine (GAL)

D-Galactosamine is a well-known experimental hepatotoxin usually used to produce acute toxicity in rat liver. Galactosamine metabolism depletes the uridine pool of hepatocytes, thus inducing transcriptional arrest and causing an increase in sensitization to cytokines such as TNF- α and an

increase in oxidative stress and GSH depletion, which lead to mitochondrial dysfunction and cell death [59]. Both oxidative and nitrosative stress play a key role in the pathogenesis of GAL-induced hepatic injury [60].

Usually rats are injected (i.p.) with GAL (400 mg/kg b.w.) as a single dose [61].

8. Cocaine

Cocaine is a psychoactive drug that has been recognized as one of the most significant examples of drug abuse. Its misuse can induce severe toxic effects, including neurotoxicity, cardiotoxicity, and hepatotoxicity. One of the main mechanisms discussed for cocaine-induced liver injury is promotion of lipid peroxidation by ROS which are produced during cocaine bioactivation to norcocaine through N-demethylation by cytochrome P 450 and flavin adenine dinucleotide containing monooxygenases [62].

A large body of evidence in both human and experimental models suggests that impairment of the antioxidant defense system by cocaine and its metabolites plays a role in the pathogenesis of cocaine hepatotoxicity [62-64]. In particular, glutathione seems to play an important protective role against cocaine-induced hepatic injury. For example, the acute administration of cocaine to rats [65] and multiple treatments of mice [63] have been shown to deplete the cellular reduced glutathione, to decrease the activity of superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx) and to increase glutathione reductase (GR) activity. The GSH depletion, induced by cocaine administration, observed in these and other studies [62, 66] might be explained by increased utilization of GSH for detoxification of ROS and lipid peroxidation products. The critical role of ROS and oxidative stress in the pathogenesis of cocaine-induced liver damage was demonstrated by the observed ameliorating effects of bioactive compounds with an antioxidant activity, administered several days before cocaine treatment [65, 67]. The bioactive compounds were found to decrease cocaine toxicity both by increasing GSH levels and antioxidant enzyme activities.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- C. Nencini, G. Giorgi, and L. Micheli, "Protective effect of silymarin on oxidative stress in rat brain," *Phytomedicine*, vol. 14, no. 2-3, pp. 129–135, 2007.
- [2] B. V. Reddy, J. S. Sundari, E. Balamurugan, and V. P. Menon, "Prevention of nicotine and streptozotocin treatment induced circulatory oxidative stress by bis-1,7-(2-hydroxyphenyl)-hepta-1,6-diene-3,5-dione in diabetic rats," *Molecular and Cellular Biochemistry*, vol. 331, no. 1-2, pp. 127–133, 2009.
- [3] S. K. Sharma, S. M. Arogya, D. H. Bhaskarmurthy, A. Agarwal, and C. C. Velusami, "Hepatoprotective activity of the *Phyllan*-

thus species on tert-butyl hydroperoxide (t-BH)-induced cytotoxicity in HepG2 cells," *Pharmacognosy Magazine*, vol. 7, no. 27, pp. 229–233, 2011.

- [4] V. M. Piñeiro-Carrero and E. O. Piñeiro, "Liver," *Pediatrics*, vol. 113, no. 4, pp. 1097–1106, 2004.
- [5] A. Subramoniam and P. Pushpangadan, "Development of phytomedicines for liver diseases," *Indian Journal of Pharmacology*, vol. 31, no. 3, pp. 166–175, 1999.
- [6] M. Joyeux, A. Rolland, J. Fleurentin, F. Mortier, and P. Dorfman, "Tert-butyl hydroperoxide-induced injury in isolated rat hepatocytes: a model for studying anti-hepatotoxic crude drugs," *Planta Medica*, vol. 56, no. 2, pp. 171–174, 1990.
- [7] H. Jaeschke, G. J. Gores, A. I. Cederbaum, J. A. Hinson, D. Pessayre, and J. J. Lemasters, "Mechanisms of hepatotoxicity," *Toxicological Sciences*, vol. 65, no. 2, pp. 166–176, 2002.
- [8] S. Basu, "Carbon tetrachloride-induced lipid peroxidation: eicosanoid formation and their regulation by antioxidant nutrients," *Toxicology*, vol. 189, no. 1-2, pp. 113–127, 2003.
- [9] L. W. D. Weber, M. Boll, and A. Stampfl, "Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as a toxicological model," *Critical Reviews in Toxicology*, vol. 33, no. 2, pp. 105–136, 2003.
- [10] Y. Masuda, "Learning toxicology from carbon tetrachlorideinduced hepatotoxicity," *Yakugaku Zasshi*, vol. 126, no. 10, pp. 885–889, 2006.
- [11] M. Bhadauria, S. K. Nirala, and S. Shukla, "Propolis protects CYP 2E1 enzymatic activity and oxidative stress induced by carbon tetrachloride," *Molecular and Cellular Biochemistry*, vol. 302, no. 1-2, pp. 215–224, 2007.
- [12] S. Sahreen, M. R. Khan, and R. A. Khan, "Hepatoprotective effects of methanol extract of *Carissa opaca* leaves on CCl₄induced damage in rat," *BMC Complementary and Alternative Medicine*, vol. 11, article 48, 2011.
- [13] M. Kondeva, M. Mitcheva, and S. Nikolov, "Effect of the Diosgenin in fresh isolated rat hepatocytes treated with carbon tetrachloride," *European Journal of Drug Metabolism and Pharmacokinetics*, vol. 28, no. 1, pp. 1–3, 2003.
- [14] R. L. Simeonova, V. B. Vitcheva, M. S. Kondeva-Burdina, I. N. Krasteva, S. D. Nikolov, and M. K. Mitcheva, "Effect of purified saponin mixture from *Astragalus corniculatus* on enzyme- and non-enzyme-induced lipid peroxidation in liver microsomes from spontaneously hypertensive rats and normotensive rats," *Phytomedicine*, vol. 17, no. 5, pp. 346–349, 2010.
- [15] V. Vitcheva, R. Simeonova, I. Krasteva, S. Nikolov, and M. Mitcheva, "Protective effects of a purified saponin mixture from *Astragalus corniculatus Bieb., in vivo* hepatotoxicity models," *Phytotherapy Research*, vol. 27, no. 5, pp. 731–736, 2013.
- [16] S. A. Ali, M. Z. Rizk, N. A. Ibrahim, M. S. Abdallah, H. M. Sharara, and M. M. Moustafa, "Protective role of *Juniperus phoe*nicea and Cupressus sempervirens against CCl₄," World Journal Gastrointestinal Pharmacology and Therapeutics, vol. 1, no. 6, pp. 123–131, 2010.
- [17] F. Sun, E. Hamagawa, C. Tsutsui, Y. Ono, Y. Ogiri, and S. Kojo, "Evaluation of oxidative stress during apoptosis and necrosis caused by carbon tetrachloride in rat liver," *Biochimica et Biophysica Acta*, vol. 1535, no. 2, pp. 186–191, 2001.
- [18] R. Simeonova, I. Krasteva, M. Kondeva-Burdina, and N. Benbassat, "Effects of extract from Astragalus Glycyphylloides on Carbon tetrachloride-induced hepatotoxicity in Wistar rats," *International Journal of Pharma and Bio Sciences*, vol. 4, no. 3, pp. 179–186, 2013.

- [19] G. Kroemer, B. Dallaporta, and M. Resche-Rigon, "The mitochondrial death/life regulator in apoptosis and necrosis," *Annual Review of Physiology*, vol. 60, pp. 619–642, 1998.
- [20] Z. Drahota, P. Křiváková, Z. Červinková et al., "Tert-butyl hydroperoxide selectively inhibits mitochondrial respiratory-chain enzymes in isolated rat hepatocytes," *Physiology Research*, vol. 54, pp. 67–72, 2005.
- [21] P. Nicotera, D. McConkey, S.-A. Svensson, G. Bellomo, and S. Orrenius, "Correlation between cytosolic Ca²⁺ concentration and cytotoxicity in hepatocytes exposed to oxidative stress," *Toxicology*, vol. 52, no. 1-2, pp. 55–63, 1988.
- [22] R. Rubin and J. L. Farber, "Mechanisms of the killing of cultured hepatocytes by hydrogen peroxide," *Archives of Biochemistry* and Biophysics, vol. 228, no. 2, pp. 450–459, 1984.
- [23] E. Kmoníčková, Z. Drahota, L. Kameníková, Z. Červinková, K. Mašek, and H. Farghali, "Modulatory effect of cyclosporin A on tert-butyl hydroperoxide-induced oxidative damage in hepatocytes," *Immunopharmacology Immunotoxicology*, vol. 23, pp. 43–54, 2001.
- [24] N. Masaki, M. E. Kyle, A. Serroni, and J. L. Farber, "Mitochondrial damage as a mechanism of cell injury in the killing of cultured hepatocytes by tert-butyl hydroperoxide," *Archives of Biochemistry and Biophysics*, vol. 270, no. 2, pp. 672–680, 1989.
- [25] K. Ollinger and U. T. Brunk, "Cellular injury induced by oxidative stress is mediated through lysosomal damage," *Free Radical Biology and Medicine*, vol. 19, no. 5, pp. 565–574, 1995.
- [26] V. O'Donnell and M. J. Burkitt, "Mitochondrial metabolism of a hydroperoxide to free radicals in human endothelial cells: an electron spin resonance spin-trapping investigation," *Biochemical Journal*, vol. 304, no. 3, pp. 707–713, 1994.
- [27] M. Mitcheva, M. Kondeva-Burdina, V. Vitcheva, I. Krasteva, and S. Nikolov, "Effect of purified saponin mixture from *Astr-agalus corniculatus* on toxicity models in isolated rat hepatocytes," *Pharmaceutical Biology*, vol. 46, no. 12, pp. 866–870, 2008.
- [28] M. Mitcheva, M. Kondeva-Burdina, I. Krasteva, and S. Nikolov, "Protective effect of purified saponin mixture from Astragalus corniculatus on toxicity models in vitro," in Medical Management of Chemical and Biological Casualties, S. Tonev, K. Kanev, and C. Dishovsky, Eds., pp. 239–251, Publishing house IRITA, Sofia, Bulgaria, 2009.
- [29] M. Adachi and H. Ishii, "Role of mitochondria in alcoholic liver injury," *Free Radical Biology and Medicine*, vol. 32, no. 6, pp. 487– 491, 2002.
- [30] G. E. Arteel, "Oxidants and antioxidants in alcohol-induced liver disease," *Gastroenterology*, vol. 124, no. 3, pp. 778–790, 2003.
- [31] A. Dey and A. I. Cederbaum, "Alcohol and oxidative liver injury," *Hepatology*, vol. 43, no. 2, pp. S63–S74, 2006.
- [32] H. Jaeschke and M. L. Bajt, "Intracellular signaling mechanisms of acetaminophen-induced liver cell death," *Toxicological Sciences*, vol. 89, no. 1, pp. 31–41, 2006.
- [33] H. Jaeschke, T. R. Knight, and M. L. Bajt, "The role of oxidant stress and reactive nitrogen species in acetaminophen hepatotoxicity," *Toxicology Letters*, vol. 144, no. 3, pp. 279–288, 2003.
- [34] A. I. Cederbaum, Y. Lu, and D. Wu, "Role of oxidative stress in alcohol-induced liver injury," *Archives of Toxicology*, vol. 83, no. 6, pp. 519–548, 2009.
- [35] F. P. Guengerich, D.-H. Kim, and M. Iwasaki, "Role of human cytochrome P-450 IIE1 in the oxidation of many low molecular weight cancer suspects," *Chemical Research in Toxicology*, vol. 4, no. 2, pp. 168–179, 1991.

- [36] D. R. Koop, "Oxidative and reductive metabolism by cytochrome P450 2E1," *The FASEB Journal*, vol. 6, no. 2, pp. 724–730, 1992.
- [37] R. Simeonova, V. Vitcheva, and M. Mitcheva, "Effect of multiple treatments with alcohol on some liver antioxidant biochemical parameters in spontaneously hypertensive rats (SHRs) versus normotensive rats (NTRs)," *Toxicology Letters*, vol. 189, 2009.
- [38] R. Simeonova, V. Vitcheva, and M. Mitcheva, "Effect of ethanol on some hepatic and brain parameters in male and female spontaneously hypertensive rats (SHRs)," *Toxicology Letters*, vol. 196, 2010.
- [39] T. Radosavljević, D. Mladenović, and D. Vucević, "The role of oxidative stress in alcoholic liver injury," *Medicinski Pregled*, vol. 62, no. 11-12, pp. 547–553, 2009.
- [40] P. J. Amar and E. R. Schiff, "Acetaminophen safety and hepatotoxicity: where do we go from here?" *Expert Opinion on Drug Safety*, vol. 6, no. 4, pp. 341–355, 2007.
- [41] L. M. Aleksunes, S. N. Campion, M. J. Goedken, and J. E. Manautou, "Acquired resistance to acetaminophen hepatotoxicity is associated with induction of multidrug resistance-associated protein 4 (Mrp4) in proliferating hepatocytes," *Toxicological Sciences*, vol. 104, no. 2, pp. 261–273, 2008.
- [42] S. U. Ruepp, R. P. Tonge, J. Shaw, N. Wallis, and F. Pognan, "Genomics and proteomics analysis of acetaminophen toxicity in mouse liver," *Toxicological Sciences*, vol. 65, no. 1, pp. 135–150, 2002.
- [43] M. Anoush, M. A. Eghbal, F. Fathiazad, H. Hamzeiy, and N. S. Kouzehkonani, "The protective effects of garlic extract against acetaminophen-induced oxidative stress and Glutathione depletion," *Pakistan Journal of Biological Sciences*, vol. 12, no. 10, pp. 765–771, 2009.
- [44] N. Kaplowitz, "Mechanisms of liver cell injury," *Journal of Hep-atology*, vol. 32, no. 1, pp. 39–47, 2000.
- [45] J. Nordberg and E. S. J. Arnér, "Reactive oxygen species, antioxidants, and the mammalian thioredoxin system," *Free Radical Biology and Medicine*, vol. 31, no. 11, pp. 1287–1312, 2001.
- [46] J. R. Mitchell, D. J. Jollow, and W. Z. Potter, "Acetaminophen induced hepatic necrosis. IV. Protective role of glutathione," *Journal of Pharmacology and Experimental Therapeutics*, vol. 187, no. 1, pp. 211–217, 1973.
- [47] T. Amimoto, T. Matsura, S.-Y. Koyama, T. Nakanishi, K. Yamada, and G. Kajiyama, "Acetaminophen-induced hepatic injury in mice: the role of lipid peroxidation and effects of pretreatment with coenzyme Q10 and α- tocopherol," *Free Radical Biology and Medicine*, vol. 19, no. 2, pp. 169–176, 1995.
- [48] A.-R. El-Zayadi, "Heavy smoking and liver," World Journal of Gastroenterology, vol. 12, no. 38, pp. 6098–6101, 2006.
- [49] K. Chattopadhyay and B. D. Chattopadyay, "Effect of Nicotine on lipid profile, peroxidation & antioxidant enzymes in female rats with restricted dietary protein," *Indian Journal of Medical Research*, vol. 127, no. 6, pp. 571–576, 2008.
- [50] B. H. Ozukutat, K. U. Ozkan, C. F. Ibrahim, E. Guldur, M. S. Kilinc, and F. Inan, "Effects of maternal nicotine exposure during on breast-fed rat pups," *Biology in Neonats*, vol. 88, no. 2, pp. 113–117, 2005.
- [51] H. Suleyman, K. Gumustekin, S. Taysi et al., "Beneficial effects of *Hippophae rhamnoides* L. on nicotine induced oxidative stress in rat blood compared with vitamin E," *Biological and Pharmaceutical Bulletin*, vol. 25, no. 9, pp. 1133–1136, 2002.
- [52] G. H. El-Sokkary, S. Cuzzocrea, and R. J. Reiter, "Effect of chronic nicotine administration on the rat lung and liver: beneficial role of melatonin," *Toxicology*, vol. 239, no. 1-2, pp. 60–67, 2007.

- [53] K. N. Srinivasan and K. V. Pugalendi, "Effect of excessive intake of thermally oxidized sesame oil on lipids, lipid peroxidation and antioxidants' status in rats," *Indian Journal of Experimental Biology*, vol. 38, no. 8, pp. 777–780, 2000.
- [54] L. Ashakumary and P. L. Vijayammal, "Additive effect of alcohol and nicotine on lipid peroxidation and antioxidant defence mechanism in rats," *Journal of Applied Toxicology*, vol. 16, pp. 305–308, 1996.
- [55] J. Zhang, S. Jiang, and R. R. Watson, "Antioxidant supplementation prevents oxidation and inflammatory responses induced by sidestream cigarette smoke in old mice," *Environmental Health Perspectives*, vol. 109, no. 10, pp. 1007–1009, 2001.
- [56] A. M. Gawish, A. M. Issa, N. S. Bassily, and S. M. Manaa, "Role of green tea on nicotine toxicity on liver and lung of mice: histological and morphometrical studies," *African Journal of Biotechnology*, vol. 11, no. 8, pp. 2013–2025, 2012.
- [57] R. Simeonova, V. Vitcheva, G. Gorneva, and M. Mitcheva, "Effects of myosmine on antioxidative defence in rat liver," *Arhiv za Higijenu Rada i Toksikologiju*, vol. 63, no. 1, pp. 7–14, 2012.
- [58] M. Micheva, M. Kondeva-Burdina, and V. Vicheva, "Study on hepatotoxicity of cytisine (Tabex) compared with nicotine in freshly isolated rat hepatocytes," *Pharmacia*, vol. 56, no. 1–4, pp. 27–32, 2009.
- [59] N. Alva, D. Cruz, S. Sanchez, J. Ma Valentin, and T. C. Bermudez, "Nitric oxide as a mediator of fructose 1, 6-bisphosphate protection in galactosamine-induced hepatotoxicity in rats," *Nitric Oxide*, vol. 28, pp. 17–23, 2013.
- [60] J. Das, J. Ghosh, A. Roy, and P. C. Sil, "Mangiferin exerts hepatoprotective activity against D-galactosamine induced acute toxicity and oxidative/nitrosative stress via Nrf2-NFκB pathways," *Toxicology and Applied Pharmacology*, vol. 260, no. 1, pp. 35–47, 2012.
- [61] G. Pushpavalli, C. Veeramani, and K. V. Pugalendi, "Effect of Piper betle on plasma antioxidant status and lipid profile against D-galactosamine-induced hepatitis in rats," *Redox Report*, vol. 14, no. 1, pp. 7–12, 2009.
- [62] R. Labib, R. Turkall, and M. S. Abdel-Rahman, "Oral cocaine produces dose-related hepatotoxicity in male mice," *Toxicology Letters*, vol. 125, no. 1–3, pp. 29–37, 2001.
- [63] R. Labib, R. Turkall, and M. S. Abdel-Rahman, "Inhibition of cocaine oxidative metabolism attenuates endotoxin potentiation of cocaine mediated hepatotoxicity," *Toxicology*, vol. 179, no. 1-2, pp. 9–19, 2002.
- [64] P. Kovacic, "Role of oxidative metabolites of cocaine in toxicity and addiction: oxidative stress and electron transfer," *Medical Hypotheses*, vol. 64, no. 2, pp. 350–356, 2005.
- [65] V. Vitcheva, R. Simeonova, I. Krasteva, M. Yotova, S. Nikolov, and M. Mitcheva, "Hepatoprotective effects of saponarin, isolated from *Gypsophila trichotoma* wend. on cocaine-induced oxidative stress in rats," *Redox Report*, vol. 16, no. 2, pp. 56–61, 2011.
- [66] T. Visalli, R. Turkall, and M. S. Abdel-Rahman, "Cocaine hepatotoxicity and its potentiation by lipopolysaccharide: treatment and gender effects," *International Journal of Toxicology*, vol. 23, no. 3, pp. 163–170, 2004.
- [67] Q.-Y. Cai, H.-B. Chen, S.-Q. Cai et al., "Effect of roots of Ficus hirta on cocaine-induced hepatotoxicity and active components," *Zhongguo Zhongyao Zazhi*, vol. 32, no. 12, pp. 1190–1193, 2007.









Autoimmune Diseases









MEDIATORS INFLAMMATION



Anesthesiology Research and Practice





Emergency Medicine International



BioMed Research International



Pain Research and Treatment





International Journal of Medicinal Chemistry



Journal of Vaccines



