

Retraction

Retracted: Influence of Two Common Polymorphisms in the *EPHX1* Gene on Warfarin Maintenance Dosage: A Meta-Analysis

BioMed Research International

Received 3 February 2019; Accepted 3 February 2019; Published 7 March 2019

Copyright © 2019 BioMed Research International. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

BioMed Research International has retracted the article titled “Influence of Two Common Polymorphisms in the *EPHX1* Gene on Warfarin Maintenance Dosage: A Meta-Analysis” [1]. This article is one of a series of very similar meta-analyses written by different authors which were published in 2014 and 2015, characterized by searching the complementary and alternative medicine database CISCOPM despite the topic not being about CAM [2]. The overlaps of language expression with these articles are concentrated in the Materials and Methods, Results, and Discussion sections. In the article, the restriction to two *EPHX1* polymorphisms was not justified and the change of one of the SNPs from rs2292566 to rs1131873 in 2013 was not mentioned.

The authors clarified that, a few years ago, they took a training course on how to write meta-analyses, in China, where they were provided with some templates. They said that they referred to these templates to write the article and sought help from a native English speaker. In addition, the corresponding author was not aware of the existence of the article before its publication. They confirmed the concerns above and asked to retract the article due to it being flawed.

References

- [1] H.-Q. Liu, C.-P. Zhang, C.-Z. Zhang, X.-C. Liu, and Z.-J. Liu, “Influence of two common polymorphisms in the *EPHX1* gene on warfarin maintenance dosage: a meta-analysis,” *BioMed Research International*, vol. 2015, Article ID 564149, 12 pages, 2015.
- [2] F. Guillaume, “A flurry of copycats on PubMed,” *The Grand Locus*, 2014, <http://blog.thegrandlocus.com/2014/10/a-flurry-of-copycats-on-pubmed>.

Review Article

Influence of Two Common Polymorphisms in the *EPHX1* Gene on Warfarin Maintenance Dosage: A Meta-Analysis

Hong-Qiang Liu,¹ Chang-Po Zhang,² Chang-Zhen Zhang,³
Xiang-Chen Liu,¹ and Zun-Jing Liu^{3,4}

¹Department of Pharmacy, Jining No. 1 People's Hospital, Jining 272011, China

²Jining Hospital of Traditional Chinese Medicine, Jining 272137, China

³Jining Municipal Authority Hospital, Jining, Shandong 272000, China

⁴Department of Neurology, China-Japan Friendship Hospital, Yinghua Road No. 2, Chaoyang District, Beijing 100029, China

Correspondence should be addressed to Zun-Jing Liu; liuzunjing905_lzj@126.com

Received 29 April 2014; Revised 5 September 2014; Accepted 5 September 2014

Academic Editor: Roberto Cirocchi

Copyright © 2015 Hong-Qiang Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We conducted a meta-analysis to investigate the influence of two common single nucleotide polymorphisms (SNPs) (rs2292566 G>A and rs4653436 A>G) in the *EPHX1* gene on warfarin maintenance dosages. Relevant literatures were searched using the *PubMed*, *Embase*, *Web of Science*, *Cochrane Library*, *CISCOM*, *CINAHL*, *Google Scholar*, *CBM*, and *CNKI* databases without any language restrictions. STATA Version 12.0 software (Stata Corporation, College Station, TX, USA) was used for this meta-analysis. Standard mean difference and its corresponding 95% confidence interval (95% CI) were calculated. Seven studies met the inclusion criteria, including 2,063 warfarin-treated patients. Meta-analysis results illustrated that *EPHX1* rs2292566 G>A polymorphism might be strongly correlated with a higher maintenance dose of warfarin. However, no interaction of *EPHX1* rs4653436 A>G polymorphism with warfarin maintenance dosage was detected. A further subgroup analysis based on stratification by ethnicity indicated that *EPHX1* rs2292566 G>A polymorphism was positively correlated with warfarin maintenance dosage among Caucasians, but not Asians. No associations were observed between *EPHX1* rs4653436 A>G polymorphism warfarin maintenance dosage among both Caucasians and Asians. Our meta-analysis provides robust and unambiguous evidence that *EPHX1* rs2292566 polymorphism may affect the maintenance dose of warfarin in Caucasians.

1. Introduction

Warfarin is an anticoagulant normally involved in preventing thrombosis and thromboembolism, which is prescribed for patients with chronic atrial fibrillation, pulmonary embolism, deep vein thrombosis, recurrent stroke, and prosthetic heart valves [1–4]. In clinical practice, warfarin anticoagulant activity should be monitored for the international normalized ratio (INRs) to ensure an appropriate, safe, and efficient dose; incorrect dosage administration may cause a high risk of potentially devastating bleeding and failure of preventing thrombosis [5–7]. Several factors have been reported to influence the variability in warfarin dose, including age, body size, vitamin K intake, interacting medications, and genetic variants [8–11]. A large number of evidences demonstrated

that genotype-guided dosing of warfarin is a widely recognized example of pharmacogenetics, and clinical utility of genetics-guided warfarin initiation would provide safe and optimal anticoagulation therapy [12, 13].

Recent studies suggested that microsomal epoxide hydroxylase 1 (*EPHX1*) may alter the pharmacokinetics and pharmacodynamics of warfarin and have a clinically significant impact on warfarin maintenance dose [14, 15]. *EPHX1* is a critical xenobiotic-metabolizing enzyme, catalyzing both detoxification and bioactivation reactions that direct the disposition of chemical epoxides including the carcinogenic metabolites of several polycyclic aromatic hydrocarbons [16, 17]. To the best of our knowledge, *EPHX1* plays an important role in the majority of xenobiotic metabolisms and ensures widespread defense against potentially genotoxic

epoxide intermediates including vitamin K epoxide, which is attributed to its exceptionally broad substrate selectivity [18].

Human *EPHX1* gene is located in the long arm of chromosome 1q42.1, consisting of 9 exons spanning approximately 35.48 kbps, and it encodes a protein of 455 amino acids [19]. *EPHX1* genetic polymorphisms affect the warfarin maintenance dose and may significantly contribute to interindividual differences in the responses to warfarin [6, 20]. A possible mechanism for the influence of *EPHX1* genetic polymorphisms on warfarin maintenance dosage is that genetic variations in the *EPHX1* gene may be conducive to the weakening of oxidized vitamin K to reduced vitamin K and the decrease of epoxide hydrolase enzyme activity, strongly impacting on the generation of active coagulation factors; therefore, it may be correlated to the efficacy and dose of warfarin [13].

Several common single nucleotide polymorphisms (SNPs) in the *EPHX1* gene have been reported previously for the effects of *EPHX1* genetic polymorphisms on the maintenance doses of warfarin; among these polymorphisms, the most frequent functional polymorphisms are rs4653436 A>G and rs2292566 G>A [14, 21, 22]. Recently, a number of studies have shown that these two common SNPs in the *EPHX1* gene might be major genetic determinants of warfarin dose [6, 23], but the results of other studies have been inconsistent [13, 21]. In view of the conflicting results from previous studies, we performed a meta-analysis of all available data to investigate the influence of *EPHX1* rs2292566 G>A and rs4653436 A>G polymorphisms on warfarin maintenance dosage.

2. Materials and Methods

2.1. Literature Search and Selection Criteria. A comprehensive search for related studies published before March 2014 was conducted on *PubMed*, *Embase*, *Web of Science*, *Cochrane Library*, *CISCOM*, *CINAHL*, *Google Scholar*, *China BioMedicine (CBM)*, and *China National Knowledge Infrastructure (CNKI)* databases. We used a series of keywords and MeSH terms as follows: ["Epoxide Hydrolases" or "EPHX1 protein, human" or "Microsomal Epoxide Hydrolase" or "Styrene Epoxide Hydrolase"] and ["single nucleotide polymorphism" or "SNP" or "polymorphism" or "mutation" or "mutant" or "variation" or "variant"] and ["Warfarin" or "Coumadin" or "Warfarin Potassium" or "Warfarin Sodium"]. There was no language restriction. We also did a manual search of reference lists from potentially relevant articles to identify other potential studies.

The studies which are in accordance with the following criteria were enrolled in the analysis: (1) clinical study focused on the influence of *EPHX1* rs2292566 G>A and rs4653436 A>G polymorphisms on warfarin maintenance dosage, and the warfarin maintenance dose was defined by the international normalized ratio (INR) measurements [24]; (2) all patients should undergo anticoagulation therapy; (3) the data of genotype frequencies and warfarin maintenance dose should be sufficient. Studies were excluded if they do not meet all of these inclusion criteria. If more than one study

by the same author using the same case series was published, not only the study with the largest sample size but also the most recent publication was included. Any disagreements were resolved by discussions and subsequent consensus.

2.2. Data Extraction and Methodological Assessment. According to the standardized form, data extraction from each included study was done by two authors. We evaluated the methodological quality of each included study based on the Newcastle-Ottawa Scale (NOS) criteria [25]. Briefly, the overall star assessed three main categories on the following: (1) subject selection: 0~4; (2) comparability of subject: 0~2; (3) clinical outcome: 0~3. NOS scores ranged from 0 to 9; and a score ≥ 7 indicates a good quality.

2.3. Statistical Analysis. Meta-analysis was performed with the use of the STATA statistical software (Version 12.0, Stata Corporation, College Station, TX, USA). Standardized mean difference (SMD, Cohen's D) and 95% confidence interval (95% CI) were calculated as estimates of relative risk for warfarin maintenance dose under different genetic models. The Z test was used to estimate the statistical significance of pooled SMDs. Heterogeneity among studies was estimated by the Cochran's Q-statistic and I^2 tests [26]. If Q-test shows a $P < 0.05$ or I^2 test exhibits $> 50\%$ which indicates significant heterogeneity, the random effects model was conducted, or else the fixed-effects model was used [27]. Meanwhile, if there was significant heterogeneity, subgroup analysis was performed to find potential explanatory variables. In order to evaluate the influence of single studies on the overall estimate, a sensitivity analysis was performed. Funnel plots and Egger's linear regression test were applied to investigate publication bias [28].

3. Results

3.1. Study Selection and Characteristics of Included Studies. Initially, the highly sensitive search strategy identified 34 articles. We reviewed the titles and abstracts of all articles and excluded 15 articles; full texts were also reviewed and 10 articles were further excluded. One study was also excluded due to the lack of data integrity. Finally, 7 cohort studies with a total of 2,063 subjects met our inclusion criteria for qualitative data analysis [6, 13, 14, 20, 21, 23, 29–38]. Figure 1 shows the selection process of eligible articles. Publication years of the eligible studies ranged from 2010 to 2013. Distribution of the number of topic-related literatures in the electronic database during the last decade is shown in Figure 2. Overall, 3 studies were conducted among Caucasians and 4 studies among Asians. LightSNiP assay, SNaPshot assay, DHPLC, TaqMan assay, and base-quenched probe were used for genotyping. NOS scores of all included studies were ≥ 5 . We summarized the study characteristics and methodological quality in Table 1.

3.2. Quantitative Data Synthesis. Meta-analysis results illustrated that *EPHX1* rs2292566 G>A polymorphism might be




FIGURE 1: Flow chart shows study selection procedure. Seven cohort studies were included in this meta-analysis.

an underlying factor for increased maintenance doses of warfarin with warfarin maintenance dosage higher in individual with *EPHX1* rs2292566 G>A mutation (GG versus AA: SMD = 0.76, 95% CI: 0.47~1.05, and $P < 0.001$; GG versus GA: SMD = 0.43, 95% CI: 0.03~0.83, and $P = 0.035$, resp.), while a similar result was not detected in the comparison of GA versus AA in the *EPHX1* rs2292566 G>A mutation (SMD = 0.31, 95% CI: -0.15~0.76, and $P = 0.186$). However, no significant interaction of *EPHX1* rs4653436 polymorphism with warfarin maintenance dosage was detected (AA versus GG: SMD = -0.01, 95% CI: -0.33~0.32, and $P = 0.974$; AA versus AG: SMD = 0.10, 95% CI: -0.05~0.24, and $P = 0.209$; and AG versus GG: SMD = -0.21, 95% CI: -0.46~0.03, and $P = 0.092$, resp.) (Figure 3).

We also conducted subgroup analyses to investigate the influence of potential factors on individual variability in warfarin dose. As shown in Figure 4, the final triangle is the weighted outcome across groups. Although no significant heterogeneity was found in most parts of those ethnic subgroups, we still continue our subgroup analysis for a secondary verification of our results. Five in all enrolled studies supplied the results of subgroup analysis based on ethnicity, indicating that *EPHX1* rs2292566 G>A polymorphism was associated with warfarin maintenance dosage among Caucasians (GG versus AA: SMD = 0.52,

95% CI: 0.07~1.03, and $P = 0.006$; GG versus GA: SMD = 0.28, 95% CI: 0.07~0.49, and $P = 0.009$, resp.), but not Asians (all $P > 0.05$) (Figure 4). Nevertheless, we observed no associations between *EPHX1* rs4653436 A>G polymorphism and warfarin maintenance dosage among both Caucasians and Asians (all $P > 0.05$). Further subgroup analyses based on sample size and genotyping method revealed significant relationships between *EPHX1* rs2292566 G>A polymorphism and an increased warfarin maintenance dosage in the large-sample-size and TaqMan assay subgroups, but not in the small-sample-size and non-TaqMan assay subgroups (as shown in Table 2). Nevertheless, we also observed no correlations between *EPHX1* rs4653436 A>G polymorphism and warfarin maintenance dosage in all these subgroups (all $P > 0.05$). Sensitivity analysis suggested that no single study could influence the pooled SMDs. Funnel plots demonstrated no evidence of obvious asymmetry existing. The Egger test also did not display strong statistical evidence for publication bias (all $P > 0.05$) (Figure 5).

4. Discussion

EPHX1 is putative subunit of the vitamin K epoxide reductase (VKOR) and suggested to be a new genetic variant




FIGURE 2: The distribution of the number of topic-related literatures in the electronic database during the last decade.

TABLE 1: Baseline characteristics and methodological quality of all included studies.


First author	Year	Country	Ethnicity	Case number	Gender (M/F)	Age (years)	Genotyping method	SNP	NOS score
Özer [20]	2013	Turkey	Asians	107	53/54	53.9 ± 13.6	LightSNiP assay	rs2292566 G>A	6
Liang [6]	2013	China	Asians	300	138/162	47.9 ± 12.5	SNaPshot assay	rs2292566 G>A rs4653436 A>G	8 8
Volcik [37]	2006	China	Asians	217	90/127	51.3 ± 15.0	DHPLC	rs4653436 A>G	7
Ciccacci [21]	2011	Italy	Caucasians	141	78/70	68.2	TaqMan	rs2292566 G>A rs4653436 A>G	6 6
Luo [23]	2010	China	Asians	197	82/115	52.9 ± 11.8	Base- quenched probe	rs2292566 G>A rs4653436 A>G	7 7
Pautas [14]	2010	France	Caucasians	290	69/231	86.7 ± 6.0	TaqMan	rs2292566 G>A	8
Carlquist [13]	2010	USA	Caucasians	168	79/89	71.0 ± 13.0	TaqMan	rs4653436 A>G	7

M: male; F: female; DHPLC: denaturing high performance liquid chromatography; NOS: Newcastle-Ottawa Scale criteria; SNP: single nucleotide polymorphisms.


affecting the warfarin maintenance dose significantly [14, 20]. Generally, through interference with the recycling of vitamin K in the liver, warfarin acts and leads to the secretion of inactive vitamin K-dependent proteins [39]. Warfarin and this vitamin K clotting factors participated in the process of biotransformation formed warfarin interactive pathways [40]. In such progression, vitamin K hydroquinone is oxidized to vitamin K epoxide which is recycled by VKOR to vitamin K and in turn is reduced to vitamin K hydroquinone stimulated by VKOR complex and epoxide hydrolase [41]. Within the vitamin K redox cycle, warfarin suppresses the action of VKOR enzyme complex, partially blocking cycle activity, resulting in the accumulated inactive hepatic vitamin K₁ 2,3-epoxide and reduced vitamin K depletion [42]. Since this form of vitamin K is an essential cofactor for γ -carboxylation of vitamin K-dependent clotting

factors, its depletion may lead to a reduction of the active clotting factors [43]. As a matter of fact, a changed dosage intake of the fat-soluble vitamin K can reverse the action of warfarin [44]. Therefore, *EPHX1* can be regarded as the other components of VKOR complex involved in the redox processes, so *EPHX1* genetic variations may be crucial candidates in the influence of warfarin anticoagulant effect [45].


In the present meta-analysis, we investigated the relationship between two common SNPs (rs2292566 G>A and rs4653436 A>G) in the *EPHX1* gene and the warfarin dose requirement. Our findings showed a strong association between *EPHX1* rs2292566 G>A polymorphism and warfarin maintenance dose, but similar association was not observed in the *EPHX1* rs4653436 A>G polymorphism, implicating that *EPHX1* rs2292566 polymorphism may be a significant



(a)



(b)



(c)

FIGURE 3: Forest plots for the relationships of *EPHX1* rs2292566 G>A and rs4653436 A>G polymorphisms with warfarin maintenance dosage. The final triangle is the weighted outcome across groups.

TABLE 2: Meta-analysis of the associations between *EPHX1* rs2292566 G>A polymorphism and warfarin maintenance dosage.

	GG versus AA			GG versus GA			GA versus AA								
	SMD	95% CI	P	P_h	I^2	SMD	95% CI	P	P_h	I^2	SMD	95% CI	P	P_h	I^2
Overall	0.76	0.47~1.05	<0.001	0.699	0.0%	0.43	0.03~0.83	0.035	<0.001	87.6%	0.31	-0.15~0.76	0.186	0.105	47.7%
Ethnicity															
Caucasians	0.52	0.07~1.03	0.006	0.716	0.0%	0.28	0.07~0.49	0.009	0.550	0.0%	0.40	-0.25~1.06	0.226	0.028	72.0%
Asians	0.84	-0.13~1.15	0.699	0.895	0.0%	0.54	-0.14~1.23	0.122	<0.001	92.6%	0.05	-0.72~0.82	0.898	0.830	0.0%
Sample size															
Small ($n < 300$)	0.71	-0.01~1.10	0.310	0.462	0.0%	0.24	-0.13~0.60	0.210	0.058	64.8%	-0.05	-0.46~0.36	0.815	0.724	0.0%
Large ($n \geq 300$)	0.82	0.40~1.24	<0.001	0.479	0.0%	0.67	0.01~1.34	0.042	<0.001	93.0%	0.62	0.17~1.07	0.007	0.334	8.7%
Genotyping method															
TaqMan assay	0.78	0.02~1.03	0.465	0.716	0.0%	0.28	0.07~0.49	0.009	0.550	92.6%	0.40	-0.25~1.06	0.226	0.028	72.0%
Non-TaqMan assay	0.84	0.00~1.15	0.051	0.895	0.0%	0.54	-0.14~1.23	0.122	<0.001	0.0%	0.05	-0.72~0.82	0.898	0.830	0.0%

EPHX1: epoxide hydrolase 1; SMD: standardized mean difference; 95% CI: 95% confidence interval; P_h : the P value for the heterogeneity test.




FIGURE 4: Continued.




FIGURE 4: Subgroup analyses for the relationships between *EPHX1* rs2292566 G>A polymorphism and warfarin maintenance dosage. The final triangle is the weighted outcome across groups.




FIGURE 5: Funnel plot of publication biases for the relationships between *EPHX1* rs2292566 G>A polymorphism and warfarin maintenance dosage. The final triangle is the weighted outcome across groups.

predictor for the interindividual variability of warfarin maintenance dose. Nevertheless, the precise mechanisms that *EPHX1* genetic polymorphisms affect the requirement of warfarin dose are still unidentified. One possible explanation could be that genetic mutations in the *EPHX1* gene may result in amino acid substitution and have some impacts on the *EPHX1* enzyme activity, with resultant impaired

warfarin metabolism and clearance, thus contributing to interindividual dose variability [13, 46]. Pautas et al. have identified *EPHX1* rs2292566 G>A polymorphism as a novel predicting factor for variable warfarin response, reporting that alteration in the *EPHX1* gene may change the pharmacokinetics and pharmacodynamics of warfarin exertion by inhibiting the activity of vitamin K epoxide reductase

(VKOR), and thereby may influence warfarin maintenance dose [14]. Loebstein et al. have revealed that VKOR was involved in vitamin K redox cycle, and VKOR plays a crucial role in promoting inactive vitamin K into active vitamin K, which is an essential cofactor for γ -carboxylation of vitamin K-dependent clotting factors in the hepatic system (II, VII, IX, and X) [42]. Furthermore, a previous study identified *EPHX1* genetic variant as a predictor of variable warfarin dose requirement because *EPHX1* has been proposed as a putative subunit of VKOR [44]. Therefore, it seems reasonable to hypothesize that *EPHX1* rs2292566 G>A polymorphism may reduce VKOR activity and consequently give rise to active vitamin K deficiency, which may lead to a decrease of the active clotting factors. Additionally, it is plausible that patients with *EPHX1* rs2292566 polymorphism may require a lower maintenance dose of warfarin. Furthermore, it should be noted that out of those six forest plots, there were only significant relationships between the comparison of GG versus AA and GG versus GA; no potential association was detected from the GA versus AA comparison in the *EPHX1* rs2292566 G>A polymorphism; one possible explanation is that the existence of possible heterogeneity sources could have an influence on the overall outcome, so no significant relationship was observed between dose variations and certain allele types for the differences in ages or ethnic backgrounds [20]. Our results are in line with a recent cohort study, which genotyped 107 patients who had stable doses and INRs at their last three consecutive visits and displayed that *EPHX1* rs2292566 G>A polymorphism was significantly associated with the warfarin maintenance dose, accounting for 1.7% of the variability in the dose [20].

Considering the possibility of existing obvious heterogeneity, which may negatively affect our association study results, stratified analyses were carefully performed based on ethnicity, genotyping method, and sample size. Subgroup analysis after the heterogeneity test was also a required step for the secondary verification of our results. The results of subgroup analysis performed by ethnicity displayed significant associations between *EPHX1* rs2292566 G>A polymorphism and a lower maintenance dose of warfarin in Caucasians, while no similar association was detected among Asians, implicating that ethnicity differences may play an important role in the effects of variants in the *EPHX1* gene on interindividual variability of warfarin maintenance dose. Although the potential mechanism of ethnicity differences is still not fully understood, we supposed that ethnicity may result in differences in alleles and genotypes among different ethnic populations. Sample size within those included papers were obviously different, and this stratified analysis revealed that a significant difference between *EPHX1* rs2292566 G>A polymorphism and warfarin maintenance dose was observed within the larger sample sizes. In summary, our findings are consistent with the previous studies that *EPHX1* rs2292566 G>A polymorphism may influence the warfarin dose requirement, suggesting that translation of this knowledge into clinical guidelines may offer a useful and informative route to improve therapeutic management during warfarin therapy.

The current meta-analysis also had several limitations that should be acknowledged. First, our results lacked sufficient statistical power to assess the correlations of *EPHX1* genetic polymorphisms with the warfarin dose requirements. Secondly, meta-analysis is a retrospective study that may inevitably induce subject selection bias and thereby have an impact on the reliability of our results. Thirdly, our meta-analysis failed to obtain original data from the included studies, which may limit further evaluation of potential role of *EPHX1* genetic polymorphisms with the warfarin dose requirements.

In conclusion, our meta-analysis provides robust and unambiguous evidence that *EPHX1* rs2292566 G>A polymorphism may affect the maintenance dose of warfarin in Caucasians, so *EPHX1* rs2292566 G>A polymorphism could be a potential and practical biomarker for the interindividual variability of warfarin maintenance dose. However, due to the limitations mentioned above, more reliable research with larger sample sizes is still required to provide a more comprehensive and representative statistical analysis precisely.

Disclosure

I would like to declare on behalf of my coauthors that the work described herein was original research that will not be submitted elsewhere and not under consideration for publication elsewhere.

Conflict of Interests

The authors declare that no competing interests exist.

Authors' Contribution

Both Hong-Qiang Liu and Xiang-Chen Liu are cofirst authors. This paper is approved by all authors for publication.

Acknowledgment

The authors would like to acknowledge the reviewers for their helpful comments on this paper.

References

- [1] C. S. Miller, S. M. Grandi, A. Shimony, K. B. Filion, and M. J. Eisenberg, "Meta-analysis of efficacy and safety of new oral anticoagulants (dabigatran, rivaroxaban, apixaban) versus warfarin in patients with atrial fibrillation," *The American Journal of Cardiology*, vol. 110, no. 3, pp. 453–460, 2012.
- [2] J. A. Heit, B. D. Lahr, T. M. Petterson, K. R. Bailey, A. A. Ashrani, and L. J. Melton III, "Heparin and warfarin anticoagulation intensity as predictors of recurrence after deep vein thrombosis or pulmonary embolism: a population-based cohort study," *Blood*, vol. 118, no. 18, pp. 4992–4999, 2011.
- [3] G. J. Hankey, M. R. Patel, S. R. Stevens et al., "Rivaroxaban compared with warfarin in patients with atrial fibrillation and previous stroke or transient ischaemic attack: a subgroup analysis of ROCKET AF," *The Lancet Neurology*, vol. 11, no. 4, pp. 315–322, 2012.

- [4] C. S. Wong, K. Batchelor, J. Bua, and F. Newall, "Safety and efficacy of warfarin in paediatric patients with prosthetic cardiac valves: a retrospective audit," *Thrombosis Research*, vol. 128, no. 4, pp. 331–334, 2011.
- [5] D. T. Le, R. T. Weibert, B. K. Sevilla, K. J. Donnelly, and S. I. Rapaport, "The international normalized ratio (INR) for monitoring warfarin therapy: reliability and relation to other monitoring methods," *Annals of Internal Medicine*, vol. 120, no. 7, pp. 552–558, 1994.
- [6] Y. Liang, Z. Chen, G. Guo et al., "Association of genetic polymorphisms with warfarin dose requirements in chinese patients," *Genetic Testing and Molecular Biomarkers*, vol. 17, no. 12, pp. 932–936, 2013.
- [7] D. Keeling, T. Baglin, C. Tait et al., "Guidelines on oral anticoagulation with warfarin—fourth edition," *British Journal of Haematology*, vol. 154, no. 3, pp. 311–324, 2011.
- [8] J. F. Carlquist and J. L. Anderson, "Using pharmacogenetics in real time to guide warfarin initiation: a clinician update," *Circulation*, vol. 124, no. 23, pp. 2554–2559, 2011.
- [9] T. T. Biss, A. J. Adamson, C. J. Seal, and F. Kamali, "The potential impact of dietary vitamin K status on anticoagulation control in children receiving warfarin," *Pediatric Hematology and Oncology*, vol. 28, no. 5, pp. 425–427, 2011.
- [10] E. Nutescu, I. Chuatrisorn, and E. Hellenbart, "Drug and dietary interactions of warfarin and novel oral anticoagulants: an update," *Journal of Thrombosis and Thrombolysis*, vol. 31, no. 3, pp. 326–343, 2011.
- [11] L. H. Cavallari, T. Y. Langaee, K. M. Momary et al., "Genetic and clinical predictors of warfarin dose requirements in African Americans," *Clinical Pharmacology and Therapeutics*, vol. 87, no. 4, pp. 459–464, 2010.
- [12] I. Y. Gong, R. G. Tirona, U. I. Schwarz et al., "Prospective evaluation of a pharmacogenetics-guided warfarin loading and maintenance dose regimen for initiation of therapy," *Blood*, vol. 118, no. 11, pp. 3163–3171, 2011.
- [13] J. F. Carlquist, B. D. Horne, C. Mower et al., "An evaluation of nine genetic variants related to metabolism and mechanism of action of warfarin as applied to stable dose prediction," *Journal of Thrombosis and Thrombolysis*, vol. 30, no. 3, pp. 358–364, 2010.
- [14] E. Pautas, C. Moreau, I. Gouin-Thibault et al., "Genetic factors (VKORC1, CYP2C9, EPHX1, and CYP4F2) are predictor variables for warfarin response in very elderly, frail inpatients," *Clinical Pharmacology and Therapeutics*, vol. 87, no. 1, pp. 57–64, 2010.
- [15] S. L. Chan, A. Thalamuthu, B. C. Goh et al., "Exon sequencing and association analysis of EPHX1 genetic variants with maintenance warfarin dose in a multiethnic Asian population," *Pharmacogenetics and Genomics*, vol. 21, no. 1, pp. 35–41, 2011.
- [16] W. Zhao, J. Luo, and X. Cai, "Association between microsomal epoxide hydrolase 1 polymorphisms and susceptibility to esophageal cancer: a meta-analysis," *Tumor Biology*, vol. 34, no. 4, pp. 2383–2388, 2013.
- [17] H. Nisa, S. Budhathoki, M. Morita et al., "Microsomal epoxide hydrolase polymorphisms, cigarette smoking, and risk of colorectal cancer: the Fukuoka colorectal cancer study," *Molecular Carcinogenesis*, vol. 52, no. 8, pp. 619–626, 2013.
- [18] T. B. Tumer, G. Sahin, and E. Arinç, "Association between polymorphisms of EPHX1 and XRCC1 genes and the risk of childhood acute lymphoblastic leukemia," *Archives of Toxicology*, vol. 86, no. 3, pp. 431–439, 2012.
- [19] J. K. Hartsfield Jr., M. J. Sutcliffe, E. T. Everett, C. Hassett, C. J. Omiecinski, and J. A. Saari, "Assignment of microsomal epoxide hydrolase (EPHX1) to human chromosome 1q42.1 by in situ hybridization," *Cytogenetics and Cell Genetics*, vol. 83, no. 1–2, pp. 44–45, 1998.
- [20] M. Özer, Y. Demirci, C. Hizel et al., "Impact of genetic factors (CYP2C9, VKORC1 and CYP4F2) on warfarin dose requirement in the Turkish population," *Basic & Clinical Pharmacology & Toxicology*, vol. 112, no. 3, pp. 209–214, 2013.
- [21] C. Ciccacci, N. Paolillo, D. Di Fusco, G. Novelli, and P. Borgiani, "EPHX1 polymorphisms are not associated with warfarin response in an Italian population," *Clinical Pharmacology and Therapeutics*, vol. 89, no. 6, p. 791, 2011.
- [22] Q. Gu, Y. Kong, J. Schneede et al., "VKORC1-1639G>A, CYP2C9, EPHX1691A>G genotype, body weight, and age are important predictors for warfarin maintenance doses in patients with mechanical heart valve prostheses in southwest China," *European Journal of Clinical Pharmacology*, vol. 66, no. 12, pp. 1217–1227, 2010.
- [23] W.-W. Luo, H.-V. Zhang, S.-M. Ma, L.-H. Xu, Z.-X. Zhang, and X.-Q. Yang, "Effect of the epoxide hydrolase 1 gene polymorphism on warfarin dose," *Chinese Pharmaceutical Journal*, vol. 45, no. 19, pp. 1489–1491, 2010.
- [24] G. Introcaso and G. Gesu, "Significance of consecutive international normalized ratio (INR) outcomes using statistical control rules in long-term anticoagulated patients. Optimization of laboratory monitoring and interpretation of borderline measurements," *Clinical Chemistry and Laboratory Medicine*, vol. 42, no. 3, pp. 294–299, 2004.
- [25] A. Stang, "Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses," *European Journal of Epidemiology*, vol. 25, no. 9, pp. 603–605, 2010.
- [26] E. Zintzaras and J. P. A. Ioannidis, "HEGESMA: genome search meta-analysis and heterogeneity testing," *Bioinformatics*, vol. 21, no. 18, pp. 3672–3673, 2005.
- [27] E. Zintzaras and J. P. A. Ioannidis, "Heterogeneity testing in meta-analysis of genome searches," *Genetic Epidemiology*, vol. 28, no. 2, pp. 123–137, 2005.
- [28] J. L. Peters, A. J. Sutton, D. R. Jones, K. R. Abrams, and L. Rushon, "Comparison of two methods to detect publication bias in meta-analysis," *Journal of the American Medical Association*, vol. 295, no. 6, pp. 676–680, 2006.
- [29] S. C. Barbaux, S. Blankenberg, H. J. Rupprecht et al., "Association between P-selectin gene polymorphisms and soluble P-selectin levels and their relation to coronary artery disease," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 21, no. 10, pp. 1668–1673, 2001.
- [30] A. M. Carter, K. Anagnostopoulou, M. W. Mansfield, and P. J. Grant, "Soluble P-selectin levels, P-selectin polymorphisms and cardiovascular disease," *Journal of Thrombosis and Haemostasis*, vol. 1, no. 8, pp. 1718–1723, 2003.
- [31] E. Elmas, P. Bugert, T. Popp et al., "The P-selectin gene polymorphism val168Met: a novel risk marker for the occurrence of primary ventricular fibrillation during acute myocardial infarction," *Journal of Cardiovascular Electrophysiology*, vol. 21, no. 11, pp. 1260–1265, 2010.
- [32] L. Ghazouani, N. Abboud, S. B. Khalifa et al., "Contribution of SELP and PSGL-1 genotypes and haplotypes to the presence of coronary heart disease in Tunisians," *Molecular Biology Reports*, vol. 38, no. 1, pp. 495–501, 2011.

- [33] C. L. Liao, M. Z. Huang, H. Y. Liu, D. Liang, and G. Z. Pam, "Association of the genotype and serum level of P-selectin with acute myocardial infarction," *Chinese Journal of Arteriosclerosis*, vol. 19, pp. 151–154, 2011.
- [34] F. Kee, G. F. Baxter, C. Morrison et al., "Polymorphisms of the P-selectin gene and risk of myocardial infarction in men and women in the ECTIM extension study," *Heart*, vol. 84, no. 5, pp. 548–552, 2000.
- [35] C. L. Liao, M. Z. Huang, H. Y. Liu, D. Liang, and G. Z. Pan, "Association P-selectin gene polymorphisms and haplotype with coronary heart disease," *Journal of Guangxi Medical University*, vol. 29, pp. 598–600, 2012.
- [36] S. A. Rosner, P. M. Ridker, R. Y. L. Zee, and N. R. Cook, "Interaction between inflammation-related gene polymorphisms and cigarette smoking on the risk of myocardial infarction in the Physician's Health Study," *Human Genetics*, vol. 118, no. 2, pp. 287–294, 2005.
- [37] K. A. Volcik, C. M. Ballantyne, J. Coresh, A. R. Folsom, K. K. Wu, and E. Boerwinkle, "P-selectin Thr715Pro polymorphism predicts P-selectin levels but not risk of incident coronary heart disease or ischemic stroke in a cohort of 14 595 participants: the Atherosclerosis Risk in Communities Study," *Atherosclerosis*, vol. 186, no. 1, pp. 74–79, 2006.
- [38] S. W. Huang, D. K. Xiang, B. L. Chen, L. Huang, B. Q. An, and G. F. Li, "Correlation between EPHX1 polymorphism and Warfarin maintenance dosage," *Tianjin Medical Journal*, vol. 39, pp. 887–889, 2011.
- [39] B. Dahlbäck, "Blood coagulation and its regulation by anti-coagulant pathways: genetic pathogenesis of bleeding and thrombotic diseases," *Journal of Internal Medicine*, vol. 257, no. 3, pp. 209–223, 2005.
- [40] M. Wadelius and M. Pirmohamed, "Pharmacogenetics of warfarin: current status and future challenges," *Pharmacogenomics Journal*, vol. 7, no. 2, pp. 99–111, 2007.
- [41] J. Oldenburg, M. Marinova, C. Müller-Reible, and M. Watzka, "The vitamin K cycle," *Vitamins and Hormones*, vol. 78, pp. 35–62, 2008.
- [42] R. Loebstein, M. Vecsler, D. Kurnik et al., "Common genetic variants of microsomal epoxide hydrolase affect warfarin dose requirements beyond the effect of cytochrome P450 2C9," *Clinical Pharmacology & Therapeutics*, vol. 77, no. 5, pp. 365–372, 2005.
- [43] M. Vecsler, R. Loebstein, S. Almog et al., "Combined genetic profiles of components and regulators of the vitamin K-dependent γ -carboxylation system affect individual sensitivity to warfarin," *Thrombosis and Haemostasis*, vol. 95, no. 2, pp. 205–211, 2006.
- [44] M. Wadelius, L. Y. Chen, N. Eriksson et al., "Association of warfarin dose with genes involved in its action and metabolism," *Human Genetics*, vol. 121, no. 1, pp. 23–34, 2007.
- [45] M. T. Lee, C. H. Chen, C. H. Chou et al., "Genetic determinants of warfarin dosing in the Han-Chinese population," *Pharmacogenomics*, vol. 10, pp. 1905–1913, 2009.
- [46] M. S. M. Issac, M. S. El-Nahid, and M. Y. Wissa, "Is there a role for MDRI, EPHX1 and Protein Z gene variants in modulation of warfarin dosage? A study on a cohort of the Egyptian population," *Molecular Diagnosis and Therapy*, vol. 18, no. 1, pp. 73–83, 2014.