Letter to the Editor

Comment on “Using Innovative Acoustic Analysis to Predict the Postoperative Outcomes of Unilateral Vocal Fold Paralysis”

Andrea Lovato and Cosimo de Filippis

Department of Neuroscience, University of Padova, Audiology Unit at Treviso Hospital, Treviso, Italy

Correspondence should be addressed to Andrea Lovato; andrea.lovato.3@hotmail.it

Received 13 July 2017; Accepted 28 December 2017; Published 15 March 2018

Academic Editor: Gjurmakch Aliev

Copyright © 2018 Andrea Lovato and Cosimo de Filippis. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We read with interest the study by Tsou et al. [1] comparing an innovative acoustic measure, that is, the number of irregular peaks (NIrrP), with jitter and shimmer as measured by the Multi-Dimensional Voice Program (MDVP); the authors wanted to verify the ability of these three acoustic measures to predict the surgical outcome of fat injection laryngoplasty in unilateral vocal fold paralysis. Using the receiver operating characteristic (ROC) curve, they found that NIrrP [Area Under the ROC Curve (AUC) = 0.98, 95% confidence interval (CI) = 0.78–0.99] was a more accurate predictor of long-term surgical outcome than jitter (AUC = 0.73, 95% CI = 0.47–0.91) and shimmer (AUC = 0.63, 95% CI = 0.37–0.85) [1].

AUC is frequently used to test the ability of clinical variables to predict the outcome in various clinical settings, especially to identify patients at a higher risk of carcinoma recurrence [2–4]. Furthermore, the efficacy of the discrimination could be measured with the scale proposed by Hosmer and Lemeshow [5, 6]. This scale is used to evaluate the discriminatory power of statistical models as follows: AUC equal to 0.5, no discrimination; AUC between 0.7 and 0.8, acceptable discrimination; AUC between 0.8 and 0.9, excellent discrimination; AUC beyond 0.90, outstanding discrimination [5, 7].

In 2016, we used the Hosmer and Lemeshow scale to compare the discriminatory power of two pieces of voice analysis software, MDVP and Praat, in distinguishing the gender of euphonic adults [8]. We found no discrimination for shimmer in dB with either MDVP (AUC = 0.658) or

Praat (AUC = 0.682); on the other hand, MDVP absolute jitter achieved an acceptable discrimination between males and females (AUC = 0.752), and Praat absolute jitter achieved an outstanding discrimination (AUC = 0.901) [8]. Using the same scale on Tsou et al.'s results [1], we could say that the innovative acoustic measure NIrrP showed outstanding discriminatory power (AUC = 0.98) in identifying unilateral vocal fold paralysis patients at a higher risk of surgical failure after fat injection laryngoplasty. Research on this and other predictors of surgical failure in phoniatry should be encouraged.

Conflicts of Interest

This was not an industry-supported study. The authors have no conflicts of interest to disclose.

References

- [1] Y.-A. Tsou, Y.-W. Liu, W.-D. Chang et al., “Using Innovative Acoustic Analysis to Predict the Postoperative Outcomes of Unilateral Vocal Fold Paralysis,” *BioMed Research International*, vol. 2016, Article ID 7821415, 2016.
- [2] J. V. Carter, J. Pan, S. N. Rai, and S. Galandiuk, “ROC-ing along: Evaluation and interpretation of receiver operating characteristic curves,” *Surgery*, vol. 159, no. 6, pp. 1638–1645, 2016.
- [3] A. Lovato, M. Lionello, A. Staffieri et al., “A higher angiogenin expression is associated with a nonnuclear maspin location in

- laryngeal carcinoma,” *Clinical and Experimental Otorhinolaryngology*, vol. 8, no. 3, pp. 268–274, 2015.
- [4] G. Marioni, S. Blandamura, M. Lionello et al., “Indications for postoperative radiotherapy in laryngeal carcinoma: A panel of tumor tissue markers for predicting locoregional recurrence in surgically treated carcinoma. A pilot study,” *Head & Neck*, vol. 36, no. 11, pp. 1534–1540, 2014.
 - [5] D. W. Hosmer and S. Lemeshow, “Applied Logistic Regression,” in *John Wiley & Sons, Inc*, pp. 156–164, 2000.
 - [6] A. Lovato, G. Marioni, E. Manzato et al., “Elderly patients at higher risk of laryngeal carcinoma recurrence could be identified by a panel of two biomarkers (nm23-H1 and CD105) and pN+ status,” *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*, vol. 272, no. 11, pp. 3417–3424, 2015.
 - [7] G. Marioni, M. Agostini, R. Cappellesso et al., “MIR-19a and SOCS-1 expression in the differential diagnosis of laryngeal (glottic) verrucous squamous cell carcinoma,” *Journal of Clinical Pathology*, vol. 69, no. 5, pp. 415–421, 2016.
 - [8] A. Lovato, W. De Colle, L. Giacomelli et al., “Multi-Dimensional Voice Program (MDVP) vs Praat for Assessing Euphonic Subjects: A Preliminary Study on the Gender-discriminating Power of Acoustic Analysis Software,” *Journal of Voice*, vol. 30, no. 6, pp. 765–765.e5, 2016.