Corrigendum

Corrigendum to “The Effect of Botulinum Toxin Type A on Expression Profiling of Long Noncoding RNAs in Human Dermal Fibroblasts”

**Ying-Ying Miao,¹ Juan Liu,¹ Jie Zhu,² Yan-Ling Tao,¹ Jia-An Zhang,¹ Dan Luo ¹,
and Bing-Rong Zhou ¹**

¹Department of Dermatology, The First Affiliated Hospital of Nanjing Medical University, Nanjing 210029, China

²Department of Dermatology, The First Affiliated Hospital of Nanjing University of TCM, Nanjing, Jiangsu 210029, China

Correspondence should be addressed to Dan Luo; daniluo2005@163.com

Received 28 October 2019; Accepted 31 October 2019; Published 26 November 2019

Copyright © 2019 Ying-Ying Miao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “The Effect of Botulinum Toxin Type A on Expression Profiling of Long Noncoding RNAs in Human Dermal Fibroblasts” [1], there was an error in Figure 4, where the second and third graphs of Figure 4(a) were mistakenly duplicated during the production process. The correct figure is shown below.

(a)

FIGURE 4: Continued.

FIGURE 4: Bioinformatic analysis of the differentially expressed genes. The p value denotes the significance of GO terms enrichment in the differentially expressed genes. The lower the p value, the more significant the GO term (p value ≤ 0.05 is recommended). We can choose the target genes for further study based on the combination of the analysis provided by GO and the biologic significance.

References

- [1] Y.-Y Miao, J. Liu, J. Zhu et al., "The effect of botulinum toxin type A on expression profiling of long noncoding RNAs in human dermal fibroblasts," *BioMed Research International*, vol. 2017, Article ID 2957941, 13 pages, 2017.