Research Article

Lactobacillus iners Is Associated with Vaginal Dysbiosis in Healthy Pregnant Women: A Preliminary Study

Nengneng Zheng ¹, Renyong Guo ², Yinyu Yao ¹, Meiyuan Jin ¹, Yiwen Cheng ³,
and Zongxin Ling ³

¹Department of Obstetrics, Tongde Hospital of Zhejiang Province, Hangzhou, Zhejiang 310012, China

²Department of Laboratory Medicine, First Affiliated Hospital, College of Medicine, Zhejiang University, Key Laboratory of Clinical In Vitro Diagnostic Techniques of Zhejiang Province, Hangzhou, Zhejiang 310003, China

³Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, National Clinical Research Center for Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou, Zhejiang 310003, China

Correspondence should be addressed to Zongxin Ling; lingzongxin@zju.edu.cn

Received 31 May 2019; Accepted 3 October 2019; Published 23 October 2019

Academic Editor: Pengjun Shi

Copyright © 2019 Nengneng Zheng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Vaginal dysbiosis has been identified to be associated with adverse pregnancy outcomes, such as preterm delivery and premature rupture of membranes. However, the overall structure and composition of vaginal microbiota in different trimesters of the pregnant women has not been fully elucidated. In this study, the physiological changes of the vaginal microbiota in healthy pregnant women were investigated. A total of 83 healthy pregnant participants were enrolled, who are in the first, second, or third pregnancy trimester. Quantitative real-time PCR was used to explore the abundant bacteria in the vaginal microbiota. No significant difference in the abundance of *Gardnerella*, *Atopobium*, *Megasphaera*, *Eggerthella*, *Leptotrichia/Sneathia*, or *Prevotella* was found among different trimesters, except *Lactobacillus*. Compared with the first pregnancy trimester, the abundance of *L. iners* decreased in the second and third trimester while the abundance of *L. crispatus* was increased in the second trimester. Moreover, we also found that vaginal cleanliness is correlated with the present of *Lactobacillus*, *Atopobium*, and *Prevotella* and leukocyte esterase is associated with *Lactobacillus*, *Atopobium*, *Gardnerella*, *Eggerthella*, *Leptotrichia/Sneathia*, and *Prevotella*. For those whose vaginal cleanliness raised or leukocyte esterase became positive, the richness of *L. iners* increased, while that of *L. crispatus* decreased significantly. Our present data indicated that the altered vaginal microbiota, mainly *Lactobacillus*, could be observed among different trimesters of pregnancy and *L. iners* could be considered as a potential bacterial marker for evaluating vaginal cleanliness and leukocyte esterase.

1. Introduction

The vaginal microbiota consist of a variety of species, including both anaerobic and aerobic microorganisms [1]. These commensal microorganisms in the vagina provide beneficial effects against opportunistic and pathogenic bacteria, constituting the first line of defense against invasive microorganisms [2]. Some of the species such as *L. crispatus*, *L. gasseri*, and *L. jensenii* take function by lowering the pH and production of large amounts of lactic acid and bactericide compounds, such as hydrogen

peroxide (H₂O₂) and bacteriocins [3, 4], which prevent possible pathogenic or opportunistic pathogenic bacteria and ensure vaginal epithelial homeostasis.

Preterm delivery is known to be one of the main causes of perinatal mortality and morbidity worldwide [5]. Generally, it will result in very low birth weight, prolonged stay in hospital, increased risk of chronic lung disease, and even cerebral palsy. Besides, premature rupture of membranes (PROM) is also a dangerous complication. In the event of PROM, the uterine cavity, placenta, and fetus are exposed to ascending infection

and increased risk of chorioamnionitis and funisitis, which are both associated with poor maternal and neonatal outcomes [6–10]. Recently, lower genital tract infection has been recognized as one of the more important risk factors associated with preterm delivery and PROM, especially in healthy nulliparous women [11]. Dysbiosis of vaginal microbiome has been recognized as a potential cause of adverse pregnancy outcomes, if they are not dominated by *Lactobacillus* [12–15]. However, the influence of vaginal dysbiosis on preterm delivery and PROM still remains unknown.

Healthy pregnancy is characterized by a temporary dynamic shift towards stable, reduced richness and low diversity in the community structures dominated by *Lactobacillus* spp. which aids in the prevention of pathogenic bacteria, different from the vaginal microbiota composition of nonpregnant women [16, 17]. Bacterial vaginosis (BV) is a condition in which the vaginal microbiota suffer a reduction in several species of probiotic *Lactobacillus* and an increase in the presence of anaerobes (*Gardnerella vaginalis*, *Atopobium vaginae*, and *Mobiluncus* sp.), which has been shown in our previous studies [18, 19]. It is hypothesized that colonization of the pathogenic bacteria in the vagina activates the local and upper (cervical and fetal membrane) innate immune system, drives an inflammatory cascade, and leads to a remodeling and disruption of membrane architecture and preterm delivery or PROM [20, 21].

Currently, cultivation-independent molecular approaches are favored for overcoming the difficult cultivation conditions such as nutrition and anaerobic requirements [22]. Thus, they have been used to demonstrate greater vaginal microbial diversity than that recognized previously. Several cultivation-independent species, such as *L. iners* and *A. vaginae*, have been identified as important elements of the vaginal microbiota as they are found across *Lactobacillus*-dominated and non-*Lactobacillus*-dominated vaginal microbiota groups [23, 24]. Thus, a deeper understanding of the composition and structure of the vaginal microbiota in healthy pregnant women is essential for fully elucidating the etiology of vaginal diseases as well as for the prevention and treatment of such diseases. Furthermore, there is an urgent need for a better understanding of when the clinical relevance of some bacteria became increasingly apparent, due to a change in either relative or absolute abundance.

The aim of this study was to investigate the vaginal microbiota in the first, second, and third pregnancy trimester in healthy pregnant women using a cultivation-independent approach. The exploration of the vaginal microbiota in these women provides a theoretical basis that may help to prevent vaginal infections during pregnancy and reduce preterm delivery and PROM.

2. Materials and Methods

2.1. Subjects' Recruitment. Healthy pregnant women, without abnormal vaginal discharge, were randomly enrolled, when they came to the Department of Obstetrics of Tongde Hospital in Zhejiang Province (Zhejiang, China) for routine medical examination between July and November 2016. A total of 83 subjects, aged 30.0 ± 4.6 years, were studied across

three time points, with 33 between 5 and 10 weeks' gestation in the first trimester, 24 between 20 and 27 weeks' gestation in the second trimester, and 26 between 37 and 39 weeks' gestation in the third trimester. The following criteria were used to exclude subjects: a history of preterm delivery; previous multiple pregnancies; placenta previa or vaginal bleeding; candidiasis, BV, or trichomoniasis; use of antibiotics, probiotics, prebiotics, or synbiotics in the previous month; cervical incompetence or suspected uterine malformation; vaginal intercourse within the last 3 days; or were immunocompromised. Informed written consent was obtained from each of the participants before enrollment. This research was conducted in agreement with the ethical principles of the Declaration of Helsinki, and the study protocol was approved by the Ethics Committee of the Tongde Hospital of Zhejiang Province.

2.2. Sample Collection. Three swabs of vaginal secretions were taken from the posterior wall of the vaginal fornix at an outpatient service when these participants underwent speculum examination. Two of the swabs were applied onto a slide for determination of vaginal pH and vaginal infection status. The remaining vaginal swab was covered, placed on ice, and used for bacterial genomic DNA extraction. Vaginal pH was measured using a pH strip (Sanaisi Company, Shanghai, China). Vaginal infection status was determined by examining a wet mount smear in potassium hydroxide for detection of candidiasis and in saline for detection of motile trichomonas and clue cells. BV diagnosis was assessed according to Amsel clinical criteria [25]. H_2O_2 and leukocyte esterase were analyzed according to the manual method of the BV union diagnosis kits (Lizhu Company, Zhuhai, Guangdong, China). Vaginal cleanliness was evaluated according to morphological observations (Table S1) [26]. The criteria of vaginal cleanliness were as follows [27]: Grade I was a large number of large Gram-positive rods (indicative of *Lactobacillus* spp.), vaginal epithelial cells, and no other bacteria observed with WBC 0–5/HP under microscopy. Grade II was some *Lactobacillus* spp. and vaginal epithelial cells, some pus cells, and other bacteria observed under microscopy with WBC 10–15/HP. Grade III was a small amount of *Lactobacillus* spp., a large number of pus cells, and other bacteria observed under microscopy with WBC 15–30/HP. Grade IV was no *Lactobacillus* spp. but pus cells and other bacteria observed under microscopy with WBC more than 30/HP. Grades I–II mean normal vaginal cleanliness, while grades III–IV mean abnormal vaginal cleanliness with inflammation. The vaginal swabs taken for bacterial genomic DNA extraction were transferred to the laboratory immediately and stored at -80°C until analyzed. Additionally, hospital records were reviewed following delivery to determine pregnancy outcome, including preterm birth, premature rupture of membranes, or preterm premature rupture of membranes.

2.3. Total Bacterial Genomic DNA Extraction. The bacterial cells retrieved on swabs were submerged in 1 mL of ice-cold phosphate-buffered saline (pH 7.5) and vigorously agitated to dislodge cells. The suspension was then transferred to a

tube containing 100 mg of glass beads (Promega Corporation, Madison, WI, USA) and placed into a FastPrep FP120 instrument (Thermo Scientific, Pittsburgh, PA, USA) to disrupt the cell membranes. The suspension was then centrifuged at 17,000 g for 10 min, resuspended in 400 μ L of lysozyme, and incubated for 1 h at 37°C. Subsequently, 500 μ L of cell lysis solution was added and warmed for an additional 30 min at 37°C. The reaction solution was then centrifuged for 10 min at 12,000 g. After removing the supernatant, 180 μ L of lysis buffer (buffer ATL) and 25 μ L of proteinase K were added to the sample and incubated for 60 min at 55°C. The QIAamp DNA Mini Extraction Kit (QIAGEN, Hilden, Germany) was then used for the isolation of the genomic DNA according to the manufacturer's instructions. One sterile cotton swab was used as a negative control alongside the patient specimens. The concentration of extracted DNA was determined using a Smart-Spec Plus spectrophotometer (Bio-Rad Laboratories, Hercules, CA, USA). Its integrity and size were checked by 1.0% agarose gel electrophoresis, containing 0.5 mg/ml ethidium bromide. All DNA was stored at -20°C before analysis.

2.4. Quantitative Real-Time PCR (qPCR) Analysis. The ABI 7900 Fast Real-Time PCR instrument and Sequence Detection Software version 1.6.3 (Applied Biosystems, Foster City, CA, USA) were used to perform qPCR on all subjects, to determine accurate copy numbers of bacteria in the vaginal specimens. The following species-specific primer sets were chosen to quantify total bacteria in the samples, *Lactobacillus* genus, *L. crispatus*, *L. jensenii*, *L. iners*, *G. vaginalis*, *A. vaginae*, *Eggerthella* sp., *Leptotrichia/Sneathia* sp. *Megasphaera* sp., and *Prevotella* sp. (Table 1) [19]. The optimal reaction conditions and parameters were obtained by adjusting the concentration of primers, reaction temperature, and the cycles of amplification. Each qPCR contained 1 μ L of template DNA, 12.5 μ L of 2 $\times$ Takara Perfect Real Time master mix, 0.3 μ L of a 10 μ M F/R primer mix, and 10.9 μ L of water. The reaction parameters were as follows: 1 cycle of predenaturation for 3 min at 95°C, 40 repeated cycles of denaturation for 30 s at 94°C, annealing for 30 s, extension for 30 s at 72°C, and a final extension step for 5 min at 72°C. At the end of the qRT-PCR, the specificity of the products produced was confirmed through a melting curve analysis, whereby the PCR products were slowly heated from 55°C to 95°C in 1.0°C increments with continuous fluorescence collection.

Serially diluted standards of plasmid DNA (1 to 10⁸ copies) containing the respective amplicon for each primer pair were run on the same plate to construct a standard curve, in order to obtain a corresponding target DNA copy number of each species in each μ L of crude DNA template. All samples including nontemplate controls were assayed three times on separate plates. Aliquots (4 μ L) of the real-time PCR products with each of the species-specific primer pairs were examined by 1.5% agarose gel electrophoresis containing 0.5 mg/mL ethidium bromide.

2.5. Statistical Analysis. The data were expressed as the mean and standard deviation for continuous variables and as

absolute and relative frequencies for categorical variables. The abundance of genus-specific bacteria was expressed as the percentage of the accurate copy numbers of the individual bacterial subgroups and total bacteria in the samples. The mean values of each item for continuous variables between two groups were compared using the Mann-Whitney *U* test, whereas differences among the three groups were analyzed using the Kruskal-Wallis *H* test. Nonparametric chi-square tests were used to compare categorical variables. Spearman's rank correlation was used to evaluate the relationship between parameters. All statistical tests were performed using SPSS, version 19.0 (SPSS Inc., Chicago, IL, USA), and any *p* value < 0.05 was considered statistically significant.

3. Results

3.1. Characteristics of the Subjects. In this study, vaginal samples of 83 healthy pregnant women were analyzed. The varying gestational ages of the women screened were compared in terms of clinical characteristics (e.g., maternal age, gravity, and parity), laboratory characteristics (vaginal pH, vaginal cleanliness, and H₂O₂ levels), and the presence of leukocyte esterase (Table 2). There was no difference in the maternal ages, gravidity, or parity among the three trimester groups, indicating that the participants in each of the different groups had similar clinical characteristics and were therefore comparable throughout this study. Comparing vaginal cleanliness and leukocyte esterase, it was found that they were significantly higher in women in the first trimester than in the second (*p* = 0.001) and third trimester (*p* < 0.001). However, vaginal pH and H₂O₂ levels showed no difference among trimesters.

3.2. Distribution of Vaginal Bacteria in Pregnant Women. In this study, using qRT-PCR, seven known abundant genera (*Lactobacillus*, *Gardnerella*, *Atopobium*, *Megasphaera*, *Eggerthella*, *Leptotrichia/Sneathia*, and *Prevotella*) were analyzed. The abundance of *Gardnerella*, *Atopobium*, *Megasphaera*, *Eggerthella*, *Leptotrichia/Sneathia*, and *Prevotella* was significantly different among the three trimesters (*p* > 0.05). The genus *Lactobacillus* constituted the major proportion of the vaginal microbiota in healthy pregnant women, which was consistent with previous studies [34, 36, 37]. *L. jensenii*, *L. iners*, and *L. crispatus* were the most frequent species [37–39]. Among them, the abundance of *L. iners* and *L. crispatus* was significantly different among the trimesters. We also found that *L. iners* decreased significantly in women in the second and third trimester when compared with women in the first trimester (*p* < 0.001), while *L. crispatus* significantly increased in the second trimester (*p* = 0.030) (Figure 1). However, the relative abundance of *L. jensenii* was unchanged obviously among the three trimesters (*p* = 0.633; Table 3).

3.3. Correlation of Vaginal Bacteria with Vaginal Cleanliness and Leucocyte Esterase. Vaginal cleanliness and leukocyte esterase varied significantly throughout the pregnancy trimester. Spearman's rank correlation was used to examine

TABLE 1: Species-specific primer sets for detection of vaginal bacteria by quantitative real-time PCR.

PCR specificity	Primer	Sequence (5'-3')	Annealing temp.	Amplicon size (bp)	Reference
All bacteria	Bac27F EUB338R-I	AGAGTTTGATCCTGGCTCAG GCTGCCTCCCGTAGGAGT	65°C	312	[28]
<i>Lactobacillus</i> genus	Bact-0011 Lab-0677	AGAGTTTGATYMTGGCTCAG CACCGCTACACATGGAG	62°C	667	[29]
<i>L. crispatus</i>	Lcris-F Lcris-R	AGCGAGCGGAACTAACAGATTTAC AGCTGATCATGCGATCTGCTT	65°C	154	[30]
<i>L. jensenii</i>	Ljens-F Ljens-R	AAGTCGAGCGAGCTTGCCTATAGA CTTCTTTCATGCGAAAGTAGC	60°C	162	[31]
<i>L. iners</i>	Liners-F Liners-R	CTCTGCCTTGAAGATCGGAGTGC ACAGTTGATAGGCATCATCTG	65°C	155	[31]
<i>Atopobium vaginae</i>	AV-F AV-R	TAGGTCAGGAGTTAAATCTG TCATGGCCCAGAAAGACCGCC	62°C	156	[32]
<i>Gardnerella vaginalis</i>	GV1-F GV3-R	TTACTGGTGTATCACTGTAAGG CCGTCACAGGCTGAACAGT	62°C	332	[33]
<i>Eggerthella</i>	Egger-621F Egger-859R	AACCTCGAGCCGGGTTCC TCGGCACGGAAGATGTAATCT	60°C	239	[34]
<i>Leptotrichia/Sneathia</i>	Lepto-395F Lepto-646R	CAATTCTGTGTGTGTGAAGAAG ACAGTTTGTAGGCAAGCCTAT	60°C	252	[34]
<i>Megasphaera type I</i>	MegaE-456F MegaE-667R	GATGCCAACAGTATCCGTCCG CCTCTCCGACACTCAAGTTCGA	64°C	212	[34]
<i>Prevotella</i>	Prevo-F Prevo-R	CCAGCCAAGTAGCGTGCA TGGACCTTCCGTATTACCGC	60°C	151	[35]

TABLE 2: Clinical and laboratory characteristics of the participants.

Parameters	Participants in 1st trimester (n = 33)	Participants in 2nd trimester (n = 24)	Participants in 3rd trimester (n = 26)	p value
Age (years)	28.30 (5.35)	29.50 (3.91)	29.35 (4.03)	0.221
Gravidity	0.88 (0.96)	1.21 (1.22)	1.31 (1.62)	0.604
Parity	0.55 (0.66)	0.50 (0.51)	0.42 (0.50)	0.832
PH	4.2 (0.3)	4.0 (0.2)	4.1 (0.2)	0.118
Cleanliness (1–2/3–4)	18/15	23/1	22/4	0.001
H ₂ O ₂ (<2/≥2)	5/28	0/24	2/24	0.125
Leukocyte esterase (P/N)	14/19	0/24	4/22	<0.001

All data are mean (standard deviation). P, positive; N, negative.

the relationship between abundant vaginal microbiota and the two indicators including vaginal cleanliness and leukocyte esterase (Table 4). We found that vaginal cleanliness was significantly correlated with the presence of *Lactobacillus*, *Atopobium*, and *Prevotella* instead of other bacteria genus. However, all abundant bacteria except for *Megasphaera* ($p < 0.05$) were altered significantly between leukocyte esterase positive and negative. Additionally, it was interesting to find that the relative abundance of *L. iners* increased with raised vaginal cleanliness grade and positive leukocyte esterase, but *L. crispatus* did not show such similar changing patterns.

3.4. Relationship between Vaginal pH and Vaginal Abundant Bacteria. In our present study, vaginal pH was maintained at a low level in these pregnant women and was not changed significantly across the three trimesters. Normally, *Lactobacilli* acidify the vagina with lactic acid, while the

overgrowth of those vaginal pathogenic bacteria may lead to the elevation of vaginal pH. We used Spearman's rank correlations to investigate the relationships between the relative abundance of genus-specific vaginal bacteria and vaginal pH. As shown in Figure 2, the vaginal pH increased when *Lactobacillus*, and more specifically *L. crispatus* decreased when the vaginal pH increased. However, the vaginal pH also was positively correlated with *L. jensenii*, *L. iners*, *A. vaginae*, and *G. vaginalis*.

4. Discussion

Vaginal cleanliness and leukocyte esterase have been proven previously to be associated with genital inflammation [27]. In this study, it was discovered that vaginal cleanliness and leukocyte esterase were significantly higher in women who were in the first trimester instead of the second and third trimester. Although this result cannot fully illustrate that these two inflammatory markers change spontaneously as

FIGURE 1: Comparison of the relative abundance of (a) *Lactobacillus*, (b) *L. crispatus*, and (c) *L. iners* in 83 asymptomatic pregnant women according to their trimester of pregnancy. Data are expressed as scatter plots, in which the horizontal lines illustrate the mean value of each genus-specific bacterium.

TABLE 3: Comparison of the relative abundance of vaginal bacteria by quantitative real-time PCR according to trimesters of pregnancy.

Species	Participants in 1st trimester (n = 33)	Participants in 2nd trimester (n = 24)	Participants in 3rd trimester (n = 26)	p value
<i>Lactobacillus</i> genus (%)	31.57 (23.94)	59.80 (21.96) b	64.14 (21.61) b	<0.001
<i>L. jensenii</i> (%) ^a	0.07 (0.32)	0.43 (2.07)	0.27 (1.31)	0.633
<i>L. crispatus</i> (%) ^a	5.74 (10.39)	13.21 (12.96) c	9.67 (10.47)	0.030
<i>L. iners</i> (%) ^a	14.86 (19.74)	4.16 (7.97) d	5.02 (7.24) d	<0.001
<i>Atopobium vaginae</i> (%)	1.21 (3.17)	2.63 (12.86)	0.03 (0.16)	0.897
<i>Gardnerella vaginalis</i> (%)	0.10 (0.24)	0.08 (0.34)	2.49 (12.67)	0.944
<i>Eggerthella</i> (%)	0.05 (0.21)	0.00 (0.00)	0.00 (0.00)	0.466
<i>Leptotrichia/Sneathia</i> (%)	0.04 (0.12)	0.00 (0.00)	0.00 (0.00)	0.432
<i>Megasphaera type</i> (%)	0.29 (0.80)	0.02 (0.10)	0.02 (0.09)	0.569
<i>Prevotella</i> (%)	0.19 (0.45)	0.00 (0.01)	0.09 (0.45)	0.169

All data are mean (standard deviation). ^aThe relative abundance of *L. crispatus*, *L. jensenii*, and *L. iners* was compared to the copy number of *Lactobacillus* genus; ^bcompared with 1st trimester, $p < 0.001$; ^ccompared with 1st trimester, $p < 0.05$; ^dcompared with 1st trimester, $p < 0.01$.

TABLE 4: Correlation of the relative abundance of vaginal bacteria with vaginal cleanliness and leucocyte esterase.

Vaginal bacteria	Vaginal cleanliness			Leucocyte esterase		
	1–2 (n = 63)	3–4 (n = 20)	p value	Negative (n = 65)	Positive (n = 18)	p value
<i>Lactobacillus</i> genus (%)	55.00 (25.14)	34.10 (27.18)	0.002	55.42 (23.86)	30.13 (29.06)	0.001
<i>L. jensenii</i> (%)	10.68 (11.91)	4.27 (8.67)	0.072	10.39 (11.84)	4.61 (9.09)	0.241
<i>L. crispatus</i> (%)	0.30 (1.53)	0.02 (0.22)	0.873	0.29 (1.51)	0.03 (0.47)	0.220
<i>L. iners</i> (%)	6.18 (13.04)	16.56 (16.49)	<0.001	6.18 (12.79)	17.73 (17.15)	<0.001
<i>Atopobium vaginae</i> (%)	0.51 (2.46)	1.09 (2.57)	0.044	0.49 (2.42)	1.23 (2.67)	0.004
<i>Gardnerella vaginalis</i> (%)	0.05 (0.22)	0.20 (0.43)	0.133	0.04 (0.22)	0.23 (0.44)	0.023
<i>Eggerthella</i> (%)	0.00 (0.01)	0.08 (0.84)	0.194	0.00 (0.01)	0.09 (1.02)	0.006
<i>Leptotrichia/Sneathia</i> (%)	0.01 (0.11)	0.04 (0.21)	0.496	0.00 (0.01)	0.06 (0.41)	0.032
<i>Megasphaera type</i> (%)	0.07 (0.23)	0.30 (0.75)	0.686	0.06 (0.43)	0.36 (1.32)	0.097
<i>Prevotella</i> (%)	0.02 (0.18)	0.36 (1.46)	0.004	0.01 (0.02)	0.45 (1.36)	0.002

All data are mean (standard deviation). The abundance of vaginal bacteria relative to total bacteria gene copy number according to vaginal cleanliness and leucocyte esterase was compared. ^aThe relative abundance of *L. crispatus*, *L. jensenii*, and *L. iners* was compared to the copy number of *Lactobacillus* genus.

pregnancy progressed, previous studies indicated that this phenomenon may be possible. Waters et al. performed a longitudinal investigation and found that most women who were BV positive in early pregnancy became BV negative in the third trimester [40]. Duff et al. have reported that BV disappeared spontaneously between 15 and 36 weeks in 11% of pregnant patients without any therapies or interventions

[41]. However, further longitudinal studies are needed to explore the definite relationship between the dynamic vaginal microbiota and the progress of pregnancy.

Vaginal pH is another known indicator of vaginal health which is commonly used for the diagnosis of genital infections. The vaginal pH significantly increases when BV or trichomoniasis vaginitis is present and it decreases when

FIGURE 2: Correlation between the relative abundance of genus-specific bacteria and vaginal pH in 83 asymptomatic pregnant women. Relationships among the vaginal pH with (a) *Lactobacillus*, (b) *L. crispatus*, (c) *L. jensenii*, (d) *L. iners*, (e) *A. vaginae*, and (f) *G. vaginalis* are shown as scatter plots and regression lines. The coefficient is taken from Spearman's correlation test.

Candida vaginitis occurs [42, 43]. However, vaginal pH did not seem to be closely associated with changes in the vagina microecosystem strongly according to the results. Furthermore, H_2O_2 is known as an important factor of vaginal health and is mostly produced by *Lactobacillus* and acts as a certain antibacterial compound which can protect against genital tract pathogens. Our data indicated that there was no significant difference in the presence of H_2O_2 among the three trimesters ($p = 0.125$), indicating that it could help to maintain vaginal homeostasis in the normal status. Of course, the bacterial interactions in the pregnancy-related vaginal microbiota are complex and are not simply related to the presence or absence of H_2O_2 -producing *Lactobacillus* [44].

The vaginal microbial community is typically characterized by abundant lactobacilli. In our study, two species of *Lactobacillus*, such as *L. iners* and *L. crispatus*, were found to be significantly different among the three trimesters. *L. iners* was found to be significantly decreased whereas *L. crispatus* was found to be significantly increased in women in the second and third trimester compared with the first trimester. Besides, we observed that vaginal cleanliness and leukocyte esterase were significantly affected by *L. iners* but not by *L. crispatus*, which was consistent with the previous study [45]. With recent advances in culture-independent community profiling, the nutritionally fastidious *L. iners* is emerging as a dominant organism, present in both healthy and *lactobacilli*-deficient aberrant vaginal environments, suggesting that *L. iners* is very flexible and can easily adapt to the fluctuating vaginal niche [46–48]. Therefore, the existence of *L. iners* may not suppress the proliferation of other potentially harmful bacteria in the vagina [49]. *L. iners* has been thought to be a precursor to being susceptible to adverse pregnant sequelae [50]. Simultaneously, *L. crispatus* has been previously suggested to be linked to healthy microbiota, a healthy pregnancy, and a term delivery and to the absence of

vaginal infection or inflammation [51]. In this work, we had identified a positive correlation between inflammatory laboratory characteristics and *L. iners*.

In fact, there is a lot of controversy over whether *L. iners* is advantageous or detrimental for the host microbiota. Gajer et al. observed that the vaginal communities dominated by *L. crispatus* changed into a community dominated by *L. iners* during menstruation [52]. Their findings indicate that *L. iners* may help in the recovery of *lactobacilli*-dominant vaginal microbiota, supporting the notion that *L. iners* is a beneficial species [53]. Most women with an *L. iners*-dominated vaginal community will deliver at term with no adverse pregnancy outcome [50, 51]. Previous studies have found that *L. iners* may be a beneficial vaginal bacterium which showed beneficial effects in several mechanisms. It has been proposed that *L. iners* activates Toll-like receptor signaling in epithelial cells, elevates the levels of heat-shock protein 70, and inhibits autophagy [54, 55]. This process could combat a non-physiological threat and maintain and promote a return to healthier conditions [56]. This would be by inducing the innate immune system in the vaginal epithelial cells, preventing harmful bacteria from obtaining important nutrients such as iron, and inhibiting their continual growth [45]. However, several studies indicated that *L. iners* offers less protection against vaginal dysbiosis [45]. In the present study, we also found that *L. iners* was more responsive to the dysbiosis of vaginal microbiota when compared to *L. crispatus*. Due to the significant complex and diverse individual discrepancies of the vaginal microbiota, more studies are required to make clear whether *L. iners* can be used as a novel biomarker to detect the presence or prognosis of vaginal inflammation and guide clinical treatment.

Our research had also explored several anaerobic microorganisms, *Gardnerella*, *Atopobium*, *Megasphaera*, *Eggerthella*, *Leptotrichia/Sneathia*, and *Prevotella*, which were considered

as the vaginal pathogenic community. Our previous studies have found that these bacteria in the vaginal pathogenic community increased significantly in patients with BV accompanying with the depletion of *Lactobacillus* [19, 57, 58], which is highly accurate for BV diagnosis. However, the abundance of these bacteria was not changed significantly among the three trimesters in our present study. Previous studies have shown that *G. vaginalis* (belonging to Actinobacteria) plays an important role in the development of BV [19, 34]. Another genus in Actinobacteria, *Atopobium*, has been suggested to be even more specific than *G. vaginalis* for the diagnosis of BV (77% and 35%, respectively) [59]. Additionally, the presence of *Eggerthella* has also been considered as an independent risk factor for BV scores (Nugent score ≥ 7) [31]. Other members such as *Prevotella* (belonging to Bacteroidetes), *Megasphaera* (especially *Megasphaera typeI*), and *Leptotrichia/Sneathia* (belonging to Fusobacteria) also appeared to have a strong positive association with BV, although previous research could not detect these genera in all BV samples [19, 34, 60, 61]. However, the clinical significance of these genera associated with BV in the vaginal ecosystem was still unknown. In this study, we found these bacteria in subjects who did not have BV, and therefore, this may suggest that these species may not be a specific marker for BV. Of course, these bacteria in the vaginal microbiota were interacted with each other closely. Previous study also found a clear negative association between *L. iners* and *L. gasseri* and between *A. vaginae* and *L. gasseri* [62]. Recently, Petrova et al. have showed that *L. iners* has more complex nutritional requirements and a Gram-variable morphology when compared to other *Lactobacillus* species [49]. Genome sequencing revealed that *L. iners* has an unusually small genome indicative of a parasitic or symbiotic lifestyle in the human vagina, which encodes inerolysin, a pore-forming toxin related to vaginolysin of *G. vaginalis* [49]. Similar to our previous studies, the *Lactobacillus* species, such as *L. iners* and *L. crispatus*, maintained the balance of the vaginal ecosystem, while these vaginal pathogenic bacteria contributed to the dysbiosis of the vaginal microbiota [19, 57, 58]. The depletion of *lactobacilli*, together with the increase of different species of anaerobes, could result in the switch from normal to a dysbiosis vaginal microbiota, which contributed to various adverse outcomes.

Our findings could have important implications when interpreting the varied results of investigations aimed at improving pregnancy outcomes. These data support the observation that the prevalence of vaginal microbiota varies significantly over the course of pregnancy, with a strong trend towards a reduction in infection by the third trimester. The limitation of this study is small sample capacity and nonlongitudinal design.

5. Conclusion

In summary, we investigated the changes of laboratory characteristics and the presence of certain bacteria in the vagina throughout pregnancy and highlighted their relationship. *L. iners* was found to be significantly decreased in the second and third trimester compared with the first trimester, while *L. crispatus* increased only in the second

trimester. It was found that *L. iners* may be highly associated with vaginal dysbiosis, and further research studies are required to identify it. Additionally, quantification of the relative and absolute numbers of *L. iners* under different conditions throughout pregnancy was needed to be able to potentially predict adverse outcomes that it may be associated with. This study was limited due to the relatively small sample size ($n=83$); however, it did provide the basis of future work to investigate the role of the microbiota in both low- and high-risk pregnancies.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Authors' Contributions

NZ and RG contributed equally to this work. NZ and ZL conceived and designed the study. NZ, RG, YY, and MJ collected the samples. YC and ZL performed qPCR. NZ, RG, YC, and ZL analyzed the data, carried out the computational analysis, interpreted the data, and drafted the manuscript.

Acknowledgments

The authors gratefully acknowledge the volunteers who participated in our study. This work was supported by the Science and Technology Foundation of Public Welfare Profession of Zhejiang Province (grant no. 2015C33198), the Medical Science and Technology Project of Zhejiang Province (grant no. 2017KY337), the National S & T Major Project of China (2018YFC2000500), and the National Natural Science Foundation of China (81771724, 31700800, 81790633, and 31870839).

Supplementary Materials

Table S1: judgment method of vaginal cleanliness. (*Supplementary Materials*)

References

- [1] B. Ma, L. J. Forney, and J. Ravel, "Vaginal microbiome: rethinking health and disease," *Annual Review of Microbiology*, vol. 66, no. 1, pp. 371–389, 2012.
- [2] R. Romero, S. S. Hassan, P. Gajer et al., "The vaginal microbiota of pregnant women who subsequently have spontaneous preterm labor and delivery and those with a normal delivery at term," *Microbiome*, vol. 2, no. 1, p. 18, 2014.
- [3] A. Machado, C. Almeida, D. Salgueiro et al., "Fluorescence in situ Hybridization method using Peptide nucleic acid probes for rapid detection of *Lactobacillus* and *Gardnerella* spp.," *BMC Microbiology*, vol. 13, no. 1, p. 82, 2013.

- [4] S. M. Husain, M. Wilks, M. Mupita et al., "Diversity and stability of cultured vaginal lactobacilli in pregnant women from a multi-ethnic urban UK population," *Journal of Applied Microbiology*, vol. 117, no. 1, pp. 258–265, 2014.
- [5] C. Tellapragada, V. K. Eshwara, P. Bhat, A. Kamath, S. Aletty, and C. Mukhopadhyay, "Screening of vulvovaginal infections during pregnancy in resource constrained settings: implications on preterm delivery," *Journal of Infection and Public Health*, vol. 10, no. 4, pp. 431–437, 2017.
- [6] R. Vigneswaran, "Infection and preterm birth: evidence of a common causal relationship with bronchopulmonary dysplasia and cerebral palsy," *Journal of Paediatrics and Child Health*, vol. 36, no. 4, pp. 293–296, 2000.
- [7] J. van Dillen, J. Zwart, J. Schutte, and J. van Roosmalen, "Maternal sepsis: epidemiology, etiology and outcome," *Current Opinion in Infectious Diseases*, vol. 23, no. 3, pp. 249–254, 2010.
- [8] D. Drassinower, A. M. Friedman, S. G. Obican, H. Levin, and C. Gyamfi-Bannerman, "Prolonged latency of preterm premature rupture of membranes and risk of cerebral palsy," *The Journal of Maternal-Fetal & Neonatal Medicine*, vol. 29, no. 17, pp. 2748–2752, 2015.
- [9] H. Lu, Q. Wang, J. Lu, Q. Zhang, and P. Kumar, "Risk factors for intraventricular hemorrhage in preterm infants born at 34 Weeks of gestation or less following preterm premature rupture of membranes," *Journal of Stroke and Cerebrovascular Diseases*, vol. 25, no. 4, pp. 807–812, 2016.
- [10] K. Puri, D. H. Taft, N. Ambalavanan, K. R. Schibler, A. L. Morrow, and S. G. Kallapur, "Association of chorioamnionitis with aberrant neonatal gut colonization and adverse clinical outcomes," *PLoS One*, vol. 11, no. 9, Article ID e0162734, 2016.
- [11] E. Faure, K. Faure, M. Figeac et al., "Vaginal mucosal homeostatic response may determine pregnancy outcome in women with bacterial vaginosis: a pilot study," *Medicine*, vol. 95, no. 5, p. e2668, 2016.
- [12] S. L. Hillier, R. P. Nugent, D. A. Eschenbach et al., "Association between bacterial vaginosis and preterm delivery of a low-birth-weight infant," *New England Journal of Medicine*, vol. 333, no. 26, pp. 1737–1742, 1995.
- [13] C. A. Flynn, A. L. Helwig, and L. N. Meurer, "Bacterial vaginosis in pregnancy and the risk of prematurity: a meta-analysis," *The Journal of Family Practice*, vol. 48, no. 11, pp. 885–892, 1999.
- [14] D. B. DiGiulio, B. J. Callahan, P. J. McMurdie et al., "Temporal and spatial variation of the human microbiota during pregnancy," *Proceedings of the National Academy of Sciences*, vol. 112, no. 35, pp. 11060–11065, 2015.
- [15] L. M. Kindinger, D. A. MacIntyre, Y. S. Lee et al., "Relationship between vaginal microbial dysbiosis, inflammation, and pregnancy outcomes in cervical cerclage," *Science Translational Medicine*, vol. 8, no. 350, p. 350ra102, 2016.
- [16] G. Reid, J. A. Younes, H. C. Van der Mei, G. B. Gloor, R. Knight, and H. J. Busscher, "Microbiota restoration: natural and supplemented recovery of human microbial communities," *Nature Reviews Microbiology*, vol. 9, no. 1, pp. 27–38, 2011.
- [17] R. Romero, S. S. Hassan, P. Gajer et al., "The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women," *Microbiome*, vol. 2, no. 1, 2014.
- [18] A. McMillan, S. Rulisa, M. Sumarah et al., "A multi-platform metabolomics approach identifies highly specific biomarkers of bacterial diversity in the vagina of pregnant and non-pregnant women," *Scientific Reports*, vol. 5, no. 1, p. 14174, 2015.
- [19] X. Z. Ling, M. J. Kong, F. Liu et al., "Molecular analysis of the diversity of vaginal microbiota associated with bacterial vaginosis," *BMC Genomics*, vol. 11, no. 1, p. 488, 2010.
- [20] A. Shobokshi and M. Shaarawy, "Maternal serum and amniotic fluid cytokines in patients with preterm premature rupture of membranes with and without intrauterine infection," *International Journal of Gynecology & Obstetrics*, vol. 79, no. 3, pp. 209–215, 2002.
- [21] K. B. Fortner, C. A. Grotegut, C. E. Ransom et al., "Bacteria localization and chorion thinning among preterm premature rupture of membranes," *PLoS One*, vol. 9, no. 1, Article ID e83338, 2014.
- [22] V. V. Demkin, S. I. Koshechkin, and A. Slesarev, "A novel real-time PCR assay for highly specific detection and quantification of vaginal lactobacilli," *Molecular and Cellular Probes*, vol. 32, pp. 33–39, 2017.
- [23] R. Verhelst, H. Verstraelen, G. Claeys et al., "Cloning of 16S rRNA genes amplified from normal and disturbed vaginal microflora suggests a strong association between *Atopobium vaginae*, *Gardnerella vaginalis* and bacterial vaginosis," *BMC Microbiology*, vol. 4, no. 1, p. 16, 2004.
- [24] D. H. Yan, Z. Lu, and J. R. Su, "Comparison of main *Lactobacillus* species between healthy women and women with bacterial vaginosis," *Chinese Medical Journal*, vol. 122, no. 22, pp. 2748–2751, 2009.
- [25] R. Amsel, P. A. Totten, C. A. Spiegel, K. C. S. Chen, D. Eschenbach, and K. K. Holmes, "Nonspecific vaginitis," *The American Journal of Medicine*, vol. 74, no. 1, pp. 14–22, 1983.
- [26] S. Kovachev, "Vaginal ecosystem," *Akush Ginekol*, vol. 50, no. 3, pp. 41–49, 2011.
- [27] X.-A. Yue, P. Chen, Y. Tang, X. Wu, and Z. Hu, "The dynamic changes of vaginal microecosystem in patients with recurrent vulvovaginal candidiasis: a retrospective study of 800 patients," *Archives of Gynecology and Obstetrics*, vol. 292, no. 6, pp. 1285–1294, 2015.
- [28] J. S. Lipp, Y. Morono, F. Inagaki, and K.-U. Hinrichs, "Significant contribution of Archaea to extant biomass in marine subsurface sediments," *Nature*, vol. 454, no. 7207, pp. 991–994, 2008.
- [29] H. G. H. J. Heilig, E. G. Zoetendal, E. E. Vaughan, P. Marteau, A. D. L. Akkermans, and W. M. de Vos, "Molecular diversity of *Lactobacillus* spp. and other lactic acid bacteria in the human intestine as determined by specific amplification of 16S ribosomal DNA," *Applied and Environmental Microbiology*, vol. 68, no. 1, pp. 114–123, 2002.
- [30] R. Byun, M. A. Nadkarni, K.-L. Chhour, F. E. Martin, N. A. Jacques, and N. Hunter, "Quantitative analysis of diverse *Lactobacillus* species present in advanced dental caries," *Journal of Clinical Microbiology*, vol. 42, no. 7, pp. 3128–3136, 2004.
- [31] R. Tamrakar, T. Yamada, I. Furuta et al., "Association between *Lactobacillus* species and bacterial vaginosis-related bacteria, and bacterial vaginosis scores in pregnant Japanese women," *BMC Infectious Diseases*, vol. 7, no. 1, p. 128, 2007.
- [32] M. J. Ferris, A. Maszta, K. E. Aldridge, J. D. Fortenberry, P. L. Fidel Jr., and D. H. Martin, "Association of *Atopobium vaginae*, a recently described metronidazole resistant anaerobe, with bacterial vaginosis," *BMC Infectious Diseases*, vol. 4, no. 5, 2004.
- [33] M. Zariffard, M. Saifuddin, B. E. Sha, and G. T. Spear, "Detection of bacterial vaginosis-related organisms by real-

- time PCR for *Lactobacilli*, *Gardnerella vaginalis* and *Mycoplasma hominis*,” *FEMS Immunology and Medical Microbiology*, vol. 34, no. 4, pp. 277–281, 2002.
- [34] D. N. Fredricks, T. L. Fiedler, and J. M. Marrazzo, “Molecular identification of bacteria associated with bacterial vaginosis,” *New England Journal of Medicine*, vol. 353, no. 18, pp. 1899–1911, 2005.
 - [35] F. E. Martin, M. A. Nadkarni, N. A. Jacques, and N. Hunter, “Quantitative microbiological study of human carious dentine by culture and real-time PCR: association of anaerobes with histopathological changes in chronic pulpitis,” *Journal of Clinical Microbiology*, vol. 40, no. 5, pp. 1698–1704, 2002.
 - [36] X. Zhou, C. J. Brown, Z. Abdo et al., “Differences in the composition of vaginal microbial communities found in healthy Caucasian and black women,” *The ISME Journal*, vol. 1, no. 2, pp. 121–133, 2007.
 - [37] H. Verstraelen, R. Verhelst, G. Claeys, E. De Backer, M. Temmerman, and M. Vaneechoutte, “Longitudinal analysis of the vaginal microflora in pregnancy suggests that *L. crispatus* promotes the stability of the normal vaginal microflora and that *L. gasseri* and/or *L. iners* are more conducive to the occurrence of abnormal vaginal microflora,” *BMC Microbiology*, vol. 9, no. 1, p. 116, 2009.
 - [38] S. I. Pavlova, A. O. Kilic, S. S. Kilic et al., “Genetic diversity of vaginal *Lactobacilli* from women in different countries based on 16S rRNA gene sequences,” *Journal of Applied Microbiology*, vol. 92, no. 3, pp. 451–459, 2002.
 - [39] M. Tärnberg, T. Jakobsson, J. Jonasson, and U. Forsum, “Identification of randomly selected colonies of *Lactobacilli* from normal vaginal fluid by pyrosequencing of the 16S rDNA variable V1 and V3 regions,” *APMIS*, vol. 110, no. 11, pp. 802–810, 2002.
 - [40] T. P. Waters, J. M. Denney, L. Mathew, R. L. Goldenberg, and J. F. Culhane, “Longitudinal trajectory of bacterial vaginosis during pregnancy,” *American Journal of Obstetrics and Gynecology*, vol. 199, no. 4, pp. 431 e431–e435, 2008.
 - [41] P. Duff, M. L. Lee, S. L. Hillier, L. M. Herd, M. A. Krohn, and D. A. Eschenbach, “Amoxicillin treatment of bacterial vaginosis during pregnancy,” *Obstetrics & Gynecology*, vol. 77, no. 3, pp. 431–435, 1991.
 - [42] J.-S. Ryu and D.-Y. Min, “Trichomonas vaginalis and trichomoniasis in the Republic of Korea,” *The Korean Journal of Parasitology*, vol. 44, no. 2, pp. 101–116, 2006.
 - [43] D. Nenadic and M. Pavlovic, “Value of bacterial culture of vaginal swabs in diagnosis of vaginal infections,” *Vojnosanitetski Pregled*, vol. 72, no. 6, pp. 523–528, 2015.
 - [44] C. Mitchell, D. Fredricks, K. Agnew, and J. Hitti, “Hydrogen peroxide-producing *Lactobacilli* are associated with lower levels of vaginal interleukin-1 β , independent of bacterial vaginosis,” *Sexually Transmitted Diseases*, vol. 42, no. 7, pp. 358–363, 2015.
 - [45] M. Vaneechoutte, “*Lactobacillus iners*, the unusual suspect,” *Research in Microbiology*, vol. 168, no. 9–10, pp. 826–836, 2017.
 - [46] J. M. Macklaim, A. D. Fernandes, J. M. Di Bella, J. A. Hammond, G. Reid, and G. B. Gloor, “Comparative meta-RNA-seq of the vaginal microbiota and differential expression by *Lactobacillus iners* in health and dysbiosis,” *Microbiome*, vol. 1, no. 1, p. 12, 2013.
 - [47] H. Borgdorff, S. D. Armstrong, H. L. Tytgat et al., “Unique insights in the cervicovaginal *Lactobacillus iners* and *L. Crispatus* proteomes and their associations with microbiota dysbiosis,” *PLoS One*, vol. 11, no. 3, Article ID e0150767, 2016.
 - [48] M. Zozaya-Hinchliffe, R. Lillis, D. H. Martin, and M. J. Ferris, “Quantitative PCR assessments of bacterial species in women with and without bacterial vaginosis,” *Journal of Clinical Microbiology*, vol. 48, no. 5, pp. 1812–1819, 2010.
 - [49] M. I. Petrova, G. Reid, M. Vaneechoutte, and S. Lebeer, “*Lactobacillus iners*: friend or foe?,” *Trends in Microbiology*, vol. 25, no. 3, pp. 182–191, 2017.
 - [50] L. M. Kindinger, P. R. Bennett, Y. S. Lee et al., “The interaction between vaginal microbiota, cervical length, and vaginal progesterone treatment for preterm birth risk,” *Microbiome*, vol. 5, no. 1, p. 6, 2017.
 - [51] L. Petricevic, K. J. Domig, F. J. Nierscher et al., “Characterisation of the vaginal *Lactobacillus* microbiota associated with preterm delivery,” *Scientific Reports*, vol. 4, no. 1, p. 5136, 2014.
 - [52] P. Gajer, R. M. Brotman, G. Bai et al., “Temporal dynamics of the human vaginal microbiota,” *Science Translational Medicine*, vol. 4, no. 132, p. 132ra52, 2012.
 - [53] S. Srinivasan, C. Liu, C. M. Mitchell et al., “Temporal variability of human vaginal bacteria and relationship with bacterial vaginosis,” *PLoS One*, vol. 5, no. 4, Article ID e10197, 2010.
 - [54] S. Y. Doerflinger, A. L. Throop, and M. M. Herbst-Kralovetz, “Bacteria in the vaginal microbiome alter the innate immune response and barrier properties of the human vaginal epithelia in a species-specific manner,” *The Journal of Infectious Diseases*, vol. 209, no. 12, pp. 1989–1999, 2014.
 - [55] Y. Feng, Z. Yao, and D. J. Klionsky, “How to control self-digestion: transcriptional, post-transcriptional, and post-translational regulation of autophagy,” *Trends in Cell Biology*, vol. 25, no. 6, pp. 354–363, 2015.
 - [56] B. Levine, N. Mizushima, and H. W. Virgin, “Autophagy in immunity and inflammation,” *Nature*, vol. 469, no. 7330, pp. 323–335, 2011.
 - [57] Z. X. Ling, X. Liu, Y. Q. Luo et al., “Associations between vaginal pathogenic community and bacterial vaginosis in Chinese reproductive-age women,” *PLoS One*, vol. 8, no. 10, Article ID e76589, 2013.
 - [58] Z. Ling, X. Liu, W. Chen et al., “The restoration of the vaginal microbiota after treatment for bacterial vaginosis with metronidazole or probiotics,” *Microbial Ecology*, vol. 65, no. 3, pp. 773–780, 2013.
 - [59] C. S. Bradshaw, S. N. Tabrizi, C. K. Fairley, A. N. Morton, E. Rudland, and S. M. Garland, “The association of *Atopobium vaginae* and *Gardnerella vaginalis* with bacterial vaginosis and recurrence after oral metronidazole therapy,” *The Journal of Infectious Diseases*, vol. 194, no. 6, pp. 828–836, 2006.
 - [60] B. B. Oakley, T. L. Fiedler, J. M. Marrazzo, and D. N. Fredricks, “Diversity of human vaginal bacterial communities and associations with clinically defined bacterial vaginosis,” *Applied and Environmental Microbiology*, vol. 74, no. 15, pp. 4898–4909, 2008.
 - [61] M. Zozaya-Hinchliffe, D. H. Martin, and M. J. Ferris, “Prevalence and abundance of uncultivated *Megasphaera*-like bacteria in the human vaginal environment,” *Applied and Environmental Microbiology*, vol. 74, no. 5, pp. 1656–1659, 2008.
 - [62] E. De Backer, R. Verhelst, H. Verstraelen et al., “Quantitative determination by real-time PCR of four vaginal *Lactobacillus* species, *Gardnerella vaginalis* and *Atopobium vaginae* indicates an inverse relationship between *L. gasseri* and *L. iners*,” *BMC Microbiology*, vol. 7, no. 1, p. 115, 2007.