

Research Article

Effect of Bitter Compounds on the Expression of Bitter Taste Receptor T2R7 Downstream Signaling Effectors in *cT2R7/pDisplay-Gα16/gust44/pcDNA3.1 (+) Cells*

Yuan Su ¹, Hang Jie ^{1,2}, Qing Zhu ¹, Xiaoling Zhao,¹ Yan Wang ¹, Huadong Yin,¹ Shailendra Kumar Mishra,¹ and Diyan Li ¹

¹Farm Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province, Sichuan Agricultural University, Chengdu 611130, China

²Chongqing Engineering Technology Research Center for GAP of Genuine Medicinal Materials, Chongqing Institute of Medicinal Plant Cultivation, Nanchuan, Chongqing 404100, China

Correspondence should be addressed to Qing Zhu; zhuqingsicau@163.com and Diyan Li; diyanli@sicau.edu.cn

Received 5 July 2019; Revised 24 September 2019; Accepted 17 October 2019; Published 31 October 2019

Academic Editor: Fengjie Sun

Copyright © 2019 Yuan Su et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bitterness is an important taste sensation for chickens, which provides useful sensory information for acquisition and selection of diet, and warns them against ingestion of potentially harmful and noxious substances in nature. Bitter taste receptors (T2Rs) mediate the recognition of bitter compounds belonging to a family of proteins known as G-protein coupled receptors. The aim of this study was to identify and evaluate the expression of *T2R7* in chicken tongue tissue and construct *cT2R7-1* and *cT2R7-2*-expressing HEK-293T cells to access the expression of *PLCβ2* and *ITPR3* after exposure with different concentrations of the bitter compounds. Using real-time PCR, we show that the relative expression level of *T2R7* mRNA in 5, 1, 0.1, and 10^{-3} mM of camphor and erythromycin solutions and 5 mM of chlorpheniramine maleate solutions was significantly higher than that in 50 mM KCL solutions. We confirmed that the bitter taste receptor *T2R7* and downstream signaling effectors are sensitive to different concentrations of bitter compounds. Moreover, *T2R7-1* (corresponding to the unique haplotype of the Tibetan chicken) had higher sensitivity to bitter compounds compared with that of *T2R7-2* (corresponding to the unique haplotype of the Jiuyuan black-chicken). These results provide great significance of taste response on dietary intake to improve chicken feeding efficiency in poultry production and have certain reference value for future taste research in other bird species.

1. Introduction

Hitherto, there are five widely known basic tastes that stimulate and are perceived by taste buds—sweet, umami, sour, bitter, and salty [1, 2]. Among them, bitter taste is an important natural defense providing useful sensory information for animals to ensure them ingestion of potentially healthy feeds rather than harmful/toxic substances [3–5]. Identification of bitter compounds is mediated by the type 2 taste receptors (T2Rs), which belong to G-protein-coupled superfamily [6, 7]. Interestingly, bitter taste receptors vary among species: a much smaller T2Rs gene repertoire exists in birds (range from 1 in the domestic

pigeon to 12 in the bar-tailed trogon) in comparison to 10–69 in mammals [8, 9], and chickens (*Gallus gallus domesticus*) have only three functional bitter taste receptors comprising T2R1, T2R2, and T2R7 [10–12]. Intriguingly, white-throated sparrows (*Zonotrichia albicollis*) were found to have 18 putatively functional T2Rs, indicating their sensitivity to wider range of bitter substances [13].

In recent years, tremendous progress has been made to elucidate different taste senses mediated by taste receptors located in oral cavity. In birds, bitter taste sensitivity varies within and between species. For instance, laying hens and turkeys have a high response to glucosinolates than broilers [14]. A behavioral study showed that T2R1 is highly sensitive

to dextromethorphan (Dex) and diphenidol (Dip) agonists affecting the behavioral responses to bitterness in chicken [15]. Recently, 6-methoxyflavanone has been identified as an antagonist for functional bitter receptors (T2R1 and T2R7) of oral epithelial cells through Ca^{2+} imaging technology [16]. Moreover, cell-based assay showed high expression levels of *T2R7* and *T2R1* in Vimentin^{-ve} taste bud cells, indicating that T2R7 and T2R1 mediate the functional bitter and umami sensation in chicken, respectively [17]. Furthermore, previous studies had reported that chickens and cockatiels (*Nymphicus hollandicus*) are similar to humans in their sensitivity to quinine, but superior to many other mammals [18, 19]. Additionally, chickens are more responsive to glucosinolates than ruminants [20]. Recently, hen protein hydrolysate (HPH) peptides are screened as the blockers of T2R4, T2R7, and T2R14 in HEK293T cell-based assay [21].

Compared with mammals, birds possess inferior taste acuity, mainly manifested as insufficient chewing ability, relatively small number of taste buds, and low saliva secretion [22]. Although T2Rs are sensitive to several bitter substances, various T2Rs exhibit different response towards the same bitter compounds (broadly “tuned”). Recently, in terms of bitterness, after screening 46 different bitter compounds, 17 bitter compounds have been proved to activate T2R7 in chicken [23]. Interestingly, one report showed a positive correlation between the number of taste buds in different breeds of chicken and their bitterness sensitivity and that the layers appear to have a high sensitivity to quinine hydrochloride than those of broilers [24]. Previous studies have shown that the bitter taste receptor gene family (*T2Rs*) and their downstream genes (*α -gustaducin*, *PLC β 2*, and *TRPM5*) are expressed in the chicken’s gustatory and extraoral gastrointestinal tract (GIT) tissues [25, 26].

By examining quinine analogs through integrated *in silico* and *in vivo* approach, three new T2R1 agonists (epiquinidine, ethylhydrocupreine, and quinidine) are identified that contribute to increase our understanding of bitter perception in chickens [27, 28]. In chickens, 2-choice test has been established to detect taste threshold for bitter, umami, and sweet tastes which have the potential to affect feed and water consumption [29]. Cheled-shoval et al. [29] found that chickens had similar thresholds to humans for quinine (bitter) but were less sensitive to sucrose and monosodium glutamate. In chickens, a study on bitter taste system demonstrated that behavioral (*in vivo*) thresholds are similar or up to two orders of magnitude higher than the heterologous (*in vitro*) ones [30].

In chickens, recent studies have specifically identified bitter taste receptors activated by various bitter compounds [15, 23]. In our previous study on *T2R7* gene, we found that haplotype HE1 corresponding to the Tibetan chicken was positively associated with high-altitude adaptation, while haplotype HE4 corresponding to the Jiuyuan black-chicken showed a negative correlation with high-altitude adaptation [31]. It shows that the elucidation of the chicken bitter taste sensitivity to different bitter compounds is extremely important in the development of chicken feed efficiency in poultry production, but the bitterness sensitivity mediated

molecular mechanism is still to be explored. Therefore, in this study, we focused on the investigation of the influence of different bitter compounds on the expression of bitter taste receptor T2R7 and downstream signaling effectors in *cT2R7/pDisplay-G α 16/gust44/pcDNA3.1 (+)* cells.

2. Materials and Methods

2.1. Chemicals. Camphor, chlorpheniramine maleate, erythromycin, benzoin, chloramphenicol, quinine, parthenolide, and KCL were purchased from Sangon Biotech Co. Ltd. Chengdu, China. For individual and cell stimulation experiments, KCL was dissolved in ultrapure water to prepare a 50 mM KCL solution. Other compounds were dissolved in 50 mM KCL solution to make a 5 mM stock solution, and the 5 mM stock solution was diluted with 50 mM KCL solution to make 1, 0.1, 10^{-3} , and 10^{-5} mM solutions. These solutions of bitter chemical compounds were stored at -20°C until use.

2.2. Animals and Individual Tests. A total of 108 Jiuyuan black-chickens (54 males and 54 females) at 120 days of age were used in this study. The same feeding management and regimens were provided to all chickens. This research was conducted under the Welfare and Management of Animals Act and approved by the Committee on the Care and Use of Laboratory Animals of the State-Level Animal Experimental Teaching Demonstration Centre of Sichuan Agricultural University (Approval ID: Decree number S20160906). Experimental procedures were performed in accordance with the Regulations for the Administration of Affairs Concerning Experimental Animals (China, 1988) for animal experiments. All efforts were made to minimize the suffering of the chickens, and no animal was injured during capture. Each independent experiment was performed in triplicate, and all three treated chickens had a similar weight. Briefly, the chemical compound test solutions (1 mL) were given with water to randomly selected chickens during the experimental period. After the exposure to the chemical compound for 24 hours, the whole tongue tissue samples were rapidly collected, frozen in liquid nitrogen, and stored at -80°C until use.

2.3. Preparation of Chicken T2R7-1 and T2R7-2 Plasmid Constructs. We fixed the point mutations using site-directed mutagenesis kit (Sangon Biotech Co. Ltd. Chengdu, China) to obtain haplotype HE1 (the unique haplotype of the Tibetan chicken) and haplotype HE4 (the unique haplotype of the Jiuyuan black-chicken), which were identified in our previous research [31]. The mutations identified in cDNA sequences of *T2R7-1* (haplotype HE1, KT377158) and *T2R7-2* (haplotype HE4, KT377161) and corresponding amino acid residues are shown in Table 1. The PCR products of the *T2R7-1* and *T2R7-2* ORF were subcloned into the pDisplay™ (Invitrogen) mammalian expression vector by using In-Fusion HD Cloning Kit (TaKaRa Biotech Corporation, Dalian, China). The endotoxin-free recombinant plasmids *cT2R7-1/pDisplay* and *cT2R7-2/pDisplay* were extracted

TABLE 1: Polymorphic sites identified in the coding region sequence of *T2R7-1* and *T2R7-2* gene in chicken.

| Nucleotide position | KT377158 (<i>T2R7-1</i>) | KT377161 (<i>T2R7-2</i>) | Mutation type | Amino acid change |
|---------------------|----------------------------|----------------------------|---------------|-------------------|
| 280 | A | C | Transversion | Met > Leu |
| 766 | A | T | Transversion | Thr > Ser |
| 802 | G | A | Transition | Val > Met |

using the E.Z.N.A. Endo-Free Plasmid Mini Kit I (Omega Bio-tek®) following the manufacturer's protocol. *Gα16/gust44/pcDNA3.1 (+)* was purchased from Sangon Biotech Co. Ltd. Chengdu, China.

2.4. Cell Cultures. Human embryonic kidney-derived 293T (HEK-293T) cells were kindly donated by Pro. Jiwen Wang's waterfowl research group (Sichuan Agricultural University). HEK-293T cells were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS at 37°C under 5% CO₂ overnight. Briefly, when the HEK-293T cells were at ~80% confluence, the cells were cotransfected with *Gα16/gust44/pcDNA3.1 (+)* and *cT2R7-1/pDisplay* or *cT2R7-2/pDisplay* (the ratio, *T2R7-pDisplay: Ga16gust44-pcDNA3.1 (+)* = 4:1) in a six-well plate using LipofectAMINE2000 (Invitrogen). About 24 hours after transfection, the cells were stimulated with 0.1 mM and 10⁻³ mM chemical compound test solutions (described above) for 1 hour. Total RNA was collected from the cells and used to evaluate the relative mRNA expression of *PLCβ2* and *ITPR3*.

2.5. RNA Extraction and Real-Time PCR. Total RNA was extracted using RNAiso Plus (TaKaRa Biotech Corporation, Dalian, China) kit and reverse transcription was performed using the PrimeScript™ RT Reagent Kit with gDNA Eraser (TaKaRa Biotech Corporation, Dalian, China). Primers for quantitative real-time PCR were designed using Primer v5.0, listed in Table 2. PCR mixture consisted of 1 μL of cDNA, 0.8 μL of 10 pmol/μL each primer, 12.5 μL of SYBR1Premix Ex Taq™ II (TaKaRa Biotech Corporation, Dalian, China), and 9.9 μL of double-distilled H₂O in a final reaction volume of 25 μL. Thermocycling parameters were 95°C for 2 min followed by 40 cycles of 95°C for 5 s, the annealing temperature for 30 s, and 60°C for 30 s.

2.6. Statistical Analysis. Statistical analysis was performed using one-way ANOVA with the SAS v9.0 statistical software package (SAS Institute Inc., Cary, NC, USA) to test the differences among treatments. All data are presented as the mean ± standard error (SE), and *P* value < 0.05 is considered statistically significant.

3. Results

3.1. Effect of Different Bitter Compounds on the Expression of *T2R7* mRNA in Tongue Tissue of Chicken. To investigate the effect of various bitter compounds in chicken, tongue tissues of chicken were stimulated with different chemical compound solutions. The relative fold changes in *T2R7* mRNA expression from the RT-PCR experiments are summarized

in Figure 1. The effect of different concentrations of camphor solution on the *T2R7* mRNA expression was significant (*P* < 0.05) (Figure 1(a)). In camphor solution, the relative fold change of 5 mM was significantly higher than that of the other concentrations (*P* < 0.05). Camphor solutions of 1, 0.1, and 10⁻³ mM concentrations had significantly higher fold changes than solutions of 10⁻⁵ mM (*P* < 0.05), but no significant difference was found among the three camphor solutions (*P* > 0.05). Moreover, there was no significant difference detected in *T2R7* mRNA expression between 10⁻⁵ mM camphor solution and 50 mM KCL solution (*P* > 0.05). The effects of chlorpheniramine maleate solution with different concentrations on the expression of *T2R7* mRNA are shown in Figure 1(b). The expression level of *T2R7* mRNA in 5 mM chlorpheniramine maleate solution was significantly increased, compared with other lower concentrations (*P* < 0.05), while there were no significant differences observed between chlorpheniramine solutions (1, 0.1, 10⁻³, and 10⁻⁵ mM) and 50 mM KCL solution (*P* > 0.05).

Interestingly, the relative mRNA abundance of *T2R7* of chicken tongue tissue was remarkably different in all different concentrations of erythromycin solutions and 50 mM KCL solution (Figure 1(c)). We found that *T2R7* mRNA expression in 5 mM erythromycin solution was significantly higher than that in other solutions (*P* < 0.05). The fold changes of 1, 0.1, and 10⁻³ mM erythromycin solutions were significantly higher than those of 50 mM KCL solutions (*P* < 0.05), but there was no difference between erythromycin solutions (*P* > 0.05). The difference between the 10⁻⁵ mM erythromycin solution and 50 mM KCL solution was not significant (*P* > 0.05). However, the relative expression levels of *T2R7* mRNA were not significantly different between the different concentration gradients of benzoin (Figure 1(d)), chloramphenicol (Figure 1(e)), quinine (Figure 1(f)), parthenolide (Figure 1(g)), and 50 mM KCL solutions (*P* > 0.05).

3.2. Effect of Different Bitter Compounds on the Expression of *PLCβ2* mRNA in *cT2R7/pDisplay-Gα16/gust44/pcDNA3.1 (+)* Cells. We determined two concentration solutions (0.1 mM and 10⁻³ mM) with reference to the results shown in Figure 2. We found that after exposure to 0.1 and 10⁻³ mM camphor solutions the relative *PLCβ2* mRNA expression was significantly higher than that of the vector-control in *cT2R7-1/pDisplay-Gα16/gust44/pcDNA3.1 (+)* cells (*P* < 0.05) (Figure 2(a)). On the other hand, the stimulus of 0.1 mM camphor solution increased expression level of *PLCβ2* mRNA significantly than that of the 10⁻³ mM camphor solution and vector-control in *cT2R7-2/pDisplay-Gα16/gust44/pcDNA3.1 (+)* cells (*P* < 0.05) (Figure 2(a)). Likewise, in *cT2R7-1/pDisplay-Gα16/gust44/pcDNA3.1 (+)*

TABLE 2: Detailed information of the primers used in real-time PCR analysis.

| Primer | Forward primer | Reverse primer | Fragment length (bp) | Annealing temperature (°C) | Accession numbers |
|----------------|-------------------------|---------------------------|----------------------|----------------------------|-------------------|
| <i>GAPDH-1</i> | CCAGAACATCATCCCAGCGTC | ACGGCAGGTCAGGTCAACAA | 136 | 60.0 | NM_204305.1 |
| <i>GAPDH-2</i> | CTTTGGTATCGTGGAAGGACTC | GTAGAGGCAGGGATGATGTTCT | 132 | 60.0 | NM_002046 |
| <i>T2R7</i> | TTCAGGCACCATTCTTCATCTAC | TGGGGCTGGTTCTGTTCTCT | 142 | 60.0 | NM_001080719.1 |
| <i>PLCβ2</i> | AAGATGCCCAAGAGCCAGAAG | GGAGACGAAGTTGTGGAAGGTG | 132 | 56.4 | NM_001284299.1 |
| <i>ITPR3</i> | TCCTGTTCTTCTCATCGTCATCA | TTGTTATCAAACCTGTCCCTCTCCA | 160 | 60.0 | NM_002224.3 |




FIGURE 1: Continued.




FIGURE 1: Relative expression levels of *T2R7* mRNA in tongue tissue of chicken stimulated by different concentrations of (a) camphor, (b) chlorpheniramine maleate, (c) erythromycin, (d) benzoin, (e) chloramphenicol, (f) quinine, and (g) parthenolide, as determined by real-time PCR. The relative gene expression was measured by qPCR and normalized to GAPDH, and a 50 mM KCL solution was the negative control. Each bar represents the mean \pm SE of the results from 2 to 3 independent experiments performed in triplicate. Different small letters on the bar indicate significant differences at $P < 0.05$.




FIGURE 2: Continued.




FIGURE 2: Relative expression levels of *PLCβ2* mRNA in *cT2R7-1*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) and *cT2R7-2*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells stimulated by different concentrations of (a) camphor, (b) chlorpheniramine maleate, (c) erythromycin, (d) benzoin, (e) chloramphenicol, (f) quinine, and (g) parthenolide, as determined by real-time PCR. The relative gene expression was measured by qPCR and normalized to GAPDH, and the vector-control was the negative control. Each bar represents the mean ± SE of the results from 2 to 3 independent experiments performed in triplicate. Different small letters on the bar indicate significant differences at $P < 0.05$.

cells, the relative *PLCβ2* mRNA expression after exposure to stimuli 0.1 and 10^{-3} mM chlorpheniramine maleate solutions was significantly higher than that of the vector-control ($P < 0.05$), whereas there was no remarkable difference measured between chlorpheniramine maleate solutions and the vector-control in *cT2R7-2*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P > 0.05$) (Figure 2(b)). The stimulus of 10^{-3} mM erythromycin solution remarkably increased relative *PLCβ2* mRNA abundance than that of the 0.1 mM erythromycin solution and vector-control ($P < 0.05$), but the 0.1 mM erythromycin solution showed no significant difference with the other two treatments in *cT2R7-1*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P > 0.05$). However, there was no significant difference in fold changes among the three treatment groups in *cT2R7-2*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P > 0.05$) (Figure 2(c)).

Furthermore, after exposure with chemical compounds benzoin (Figure 2(d)) and chloramphenicol (Figure 2(e)), a significant increase occurred in the relative expression level of *PLCβ2* mRNA in the 10^{-3} mM solution compared with that in the other concentration in *cT2R7-1*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P < 0.05$), whereas the all three treatment groups were not significantly different in *cT2R7-2*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P > 0.05$). As

illustrated in Figure 2(f), the relative expression levels of *PLCβ2* mRNA were not significantly influenced by different concentrations of quinine compared with the vector-control ($P > 0.05$). The change in relative expression levels of *PLCβ2* mRNA caused by parthenolide is summarized in Figure 2(g). The stimulus of 10^{-3} mM solution had a significantly higher relative expression level than that of the 0.1 mM solution and vector-control, and the level in the 0.1 mM solution was significantly higher than that of the vector-control in *cT2R7-1*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P < 0.05$). By contrast, all the three treatment groups were not significantly different in *cT2R7-2*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells ($P > 0.05$).

3.3. Effect of Different Bitter Compounds on the Expression of *ITPR3* mRNA in *cT2R7*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) Cells. In addition to the above analysis, *cT2R7*/pDisplay-*Gα16*/*gust44*/pcDNA3.1 (+) cells were treated with 0.1 mM and 10^{-3} mM compound solutions, and the expression of *ITPR3* mRNA was measured by real-time PCR. The relative expression level is summarized in Figure 3. After exposure with camphor (Figure 3(a)) and quinine (Figure 3(f)) compounds, a significant increase occurred in the relative expression levels of *ITPR3* mRNA in the 0.1 and 10^{-3} mM




FIGURE 3: Continued.




FIGURE 3: Relative expression levels of *ITPR3* mRNA in *cT2R7-1*/pDisplay-G α 16/gust44/pcDNA3.1 (+) and *cT2R7-2*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells stimulated by different concentrations of (a) camphor, (b) chlorpheniramine maleate, (c) erythromycin, (d) benzoin, (e) chloramphenicol, (f) quinine, and (g) parthenolide, as determined by real-time PCR. The relative gene expression was measured by qPCR and normalized to GAPDH, and the vector-control was the negative control. Each bar represents the mean \pm SE of the results from 2 to 3 independent experiments performed in triplicate. Different small letters on the bar indicate significant differences at $P < 0.05$.

solutions compared with that of the vector-control in *cT2R7-1*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P < 0.05$). The expression level in the stimulus of 10^{-3} mM camphor solution was significantly higher than that in the 0.1 mM camphor solution and vector-control, and the expression level of the 0.1 mM solution was significantly higher than that of the vector-control in *cT2R7-2*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P < 0.05$). On the other hand, there was no significant difference detected among the three treatment groups for quinine ($P > 0.05$). After exposure with chlorpheniramine maleate, the stimuli of 10^{-3} mM solution was significantly increased in the relative expression of *ITPR3* mRNA compared with that of 0.1 mM and vector-control in *cT2R7-1*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P < 0.05$) (Figure 3(b)); however, no significant differences were found in expression level of *ITPR3* mRNA between these stimuli in *cT2R7-2*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P > 0.05$). After exposure with erythromycin (Figure 3(c)), benzoin (Figure 3(d)), and chloramphenicol (Figure 3(e)) compounds, a significant increase occurred in the relative expression level with the stimulus of 0.1 mM solution compared with that in the other concentrations ($P < 0.05$). Moreover, the level in the 10^{-3} mM erythromycin solution was significantly higher than that in the vector-control in *cT2R7-1*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P < 0.05$). By contrast, all the three treatment groups for chloramphenicol were not significantly different in *cT2R7-2*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P > 0.05$). Similar patterns of expression level were observed in both the *cT2R7-1*/pDisplay-G α 16/gust44/pcDNA3.1 (+) and *cT2R7-2*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells after exposure with parthenolide solution. We determined that the 10^{-3} mM parthenolide solution had a significantly higher relative expression level than that of the 0.1 mM solution and vector-control ($P < 0.05$). Further, the level of mRNA expression in the 0.1 mM parthenolide solution was significantly higher than that of the vector-control in *cT2R7-1*/pDisplay-G α 16/gust44/pcDNA3.1 (+) cells ($P < 0.05$) (Figure 3(g)).

4. Discussion

Tibetan chickens living in Tibet plateau have adapted to the special environment and developed unique genetic predispositions and dietary habits [32]. Jiuyuan black-chickens used in the study are important as egg layers. It has been demonstrated that the broiler chickens (broiler-type) are more sensitive to bitter taste than White Leghorn (layer-type) [33]. Previously, researchers found a relationship between the number of taste buds and taste sensitivity [33, 34]. Furthermore, optimizing the consumption of balanced diets plays a central role in the welfare, development, health, and productivity of animals, especially when raised or preserved in captivity [35].

In this study, we ultimately explore the sensibility of taste receptor proteins to bitter compounds in order to ascertain whether chickens are sensitive to several bitter compounds with concentrations gradient. Significant changes in mRNA expression of *T2R7* were found in a dose-dependent fashion after induction with camphor, chlorpheniramine maleate ($P < 0.05$), and erythromycin solutions, but there was no significant difference found between the different concentrations in the other four compounds ($P > 0.05$). These results implied that the sensitivity of chickens to bitter compounds (camphor, chlorpheniramine maleate, and erythromycin) can be determined by the relative expression of *T2R7* mRNA in tongue tissues. The postnatal administration of the bitter tasting quinine to chicks increases the mRNA expression level of *T2R1* and *T2R7* genes in the chicken's tongue [25], and four chemical substances chloramphenicol, chlorpheniramine, diphenidol, and quinine sulfate activate all three bitter taste receptors in chickens [23]. However, the expression of *T2R7* mRNA was not related to the intake of quinine and chloramphenicol at different concentrations. Recent behavioral experiments reveal that baby chicks have a greater aversion to salt and sour taste qualities than in adults [36]. A latest research on chickens shows that younger chicks are more sensitive to

bitter compounds compared with older chicks [37]. Previously, it was reported that NaCl solutions of 85 mM and 100 mM are the preferred solutions for chickens, while chickens refused to accept NaCl solutions of 250 mM or higher concentration due to aversive taste [38]. In this study, we used Jiuyuan black-chickens at age of 120 days, which were fed the normal commercial feed formulation. Thus, we speculate that the age of chickens resulted in a reduction in their sensitivity to bitter compounds. Additionally, normal commercial feed does not contain strongly bitter substances [37]; therefore, the bitter taste experience of these chickens was not extensive under the experimental conditions of this study. Moreover, the long-term feed formulation and highly selective breeding of farm-raised chickens expose them to less toxic bitter substances that exist in nature. Therefore, we speculate that this series of artificial interventions resulted in the degeneration of the *T2R7* gene during the long-term breeding process of chicken. Certainly, additional studies are necessary to confirm this hypothesis.

Furthermore, quinine stimulation elevated the mRNA level of *ITPR3* without affecting the *T2R7* and *PLCβ2* mRNA levels, and these results are consistent with earlier studies which show that chickens are less sensitive to quinine compared to the other bitter compounds [30, 39]. Our findings are in accordance with a recent study which demonstrated that *IP3R3*-mediated Ca^{2+} flux is strongly inhibited by quinine using heterologous systems and/or cell lines [40]. The gustducin activates phospholipase C (*PLCβ2*) and catalyzes phosphatidylinositol phosphate to produce the second messengers inositol 1,4,5 trisphosphate (*IP₃*) and diacylglycerol (*DAG*), leading to calcium release [41, 42]. A prior study using knockout mice reported that the elimination of *PLCβ2*, *TrpM5*, or *IP₃* receptor proteins leads to a reduction or complete loss of sensitivity to bitter, sweet, or umami taste suggesting that these taste sensations strongly favor a conserved, streamline signaling cascade [43]. Thus, a bitter substance can induce the activation of the bitter receptor protein and enhance the relative *PLCβ2* and *ITPR3* mRNA expression. These results provide favorable evidence for reducing the sensitivity of different bitter compounds to bitter taste, although further research is required to confirm this hypothesis.

Bitter compounds elicit innate aversive response across species, which is typically considered to prevent the ingestion of toxic substances [44, 45]. Hirai et al. established correlation between patients' taste sensitivity, perception, and its mRNA expression, which increases the expression of taste receptor genes linked to phantogeusia pathogenesis by enhancing bitter taste sensation [46]. We screened several bitter compounds identified through inquiry of the BitterDB probe (<http://bitterdb.agri.huji.ac.il/bitterdb/>, [47]) to explore the relative mRNA expression levels in downstream signaling genes (*PLCβ2* and *ITPR3*) of chicken. Previous reports have described *IP3R3* as the dominant form of the *IP₃* receptor which plays an important role in bitter taste transduction [48, 49]. In addition, knocking out of *PLCβ2* and *ITPR3* reduces the ability of mice to recognize most bitter compounds [50]. We found that the relative mRNA expression levels of *PLCβ2* and *ITPR3* in *cT2R7-1/pDisplay-Gα16/gust44/pcDNA3.1 (+)* cells induced by different bitter

compounds, such as 0.1 and 10^{-3} mM camphor, 10^{-3} mM chlorpheniramine maleate, 10^{-3} mM erythromycin, 10^{-3} mM quinine, and 0.1 mM and 10^{-3} mM parthenolide, were significantly higher than that of the vector-control ($P < 0.05$). Previously, *PLCβ2* and *ITPR3* were identified as the key genes in the signal transduction pathway of bitter taste [49–51]. Bitter taste signaling is believed to be complex, and the pathways that transduce the taste response to bitter compounds lead to cation channel opening, depolarization of the taste receptor cells (TRCs), and subsequent neurotransmitter release [50]. In particular, chemical bitter compounds (quinine, garlic oil, almond oil, clove oil, and magnesium chloride) play a crucial role in reducing feather pecking incidence in laying hens [52–54].

By contrast, the mRNA expression of *PLCβ2* and *ITPR3* in *cT2R7-2/pDisplay-Gα16/gust44/pcDNA3.1 (+)* cells stimulated by 0.1 mM camphor was significantly higher than that of the vector-control ($P < 0.05$). Primarily, chickens appear to have an acute sense of taste allowing the discrimination of five primary tastes including fatty, umami, salty, sour, and bitter [55]. Interestingly, dietary preferences are suppressed when the tongue becomes paralyzed, suggesting the role of olfaction in dietary fat preferences [56]. Meanwhile, T2Rs have been identified in nonoral tissues, including the respiratory tract, the gut, the spleen, the lung, the heart, and bursa Fabricius, suggesting its involvement in other physiological functions such as appetite regulation, innate immunity, and internal organs, besides taste [57, 58]. In addition, preferences for a balanced diet containing synthetic AA (potentially related to umami taste) suggest that lysine, methionine, and tryptophan in a diet increase feed intake [59]. It has been reported that PKD1L3 and PKD2L1 ion channels function as a likely candidate for a mammalian sour taste receptor related to the generation of acidic taste [60]. Bitter taste receptor harbors multiple single nucleotide polymorphisms (SNPs) in the coding region, which are associated with dietary preferences, metabolic traits, and body mass index [61].

Our results reveal that bitter taste receptor (*T2R7*) is sensitive to different concentrations of bitter compounds and *T2R7-1* (corresponding to the unique haplotype of the Tibetan chicken) had higher sensitivity to bitter compounds compared with that of *T2R7-2* (corresponding to the unique haplotype of the Jiuyuan black-chicken). Similarly, recent studies have shown an important link between the haplotypes and the ability to identify bitter compounds [62]. They showed that the sequence variants of *T2R38* gene had a direct influence on phenylthiocarbamide (PTC) taste sensitivity, in which the haplotype PAV had greater PTC sensitivity compared with that of the haplotype AVI. In addition, Tibetan chickens raised outside cage may absorb or detect more natural bitter substances than Jiuyuan black-chickens bred in cages. The long-term feed formulation and highly selective breeding may reduce the sensitivity of Jiuyuan black-chickens to bitter substances. Overall, chicken chemosensory research has been applied to chicken initial choice of feed and the level of feed consumption, which is critical to the understanding of molecular mechanisms of chicken taste.

5. Conclusions

In summary, we evaluated the sensitivity of bitter taste receptor (T2R7) and its downstream signaling molecule on synthetic bitter compounds for chicken. The bitter taste receptor T2R7 was found sensitive to different concentrations of bitter compounds. Furthermore, bitter taste receptor T2R7-1 (corresponding to the unique haplotype of the Tibetan chicken) has higher sensitivity to bitter compounds compared with that of T2R7-2. Our results will be helpful in the enhancement of chicken feed efficiency as well as beneficial to future taste sensation research in context of improving feeding strategies among Aves.

Data Availability

All data are available within the article or available from the authors upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Yuan Su and Hang Jie contributed equally to this work. Diyan Li and Qing Zhu conceived and designed the experiments. Yuan Su and Hang Jie drafted the manuscript. Yuan Su and Yan Wang conducted the experiments. Xiaoling Zhao and Huadong Yin analyzed the data. Proofreading and editing is done by Shailendra Kumar Mishra and Yuan Su. All authors read and approved the final draft of the manuscript.

Acknowledgments

This study was financially supported by the National Natural Science Foundation of China (31402063). The authors thank Prof. Jiwen Wang (Sichuan Agricultural University) for providing the HEK-293T cells.

References

- [1] S. C. Kinnamon and T. A. Cummings, "Chemosensory transduction mechanisms in taste," *Annual Review of Physiology*, vol. 54, no. 54, pp. 715–731, 1992.
- [2] B. Lindemann, "Taste reception," *Physiological Reviews*, vol. 76, no. 3, pp. 719–766, 1996.
- [3] J. B. Epstein and A. Barasch, "Taste disorders in cancer patients: pathogenesis, and approach to assessment and management," *Oral Oncology*, vol. 46, no. 2, pp. 77–81, 2010.
- [4] J. B. Epstein, G. Smutzer, and R. L. Doty, "Understanding the impact of taste changes in oncology care," *Supportive Care in Cancer*, vol. 24, no. 4, pp. 1917–1931, 2016.
- [5] A. J. Hovan, P. M. Williams, P. Stevenson-Moore et al., "A systematic review of dysgeusia induced by cancer therapies," *Supportive Care in Cancer*, vol. 18, no. 8, pp. 1081–1087, 2010.
- [6] H. Matsunami, J.-P. Montmayeur, and L. B. Buck, "A family of candidate taste receptors in human and mouse," *Nature*, vol. 404, no. 6778, pp. 601–604, 2000.
- [7] D. Li and J. Zhang, "Diet shapes the evolution of the vertebrate bitter taste receptor gene repertoire," *Molecular Biology and Evolution*, vol. 31, no. 2, pp. 303–309, 2014.
- [8] K. Wang and H. Zhao, "Birds generally carry a small repertoire of bitter taste receptor genes," *Genome Biology and Evolution*, vol. 7, no. 9, pp. 2705–2715, 2015.
- [9] C. Conte, M. Ebeling, A. Marcuz, P. Nef, and P. J. Andres-Barquin, "Evolutionary relationships of the Tas2r receptor gene families in mouse and human," *Physiological Genomics*, vol. 14, no. 1, pp. 73–82, 2003.
- [10] L. Clark, J. Hagelin, and S. Werner, "The chemical senses in birds," in *Sturkie's Avian Physiology*, pp. 89–111, Elsevier, Amsterdam, The Netherlands, 6th edition, 2015.
- [11] Y. Go, "Lineage-specific expansions and contractions of the bitter taste receptor gene repertoire in vertebrates," *Molecular Biology and Evolution*, vol. 23, no. 5, pp. 964–972, 2006.
- [12] D. Dong, G. Jones, and S. Zhang, "Dynamic evolution of bitter taste receptor genes in vertebrates," *BMC Evolutionary Biology*, vol. 9, no. 1, p. 12, 2009.
- [13] J. K. Davis, J. J. Lowman, P. J. Thomas et al., "Evolution of a bitter taste receptor gene cluster in a new world sparrow," *Genome Biology and Evolution*, vol. 2, pp. 358–370, 2010.
- [14] G. R. Fenwick and R. F. Curtis, "Rapeseed meal and its use in poultry diets. A review," *Animal Feed Science and Technology*, vol. 5, no. 4, pp. 255–298, 1980.
- [15] N. Hirose, Y. Kawabata, F. Kawabata, S. Nishimura, and S. Tabata, "Bitter taste receptor T2R1 activities were compatible with behavioral sensitivity to bitterness in chickens," *Biochemical and Biophysical Research Communications*, vol. 460, no. 2, pp. 464–468, 2015.
- [16] B. Dey, "Identification of functional bitter taste receptors and their antagonist in chickens," *Biochemical & Biophysical Research Communications*, vol. 482, no. 4, pp. 693–699, 2016.
- [17] Y. Yoshida, Z. Wang, K. F. Tehrani et al., "Bitter taste receptor T2R7 and umami taste receptor subunit T1R1 are expressed highly in Vimentin-negative taste bud cells in chickens," *Biochemical and Biophysical Research Communications*, vol. 511, no. 2, pp. 280–286, 2019.
- [18] H. Ueda and S. Kainou, "Aversion to quinine is associated with taste sensation in chicks," *The Journal of Poultry Science*, vol. 42, no. 3, pp. 254–262, 2005.
- [19] K. D. Matson, J. R. Millam, and K. C. Klasing, "Cockatiels (*Nymphicus hollandicus*) reject very low levels of plant secondary compounds," *Applied Animal Behaviour Science*, vol. 85, no. 1–2, pp. 141–156, 2004.
- [20] M. K. Tripathi and A. S. Mishra, "Glucosinolates in animal nutrition: a review," *Animal Feed Science and Technology*, vol. 132, no. 1–2, pp. 1–27, 2007.
- [21] Q. Xu, N. Singh, H. Hong et al., "Hen protein-derived peptides as the blockers of human bitter taste receptors T2R4, T2R7 and T2R14," *Food Chemistry*, vol. 283, pp. 621–627, 2019.
- [22] E. Roura, M. W. Baldwin, and K. C. Klasing, "The avian taste system: potential implications in poultry nutrition," *Animal Feed Science & Technology*, vol. 180, no. 1–4, pp. 1–9, 2013.
- [23] M. Behrens, S. I. Korsching, and W. Meyerhof, "Tuning properties of avian and frog bitter taste receptors dynamically fit gene repertoire sizes," *Molecular Biology and Evolution*, vol. 31, no. 12, pp. 3216–3227, 2014.
- [24] K.-I. Kudo, J.-I. Shiraishi, S. Nishimura, T. Bungo, and S. Tabata, "The number of taste buds is related to bitter taste sensitivity in layer and broiler chickens," *Animal Science Journal*, vol. 81, no. 2, pp. 240–244, 2010.
- [25] S. L. Cheled-Shoval, M. Behrens, W. Meyerhof, M. Y. Niv, and Z. Uni, "Perinatal administration of a bitter tastant influences

- gene expression in chicken palate and duodenum," *Journal of Agricultural and Food Chemistry*, vol. 62, no. 52, pp. 12512–12520, 2014.
- [26] S. L. Cheled-Shoval, S. Druyan, and Z. Uni, "Bitter, sweet and umami taste receptors and downstream signaling effectors: expression in embryonic and growing chicken gastrointestinal tract," *Poultry Science*, vol. 94, no. 8, pp. 1928–1941, 2015.
- [27] P. A. Di, "Ligand binding modes from low resolution GPCR models and mutagenesis: chicken bitter taste receptor as a test-case," *Scientific Reports*, vol. 7, no. 1, p. 8223, 2017.
- [28] P. A. Di, "Molecular features underlying selectivity in chicken bitter taste receptors," *Frontiers in Molecular Biosciences*, vol. 5, p. 6, 2018.
- [29] S. L. Cheled-Shoval, N. Reicher, M. Y. Niv, and Z. Uni, "Detecting thresholds for bitter, umami, and sweet tastants in broiler chicken using a 2-choice test method," *Poultry Science*, vol. 96, no. 7, pp. 2206–2218, 2017.
- [30] S. Cheled-Shoval, M. Behrens, A. Korb et al., "From cell to beak: in-vitro and in-vivo characterization of chicken bitter taste thresholds," *Molecules*, vol. 22, no. 5, p. 821, 2017.
- [31] Y. Su, D. Li, U. Gaur et al., "Genetic diversity of bitter taste receptor gene family in Sichuan domestic and Tibetan chicken populations," *Journal of Genetics*, vol. 95, no. 3, pp. 675–681, 2016.
- [32] M.-S. Wang, Y. Li, M.-S. Peng et al., "Genomic analyses reveal potential independent adaptation to high altitude in Tibetan chickens," *Molecular Biology and Evolution*, vol. 32, no. 7, pp. 1880–1889, 2015.
- [33] D. Ganchrow and J. R. Ganchrow, "Number and distribution of taste buds in the oral cavity of hatchling chicks," *Physiology & Behavior*, vol. 34, no. 6, pp. 889–894, 1985.
- [34] J. R. Ganchrow and D. Ganchrow, "Taste bud development in chickens (*Gallus gallus domesticus*)," *The Anatomical Record*, vol. 218, no. 1, pp. 88–93, 1987.
- [35] S. Niknafs and E. Roura, "Nutrient sensing, taste and feed intake in avian species," *Nutrition Research Reviews*, vol. 31, no. 2, pp. 256–266, 2018.
- [36] H. Berkhoudt, *Avian Taste Buds: Topography, Structure and Function*, Springer, Boston, MA, USA, 1992.
- [37] B. Dey, F. Kawabata, Y. Kawabata, S. Nishimura, and S. Tabata, "Bitter taste sensitivity and the expression of bitter taste receptors at different growth stages of chicks," *The Journal of Poultry Science*, vol. 55, no. 3, pp. 204–209, 2018.
- [38] C. J. Duncan, "Salt preferences of birds and mammals," *Physiological Zoology*, vol. 35, no. 2, pp. 120–132, 1962.
- [39] Y. Yoshida, F. Kawabata, Y. Kawabata, S. Nishimura, and S. Tabata, "Expression levels of taste-related genes in palate and tongue tip, and involvement of transient receptor potential subfamily M member 5 (TRPM5) in taste sense in chickens," *Animal Science Journal*, vol. 89, no. 2, pp. 441–447, 2018.
- [40] M. Gees, Y. A. Alpizar, T. Luyten et al., "Differential effects of bitter compounds on the taste transduction channels TRPM5 and IP3 receptor type 3," *Chemical Senses*, vol. 39, no. 4, pp. 295–311, 2014.
- [41] B. Lindemann, "Receptors and transduction in taste," *Nature*, vol. 413, no. 6852, pp. 219–225, 2001.
- [42] R. F. Margolskee, "Molecular mechanisms of bitter and sweet taste transduction," *Journal of Biological Chemistry*, vol. 277, no. 1, pp. 1–4, 2002.
- [43] Y. Zhang, M. A. Hoon, J. Chandrashekar et al., "Coding of sweet, bitter, and umami tastes," *Cell*, vol. 112, no. 3, pp. 293–301, 2003.
- [44] J. Chandrashekar, M. A. Hoon, N. J. P. Ryba, and C. S. Zuker, "The receptors and cells for mammalian taste," *Nature*, vol. 444, no. 7117, pp. 288–294, 2006.
- [45] J. E. Steiner, D. Glaser, M. E. Hawilo, and K. C. Berridge, "Comparative expression of hedonic impact: affective reactions to taste by human infants and other primates," *Neuroscience & Biobehavioral Reviews*, vol. 25, no. 1, pp. 53–74, 2001.
- [46] R. Hirai, K. Takao, K. Onoda, S. Kokubun, and M. Ikeda, "Patients with phantogeusia show increased expression of T2R taste receptor genes in their tongues," *Annals of Otolaryngology & Laryngology*, vol. 121, no. 2, pp. 113–118, 2012.
- [47] A. Wiener, M. Shudler, A. Levit, and M. Y. Niv, "BitterDB: a database of bitter compounds," *Nucleic Acids Research*, vol. 40, no. D1, pp. D413–D419, 2012.
- [48] T. H. Meuwissen, B. J. Hayes, and M. E. Goddard, "Prediction of total genetic value using genome-wide dense marker maps," *Genetics*, vol. 157, no. 4, pp. 1819–1829, 2001.
- [49] T. R. Clapp, L. M. Stone, R. F. Margolskee, and S. C. Kinnamon, "Immunocytochemical evidence for co-expression of Type III IP3 receptor with signaling components of bitter taste transduction," *Bmc Neuroscience*, vol. 2, no. 1, p. 6, 2001.
- [50] S. Damak, M. Rong, K. Yasumatsu et al., "Trpm5 null mice respond to bitter, sweet, and umami compounds," *Chemical Senses*, vol. 31, no. 3, pp. 253–264, 2006.
- [51] M. A. Miyoshi, K. Abe, and Y. Emori, "IP3 receptor type 3 and PLCbeta2 are co-expressed with taste receptors T1R and T2R in rat taste bud cells," *Chemical Senses*, vol. 26, no. 3, pp. 259–265, 2001.
- [52] A. Harlander-Matauschek, P. Beck, and H.-P. Piepho, "Taste aversion learning to eliminate feather pecking in laying hens, *Gallus gallus domesticus*," *Animal Behaviour*, vol. 78, no. 2, pp. 485–490, 2009.
- [53] A. H. Matauschek, P. Beck, and T. B. Rodenburg, "Effect of an early bitter taste experience on subsequent feather-pecking behaviour in laying hens," *Applied Animal Behaviour Science*, vol. 127, no. 3–4, pp. 108–114, 2010.
- [54] A. H. Matauschek and T. B. Rodenburg, "Applying chemical stimuli on feathers to reduce feather pecking in laying hens," *Applied Animal Behaviour Science*, vol. 132, no. 3–4, pp. 146–151, 2011.
- [55] A. R. Y. E. Boushy and A. F. B. V. D. Poel, *Palatability and Feed Intake Regulations*, Springer, Boston, MA, USA, 1994.
- [56] R. T. Mabayo, J.-I. Okumura, A. Hirao, S. Sugita, K. Sugahara, and M. Furuse, "The role of olfaction in oil preference in the chicken," *Physiology & Behavior*, vol. 59, no. 6, pp. 1185–1188, 1996.
- [57] E. Hamdard, Z. Lv, J. Jiang et al., "Responsiveness expressions of bitter taste receptors against denatonium benzoate and genistein in the heart, spleen, lung, kidney, and bursa Fabricius of Chinese fast yellow chicken," *Animals*, vol. 9, no. 8, p. 532, 2019.
- [58] B. Avau and I. Depoortere, "The bitter truth about bitter taste receptors: beyond sensing bitter in the oral cavity," *Acta Physiologica*, vol. 216, no. 4, pp. 407–420, 2016.
- [59] M. L. Picard, G. Uzu, E. A. Dunnington, and P. B. Siegel, "Food intake adjustments of chicks: short term reactions to deficiencies in lysine, methionine and tryptophan," *British Poultry Science*, vol. 34, no. 4, pp. 737–746, 1993.
- [60] Y. Ishimaru, H. Inada, M. Kubota, H. Zhuang, M. Tominaga, and H. Matsunami, "Transient receptor potential family members PKD1L3 and PKD2L1 form a candidate sour taste receptor," *Proceedings of the National Academy of Sciences*, vol. 103, no. 33, pp. 12569–12574, 2006.

- [61] V. B. Duffy, "Variation in oral sensation: implications for diet and health," *Current Opinion in Gastroenterology*, vol. 23, no. 2, pp. 171–177, 2007.
- [62] S. V. Lipchock, J. A. Mennella, A. I. Spielman, and D. R. Reed, "Human bitter perception correlates with bitter receptor messenger RNA expression in taste cells," *The American Journal of Clinical Nutrition*, vol. 98, no. 4, pp. 1136–1143, 2013.



Hindawi

Submit your manuscripts at
www.hindawi.com

