

## Research Article

# Decreased Levels of Serum IL-34 Associated with Cognitive Impairment in Vascular Dementia

Yang Wang,<sup>1</sup> Wei Lu,<sup>2</sup> Wenjing Ning,<sup>3,4</sup> Yan Chen ,<sup>5</sup> and Lingxing Li <sup>6</sup>

<sup>1</sup>Department of Neurology, Qingpu Branch of Zhongshan Hospital, Fudan University, Shanghai 201700, China

<sup>2</sup>Department of Intensive Rehabilitation, Shandong Provincial Third Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong Province 250000, China

<sup>3</sup>Department of Geriatric, Shandong Provincial Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong Province 250021, China

<sup>4</sup>Department of Neurology, Jinan First People's Hospital, Shandong Traditional Chinese Medicine University, Jinan, Shandong Province 250013, China

<sup>5</sup>Department of Geriatric Medicine, Shandong Provincial Third Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong Province 250000, China

<sup>6</sup>Department of Cardiovascular Medicine, Taian City Central Hospital, Shandong First Medical University & Shandong Academy of Medical Sciences, Taian, Shandong Province. 271000, China

Correspondence should be addressed to Yan Chen; [chenyan2004925@163.com](mailto:chenyan2004925@163.com) and Lingxing Li; [luckykeyan@163.com](mailto:luckykeyan@163.com)

Received 16 April 2021; Revised 5 May 2021; Accepted 13 May 2021; Published 21 May 2021

Academic Editor: Juan Yang

Copyright © 2021 Yang Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Objective.** Interleukin- (IL-) 34 is a new type of cytokine with neuroprotective effects discovered in recent years. However, the relationship between IL-34 and vascular dementia (VaD) has not yet been elucidated. The purpose of this study is to determine whether IL-34 is involved in cognitive impairment of VaD. **Methods.** From January 2017 to December 2020, 84 VaD patients and 60 healthy controls who attended Qingpu Branch of Zhongshan Hospital were prospectively included in the study. Once included in the study, demographic features of all research subjects are collected. They include age, gender, education, white blood cells (WBC), neutrophil, lymphocyte, systolic blood pressure (SBP), diastolic blood pressure (DBP), fasting blood glucose (FBG), triglycerides (TG), and total cholesterol (TC). Meanwhile, the Montreal Cognitive Assessment (MoCA) scale was used to assess the cognitive function of participants. The serum IL-34 level was determined by enzyme-linked immunosorbent assay (ELISA). **Results.** There was no significant difference between the demographic features of VaD patients and healthy controls ( $p > 0.05$ ). However, the serum IL-34 levels of VaD patients and healthy controls are  $27.6 \pm 3.9$  pg/ml and  $41.8 \pm 6.0$  pg/ml, respectively, and there is a significant statistical difference between them ( $p < 0.001$ ). The results of bivariate correlation analysis showed that serum IL-34 levels were significantly positively correlated with MoCA scores ( $r = 0.371$ ,  $p = 0.023$ ). Further regression analysis showed that IL-34 was still correlated with MoCA after adjusting for demographic features ( $\beta = 0.276$ ,  $p = 0.038$ ). **Conclusions.** Serum IL-34 levels in VaD patients were significantly reduced, which may be an independent predictor of cognitive impairment in VaD patients.

## 1. Introduction

Vascular dementia (VaD) is a common type of dementia. The main cause of VaD is cerebrovascular disease, which leads to impaired cerebral blood flow and further damages neurons in the hippocampus or cortex [1, 2]. According to statistics from the World Health Organization (WHO), there are cur-

rently about 36 million people with dementia in the world. By 2050, this number will reach to 115 million, and VaD accounts for about 20% of them, second only to Alzheimer's disease (AD) [3]. It is reported that the prevalence of VaD in developed countries ranges from 1% to 4%, while the prevalence of VD in developing countries is about 8%, and the prevalence of VD in the elderly is rising rapidly [4]. The annual

cost of VaD is as high as 200 billion US dollars, and with the extension of human life expectancy, VaD has gradually become one of the main health problems that need to be solved urgently in countries all over the world [5]. Our existing treatment may alleviate some of the symptoms of VaD in the short term, but it cannot fundamentally slow down the disease progression or reverse the disease process [6]. Therefore, it is particularly urgent to continue to explore the potential pathogenic mechanism and therapeutic targets of VaD.

Interleukin- (IL-) 34 was first discovered by Lin's team in 2008 and is considered to be a nonspecific ligand for colony stimulating factor- (CSF-) 1 receptor (CSF-1R) [7]. In addition to CSF-1R, IL-34 can also bind to protein-tyrosine phosphatase- (PTP-)  $\zeta$  and syndecan-1. IL-34 is a homoglycan protein secreted mainly by brain neuron cells and skin keratinocytes, and it is highly conserved in vertebrates [8]. The human IL-34 gene is located on the 16q22.1 chromosome, while the mouse IL-34 gene is located on the 8E1 chromosome, which has high homology among different species [9]. More and more evidences show that IL-34 is involved in the etiology of various diseases, such as inflammation, infection, autoimmune diseases, and tumors [10].

Recent studies have shown that IL-34 has a certain neuroprotective effect in certain special diseases. However, it is not clear whether IL-34 is involved in the disease process of VaD. Therefore, the main purpose of our study is to explore the relationship between IL-34 and cognitive impairment in VaD patients, in order to provide potential targets and drugs for the treatment of VaD.

## 2. Methods

**2.1. Study Participants.** From January 2017 to December 2020, 84 VaD patients treated at Qingpu Branch of Zhongshan Hospital were included in the study. At the same time, 60 age- and gender-matched healthy controls were included. The diagnosis of VaD was completed by experienced physicians with reference to the National Institute of Nervous System Diseases and Stroke (NINDSAIREN) and the 11th edition of the International Classification of Diseases (ICD-11) [11, 12]. The exclusion criteria are as follows: combined with AD or Lewy body dementia; severe infectious diseases; taking anti-infective drugs; severe trauma; recent history of surgery; suffering from severe liver disease, kidney disease, or tumor; alcoholism; and severe mental illness. The patients or family members are written informed of the current research and agree to participate in the research. The study was approved by the ethics committee of our hospital and abides by the Declaration of Helsinki.

**2.2. Demographic Features of the Participants.** Of the 144 study participants, 84 were VaD patients, and the other 60 were healthy controls. The demographic features of all participants were carefully recorded by dedicated personnel, including age, gender, education, white blood cells (WBC), neutrophil, lymphocyte, systolic blood pressure (SBP), diastolic blood pressure (DBP), fasting blood glucose (FBG), triglycerides (TG), and total cholesterol (TC). Demographic feature information is obtained in the form of questionnaires.

**2.3. MoCA Scale Evaluation.** The Montreal Cognitive Assessment (MoCA) is used as a tool to evaluate cognitive function. Dr. Ziad Nasreddine of Canada invented the MoCA scale in 1996, which was later translated into multiple languages and widely used because of its good accuracy. MoCA evaluated the following cognitive areas: short-term memory, visuospatial abilities, executive functions, attention, language, abstract reasoning, and orientation. The MoCA score ranges from 0 to 30, and a score of 26 or more is considered normal [13, 14].

**2.4. Serum ELISA Test.** Enzyme-linked immunosorbent assay (ELISA) is a popular immunoassay method that can quickly and easily analyze the same indicator in a large number of samples. In this experiment, we used an ELISA kit (MyBioSource, San Diego, CA, USA) to detect the concentration of IL-34 in the serum. Fasting venous blood of all participants was collected within 24 hours of enrollment. The collected whole blood was placed in the refrigerator overnight at 4°C and then centrifuged at 3000 rpm for 10 minutes, and the supernatant was immediately taken to be tested and stored in the refrigerator at -80°C. The specific experimental steps refer to previous reports and product manuals [15].

**2.5. Statistical Analysis.** In this study, categorical variables are represented by numbers, and continuous variables are represented by mean  $\pm$  standard deviation. The comparison between the two groups used *t*-test or chi-square test. Spearman's rank was used to assess the bivariate correlation, and multivariate regression analysis was used to clarify the causal relationship between the variables. All data were analyzed using the SPSS 22.0 software (SPSS Inc., IL, USA), and a two-tailed *p* value  $< 0.05$  was considered statistically significant.

## 3. Results

**3.1. Demographic Features of the Participants.** Of the 144 study participants, 84 were VaD patients, and the other 60 were healthy controls. The demographic features of the participants are shown in Table 1. There was no significant difference between the demographic features (age, gender, education, WBC, neutrophil, lymphocyte, SBP, DBP, FBG, TG, and TC) of VaD patients and healthy controls ( $p > 0.05$ ). However, the serum IL-34 levels of VaD patients and healthy controls are  $27.6 \pm 3.9$  pg/ml and  $41.8 \pm 6.0$  pg/ml, respectively, and there is a significant statistical difference between them ( $p < 0.001$ ). The comparison of the MoCA score and serum IL-34 levels of VaD patients and healthy controls is shown in Figure 1.

**3.2. Bivariate Correlation Analysis.** In order to clarify the potential factors affecting the cognitive function of VaD patients, we conducted a bivariate correlation analysis of the MoCA score and the variables of the demographic features. The results of the bivariate correlation analysis are summarized in Table 2. From the results in Table 2, it can be seen that the MoCA score of VaD patients is significantly positively correlated with serum IL-34 levels

TABLE 1: Demographic features of the participants.

	VaD (N = 84)	Controls (N = 60)	p value
Age (year, mean $\pm$ SD)	70.3 $\pm$ 8.9	70.5 $\pm$ 9.2	0.896
Gender (male/female)	50/34	38/22	0.644
Education (year, mean $\pm$ SD)	8.4 $\pm$ 1.5	8.6 $\pm$ 1.8	0.469
WBC ( $\times 10^9/L$ , mean $\pm$ SD)	7.6 $\pm$ 1.3	7.4 $\pm$ 1.6	0.410
Neutrophil ( $\times 10^9/L$ , mean $\pm$ SD)	5.2 $\pm$ 1.4	5.0 $\pm$ 1.1	0.358
Lymphocyte ( $\times 10^9/L$ , mean $\pm$ SD)	1.8 $\pm$ 0.6	1.9 $\pm$ 0.7	0.359
SBP (mmHg, mean $\pm$ SD)	140.6 $\pm$ 12.2	139.8 $\pm$ 12.9	0.705
DBP (mmHg, mean $\pm$ SD)	87.7 $\pm$ 8.8	87.1 $\pm$ 8.5	0.683
FBG (mmol/L, mean $\pm$ SD)	7.3 $\pm$ 0.6	7.1 $\pm$ 0.8	0.089
TG (mmol/L, mean $\pm$ SD)	1.6 $\pm$ 0.2	1.6 $\pm$ 0.1	1.000
TC (mmol/L, mean $\pm$ SD)	4.6 $\pm$ 0.5	4.5 $\pm$ 0.4	0.202
MoCA (points, mean $\pm$ SD)	22.5 $\pm$ 2.6	28.1 $\pm$ 1.2	<0.001
IL-34 (pg/ml, mean $\pm$ SD)	27.6 $\pm$ 3.9	41.8 $\pm$ 6.0	<0.001

Abbreviations: VaD, vascular dementia; WBC, white blood cells; SBP, systolic blood pressure; DBP, diastolic blood pressure; FBG, fasting blood glucose; TG, triglycerides; TC, total cholesterol; MoCA, Montreal Cognitive Assessment; IL-34, interleukin-34.




FIGURE 1: The MoCA scores and serum IL-34 levels of VaD patients and healthy controls. (a) MoCA scores; (b) serum IL-34 levels. Compared with the controls, \* $p < 0.05$ .

( $r = 0.371$ ,  $p = 0.023$ ). However, there is no significant correlation between MoCA score and age, gender, education, WBC, neutrophil, lymphocyte, SBP, DBP, FBG, TG, and TC ( $p > 0.05$ ).

**3.3. Multivariate Regression Analysis.** In order to understand the relationship between IL-34 levels and the cognitive function of VaD patients, we conducted a multivariate logistic regression analysis between the MoCA score and demographic features including IL-34. The results of the multivariate logistic regression analysis are summarized in Table 3. After adjusting for various demographic features, the relationship between IL-34 and the cognitive impairment of VaD patients is still significant, with an OR of 2.171 ( $\beta = 0.276$ ,  $p = 0.0038$ ). These results indicate that IL-34 levels are positively correlated with VaD MoCA scores. In other words, VaD patients with low serum IL-34 levels may have more severe cognitive impairment.

## 4. Discussion

This study mainly explored the correlation between the cognitive function of VaD patients and the level of serum IL-34. The preliminary results showed that the serum IL-34 level and MoCA score of VaD patients were significantly lower than those of healthy controls. The results of correlation analysis showed that the MoCA score of VaD patients was significantly positively correlated with serum IL-34 levels and was not related to age, gender, education, WBC, neutrophil, lymphocyte, SBP, DBP, FBG, TG, and TC and other demographic features. Regression analysis, after correcting for interference factors, further confirmed the correlation between serum IL-34 and MoCA in VaD patients. As far as we know, this is the first research report that uses serum IL-34 as an index to predict cognitive impairment in VaD patients.

Cytokines are a group of secreted proteins, and its members include interleukins, lymphokines, chemokines, tumor

TABLE 2: Correlation between demographic features and MoCA in VaD.

	<i>r</i>	<i>p</i> value
Age (year, mean ± SD)	-0.287	0.134
Gender (male/female)	0.329	0.399
Education (year, mean ± SD)	-0.173	0.097
WBC ( $\times 10^9/L$ , mean ± SD)	0.418	0.283
Neutrophil ( $\times 10^9/L$ , mean ± SD)	0.556	0.431
Lymphocyte ( $\times 10^9/L$ , mean ± SD)	0.314	0.262
SBP (mmHg, mean ± SD)	-0.205	0.305
DBP (mmHg, mean ± SD)	-0.231	0.338
FBG (mmol/L, mean ± SD)	-0.270	0.196
TC (mmol/L, mean ± SD)	-0.402	0.583
TG (mmol/L, mean ± SD)	-0.357	0.677
IL-34 (ng/ml, mean ± SD)	0.371	0.023

Abbreviations: MoCA, Montreal Cognitive Assessment; VaD, vascular dementia; WBC, white blood cells; SBP, systolic blood pressure; DBP, diastolic blood pressure; FBG, fasting blood glucose; TG, triglycerides; TC, total cholesterol; IL-34, interleukin-34.

TABLE 3: Multivariable analyses of demographic features and MoCA in VaD.

	Regression coefficient	<i>p</i> value
Age (year, mean ± SD)	0.213	0.569
Gender (male/female)	0.217	0.117
Education (year, mean ± SD)	0.139	0.209
WBC ( $\times 10^9/L$ , mean ± SD)	0.296	0.318
Neutrophil ( $\times 10^9/L$ , mean ± SD)	0.245	0.167
Lymphocyte ( $\times 10^9/L$ , mean ± SD)	0.197	0.388
SBP (mmHg, mean ± SD)	0.115	0.271
DBP (mmHg, mean ± SD)	0.168	0.226
FBG (mmol/L, mean ± SD)	0.103	0.309
TC (mmol/L, mean ± SD)	0.186	0.275
TG (mmol/L, mean ± SD)	0.153	0.224
IL-34 (ng/ml, mean ± SD)	0.276	0.038

Abbreviations: MoCA, Montreal Cognitive Assessment; VaD, vascular dementia; WBC, white blood cells; SBP, systolic blood pressure; DBP, diastolic blood pressure; FBG, fasting blood glucose; TG, triglycerides; TC, total cholesterol; IL-34, interleukin-34.

necrosis factor, and interferon. Cytokines produced by a variety of cells can mediate cell-cell signal transduction through autocrine and paracrine and play a key role in the communication between different cells, tissues, and organs. The combination of cytokines and specific receptors not only controls the growth, proliferation, and migration of main cells at the cellular level but also participates in the body's inflammation, tumor, immunity, and angiogenesis [16]. Therefore, research on cytokines as diagnostic markers and therapeutic targets has been in the ascendant for a long time, and some progress has been made in predicting the severity and prognosis of the disease. IL-34 was identified as the second ligand of CSF1R in 2008. The full length of human IL-34

contains 242 amino acids. The first 182 amino acids are essential for the stability and proper folding of IL-34, and the last 50 residues are a large number of disordered and Pro-Ser-Thr-rich amino acids [17]. It has a similar active region to CSF-1, but their sequence does not have homology. IL-34 is a hematopoietic cytokine, which is widely distributed in the neurons, skin, heart, lung, liver, kidney, testes, and lymph nodes. It is a key regulator of the survival, proliferation, and differentiation of myeloid cells [18].

Recent studies have found that IL-34 is involved in the process of many diseases. Scholars from Dalian Medical University found that IL-34 increases the expression of IL-6 and then upregulates Th17 to participate in the course of rheumatoid arthritis [19]. Researchers from Chongqing Medical University found that serum IL-34 is elevated in patients with systemic lupus erythematosus (SLE), which may be used as a diagnostic marker for SLE [20]. In addition, Polish studies have shown that IL-34 can be used as a potential inflammatory biomarker for predicting the risk of diabetes complications [21]. Interestingly, the scientific research team of Shanghai Jiao tong University found that increased levels of IL-34 in the acute phase are associated with increased risk of heart failure and poor prognosis after myocardial infarction [22]. IL-34 is also believed to be related to psoriasis, psoriatic arthritis, obesity, liver disease, kidney disease, and inflammatory bowel disease [23–28].

The role of IL-34 in neurological diseases has gradually attracted attention in recent years. IL-34 can act on a variety of cells such as neurons, microglia, and endothelial cells to maintain the homeostasis of the central nervous system [29, 30]. A professor at Stanford University found that IL-34 provides a powerful neuroprotective and survival signal for neurons in brain injury and neurodegeneration [31]. Recently, Iranian scientists have found that serum IL-34 levels in patients with acute and chronic inflammatory demyelinating polyneuropathies (AIDP and CIDP) are elevated, suggesting that IL-34 may participate in the pathogenesis of autoimmune diseases [32]. In addition, West Nile virus infection can cause cognitive impairment, and IL-34 knockout mice can reduce the synaptic loss caused by this virus infection, suggesting that IL-34 may participate in the course of cognitive impairment [33]. Canadian researchers initially revealed the mechanism by which IL-34 is involved in the pathogenesis of AD [34], but its relationship with VaD has not been reported yet.

Our research has some limitations. First of all, we are a single-center small sample study; secondly, our VaD patients may have mixed types of dementia; and finally, we have not done interventional mechanism research. However, we have studied the relationship between serum IL-34 and VaD for the first time, which is the advantage of our research.

## 5. Conclusions

In summary, we found that serum IL-34 levels in VaD patients were significantly reduced. We report for the first time the role of serum IL-34 in VaD. IL-34 may be used in clinical services as a diagnosis and treatment target of VaD.

## Data Availability

The data used to support the findings of this study are available from the corresponding authors upon request.

## Conflicts of Interest

All the authors declare no conflict of interests.

## Authors' Contributions

Yang Wang and Wei Lu are co-first authors, and they contributed equally to this work.

## References

- [1] Q. Wang, W. Yang, J. Zhang, Y. Zhao, and Y. Xu, "TREM2 overexpression attenuates cognitive deficits in experimental models of vascular dementia," *Neural Plasticity*, vol. 2020, Article ID 8834275, 10 pages, 2020.
- [2] M. Xu, X. Y. He, and P. Huang, "The relationship between serum amyloid A level and cognitive dysfunction in patients with vascular dementia: preliminary findings," *BioMed Research International*, vol. 2021, Article ID 6676144, 6 pages, 2021.
- [3] J. Li, S. Li, Y. Song et al., "Association of serum FAM19A5 with cognitive impairment in vascular dementia," *Disease Markers*, vol. 2020, Article ID 8895900, 5 pages, 2020.
- [4] K. Yang, L. Zeng, A. Ge, Y. Yi, S. Wang, and J. Ge, "Exploring the oxidative stress mechanism of Buyang Huanwu decoction in intervention of vascular dementia based on systems biology strategy," *Oxidative Medicine and Cellular Longevity*, vol. 2021, Article ID 8879060, 29 pages, 2021.
- [5] E. E. Smith, "Clinical presentations and epidemiology of vascular dementia," *Clinical Science*, vol. 131, no. 11, pp. 1059–1068, 2017.
- [6] L. Wang, J. W. Yang, L. T. Lin et al., "Acupuncture attenuates inflammation in microglia of vascular dementia rats by inhibiting miR-93-mediated TLR4/MyD88/NF- $\kappa$ B signaling pathway," *Oxidative Medicine and Cellular Longevity*, vol. 2020, Article ID 8253904, 15 pages, 2020.
- [7] H. Lin, E. Lee, K. Hestir et al., "Discovery of a cytokine and its receptor by functional screening of the extracellular proteome," *Science*, vol. 320, no. 5877, pp. 807–811, 2008.
- [8] T. L. Tay, J. C. Savage, C. W. Hui, K. Bisht, and M. Tremblay, "Microglia across the lifespan: from origin to function in brain development, plasticity and cognition," *The Journal of Physiology*, vol. 595, no. 6, pp. 1929–1945, 2017.
- [9] E. Franzè, C. Stolfi, E. Troncone, P. Scarozza, and G. Monteleone, "Role of interleukin-34 in cancer," *Cancers*, vol. 12, no. 1, p. 252, 2020.
- [10] Y. Nakamichi, N. Udagawa, and N. Takahashi, "IL-34 and CSF-1: similarities and differences," *Journal of Bone and Mineral Metabolism*, vol. 31, no. 5, pp. 486–495, 2013.
- [11] O. Lopez, M. Larumbe, J. Becker et al., "Reliability of NINDS-AIREN clinical criteria for the diagnosis of vascular dementia," *Neurology*, vol. 44, no. 7, pp. 1240–1240, 1994.
- [12] M. Zaudig, "Behavioral and psychological symptoms of dementia in the International Classification of Diseases (ICD)-10 and Beyond (ICD-11)," *International Psychogeriatrics*, vol. 12, p. 29, 2000.
- [13] Y. Chen, L. Liang, S. Cao et al., "Serum CCL21 as a potential biomarker for cognitive impairment in spinal cord injury," *BioMed Research International*, vol. 2020, Article ID 6692802, 5 pages, 2020.
- [14] X. Wang, F. Zhang, W. Ma, D. Feng, J. Zhang, and J. Xu, "Increased levels of serum neuregulin 1 associated with cognitive impairment in vascular dementia," *BioMed Research International*, vol. 2020, Article ID 6683747, 5 pages, 2020.
- [15] L. Wang, Y. Chen, D. Feng, and X. Wang, "Serum ICAM-1 as a predictor of prognosis in patients with acute ischemic stroke," *BioMed Research International*, vol. 2021, Article ID 5539304, 6 pages, 2021.
- [16] M. Baghdadi, H. Endo, Y. Tanaka, H. Wada, and K. I. Seino, "Interleukin 34, from pathogenesis to clinical applications," *Cytokine*, vol. 99, pp. 139–147, 2017.
- [17] M. Baghdadi, Y. Umeyama, N. Hama et al., "Interleukin-34, a comprehensive review," *Journal of Leukocyte Biology*, vol. 104, no. 5, pp. 931–951, 2018.
- [18] I. Lelios, D. Cansever, S. G. Utz, W. Mildenerberger, S. A. Stifter, and M. Greter, "Emerging roles of IL-34 in health and disease," *The Journal of Experimental Medicine*, vol. 217, no. 3, 2020.
- [19] B. Wang, Z. Ma, M. Wang et al., "IL-34 upregulated Th17 production through increased IL-6 expression by rheumatoid fibroblast-like Synoviocytes," *Mediators of Inflammation*, vol. 2017, Article ID 1567120, 10 pages, 2017.
- [20] H. Wang, J. Cao, and X. Lai, "Serum interleukin-34 levels are elevated in patients with systemic lupus erythematosus," *Molecules*, vol. 22, 2017.
- [21] K. Zorena, O. Jachimowicz-Duda, and P. Wąż, "The cut-off value for interleukin 34 as an additional potential inflammatory biomarker for the prediction of the risk of diabetic complications," *Biomarkers: biochemical indicators of exposure, response, and susceptibility to chemicals*, vol. 21, no. 3, pp. 276–282, 2016.
- [22] Q. Fan, R. Tao, H. Zhang et al., "Interleukin-34 levels were associated with prognosis in patients with acute myocardial infarction," *International Heart Journal*, vol. 60, no. 6, pp. 1259–1267, 2019.
- [23] J. Li, L. Liu, W. Rui et al., "New interleukins in psoriasis and psoriatic arthritis patients: the possible roles of interleukin-33 to interleukin-38 in disease activities and bone erosions," *Dermatology*, vol. 233, pp. 37–46, 2017.
- [24] N. Li, L. Jiang, Y. Cai et al., "The correlation between interleukin-34 and bone erosion under ultrasound in rheumatoid arthritis," *Modern Rheumatology*, vol. 30, no. 2, pp. 269–275, 2020.
- [25] E. J. Chang, S. K. Lee, Y. S. Song et al., "IL-34 is associated with obesity, chronic inflammation, and insulin resistance," *The Journal of Clinical Endocrinology and Metabolism*, vol. 99, no. 7, pp. E1263–E1271, 2014.
- [26] F. Al-Shaebi, L. Wenzhang, K. Hezam, M. Almezgagi, and L. Wei, "Recent insights of the role and signalling pathways of interleukin-34 in liver diseases," *International Immunopharmacology*, vol. 89, p. 107023, 2020.
- [27] J. H. Baek, R. Zeng, J. Weinmann-Menke et al., "IL-34 mediates acute kidney injury and worsens subsequent chronic kidney disease," *The Journal of Clinical Investigation*, vol. 125, no. 8, pp. 3198–3214, 2015.
- [28] S. Zwicker, G. L. Martinez, M. Bosma et al., "Interleukin 34: a new modulator of human and experimental inflammatory

- bowel disease,” *Clinical science*, vol. 129, no. 3, pp. 281–290, 2015.
- [29] S. Jin, Y. Sonobe, J. Kawanokuchi et al., “Interleukin-34 restores blood-brain barrier integrity by upregulating tight junction proteins in endothelial cells,” *PLoS One*, vol. 9, no. 12, article e115981, 2014.
- [30] P. W. Lee, A. Selhorst, S. G. Lampe, Y. Liu, Y. Yang, and A. E. Lovett-Racke, “Neuron-specific vitamin D signaling attenuates microglia activation and CNS autoimmunity,” *Frontiers in Neurology*, vol. 11, p. 19, 2020.
- [31] J. Luo, F. Elwood, M. Britschgi et al., “Colony-stimulating factor 1 receptor (CSF1R) signaling in injured neurons facilitates protection and survival,” *The Journal of Experimental Medicine*, vol. 210, no. 1, pp. 157–172, 2013.
- [32] S. Rezaei, S. Ghafouri-Fard, A. Komaki, M. Mazdeh, M. Taheri, and M. M. Eftekharian, “Increased levels of IL-34 in acquired immune-mediated neuropathies,” *Journal of molecular neuroscience*, vol. 71, no. 1, pp. 137–141, 2021.
- [33] M. J. Vasek, C. Garber, D. Dorsey et al., “A complement-microglial axis drives synapse loss during virus-induced memory impairment,” *Nature*, vol. 534, no. 7608, pp. 538–543, 2016.
- [34] V. Pons, P. Lévesque, M. M. Plante, and S. Rivest, “Conditional genetic deletion of CSF1 receptor in microglia ameliorates the pathophysiology of Alzheimer’s disease,” *Alzheimer’s Research & Therapy*, vol. 13, no. 1, p. 8, 2021.