Review Article

Traditional Medicinal Plants as a Source of Antituberculosis Drugs: A System Review

Yuhui Xu,¹ Bowen Liang,² Chengcheng Kong,^{3,4} and Zhaogang Sun^{3,4}

¹Institute of Chinese Materia Medica, China Academy of Chinese Medical Science, Beijing 100700, China

²Department of Traditional Chinese Medicine, Beijing Chest Hospital, Capital Medical University, Beijing 101149, China

³Translational Medicine Center, Beijing Chest Hospital, Capital Medical University, Beijing 101149, China

⁴Beijing Key Laboratory in Drug Resistant Tuberculosis Research, Beijing Tuberculosis & Thoracic Tumor Research Institute, Beijing 101149, China

Correspondence should be addressed to Zhaogang Sun; sunzg75@163.com

Received 16 March 2021; Accepted 9 August 2021; Published 10 September 2021

Academic Editor: Jane Hanrahan

Copyright © 2021 Yuhui Xu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Medicinal plants are the chief components in the different oriental formulations in different traditional medical systems worldwide. As a thriving source of medicine, the medicinal plants with antituberculosis (TB) properties inspire the pharmacists to develop new drugs based on their active components or semimetabolites. In the present review, the anti-TB medicinal plants were screened from the scientific literatures, based on the botanical classification and the anti-TB activity. The obtained anti-TB medicinal plants were categorized into three different categories, viz., 159 plants critically examined with a total 335 isolated compounds, 131 plants with their crude extracts showing anti-TB activity, and 27 plants in literature with the prescribed formula by the traditional healers. Our systemic analysis on the medicinal plants can assist the discovery of novel and more efficacious anti-TB drugs.

1. Introduction

Globally, traditional medicines (TM) make a vital contribution to the health care industry. In some countries, TM is the main source of health care or even the sole health care service available, especially in the rural sector [1]. The popularity of TM is also increasing in the developed countries for many different reasons, one of which is that the effectiveness of these TM was proved by the ethnopharmacological research. Early in 1972, the World Health Organization (WHO) established a Department of Traditional Medicine (DTM). Later, WHO (2013) called on to strengthen its public services of the traditional medicine [1]. Recently, the International Classification of Traditional Medicine (ICTM) was added as a new chapter into the International Classification of Diseases—11 (ICD-11) [2]. This achievement currently refers to the Traditional Chinese Medicine (TCM) alone, which opens its doors to accommodate many other thriving traditional health care philosophies prevailing glob-

ally, such as Ayurveda and Traditional African medicine (TAM).

The spread of tuberculosis (TB) occurred from East Africa to the rest of the world with the migration of Homo sapiens, especially along the established trade routes with increased mingling and crowding of populations [3, 4]. Currently, there exist more than 10 million new cases of active disease and nearly 1.3 million deaths annually [5, 6]. In response to this spreading route, different countries developed their own traditional anti-TB formulations during the long courses in fighting this old plaque. Reports relating TB can be found in many ancestral data of the TM medical system, especially the TCM, Ayurveda, and TAM for its long history coexisting with human kinds for an estimated 70,000 years [7]. Investigations on the TM formulations show that the plants or herbs are the main composition of the traditional anti-TB formula, from which the active components or semimetabolites present a thriving source of new drugs. In the last 20 years, nearly 50% of drugs approved by the

FDA in the United States of America have been derivatives of the natural products, including natural plant products [8]. Among the 435,000 plant species reported worldwide [9], an estimated 70,000 species of plants are used for medicinal purposes [10]. Thus, selecting plants based on ethnobotanical knowledge can enhance the probability to find new compounds with anti-TB activity.

Before this review, some articles summarized the role of local medical plants but only few with anti-TB purpose [11, 12]. In this review, the anti-TB medicinal plants in different countries or regions are included to analyze their botanical classification, active botanical parts, extract method, and *in vitro* anti-TB activities in brief. Subsequently, the effective anti-TB plants are described with the following three branches: those with the isolated effective compounds, those with their crude plant extracts showing anti-TB effect, and those only found in the formula prescribed by traditional healers. Finally, we discuss the influencing factors on the development of traditional medicine and its future trend. This review is to inspire the development of possible new anti-TB agents derived from plants.

2. Brief Description of the Overall Anti-TB Medicinal Plants

We present the data by searching the main three databases: Wangfang Med, Chinese National Knowledge Infrastructure, and PubMed. Combinations of the following search terms are used: "tuberculosis," "plant," "herb," "Chinese and western medicine," and "random." In the present review, only the nonrepetitive plant species with good *in vivo* or *in vitro* anti-TB effect were accepted, although the criterion of the effectiveness was quite different with the inhibition concentration expressed in several different ways in different *Mycobacteria*, especially the *M. tuberculosis* H37Rv and the clinical isolates. The plants employed for treating the fever in traditional medicine have not been included, as fever is taken to be a nonspecific indication of many infections that are not restricted to TB. The exception to this is where fever is treated in conjunction with other TB-related symptoms like coughing.

The classification of the traditional anti-TB medicinal plants in the present review belongs to 90 families including 230 genus and 277 species (Figure 1). The top 11 families with more than 7 plant species include Fabaceae (21 species in 18 genus), Asteraceae (20 in 16 genus), Euphorbiaceae (14 in 11 genus), Lamiaceae (13 in 11 genus), Rutaceae (14 in 10 genus), Combretaceae (9 in 4 genus), Piperaceae (9 in 1 genus), Zingiberaceae (8 in 3 genus), Annonaceae (7 in 6 genus), and Apiaceae (7 in 7 genus). Forty plant families are only reported once. A total of 6 *Terminalia* genus that belongs to the Combretaceae family have up to 6 anti-TB plant species, and about 9 anti-TB plant species belong to only one genus *Piper*.

The literatures that we studied reported the anti-TB properties of the plant species from different plant parts (aerial parts, almonds, bark, bulbs, branches, fruits, flowers, heart woods, leaves, rhizomes, roots, stems, seeds, shoots, twigs, tubers, wood, whole plants, and even the ethnomedic-

inal recipes). With the leaves (83 cases), roots (61), aerial parts (32), barks (30), stems (14), whole plants (9), seeds (9), fruits (8), rhizomes (8), and flowers (7) are the top 10 most used anti-TB plant parts. For the same plant species, different parts of the plant presented a varied anti-TB effect. The useful plant parts of the genus *Lantana*, *Piper*, and *Terminalia* mainly focused on the leaves, leaves, and both leaves and roots, respectively.

It was observed that the extraction methods of the medicinal plants available in the literature significantly affected the anti-TB results. The general problems concerning the antibacterial screening of medicinal plant extracts have already been discussed in the literature [13]. There is still no single extraction method that is regarded as a standard for extracting the bioactive compounds from medicinal plants. One or more of the following solvents were mainly used in the studies: dichloromethane (268 times), methanol (65), ethanol (45), hexane (29), chloroform (18), ethyl acetate (11), water (11), and acetone (10), while diethylether, acetate, and hydroalcoholic solutions were seldom used. Since the extraction process deeply influences the results of the bioactivity tests and the subsequent isolation of bioactive compounds, selection of the best extraction method by consulting the traditional knowledge about the preparation of the herbal remedy remains crucial [14].

As noted earlier and evidenced by this review, many reports lack adequate statistical analysis of their results and appropriate controls for their anti-TB activity, while some studies lack the generic extraction schemes or tests against a panel of various species of *Mycobacteria* to avoid false positive results. In this review, the parallel cytotoxic evaluation on mammalian cell lines has not been provided, since our main aim was to summarize the crude extracts or compound precursors of the anti-TB medicinal plants, although this needs to be overcome in the future.

3. Compounds from the Plants with Anti-TB Activity

Different from the conventional process of drug discovery involving the screening of large molecular libraries for biological activities and/or *in silico* data mining approaches based on cheminformatics modeling, the bioactivity-guided fractionation was mostly employed in medicinal plants to isolate the bioactive compounds. They were extracted first from the specified parts of the plants, then fractionized and characterized by infrared spectroscopy, mass spectrometry, and nuclear magnetic resonance spectroscopy, to obtain the structural data. Finally, their bioactivities were verified in different mycobacteria.

Several groups summarized the active anti-TB natural products from the different organs and regions. Early in 2007, Copp and Pearce [15] summarized a total of 353 natural products (secondary metabolites) with reported growth inhibitory activity towards *M. tuberculosis* or related organisms from terrestrial and marine plants and animals and microorganisms. Abedinzadeh et al. [16] stressed the natural antimycobacterial peptides from bacteria, fungi, plants, and animals. Chinsebu [17] described the natural

FIGURE 1: Classification of traditional anti-TB medicinal plants with effective crude extracts and the compounds. (a) Botanical families consisting of the anti-TB medicinal plants. There are 108 families including 230 genus and 277 species in this summary. (b) Genus number (>2) of the anti-TB medicinal plant families.

antimycobacterial agents from endophytes and medicinal plants in different regions of Africa, Europe, Asia, South America, and Canada. The present review only focused on the medicinal plants and the plants with anti-TB components belonged to the 156 species, 123 genus, and 64 families, of which Fabaceae (13 species, 10 genus), Rutaceae (10 species, 7 genus), and Lamiaceae (9 species, 7 genus) were the top three families; accordingly, more genera belong to those family with anti-TB activity (Table 1).

Many plants consisted several components with anti-TB activity, and only the active compounds that were reported are listed in this review. Table 1 presents the list of 335 compounds, which were tested for their anti-TB activities. Those 335 compounds could be divided into mainly 11 classes, such as terpenes (37 types), ketones (31), acids (14), alcohols (10), esters (9), hydrocarbons (9), quinones (8), furans (7), phenols (6), and quinolones (3). The typical structures of

the 335 compounds are sorted out in Figure 2. Of all the anti-TB natural compounds, the derivatives and analogs of phytol, flavones, and terpenoids were critically reviewed by Singh et al. [173] and Cantrell et al. [174] for their pharmacological activities of various diseases. These 335 compounds were natural products or secondary metabolites, and few of their synthetic modified derivatives have been mentioned in this review.

In fact, many semisynthetic derivatives proved to be more active than the parent compounds; for example, the methylation of natural compounds of mulinenic acid and 13-hydroxy-mulin-11-en-20-oic acid methyl ester decreased the minimum inhibitory concentration (MIC) by 8 times [42]; n-propyl ester and n-butyl ester of isomulinic acid decreased the MIC by 4 times [42]. The triacetylated methyl gallate decreased the MIC 2-4 times, since the acetylation increased the lipophilic nature of methyl gallate [19]. The

TABLE 1: Medicinal plants and natural products showing *in vitro* anti-TB activity.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Abrus precatorius</i>	Fabaceae	Aerial parts	Dichloromethane fraction	Isoflavanquinone	Abruquinone B (1) against <i>M. tuberculosis</i> H37Ra with MIC of 12.5 $\mu\text{g/ml}$ by MABA ^{a,b}	[18]
<i>Acacia farnesiana</i>	Mimosaceae	Fruit	Methanolic extract	Parabens, flavanones	Methyl gallate (2) showed activity against the sensible strain <i>M. tuberculosis</i> H37Rv with MIC of 50 $\mu\text{g/ml}$, respectively. The (2S)-Naringenin 7-O- β -galloylglucopyranoside (3) showed activity against multidrug resistant <i>M. tuberculosis</i> G122 with MIC of 50 $\mu\text{g/ml}$ by MABA	[19]
<i>Aglaia forbesii</i>	Meliaceae	Leaf	Dichloromethane fraction	Benzopyran flavaglines	Desacetylpyramidaglain D (4) against <i>M. tuberculosis</i> Ra with MIC of 25 $\mu\text{g/ml}$ by MABA	[20]
<i>Allanblackia floribunda</i>	Guttiferae	Root bark	Successively macerated in dichloromethane-methanol (1:1) and methanol for 4 h	Biflavonoids	Morelloflavone (5) with the MIC of 19.53 and 39.06 $\mu\text{g/ml}$ against <i>M. smegmatis</i> and <i>M. tuberculosis</i> , respectively, by MABA	[21]
<i>Allium neapolitanum</i>	Alliaceae	Bulb	Chloroform extract	Canthinone	Canthin-6-one (6), 8-hydroxy-canthin-6-one (7), and 5(ζ)-hydroxy-octadeca-6(E)-8(Z)-dienic acid (8) with MICs in the range 8–32 $\mu\text{g/ml}$ against a panel of fast-growing Mycobacterium species by dilution method	[22]
<i>Allium sativum</i>	Liliaceae	Bulb	Petroleum ether extract	Fatty acids	Lauric acid (9) and myristic acid (10) with MIC of 22.2 and 66.7 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[23]
<i>Allophylus edulis</i>	Sapindaceae	Leaf	Hydrodistillation	Cycloprop[<i>e</i>]azulen-4-ol	Viridiflorol (11) against <i>M. tuberculosis</i> H37Rv (ATCC27294) with MIC of 190.0 $\mu\text{g/ml}$ by the microplate resazurin assay	[24]
<i>Alnus incana</i>	Betulaceae	Bark	Methanol extract	Triterpenes	Betulin (12), betulinic acid (13), and betulone (14) with MIC of 12.5, 84, and 57 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by the microplate resazurin assay	[25]
<i>Alpinia katsumadai</i>	Zingiberaceae	Seed	n-Hexane	Diarylheptanoids	Trans,trans-1,7-diphenylhepta-4,6-dien-3-one (15) as efflux inhibitors against <i>M. smegmatis</i> mc ² 155 by thiazolyl blue tetrazolium bromide method	[26]
<i>Amphipterygium adstringens</i>	Anacardiaceae	Stem bark	Dichloromethane/methanol (1:1)	Tirucallanes	(14 β , 24E)-3-oxolanosta-7,24-dien-26-oic acid (16) and (14 β ,24E)-3-hydroxyalanosta-7,24-dien-26-oic acid (17) with MIC of 64 and 32 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by Bactec 460-TB apparatus	[27]
<i>Amyris elemifera</i>	Rutaceae	Leaf	Chloroform extract	Texalin	Texalin (18) with MIC of 25 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv by Bactec 460-TB radiometric methodology	[28]
<i>Androsace umbellata</i>	Primulaceae	Whole plants	Ethanol extract	Saxifragifolin	Saxifragifolin D (19) reduced the intracellular replication of <i>M. tuberculosis</i> in THP-1-derived macrophages but not in A549 cells	[29]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Angelica sinensis</i>	Apiaceae	Root	Chloroform extract	Fatty diol	Falcarindiol (20), 9Z,17-octadecadiene-12,14-diyne-1,11,16-triol,1-acetate (21), and oplopandiol (22) with MIC of 26.7, 25.3, and 50.2 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv by MABA	[30]
<i>Anisochilus harmandii</i>	Lamiaceae	Aerial parts	Water	Diterpenes	One pimarane-type diterpene named pimaric acid (23) and two abietane-type diterpenes named 9a,13a-epidioxyabiet-8 (14)-en-18-oic acid (24) and 15-hydroxydehydroabietic acid (25) with MIC of 50 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[31]
<i>Aralia nudicaulis</i>	Araliaceae	Rhizome	Methanolic extract	Polyacetylene	(3R)-falcarinol (26) and (3R, 9R, 10S)-panaxydol (27) with MICs of 25.6 μM and 36.0 μM and IC50's of 15.3 μM and 23.5 μM against <i>M. tuberculosis</i> H37Ra by microplate resazurin assay	[32]
<i>Ardisia gigantifolia</i>	Primulaceae	Leaves and stems	Chloroform extract	Resorcinol	5-alkylresorcinols (28), 5-(8Z-heptadecenyl) resorcinol (29), and 5-(8Z-pentadecenyl) resorcinol (30) with MIC of 34.4 and 79.2 μM against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA assay, respectively	[33]
<i>Argyreia speciosa</i>	Convolvulaceae	Root	Methanolic extract	Flavonoid sulphates	Quercetin 3'-O methyl 3-sulphate (31) and kaempferol 7-O methyl 3-sulphate (32) against <i>M. tuberculosis</i> H37Rv with MIC values of 25 $\mu\text{g/ml}$, respectively, by MABA	[34]
<i>Arisaema sinii</i>	Araceae	Whole plant	80% ethanol	Fatty acid ester	(E)-2-(methyl (phenyl) amino) ethyl 2-(2-hydroxyundecanamido)-7 (33), 11-dimethyl-3-oxotetradec-4-enoate (34), and compound 1 inhibit mycobacterial biofilm formation, disperse the preformed biofilms, and disrupt the mature biofilms at concentration of 4, 8, and 32 $\mu\text{g/ml}$, respectively	[35]
<i>Aristolochia brevipes</i>	Aristolochiaceae	Rhizome	Dichloromethane extract	Benzo[f]-1,3-benzodioxolo[6,5,4-cd]indol-5(6H)-one	Aristolactam I (35) with MIC of 12.5-25 $\mu\text{g/ml}$ against drug resistant <i>M. tuberculosis</i> by fluorimetrit MABA	[36]
<i>Aristolochia taliscana</i>	Aristolochiaceae	Hook roots	Hexanic extract	Neolignans	Licarin A (36), licarin B (37), and eupomatenoide-7 (38) with MIC of 3.12-12.5 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> strains: H37Rv, four mono-resistant H37Rv variants and 12 clinical MDR isolates by MABA	[37, 38]
<i>Arracacia toluensis</i>	Umbelliferae	Aerial parts	Dichloromethane-methanol (1:1)	Coumarins	Isoimperatorin (39), osthol (40), suberosin (41), and 8-methoxyxyporalen (42) with MIC of 64, 32, 16, and 128 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[39]
<i>Artemisia capillaris</i>	Asteraceae	Aerial parts	Methanol extracts	Stilbene derivatives	Ursolic acid (43) and hydroquinone (44) with MIC of 12.5-25 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> MDR/XDR strains by MABA	[40]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
Artocarpus lakoocha	Moraceae	Root	Dichloromethane extract	Benzofuran	Lakoochin A (45) and lakoochin B (46) with MIC of 12.5 and 50 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by MABA	[41]
Azorella compacta	Umbelliferae	Aerial parts	n-Hexane	Diterpenoid	Azorellanol (47) and 17-acetoxy-13-alpha-hydroxy-azorellane (48) showed the strongest activity, with MIC of 12.5 $\mu\text{g/ml}$ against both strains (<i>M. tuberculosis</i> H37Rv (ATTC 27294) and a clinical isolate CIBIN/UMF1599) by MABA	[42, 43]
Bauhinia purpurea	Fabaceae	Root	Dichloromethane extract	Dibenz[b,f]oxepin	Bauhinoxepin J (5,6-dihydro-3-methoxy-1,4-dione)dibenz[b,f]oxepin (49) against <i>M. tuberculosis</i> H37Ra with a MIC of 24.4 μM by MABA	[44]
Bauhinia saccocalyx	Fabaceae	Root	Dichloromethane extract	Dibenz[b,f]oxepin	Bauhinoxepins A (=3,3,5-trimethylbenzo[b]pyrano[g]benzoxepin-6,11-diol) (50) and B (=6-methoxy-7-methyl-2-(3-methylbut-2-enyl)dibenzo[b,f]oxepine-1,8-diol) (51) against <i>M. tuberculosis</i> H37Rv with MIC of 6.25 and 12.5 $\mu\text{g/ml}$, respectively, by MABA	[45]
Beilschmiedia erythrophloia	Lauraceae	Root	Methanol	Endiandric acid	Erythrophloin C (52) and suberosol B (53) against <i>M. tuberculosis</i> H37Rv with MICs of 50 and 28.9 $\mu\text{g/ml}$, respectively, by MABA	[46]
Beilschmiedia tsangii	Lauraceae	Leaf	Methanol	Epoxyfuranoid lignans	Three new epoxyfuranoid lignans, 4a,5a-epoxybeilschmin A (54) and B (55), and beilschmin D (56), together with known beilschmin A (57) and B (58) with MICs of 30, 40, 50, 2.5, and 7.5 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> 90-211378 by proportion method on agar	[47]
Blepharodon nitidum	Asclepiadaceae	Whole plant	Ethanol extract	Hydroperoxycycloartanes	24-Hydroperoxycycloart-25-en-3 β -ol (59) and 25-hydroperoxycycloart-23-en-3 β -ol (60) with MIC of 12.5 and 25 $\mu\text{g/ml}$ against drug-resistant clinical isolates by MABA	[48]
Bocconia arborea	Papaveraceae	Aerial parts	Chloroform extract	Dihydrochelirubine	Alkaloids 6-methoxydihydrochelirubine (61) and 6-methoxydihydrochelirubine (62) against <i>M. tuberculosis</i> H37Rv and MDR tuberculosis with MIC of 12.5-50 $\mu\text{g/ml}$ by MABA	[49]
Caesalpinia pulcherrima	Fabaceae	Root	Dichloromethane	Cassane-furanoditerpenoids	6 Beta-benzoyl-7 beta-hydroxyvouacapen-5 alpha-ol (63) and 6 beta-cinnamoyl-7beta-hydroxyvouacapen-5 alpha-ol (64) with MIC of 25 and 6.25 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by MABA	[50]
Caesalpinia sappan	Fabaceae	Heartwood	Methanol	Chalcone	3-Deoxysappanchalcone (65) against both drug-susceptible and drug-resistant strains of <i>M. tuberculosis</i> at MIC50s of 3.125-12.5 $\mu\text{g/ml}$ in culture broth and MIC50s of 6.25-12.5 $\mu\text{g/ml}$ inside macrophages and pneumocytes	[51]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
Calliandra californica	Fabaceae	Root	Ethyl acetate	Cassane-type diterpenes	Escobarine A (66) and B (67) with MIC of 25-50 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv	[52]
Callicarpa pilosissima	Verbenaceae	Leaves and twigs	Methanol extract	Diterpenoids	12-Deoxy-11,12-dihydro-seco-hinokiol methyl ester (68), callicarpic acid B (69), and alpha-tocopherol trimer B (70) with MICs $\leq 63.6 \mu\text{M}$ against <i>M. tuberculosis</i> H37Rv in vitro by MABA	[53]
Calophyllum lanigerum	Clusiaceae	All plant materials	Methanol	Coumarin	(+)-Calanolide A (71) against <i>M. tuberculosis</i> with H37Rv with MIC of 3.13 $\mu\text{g/ml}$ by the radiometric BACTEC	[54, 55]
Camchaya calcarea	Asteraceae	Whole plant	Dichloromethane	Cyclodeca[b]furan	Goyazensolide (72), centratherin (73), lychnophorolide B (74), isogoyazensolide (75), isocentratherin (76), 5-epi-isogoyazensolide (77), and 5-epi-isocentratherin (78) with MICs of 3.1, 3.1, 6.2, 1.5, 3.1, 3.1, and 3.1 $\mu\text{g/ml}$, against <i>M. tuberculosis</i> H37Rv, respectively, by MABA	[56]
Celastrus vulcanicola	Celastraceae	Leaf	Dichloromethane extract	Sesquiterpenes	1 α -Acetoxy-6 β , 9 β -dibenzoyloxy-dihydro-b-agarofuran (79) with MIC value of 6.2 $\mu\text{g/ml}$ against sensitive and resistant <i>M. tuberculosis</i> strains by MTT method	[57]
Chamaedorea tepejilote	Areaceae	Leaf	Hexane extracts	Pentacyclic triterpenes, fatty alcohols	Ursolic acid (43) and farnesol (80) against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 50 $\mu\text{g/ml}$ and 8 $\mu\text{g/ml}$, respectively, by MABA	[58, 59]
Chrysanthemum morifolium	Asteraceae	Flower	Methanol extract	3-Hydroxy triterpenoids	Maniladiol (81), 3-epilupeol (82), and 4,5a-epoxyhelianol (83) with MIC of 4 $\mu\text{g/ml}$, 4 $\mu\text{g/ml}$, and 6 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv by MABA	[60]
Citrullus colocynthis	Cucurbitaceae	Fruit	Methanolic extract	Triterpenes	Ursolic acid (43) and cucurbitacin E 2-0- β -D-glucopyranoside (84) against <i>M. tuberculosis</i> H37Rv with MICs of 50 and 25 $\mu\text{g/ml}$, respectively, by BACTEC 460TB system	[61]
Citrus aurantiifolia	Rutaceae	Fruit peels	Hexane extract	Furo[3,2-G]coumarin, fatty acid	Both 5,8-dimethoxypsoralen (85) and palmitic acid (86) with the MIC of 25 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv (ATCC 27294) and <i>M. tuberculosis</i> H10 by MABA	[62]
Clausena excavata	Rutaceae	Rhizome	Chloroform extract	Coumarins, carbazole derivatives	Dentatin (87), nor-dentatin (88), clausenidin (89), 3-formylcarbazole(90), mukonal (91), 3-methoxycarbonylcarbazole (92), 2-hydroxy-3-formyl-7-methoxycarbazole (93), and clausoline J (94) with MIC of 50, 100, 200, 100, 200, 50, 100, and 100 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by MABA	[63]
Clavija procera	Theophrastaceae	Stem and bark	Ethanol extract	Oleanane triterpene	Oleanane triterpenoid aegicerin (95) with MIC values ranged between 1.6 and 3.12 $\mu\text{g/ml}$ against 37 different sensitive and resistant MTB strains by thiazolyl blue tetrazolium bromide method	[64]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Clinacanthus siamensis</i>	Acanthaceae	Leaf	Ethanol	Amide	Trans-3-methylthioacrylamide (96) with the MIC of 200 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by MABA	[65]
<i>Cnidocolus chayamansa</i>	Euphorbiaceae	Leaf	Chloroform : methanol (1 : 1)	Pentacyclic triterpenes	Moretenol (97) and moretenyl acetate (98) showed MIC of 25 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv and four monoresistant strains	[66]
<i>Combretum molle</i>	Combretaceae	Stem bark	Acetone	Polyphenol	Punicalagin (99) with MIC of 600 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv by the agar proportional method	[67, 68]
<i>Cordia globifera</i>	Ehretiaceae	Root	A hexane-soluble extract	Quinones	Globiferin (100) and cordiachrome C (101) with MICs of 6.2 and 1.5 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[69]
<i>Croton kongensis</i>	Euphorbiaceae	Leaf	Dichloromethane	Diterpenedione	16-Dien-9,15-dione (102), ent-8,9-seco-8,14-epoxy-7R-hydroxy-11a-acetoxy-16-kauren-9,15-dione (103), ent-8,9-seco-7R-hydroxy-11a-acetoxykaura-8 (14) with MICs of 25.0, 6.25, and 6.25 $\mu\text{g/ml}$, respectively, and possessed antimalarial activity with IC_{50} ranges of 1.0-2.8 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by MABA	[70]
<i>Curcuma longa</i>	Zingiberaceae	Rhizomes	Chloroform extracts	Curcumin	Curcuminoid demethoxycurcumin (104) with MIC of 200 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv by BACTEC 460	[71, 72]
<i>Curtisia dentata</i>	Curtisiaceae	Leaf	Ethanol extracts	Triterpenes	Ursolic acid acetate (105) and betulinic acid acetate (106) with MIC of 3.4 $\mu\text{g/ml}$ and 19.8 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[73]
<i>Cynanchum atratum</i>	Asclepiadaceae	Roots	Ethanol extract	Isoquinolin	(-)-Deoxypergularinine (107) with MIC of 12.5 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra, H37Rv, MDR, and XDR strains by Bactec MGIT 960TM	[74]
<i>Derris indica</i>	Fabaceae	Stem, root	Hexane : dichloromethane (1 : 1)	Flavonoids	3-Methoxy-(3''-4''-dihydro-3''-4''-diacetoxy)-2'',2''-dimethylpyrano-(7,8:5'',6'')-flavone (108), 3,4-methylenedioxy-10-methoxy-7-oxo[2]benzopyrano[4,3-b]benzopyran (109), karanjachromene (110), and pinnatin (111) against <i>M. tuberculosis</i> H37Ra with MICs of 25, 6.25, 12.5, and 12.5 $\mu\text{g/ml}$ by MABA	[75, 76]
<i>Diospyros anisandra</i>	Ebenaceae	Stem bark	n-Hexane extract	Naphthalene	Plumbagin (112) and 3,3'-biplumbagin(113) against <i>M. tuberculosis</i> H37Rv with MIC of 1.56 and 3.33 $\mu\text{g/ml}$ by MABA	[7]
<i>Diospyros decandra</i>	Ebenaceae	Bark	Hexane	Triterpenes	2-Oxo-3b,19a-dihydroxy-24-nor-urs-12-en-28-oic acid (114) with MIC of $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Ra by MABA	[77]
<i>Diospyros montana</i>	Ebenaceae	Stem bark	Chloroform extract	Quinonoids	Plumbagin (112) > emodin (115) > menadione (116) > thymoquinone (117) > diospyrin (118) against <i>M. tuberculosis</i> H37Ra (ATCC 25177) and MDR-TB by MABA	[78, 79]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Distemonanthus benthamianus</i>	Fabaceae	Stem bark	Methanol extract	Flavonoids	Distemonanthoside (119), sitosterol 3-O- -d-glucopyranoside, 4-methoxygallic acid (118), and quercetin (121), against a clinical isolate strain of <i>M. tuberculosis</i> AC 45 with MIC ranged from 31.25 to 125 µg/ml by MABA	[80]
<i>Dracaena angustifolia</i>	Dracaenaceae	Leaf	Dichloromethane extract	Triterpenes, fatty acids, fatty alcohols	Ergosterol-5,8-endoperoxide (122), linoleic acid (123), and E-phytol (124) with MICs ≤ 2 µg/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[81]
<i>Ehretia longiflora</i>	Boraginaceae	Root	Methanolic extract	Quinone	Ehretiquinone (125) and prenylhydroquinone (126) with MIC of 25 and 26.2 µg/ml against <i>M. tuberculosis</i> strain H37Rv by the agar proportion method	[82]
<i>Engelhardia roxburghiana</i>	Juglandaceae	Root	Methanol	Quinone	Engelharquinone (127), 3-methoxyjuglone (128), and (-)-4-hydroxy-1-tetralone (129) with MIC of 3.125, 3.125, and 6.25 µg/ml against <i>M. tuberculosis</i> 90-221387 and 0.2, 0.2, and 4.0 µg/ml against <i>M. tuberculosis</i> H37Rv, respectively, by agar proportion method	[83, 84]
<i>Eriosema chinense</i>	Fabaceae	Root	Exane and dichloromethane extracts	Flavonoids	Khonklonginol A (130), B (131), H (132), and lupinifolinol (133), dehydrolupinifolinol (134), flemichin (135), eriosemaone A (136), and lupinifolin (137), with MICs of 25, 50, 25, 25, 12.5, 12.5, 12.5, and 12.5 µg/ml, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[85]
<i>Erythrina gibbosa</i>	Fabaceae	Ethnomedicinal recipes	70% aqueous methanol	Isoflavonoids	Isoflavonoidspasellidin (138) and erythobissin (139) with MICs, respectively, lying between 8 µg/ml and 25 µg/ml against MTB; 3-phenyl coumarin derivative indicanine (140) with MIC of 18.5 µg/ml against <i>M. smegmatis</i> by agar proportion method	[86]
<i>Erythrina indica</i>	Fabaceae	Root bark	Successively extracted with dichloromethane-methanol (1:1) and methanol	Isoflavonoids	Indicanine B (141) with MIC of 18.5 µg/ml against <i>M. smegmatis</i> by agar proportion method	[86]
<i>Erythrina senegalensis</i>	Fabaceae	Ethnomedicinal recipes	70% aqueous methanol	Isoflavonoids	Isoflavonoidspasellidin (138) and erythobissin (139) with MICs between 8 µg/ml and 25 µg/ml against <i>M. tuberculosis</i> H37Rv; 3-phenyl coumarin derivative indicanine (138) with MIC of 18.5 µg/ml against <i>M. smegmatis</i> by agar proportion method	[86]
<i>Eucalyptus torelliana</i>	Myrtaceae	Leaf	Hexane extract	Fatty acid ester	Hydroxyomyristic acid methylester (142) and a substituted pyrenyl ester (143), a sterol with MIC of 49.45 and 46.99 µg/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[87]
<i>Eudea natalensis</i>	Ebenaceae	Root	Acetone extract	Naphthoquinone	Diospyrin (118) and 7-methyljuglone (144) against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by BACTEC 460 with four- to sixfold reduction of MIC	[88]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Euphorbia ebracteolata</i>	Euphorbiaceae	Roots	80% ethanol	Diterpenoids	Rosane-type diterpenoids 3 (145) and 8 (146) displayed moderate inhibitory effects on with the MIC of 18 μ g/ml and 25 μ g/ml, respectively, by MABA	[89, 90]
<i>Euphorbia lagascae</i>	Euphorbiaceae	Air-dried powdered plant	Methanol extract	Steroids	Ergosterol peroxide (147), cycloart-23-en-3 β ,25-diol (148), vanillin (149), and 4-hydroxybenzaldehyde (150) against <i>M. tuberculosis</i> H37Rv ATCC 27294 strain using two different systems: BACTEC 460TB (Bactec 460)	[91]
<i>Exocarpos latifolius</i>	Santalaceae	Stem	Methanol extract	Fatty acid	Exocarpic acid (E-octadeca-13-ene-9,11-diynoic-acid) (151) with MIC of 20 μ g/ml against <i>M. tuberculosis</i> H37Ra (ATCC 25177) by the thiazolyl blue tetrazolium bromide method	[92]
<i>Fatoua pilosa</i>	Moraceae	Whole plant	Methanol extract	Coumarin; chalcones	Scopoletin (152), isobavachalcone (153) and (E)-1-[2,4-dihydroxy-3-(3-methylbut-2-en-1-yl)phenyl]-3-(2,2-dimethyl-8-hydroxy-2H-benzopyran-6-yl)prop-2-en-1-one (154), copoletin (155), and umbelliferone (156) with MICs of 42, 18, 30, 42, and 58.3 μ g/ml against <i>M. tuberculosis</i> H37Rv by the agar proportion method	[93]
<i>Ferula hermonis</i>	Apiaceae	Root	Ethanol extraction	Octahydroazulen	jaeschkeanadiol benzoate (teferidin) (157) and jaeschkeanadiol p-hydroxybenzoate (ferutin) (158) with MIC values of 3.125 and 1.56 μ g/ml against <i>M. bovis</i> BCG Pasteur 1173P2 and 0.69 and 2 μ g/ml against <i>M. tuberculosis</i> H37Rv, respectively, by fluorescence assay	[94]
<i>Garcinia livingstonei</i>	Guttiferae	Leaf	Acetone extract	Flavonoids	Amentoflavone (159) and 4-monomethoxy amentoflavone (160) with MIC of 0.6 and 1.4 mg/ml against <i>M. smegmatis</i> (ATCC 1441), with the positive control isoniazid (MIC = 1.70 mg/ml) by tetrazolium violet indicator	[95]
<i>Garcinia multiflora</i>	Clusiaceae	Heartwood	Methanol	Biflavones	6,6''-Biapigenin hexamethylether (161), volkensiflavone hexamethylether (162), and hexamethylether of GB-1a (163) against <i>M. tuberculosis</i> H37Rv with inhibition of 96%, 95%, and 87% at 1.25 μ g/ml by BACTEC 460TB	[96]
<i>Garcinia nobilis</i>	Clusiaceae	Stem bark	Methanol	Flavonoids	4-Prenyl-2-(3,7-dimethyl-2,6-octadienyl)-1,3,5,8-tetrahydroxyxanthone (164) with MIC of 8 μ g/ml against <i>M. tuberculosis</i> H37Rv ATCC 27294 and <i>M. tuberculosis</i> clinical MTCS2 by MABA	[97]
<i>Goniothalamus laoticus</i>	Annonaceae	Flower	Ethyl acetate extract	Lactone derivative	(+)-Altholactone (165) and howinin A (166) with MIC of 6.25 μ g/ml, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[98]
<i>Harrisonia perforata</i>	Simaroubaceae	Branches	Ethanol extract	Flavonoids	Perforamone B (5-hydroxy-7-methoxy-2-methyl-8-(1-hydroxy-3-methyl-3-butenyl)chromone) (167) and D (2-hydroxymethylalloptaeroxylin) (168), peucin-7-methyl	[99]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
Helichrysum melanacme	Asteraceae	Shoots	Acetonic extract	Chalcone	ether (169), and greveichromenol (170) with MICs of 25, 25, 50, and 50 µg/ml against <i>M. tuberculosis</i> H37Ra by MABA 2',4',6'-Trihydroxy-3'-prenylchalcone (171) and 4',6',5'',-trihydroxy-6'',6''-dimethyldihydropyrano-[2'',3''-2',3'] chalcone (172) with MICs of 0.05 µg/ml against <i>M. tuberculosis</i> H37Rv by BACTEC radiometric method (3R,8S)-Falarindiol (173) and 6-isopentenylisobergapten (174) with MICs of 24 µM and 167 µM and IC ₅₀ s of 6 µM and 27 µM against <i>M. tuberculosis</i> H37Ra, respectively	[100]
Heracleum maximum	Apiaceae	Root	Methanolic extract	Furanocoumarins	Unsaturated fat oleic and linoleic acids (175) with MIC of 4 and 16 µg/ml against <i>M. fortuitum</i> by thiazolyl blue tetrazolium bromide method	[101]
Humulus lupulus	Cannabaceae	Strobile hops	Hexane extract	Fatty acid	Anthelminticins A (176), B (177), C (178) (11-(cyclopent-1-en-1-yl)-11-oxoundecanoic acid, 2,3-dihydroxypropyl 9-[(R)-cyclopent-2-en-1-yl]nonanoate, 2,3-dihydroxypropyl 13-[(R)-cyclopent-2-en-1-yl]tridecanoate), and two known ones, namely, chaulmoogric acid (179) and ethyl chaulmoograte (180) with MIC of 5.54, 16.70, 4.38, 9.82, and 16.80 µM, respectively, by GFP-expressed <i>M. tuberculosis</i> H37Rv	[102]
Hydnocarpus anthelmintica	Flacourtiaceae	Seeds	95% ethanol	Fatty acid	Hyperenone A (181) against <i>M. tuberculosis</i> H37Rv and <i>M. bovis</i> BCG with MIC of 75 µg/ml and 100 µg/ml, respectively, by thiazolyl blue tetrazolium bromide method	[103]
Hypericum perforatum	Guttiferae	Aerial parts	Hexane and chloroform extracts	Pyranone	Diterpene 12-methyl-5-dehydroacetylthorminone (182) with MIC of 0.38 µg/ml against <i>M. tuberculosis</i> H37Rv by proportion method	[104]
Indigofera longeracemosa	Fabaceae	Stem	Methanol extract	Diterpene	3â,25-Epoxy-3R,21R-dihydroxy-22â-angeloyloxyolean-12-ene-28-oic acid (183), camaric acid (184), and rehmannic acid (185) with MICs of 64, 64, and 32 µM, respectively, against <i>M. tuberculosis</i> H37Rv by BACTEC 460 radiometric system	[105]
Ipomoea leptophylla	Convolvulaceae	Leaf	Organic soluble extract	Triterpenes	Isocupressic acid (186) and communic acid (187) displayed MICs of 78 µM and 31 µM against <i>M. tuberculosis</i> H37Ra, respectively, by microplate resazurin assay	[106]
Juniperus communis	Cupressaceae	Needles and branches	Methanolic extracts	Diterpene	Plumbagin (112) and 7β-hydroxyabieta-8,13-dien-11,12-dione (188) with MIC < 12.5 µg/ml against <i>M. tuberculosis</i> H37Rv by visible growth	[107]
Juniperus procera	Cupressaceae	Leaf and bark	95% ethanol	Diterpene		[108]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
Kaempferia galanga	Zingiberaceae	Rhizomes	Absolute ethanol	Aromatic acid	Ethyl p-methoxycinnamate (189) inhibited the growth of <i>M. tuberculosis</i> H37Ra and H37Rv with MICs of 48.5 and 24.2 mM by Bactec 460 system	[109]
Kaempferia parviflora	Zingiberaceae	Rhizomes	Water	Flavonoids	3,5,7,4'-Tetramethoxyflavone (190) and 5,7,4'-trimethoxyflavone (191) with MIC of 200 and 50 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv by MABA	[110]
Lantana hispida	Verbenaceae	Aerial parts	Hexane extract	Pentacyclic triterpenoids	Three pentacyclic triterpenoids of 3-acetoxy-22-(2'-methyl-2Z-butenyloxy)-12-oleanen-28-oic acid (192), 3-hydroxy-22β-(2'-methyl-2Z-butenyloxy)-12-oleanen-28-oic acid (reduced lantadene A) (193), and oleanolic acid (194) with MIC of 50, 50, and 25 µg/ml against <i>M. tuberculosis</i> H37Rv by MABA	[111]
Lantana horrida	Verbenaceae	Aerial parts	Hexanic extracts	Triterpenes	Ursolic acid (43) and oleanolic acid (194) against <i>M. tuberculosis</i> H37Rv by MABA	[58]
Larrea divaricata	Zygophyllaceae	Aerial parts	Dichloromethane/methanol (1:1)	Flavone, polyphenols	5,7-Dihydroxy-3-methoxy-2-(4-methoxy-phenyl)-4H-chromen-4-one (195); 5,6,7-trihydroxy-3-methoxy-2-(4-methoxyphenyl)-H-chromen-4-one (196); β,γ-dimethyl-α,δ-bis(3,4-dihydroxyphenyl)butane (17) with MIC of 50 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by Bactec 460-TB apparatus	[27]
Ligusticum officinale	Apiaceae	Root	n-Hexane extract	Fatty alcohol	Falcarindiol (198) with MIC of 20 µg/ml and sesquiterpene alcohol a-prethapsenol (199) with MIC of 60 µg/ml against <i>M. tuberculosis</i> H37Rv by spot culture growth inhibition assay	[112]
Limnophila geoffrayi	Scrophulariaceae	Aerial parts	Chloroform extract	Flavonoids	Lavonesnevadensin (5,7-dihydroxy-6,8,4'-trimethoxyflavone) (200) and isothymusin (6,7-dimethoxy-5,8,4'-trihydroxyflavone) (201) against <i>M. tuberculosis</i> H37Ra by MABA	[113]
Lippia turbinata	Verbenaceae	Aerial parts	Methanol-dichloromethane	Triterpenoids	3β,25-Epoxy-3α,21α-dihydroxy-22β-(3-methylbut-2-en-1-oyloxy) olean-12-ene-28-oic acid (202); 3β,25-epoxy-3α,21α-dihydroxy-22β-angeloyloxyolean-12-ene-28-oic acid (203); 3β,25-epoxy-3α,21α-dihydroxy-22β-tigloyloxyolean-12-ene-28-oic acid (204); and 3β,25-epoxy-3R-hydroxy-22β-(2-methylbutan-1-oyloxy)olean-12-ene-28-oic acid (205), lantanilic acid (206), camaric acid (184), lantanolic acid (207), and rehmmanic acid (185) against <i>M. tuberculosis</i> H37Rv (ATCC 27294) using the radiorespirometric BACTEC 460 system	[106]
Litsea hypophaea	Lauraceae	Root	Methanol	Lactone, phenol	Litseekolide L (208) and N-trans-feruloylmethoxytyramine (209) with MIC values of 25	[76]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Lophira lanceolata</i>	Ochnaceae	Roots	Methanol extract	Flavonoids	and 1.6 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv by agar proportion method Dihydroflavone A (210) and lophirone A (211) with MIC of 31.25 and 15.75 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> by MABA	[114]
<i>Lunasia amara</i>	Rutaceae	Leaf	Ethanol	Phenyl quinoline	4-Methoxy-2-phenylquinoline (212) and 4-methoxy-2-(3,4-ethylenedioxy) phenyl quinoline (213) with MICs of 16 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv by BACTEC radiometric method	[115]
<i>Marsypopetalum modestum</i>	Annonaceae	Stem	Ethanol	Dithiopyridine	Dipyrithione (214) with MIC < 0.039 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv by MABA	[116]
<i>Micromelum hirsutum</i>	Rutaceae	Stem bark	Dichloromethane extract	Carbazole	Micromolide ((-)-Z-9-octadecene-4-olide) (215) and five known alkaloids: lansine (216), 3-formylcarbazole (90), and 3-formyl-6-methoxycarbazole (217) with MIC of 1.5, 31.5, 14.3, 42.3, and 15.6 $\mu\text{g/ml}$, against <i>M. tuberculosis</i> H37Rv by MABA	[117]
<i>Microtropis fokenensis</i>	Celastraceae	Root	Methanol	Sesquiterpenes	1 α -Acetoxy-2 α -hydroxy-6 β ,9 β ,15-tribenzoyloxy- β -dihydroarogofuran (218), 2 α -acetoxy-1 α -hydroxy-6 β ,9 β ,15-tribenzoyloxy- β -dihydroarogofuran (219), ricolin G (220), and triptogelin G-2 (221) with MICs of 19.5, 15.8, 14.6, and 26 μM against <i>M. tuberculosis</i> 90-221387 by agar proportion method	[118]
<i>Microtropis japonica</i>	Celastraceae	Stem	Methanol	Sesquiterpenes	15-Acetoxyrubiculin G (222), celahin C (223), salasol A (224), and 8-acetoxymutangan (225) with a MIC of 39.6 μM , 15 μM , and 10 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv by agar proportion method	[119, 120]
<i>Morinda citrifolia</i>	Rubiaceae	Leaves	Hexane fraction	Steroid	E-Phytol (mixture of the two ketosteroids, stigmasta-4-en-3-one (226) and stigmasta-4-22-dien-3-one (227)) and the epidioxysterol campesta-5,7,22-trien-3 β -ol (228) with MIC of 2.5 $\mu\text{g/ml}$ and minus 2.0 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by the growth index	[121, 122]
<i>Mulinum crassifolium</i>	Apiaceae	Aerial parts	n-Hexane	Diterpenoid	13-Hydroxy-mulin-11-en-20-oic-acid methyl ester (229), isomulinic acid n-propyl ester (230), and isomulinic acid n-butyl ester (231) with MIC of 6.25 $\mu\text{g/ml}$, respectively, against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[43]
<i>Myrtus communis</i>	Myrtaceae	Leaf	Hydrodistillation	Monoterpene	Essential oil of limonene (232), 1-8 cineole (233), and α -pinene (234) against <i>M. tuberculosis</i> H37Rv with MIC of 2% (v/v)	[123]
<i>Nigella sativa</i>	Ranunculaceae	Seed	Methanolic extract	Quinones	Thymoquinone (TQ; 2-isopropyl-5-methyl-1, 4-benzoquinone) (117) with MIC of 12.5 $\mu\text{g/ml}$	[124]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Ocimum basilicum</i>	Lamiaceae	Aerial parts (leaves, fruits, and flowers)	Methanolic extract	Fatty acid ester	intracellular <i>M. tuberculosis</i> H37Rv and extensively drug-resistant tuberculosis (XDR-TB) 72 h postinfection in RAW 264.7 cells (E)-3'-Hydroxy-4'-(1''-hydroxyethyl)-phenyl-4-methoxycinnamate (235) against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[125]
<i>Ocimum sanctum</i>	Lamiaceae	Leaf	Ethanol extract	Triterpenes	Ursolic acid (UA, 3b-hydroxy-urs-12-en-28-oic-acid) (43) with MIC of 12.5 µg/ml against <i>M. tuberculosis</i> by MABA	[40, 126]
<i>Ocotea macrophylla</i>	Lauraceae	Wood, leaf	Ethanol extract	Apomorphine	(S)-3-Methoxynordomesticine hydrochloride (236) with MIC of 4 µg/ml against <i>M. tuberculosis</i> H37Rv by thiazolyl blue tetrazolium bromide method	[127]
<i>Oplopanax horridus</i>	Araliaceae	Inner stem bark	Consecutively extracted with dichloromethane, methanol, and 50% aqueous methanol	Sesquiterpenes	Oplopandiol (237) and faltarindiol (198) with MI of 61.5 and 6.2 µg/ml against <i>M. tuberculosis</i> (ATCC 35801) by MABA	[128]
<i>Pedilanthus tithymaloides</i>	Euphorbiaceae	Root	Successively using hexane, dichloromethane, and methanol	Diterpenoid	Caniojane (238) against <i>M. tuberculosis</i> H37Ra with MIC of 25 µg/ml by MABA	[129]
<i>Phoradendron robinsonii</i>	Santalaceae	Whole plant	Dichloromethane/methanol (1:1)	Flavanone	5-Hydroxy-2-(4'-hydroxyphenyl)-7-methoxy-2,3-dihydro-4H-chromen-4-one(239) with MIC of 128 µg/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by Bactec 460-TB apparatus	[27]
<i>Physalis angulata</i>	Solanaceae	Aerial parts	Chloroform	Physalin	Physalin D (240) with MIC of 32 µg/ml against <i>M. tuberculosis</i> H37Rv ATCC 27294 by MABA	[130]
<i>Piper chaba</i>	Piperaceae	Root	95% ethanol	Amide	Pellitorine (241), guineensine (242), sarmentine (243), brachyamide B (244), 1-piperetyl pyrrolidine (245), sarmentosine (246), and 1-(3,4-methylenedioxyphenyl)-1E-tetradecene (247) with MIC values in the range of 25–200 µg/ml against <i>M. tuberculosis</i> H37Ra by MABA	[99, 131]
<i>Piper regnellii</i>	Piperaceae	Leaves and stems	Supercritical fluid extraction (CO2 extracts)	Benzofuran	Eupomatenoid-3 (248) and 5-(3-methyl-5-propylbenzofuran-2-yl) benzo[d][1,3]dioxole (249) against <i>M. tuberculosis</i> H37Rv by colorimetric resazurin microtiter assay plate	[132]
<i>Piper solmsianum</i>	Piperaceae	Leaf	Methanolic extract	Benzofuran	Synergism was observed in <i>M. tuberculosis</i> H37Rv and eight clinical isolates with eupomatenoid-5 (250) + rifampicin and in <i>M. tuberculosis</i> H37Rv and 17 clinical isolates with eupomatenoid-5+ ethambutol combinations	[133]
<i>Plectranthus grandidentatus</i>	Lamiaceae	Aerial parts	Chloroform	Royleanone	Abietanes 6β,7α-dihydroxyroyleanone (251), horminone, and 6,7-dehydroroyleanone (252) with MIC of 25 and	[134]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
Plectranthus ornatus	Lamiaceae	Aerial parts	Chloroform	Royleanone	12.5 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv by MTT 7α-Acetoxy-6β-hydroxyroyleanone (MIC 0.0083 µM) (253) and 6,7-dehydroroyleanone (MIC 0.039 µM) (254) against <i>M. tuberculosis</i> H37Rv by thiazolyl blue tetrazolium bromide method	[134, 135]
Plumbago indica	Plumbaginaceae	Root	Petroleum ether	Naphthalene	Plumbagin (112) with MIC of 0.25 and 8 µg/ml against MDR-TB and 2 and 4 µg/ml against the XDR-TB isolates by thiazolyl blue tetrazolium bromide method	[78]
Plumeria bicolor	Apocynaceae	Leaves	Chloroform extract	Naphthalene	Plumericin (112) against active and MDR TB with MIC of 0.12, 0.15, 0.07, 0.13, and 0.14 µg/ml, respectively, better than isoplugmercin	[136]
Polyalthia cerasoides	Annonaceae	Root	Extracted successively with hexane, ethyl acetate, and methanol	Apomorphine, fatty acid, sesquiterpenes	Bidebilin E (255), octadeca-9,11,13-triynoic acid (256), and α-humulene (257), with 6.25 µg/ml against <i>M. tuberculosis</i> H37Ra MABA	[137]
Polyalthia debilis	Annonaceae	Root	Methanol	Lactone derivative	Debilisones B (258), C (259), and E (260) with MIC of 25, 12.5, and 25 µg/ml, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[138]
Polyalthia evecta	Annonaceae	Root	Extracted successively with hexane, dichloromethane, and methanol	Furan	Furanoid polyacetylene (261) with MIC of 3.1 µg/ml against <i>M. tuberculosis</i> H37Ra by MABA	[139]
Potamogeton malaianus	Potamogetonaceae	Whole plant	Dichloromethane	Diterpenes	Potamogetonyde (262), potamogetonol (263), potamogetonin (264), and 15,16-epoxy-12-oxo-8 (17),13 (16),14-labdatrien-20,19-olide (265) with MIC of 50–100 µg/ml against <i>M. tuberculosis</i> H37Ra by MABA	[140]
Pourthiaea lucida	Rosaceae	Leaf	Methanol	Alcohol	a-Tocospino A (266) and B (267), a-tocopherylquinone (268), and (E)-phytol (124) with MICs of 30, 50, 25, and 12.5 µg/ml against <i>M. tuberculosis</i> H37Rv by agar proportion method	[141]
Premna odorata	Lamiaceae	Leaf	Methanol	Aldehydes	1-Heneicosyl formate (269) with MIC of 8 µg/ml against <i>M. tuberculosis</i> H37Rv (ATCC27294) by MABA	[142]
Prunus cerasoides	Rosaceae	Root	Successively with hexane, ethyl acetate, and methanol	Fatty acid	Octadeca-9,11,13-triynoic acid (256) with MIC of 6.25 µg/ml against <i>M. tuberculosis</i> H37Ra by MABA	[137]
Punica granatum	Punicaceae	Peel of the fruit	Water	Polyphenol	Epigallocatechin-3-gallate (EGCG) (270) and quercetin (271) with MIC of 32–256 µg/ml against nine <i>M. tuberculosis</i> isolates by thiazolyl blue tetrazolium bromide method	[143]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
Radermachera boniana	Bignoniaceae	Leaves and twigs	Ethyl acetate	Sterol	Ergosterol peroxide (147) and b-sitostenone (272) with MICs of 34.8 and 9.9 μ M, respectively, against <i>M. tuberculosis</i> H37Rv by MABA	[144]
Ranunculus ternatus	Ranunculaceae	Roots	Ethanol	Benzophenones	(benzophenones) Methyl (R)-3-[2-(3,4-dihydroxybenzoyl)-4,5-dihydroxyphenyl]-2-hydroxypropanoate (273) with MIC of 41.67 μ g/ml against <i>M. tuberculosis</i> H37Rv by MABA	[145]
Rumex hymenosepalus	Polygonaceae	Root	Dichloromethane/methanol (1:1)	Diphenylethylene	5-[(E)-2-(4-Acetoxyphenyl) ethenyl]-1,3-benzenediol(1a) (274) with MIC of 32 μ g/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by Bactec 460-TB apparatus	[27]
Rumex nepalensis	Polygonaceae	Root	Ethanol extracts	Glycoside	Rumexneposide A (275), torachrysone (276), nepodin-8-O-b-D-glucopyranoside (277), torachrysone-8-O-b-D-glucopyranoside (278), and chrysophanol-8-O-b-D-glucopyranoside (279), which showed MICs of 20.7, 6.1, 26.6, 8.9, and 4.1 μ M, respectively, by fluorescence assay	[146]
Salvia africana lutea	Lamiaceae	Aerial parts	Ethanol extract	Diterpene	Abietane-type diterpene carnosic acid (280) with MIC of 28 μ M against <i>M. tuberculosis</i> H37Rv (ATCC27294) by a rapid radiometric method	[147]
Salvia miltiorrhiza	Lamiaceae	Roots	Acetone	Tanshinones	Tanshinone I (281), tanshinone IIA (282), and cryptotanshinone (283) with MIC in the range of 1.17–26.57 μ g/ml against <i>M. tuberculosis</i> H37Rv by agar proportion method	[148, 149]
Sapium indicum	Euphorbiaceae	Fruit	Hexane extract	Phorbol ester	Sapintoxin A (284), sapintoxin B (285), 12-(2 α -N-methylaminobenzoyl)-4R-deoxy-5,20-dihydroxyphorbol-13-acetate (286), and 12-(2-methylaminobenzoyl)-4-deoxyphorbolaldehyde-13-acetate (287) with MIC of 3.12, 12.5, 25, and 25 μ g/ml, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[150]
Saussurea lappa	Asteraceae	Bark	Ethanol extract	Sesquiterpenoids	Saussureamine C (methyl 3-O-feruloylquininate) (288) against <i>M. tuberculosis</i> by inhibiting folC	[151]
Scleropyrum wallichianum	Santalaceae	Twig	Successively with n-hexane, chloroform, and methanol	Fatty acid	Scleropyric acid (289) with MIC of 25 μ g/ml against <i>M. tuberculosis</i> H37Ra by MABA	[152]
Solanum torvum	Solanaceae	Fruit	Methanol extracts	Xanthine	Methyl caffeate (290) with MIC of 8 μ g/ml against <i>M. tuberculosis</i> by agar proportion method	[153]
Stephania dinklagei	Menispermaceae	Aerial parts	Chloroform extract	Flavanone	Flavanone pinostrobin (291) against <i>M. tuberculosis</i> H37Rv with MIC of 3.125 μ g/ml by MABA	[49]
Strobilanthes cusia	Acanthaceae	Leaf	Methanol extracts	Quinazoline	Tryptanthrin (292) with MIC of 1 mg/ml against <i>M. tuberculosis</i> by BACTECH	[154, 155]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Tabernaemontana citrifolia</i>	Apocynaceae	Leaf	Chloroform extract	Voacangine	Ibogaïne (293) and voacangine (294) with MIC of 50 µg/ml against <i>M. tuberculosis</i> H37Rv by Bactec 460-TB radiometric methodology	[28]
<i>Teloxys graveolens</i>	Chenopodiaceae	Aerial parts	Acetone extract	Flavanone	Flavanone pinostrobin (291) against <i>M. tuberculosis</i> H37Rv with MIC of 12.5 µg/ml by MABA	[49]
<i>Terminalia avicennioides</i>	Combretaceae	Root bark	Successively with petroleum ether, ethyl acetate, chloroform, and methanol	Triterpenes	Arjunolic acid (295) and friedelin (296) which, respectively, had MICs against BCG of 156 µg/ml and 4.9 µg/ml, respectively, by broth microdilution method	[156, 157]
<i>Terminalia brownii</i>	Combretaceae	Root	Methanol extract	Flavones, ellagic acid	Methyl (S)-flavogallionate (297), ellagic acid xyloside (298), and methyl ellagic acid xyloside (299) against <i>M. smegmatis</i> by measured spectrophotometrically at 620 nm	[158]
<i>Byrsonima fagifolia</i>	Malpighiaceae	Leaf	Chloroform extract	Amyrin	α- and β-Amyrin (300), α-amyrin acetate (301), and dotriacontane (302) with MIC of 31.25, 62.5, and 62.5 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[159]
<i>Terminalia laxiflora</i>	Combretaceae	Root	Methanol extract	Triterpenes, fatty alcohol, sterol	Friedelin (296), triacontanol (303), β-sitosterol (34), and sitosterone (35) with MIC of 250 µg/ml, 250 µg/ml, 500 µg/ml, and 500 µg/ml, respectively, against <i>M. smegmatis</i> by measured spectrophotometrically at 620 nm	[158]
<i>Terminalia superba</i>	Combretaceae	Stem bark	Methanol extract	Ellagic acid	3,4'-Di-O-methyllellagic acid 3'-O-β-D-xylopyranoside (306) and 4'-O-galloyl-3,3'-di-O-methyllellagic acid 4-O-β-D-xylopyranoside (307) with MIC of 4.88 µg/ml and 9.76 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv (ATCC 27294) by MABA	[160]
<i>Tetracera potatoria</i>	Dilleniaceae	Stem bark	Methanol/ dichloromethane (1 : 1)	Alcohol	Tetraceranoate (308) and N-hydroxy imidate-tetracerane (309) with MIC of 7.8 µg/ml and 15 µg/ml, respectively, against <i>M. smegmatis</i> (ATCC 23246) by tetrazolium method	[161]
<i>Tetradenia riparia</i>	Lamiaceae	Leaf	Hydrodistillation	Royleanone	6,7-Dehydoroyleanone (254) with MIC of 31.2 µg/ml against <i>M. tuberculosis</i> H37Rv by resazurin microtiter assay	[162]
<i>Thalia multiflora</i>	Marantaceae	Aerial parts	Dichloromethane-methanol (1 : 1) followed by a 100% methanol	Steroids	Stigmast-5-en-3β-ol-7-one (310), stigmast-4-ene-6β-ol-3-one (311), stigmast-5,22-dien-3β-ol-7-one (312), and stigmast-4,22-dien-6β-ol-3-one (313) were found to be the most active compounds with MIC of 1.9, 4, 1.0, and 2.2 µg/ml, respectively, against <i>M. tuberculosis</i> by fluorescence assay	[163]

TABLE 1: Continued.

Plant species	Plant family	Part used	Extracts	Compound class	Active constituents	References
<i>Tiliacora triandra</i>	Menispermaceae	Root	Dichloromethane	Bisbenzylisoquinoline alkaloids	Tiliacoronine (314), 2'-nortiliacoronine (315), and tiliacoronine (316) are bisbenzylisoquinoline alkaloids with MIC of 3.1–6.2 mg/ml against <i>M. tuberculosis</i> different strains by MABA	[164]
<i>Tussilago farfara</i>	Asteraceae	Aerial parts	Soxhlet extracted	Aromatic acid	p-Coumaric acid (317) and 4-hydroxybenzoic acid (318) with MIC 31.3 µg/ml of 62.5 µg/ml against <i>M. tuberculosis</i> H37Rv by high throughput spot culture growth inhibition assay	[165]
<i>Ventilago madraspatana</i>	Rhamnaceae	Stem bark	Methanol	Anthraquinone	Emodin (115) with MIC of 4 µg/ml against <i>M. tuberculosis</i> H37Rv by MABA	[78, 166]
<i>Zanthoxylum capense</i>	Rutaceae	Leaf	80% ethanol	Phenol, amide	Decarine (319) and an N-isobutylamide, N-isobutyl-(2E,4E)-2,4-tetradecadienamide (320) with MIC of 1.6 µg/ml against <i>M. tuberculosis</i> H37Rv by measuring the optical density at 600 nm in a Tecan M200 plate spectrophotometer	[167]
<i>Zanthoxylum leprieurii</i>	Rutaceae	Stem bark	Methanol extract	Dihydroacridine	Hydroxy-1, 3-dimethoxy-10-methyl-9-acridone (321), 3-hydroxy-1, 5, 6-trimethoxy-9-acridone (322) with MIC of 5.1 and 1.5 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv by MABA	[168]
<i>Zanthoxylum schinifolium</i>	Rutaceae	Leaf	Methanolic extracts	Coumarin	7-[(2E)-3,7-dimethylocta-2,6-dienoxy]-8-methoxychromen-2-one (collinin) (323) with MIC50 of 3.13–6.25 µg/ml against both drug-susceptible and -resistant strains of <i>M. tuberculosis</i> by luminescent viability assay kit	[169]
<i>Zanthoxylum wutaense</i>	Rutaceae	Root	Methanol extract	Benzofuran, furo[3,2-b]quinoline	7-Methoxyanodendroate (324), 7-methoxywutaifuranal (325), wutaensal (326), dictamine (327), and γ-fagarine (328), with MIC of 35, 35, 30, 30, and 30 µg/ml, respectively, against <i>M. tuberculosis</i> H37Rv by the agar proportion method	[170]
<i>Zingiber cassumunar</i>	Zingiberaceae	Root	Methanol	Three fatty acid esters	(E)-4-(3,4-Dimethoxyphenyl)but-3-en-1-yl linoleate (329), (E)-4-(3,4-dimethoxyphenyl)but-3-en-1-yl oleate (330), and (E)-4-(3,4-dimethoxyphenyl) but-3-en-1-yl palmitate (331), with MIC of 200, 100, and 200 µg/ml, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[171]
<i>Ziziphus cambodiana</i>	Rhamnaceae	Root bark	Acetate extract	Triterpenes	3-O-Vanillylceanothic acid (332), betulinolaldehyde (333), betulinic acid (334), and 2-O-E-p-coumaryl aliphatic acid (335) with MIC of 25, 25, 25, and 12.5 µg/ml, respectively, against <i>M. tuberculosis</i> H37Ra by MABA	[172]

MABA: microplate alamar blue assay; MIC: minimum inhibitory concentration.

FIGURE 2: Typical structures sorted out of the 335 anti-TB compounds from the reported medicinal plants, which included sesquiterpene, diterpene, triterpene, quinone, polyphenol, fatty alcohol, flavanone, flavonoid, chalcone, benzofuran, fatty acid ester, fatty acid, steroid, amide, and coumarin.

abietane diterpenoid had an MIC of 1.2 $\mu\text{g/ml}$, while its C-12 acetate analogue was more active with an MIC of 0.89 $\mu\text{g/ml}$ [175]. One of the most impressive natural products was (+)-calanolide A, a novel dipyrano coumarin from the Malaysian tree *Calophyllum lanigerum* var. *austrocoriaceum*. This distinct compound was first reported with good activity against the strains of HIV-1, which was resistant to diverse other nonnucleosides as well as nucleoside (AZT) reverse transcriptase inhibitors [176, 177]. Later, the novel calanolides with the ring-D-modification were synthesized with selective activity against the replication and/or nonreplicating *M. tuberculosis* by targeting the Rv2466c [55]. In particular, analogues bearing 2-nitrofurano group at the ring D position markedly improved the *in vitro* efficacy and reduced the mammalian cell toxicity, when compared with the parent compound (+)-calanolide A [55]. Recently, Mu et al. demonstrated that the nitrofuranyl calanolides could be employed as novel fluorescent probes that can serve as a much needed high-throughput and low-cost detection method for detection of living *M. tuberculosis* and can precisely determine the MIC values for a full range of available drugs [178]. Thus, different modifications of the calanolide derivatives demonstrated three aspects (anti-HIV, anti-TB, and TB diagnosis) of potent usage in TB disease.

Of all the 335 natural plant compounds and its semisynthetic analogues, only few were found for their mechanistic role of their anti-TB activities. The calanolides target the Rv2466c, and hyperenone A inhibits the ATP-dependent MurE ligase, which involves in the cytoplasmic steps of peptidoglycan biosynthesis [104]. It was reported that saussureamine C (methyl 3-O-feruloylquininate) targets the folC [151] and eupractenoid B targets the acetyl transfer activity of GlmU [90]. The trans, trans-1, 7-diphenylhepta-4, 6-dien-3-one target the efflux pumps [26]. *In silico* analysis revealed that some fatty acids could bind at the cleft between the N-terminal and C-terminal lobes of the cytosolic domain of serine/threonine protein kinase B (PknB) [23]. The anti-TB plant medicinal compounds included in this review lack the molecular basis of the action and mechanisms of modulation on the metabolism of *M. tuberculosis* nor the immunomodulatory activities of those compounds.

4. Plants Showing Anti-TB Effect in Form of Crude Extracts

The plants whose active components were isolated and tested for their anti-TB activity as described in Section 3. This section summarizes the reported plants for their anti-TB activity only in the form of crude extracts. They are listed in Table 2 with the total amount of 128 plant species. The top five plant families were *Asteraceae*, *Euphorbiaceae*, *Fabaceae*, *Piperaceae*, and *Acanthaceae*, and the plant parts mainly used for extraction were root for *Fabaceae* family and leaf for *Asteraceae*, *Piperaceae*, and *Acanthaceae*, respectively. For *Euphorbiaceae* family, the plant parts of bark, fruit, leaf, root, and seed were reported with the anti-TB function.

Among the extraction methods, ethanol, hexane, and methanol were found to be the top three frequently used

TABLE 2: Medicinal plants and their crude extracts showing *in vitro* anti-TB activity.

Plant species	Plant family	Part used	Extracts	References
Acacia catechu	Liliaceae	Root	Hexane extracts with inhibition of mycobacterial (standard and clinical) growth	[179]
Acacia senegal	Fabaceae	Root	Aqueous extraction has potential antimycobacterial activity	[180]
Acalypha indica	Euphorbiaceae	Leaf	Aqueous extracts with inhibition of 95% at 4 percent <i>v/v</i> concentration in L-J medium for sensitive <i>M. tuberculosis</i> H37Rv	[181]
Achyrocline alata	Asteraceae	Leaf, stem	Aqueous extracts against <i>M. tuberculosis</i> H37Rv strain (ATCC 27294) with MIC ^a of 62.5 $\mu\text{g/ml}$	[182]
Acorus calamus	Acoraceae	Root	Aqueous extracts against <i>M. bovis</i> BCG by OD units	[183]
Adhatoda vasica	Acanthaceae	Leaf	Methanol extracts with the oils inhibiting the growth of MTB B19-4 at 2 $\mu\text{g/ml}$	[180]
Allium sativum	Liliaceae	Leaf	Aqueous extract was found to be 63% at 4 percent <i>v/v</i> concentration in L-J medium for sensitive <i>M. tuberculosis</i> H37Rv	[181]
Aloe vera	Xanthorrhoeaceae	Leaf	Aqueous extract was found to be 41% at 4 percent <i>v/v</i> concentration in L-J medium for sensitive <i>M. tuberculosis</i> H37Rv	[181, 184]
Alstonia scholaris	Apocynaceae	Leaf	Methanol extracts have potential antimycobacterial activity and the synergistic group consisting of rifampicin in murine model	[185]
Amborella trichopoda	Amborellaceae	Fruit	Methanol extracts against <i>M. bovis</i> BCG (strain 11-73 P2) with MIC of 2.5 $\mu\text{g/ml}$	[186]
Ambrosia ambrosioides	Asteraceae	Aerial parts	Methanolic extracts against <i>M. tuberculosis</i> H37Rv with MIC of 790 $\mu\text{g/ml}$	[187]
Ambrosia confertiflora	Asteraceae	Aerial parts	Methanol, chloroform, dichloromethane, and ethyl acetate extracts against <i>M. tuberculosis</i> H37Rv with MIC of 200, 90, 120, and 160 $\mu\text{g/ml}$, respectively	[187]
Amphipterygium simplicifolium	Julianaceae	Leaf	Dichloromethane-methanol extracts (1:1) inhibit the <i>M. tuberculosis</i> H37Rv at 50 $\mu\text{g/ml}$ with $90.5 \pm 1.0\%$	[188]
Andrographis paniculata	Acanthaceae	Aerial parts	Hexane and methanol (1:5) extracts with maximum antimycobacterial activity at 250 $\mu\text{g/ml}$ against all the tested strains of <i>M. tuberculosis</i> (H37Rv, MDR, and drug sensitive)	[189]
Andrographis paniculata	Acanthaceae	Leaf	Ethanol extracts with inhibition of mycobacterial (standard and clinical) growth	[7, 190]
Angiopteris evecta	Marattiaceae	Leaf	80% methanol extract against <i>M. tuberculosis</i> H37Rv ATCC 25618 with an MIC of 400 $\mu\text{g/ml}$	[191]
Apodytes dimidiata	Icacinaeae	Leaf	Hexane extractions against the field strain of MDR-TB and against the <i>M. tuberculosis</i> H37Rv with MIC of 0.47 and 0.31 mg/ml, respectively	[192]
Artemisia ludoviciana	Asteraceae	Bark, leaf	Hexane extracts against MDR-TB clinical isolates with MIC of 25-100 $\mu\text{g/ml}$	[58, 193]
Artemisia nilagirica	Asteraceae	Leaf	Ethanol extracts against <i>M. smegmatis</i> with IC ₅₀ ^c of 300 $\mu\text{g/ml}$	[194]
Beilschmiedia obscura	Lauraceae	Root	Ethyl acetate extracts against <i>M. tuberculosis</i> H37Rv with MIC of 31.25 $\mu\text{g/ml}$ by MABA ^b	[195]
Bidens odorata	Asteraceae	Aerial parts	Hexane, dichloromethane, ethyl acetate, and ethanolic extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 100, 12.5, 12.5, and 12.5 $\mu\text{g/ml}$	[196]
Bridelia micrantha	Euphorbiaceae	Bark	Acetone extracts against <i>M. tuberculosis</i> H37Ra with MIC of 25 $\mu\text{g/ml}$	[197]
Calluna vulgaris	Ericaceae	Aerial parts	Ethyl acetate extracts with 97% inhibition at 100 $\mu\text{g/ml}$ against <i>M. tuberculosis</i> H37Rv (ATCC 27294)	[198]
Calophyllum brasiliense	Clusiaceae	Leaf	Dichloromethane-methanol extracts (1:1) inhibit the <i>M. tuberculosis</i> H37Rv at 50 $\mu\text{g/ml}$ with $82.8 \pm 0.4\%$	[188]
Capparis zeylanica	Capparidaceae	Leaf	Ethyl acetate extracts against <i>M. tuberculosis</i> H37Rv with the 32 mm minimum zone of inhibition	[199, 200]
Carya illinoensis	Juglandaceae	Bark	Hexane extracts against <i>M. tuberculosis</i> H37Rv with MIC of 31 $\mu\text{g/ml}$	[193]
Cassia sophera	Caesalpiniaceae	Seed	Methanol extracts against <i>M. smegmatis</i> with MIC of 125 $\mu\text{g/ml}$	[201]

TABLE 2: Continued.

Plant species	Plant family	Part used	Extracts	References
<i>Chenopodium ambrosioides</i>	Amaranthaceae	Leaf	80% ethanol crude extracts against <i>M. tuberculosis</i> H37Ra (ATCC 25177™) with MIC of 5000 µg/ml	[184]
<i>Chrysactinia mexicana</i>	Asteraceae	Root	Ethyl ether extract against a drug-resistant strain of <i>M. tuberculosis</i> CIBIN/UMF15:99 with MIC of 62.5 µg/ml	[202]
<i>Citrullus colocynthis</i>	Cucurbitaceae	Leaf	Chloroform extracts against <i>M. tuberculosis</i> H37Rv with MIC of 2.5 mg/ml by MABA	[203]
<i>Citrus sinensis</i>	Rutaceae	Fruit peel	Hexane extracts against two drug-resistant strains of <i>M. tuberculosis</i> with MIC of 25 and 50 µg/ml	[204]
<i>Cladonia arbuscula</i>	Cladoniaceae	Root	Hexane and ethyl acetate extracts with 96% and 99% inhibition at 100 µg/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294), respectively	[198]
<i>Cocculus hirsutus</i>	Menispermaceae	Leaf	Ethanol extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 500 µg/ml	[196]
<i>Codiaeum peltatum</i>	Euphorbiaceae	Stem	Methanol extracts against <i>M. bovis</i> BCG (strain 11-73 P2) with MIC of 100 µg/ml	[186]
<i>Combretum aculeatum</i>	Combretaceae	Aerial part	Aqueous extracts inhibiting <i>M. marinum</i> with MIC of 0.2 mg/ml	[205]
<i>Costus speciosus</i>	Zingiberaceae	Stem, flower	Hexane partition from methanol extracts against <i>M. tuberculosis</i> H37Rv with MIC of 100 µg/ml	[206]
<i>Cremaspora triflora</i>	Rubiaceae	Leaf	Acetone extracts decreased 16-fold of MIC in combination with rifampicin against <i>M. aurum</i> and reduction of the MICs of the anti-TB drug ranged from 2-fold to 4-fold, 2-fold to 64-fold, and 2-fold to 64-fold for <i>M. smegmatis</i> , <i>M. aurum</i> , and <i>M. tuberculosis</i> , respectively	[207]
<i>Croton sylvaticus</i>	Euphorbiaceae	Leaf, root, stem bark	Decoction, not known	[208]
<i>Curcuma caesia</i>	Zingiberaceae	Rhizome	Ethanol extract against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 31.25 µg/ml	[196]
<i>Cymbopogon citratus</i>	Poaceae	Stem, rhizome	Hexane partition from methanol extracts of 200 µg/ml against <i>M. tuberculosis</i> H37Rv	[206]
<i>Cyperus rotundus</i>	Cyperaceae	Root	Ethanol extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 62.5 µg/ml	[209]
<i>Davilla elliptica</i>	Dilleniaceae	Leaf	Chloroform extracts showed a promising antimycobacterial activity with a MIC of 62.5 µg/ml by MABA	[209]
<i>Dissotis rotundifolia</i>	Melastomataceae	Leaf	80% ethanol crude extracts against <i>M. tuberculosis</i> H37Ra (ATCC 25177™) with MIC of 5000 µg/ml	[184]
<i>Dryopteris stewartii</i>	Dryopteridaceae	Whole plant	Decoction, not known	[203]
<i>Echinops giganteus</i>	Asteraceae	Root	Methanol extracts against <i>M. tuberculosis</i> H37Ra and H37Rv with MIC of 32 and 16 µg/ml, respectively	[210]
<i>Empetrum nigrum</i>	Empetraceae	Root	Hexane extracts with 95% inhibition at 100 µg/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294)	[198]
<i>Erythrina abyssinica</i>	Fabaceae	Root bark	Methanol extracts showed the highest activity on <i>M. tuberculosis</i> H37Rv (MIC 390 µg/ml)	[211]
<i>Eulophia nuda</i>	Orchidaceae	Tubers	Ethanol extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 500 µg/ml	[196]
<i>Euphorbia albomarginata</i>	Euphorbiaceae	Shoots	Extracts by n-hexane, dichloromethane, ethyl acetate, and methanol individually against <i>M. tuberculosis</i> H37Rv with MIC of 250-1000 µg/ml	[212]
<i>Euphorbia hirta</i>	Euphorbiaceae	Leaf	Ethyl acetate extracts showed better activity with maximum of 64.73% reduction in relative light units against <i>M. tuberculosis</i> H37R	[213]
<i>Evodia elleryana</i>	Rutaceae	Bark	Ethyl acetate extracts with 95% inhibition of <i>M. tuberculosis</i> H37Ra grown in vitro (ATCC 25177) at 50 µg/ml	[214]
<i>Ficus sur</i>	Moraceae	Root	80% ethanol against <i>M. tuberculosis</i> H37Ra (ATCC 25177) with MIC of 0.78 mg/ml	[215]
<i>Ficus citrifolia</i>	Moraceae	Leaf	95% ethanol extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with 91% inhibition at 100 µg/ml	[216]

TABLE 2: Continued.

Plant species	Plant family	Part used	Extracts	References
<i>Flourensia cernua</i>	Asteraceae	Leaf	Hexane extracts against sensitive and resistant strains, respectively, with MIC of 25-50 $\mu\text{g/ml}$	[217]
<i>Foeniculum vulgare</i>	Umbelliferae	Aerial parts	Hexane extracts against <i>M. tuberculosis</i> H37Rv with MIC of 100 $\mu\text{g/ml}$	[204]
<i>Globularia alypum</i>	Globulariaceae	Leaf	Petroleum ether extracts against <i>M. tuberculosis</i> H37Rv with IC_{50} of 77 $\mu\text{g/ml}$	[218]
<i>Glycyrrhiza glabra</i>	Fabaceae	Root	Ethanol extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 250 $\mu\text{g/ml}$	[219]
<i>Guaiacum coulteri</i>	Zygophyllaceae	Flower	Methanol extracts against <i>M. tuberculosis</i> H37Rv with MIC of 1000 $\mu\text{g/ml}$	[188]
<i>Guiera senegalensis</i>	Combretaceae	Aerial parts	Aqueous extracts inhibiting <i>M. marinum</i> with MIC of 200 $\mu\text{g/ml}$	[205]
<i>Gymnosperma glutinosum</i>	Asteraceae	Leaf	Hexane extracts against <i>M. tuberculosis</i> H37Ra and H37Rv both at 31.2 $\mu\text{g/ml}$	[220]
<i>Helianthus annuus</i>	Asteraceae	Stem	Extracts by n-hexane, dichloromethane, ethyl acetate, and methanol individually against <i>M. tuberculosis</i> H37Rv with MIC of 250-500 $\mu\text{g/ml}$	[212]
<i>Heracleum maximum</i>	Apiaceae	Root	Aqueous extracts against <i>M. bovis</i> BCG by OD units	[183]
<i>Heteromorpha trifoliata</i>	Apiaceae	Leaf	Ethanol extracts against <i>M. tuberculosis</i> H37Rv with MIC of 80 $\mu\text{g/ml}$	[221]
<i>Hygrophila auriculata</i>	Acanthaceae	Root, leaf	Acetone extract against <i>M. tuberculosis</i> H37Rv by the disc diffusion method	[222]
<i>Juglans mollis</i>	Juglandaceae	Bark	Hexane extracts against <i>M. tuberculosis</i> H37Rv with MIC of 50 $\mu\text{g/ml}$	[193]
<i>Juglans regia</i>	Juglandaceae	Bark, leaf	Hexane extracts against <i>M. tuberculosis</i> strain H37Rv with MIC of 100 $\mu\text{g/ml}$	[193, 194]
<i>Justicia adhatoda</i>	Acanthaceae	Leaf	Ethanol extract against <i>M. tuberculosis</i> H37Rv by the disc diffusion method	[223]
<i>Khaya senegalensis</i>	Meliaceae	Bark, leaf	Ethanol extracts against <i>M. tuberculosis</i> H37Ra with MIC of 6.25 $\mu\text{g/ml}$	[224]
<i>Lantana camara</i>	Verbenaceae	Leaf	Methanol extracts against <i>M. tuberculosis</i> H37Rv with MIC of 20 $\mu\text{g/ml}$	[17]
<i>Lantana hispida</i>	Verbenaceae	Leaf	Hexane extracts against drug-resistant clinical isolates of <i>M. tuberculosis</i> with MIC of 100-200 $\mu\text{g/ml}$	[193, 194]
<i>Laurelia novae-zelandiae</i>	Monimiaceae	Leaf, flower	Aqueous extract against <i>M. smegmatis</i> with IC_{50} of 0.02 mg/ml	[225]
<i>Leucophyllum frutescens</i>	Scrophulariaceae	Root, leaf	Methanol extracts against a drug-resistant strain of <i>M. tuberculosis</i> CIBIN/UMF15:99 with MIC of 62.5 $\mu\text{g/ml}$	[202]
<i>Maerua edulis</i>	Capparaceae	Root	Hexane extracts against <i>M. bovis</i> BCG, <i>M. tuberculosis</i> H37Ra with MIC 31.2–62.5 $\mu\text{g/ml}$	[226]
<i>Mallotus philippensis</i>	Euphorbiaceae	Leaf, fruit	Ethanol extracts of fruit and leaves against <i>M. tuberculosis</i> H37Rv (ATCC 27294) both with MIC of 250 $\mu\text{g/ml}$	[227]
<i>Metrosideros excelsa</i>	Myrtaceae	Leaf	Methanol extracts against <i>M. smegmatis</i> with IC_{50} of 0.11 mg/ml	[226]
<i>Millettia stuhlmannii</i>	Fabaceae	Leaf	Acetone extracts against <i>M. smegmatis</i> with MIC of 0.13 mg/ml	[228]
<i>Morinda citrifolia</i>	Rubiaceae	Leaf	Aqueous extract has an inhibition rate of 89% against <i>M. tuberculosis</i> H37Rv	[180]
<i>Mucuna imbricata</i>	Fabaceae	Seed	Methanol extracts have potential antimycobacterial activity and the synergistic group consisting of rifampicin in murine model	[185]
<i>Murraya koenigii</i>	Rutaceae	Leaf	Ethanol extracts against <i>M. smegmatis</i> with IC_{50} of 300 $\mu\text{g/ml}$	[194]
<i>Musa acuminata</i>	Musaceae	Stem	Methanol extracts against drug-resistant variants of <i>M. tuberculosis</i> with MIC of 200 $\mu\text{g/ml}$	[204]
<i>Myoporum crassifolium</i>	Scrophulariaceae	Wood	Hydrodistillation with essential oils against <i>M. bovis</i> BCG (strain 11-73 P2) with MIC of 50 $\mu\text{g/ml}$	[186]

TABLE 2: Continued.

Plant species	Plant family	Part used	Extracts	References
<i>Myrica gale</i>	Myricaceae	Root, stem	Ethyl acetate extracts with 96% inhibition at 100 μ g/ml against <i>M. tuberculosis</i> H37Rv (ATCC 27294)	[198]
<i>Myristica fatua</i>	Myricaceae	Almond	Dichloromethane soluble extracts against <i>M. bovis</i> BCG (strain 11-73 P2) with MIC of 50 μ g/ml	[186]
<i>Nasturtium officinale</i>	Cruciferae	Aerial parts	Chloroform extracts against two drug-resistant strains of <i>M. tuberculosis</i> with MIC of 50-100 μ M	[204]
<i>Olea europaea</i>	Oleaceae	Leaf	Hexane extracts against the drug-resistant variants of <i>M. tuberculosis</i> with MIC of 25-100 μ M	[204]
<i>Otostegia integrifolia</i>	Lamiaceae	Root	Chloroform extract of roots was the most active on <i>M. tuberculosis</i> H37Rv (MIC 156 μ g/ml) and AOZ8W-4 (MDR-TB clinical isolate) (MIC 0.078 mg/ml)	[229]
<i>Pelargonium graveolens</i>	Geraniaceae	Seed	Hydrodistillation for essential oil against tested isolates ranged from 19.5 μ g/ml to 78 μ g/ml	[230]
<i>Pelargonium sidoides</i>	Geraniaceae	Root	Aqueous extracts inhibiting the growth of <i>M. tuberculosis</i> H37Rv (ATCC 27294) by 96% at a sample concentration of 12.5 μ g/ml	[231]
<i>Pentanisia prunelloides</i>	Rubiaceae	Root	80% ethanol against <i>M. tuberculosis</i> H37Ra (ATCC 25177) with MIC of 0.78 mg/ml	[215]
<i>Persea americana</i>	Lauraceae	Leaf, seed	Methanolic extracts against <i>M. tuberculosis</i> H37Ra with MIC of 31.2 μ g/ml and H37Rv; chloroformic extract of seeds against <i>M. tuberculosis</i> H37Rv MIC less than 50 μ g/ml	[229, 232]
<i>Phymaspermum acerosum</i>	Asteraceae	Root, leaf	Ethanol and water extracts had the best MIC value of 20 μ g/ml against five <i>M. tuberculosis</i> strains	[221]
<i>Piper cernuum</i>	Piperaceae	Leaf	Hydrodistillation with water displayed moderate activity against the <i>M. tuberculosis</i> H37Rv with MIC of 125 μ g/ml	[233]
<i>Piper diospyrifolium</i>	Piperaceae	Leaf	Hydrodistillation with water displayed moderate activity against the <i>M. tuberculosis</i> H37Rv with MIC of 125 μ g/ml	[233]
<i>Piper guineense</i>	Piperaceae	Seed	Methanol extracts against <i>M. tuberculosis</i> H37Ra and H37Rv with MIC of 256 μ g/ml	[210]
<i>Piper imperiale</i>	Piperaceae	Flower	Ethanol extracts against <i>M. tuberculosis</i> H37Rv with MIC of 75 μ g/ml	[234]
<i>Piper rivinoides</i>	Piperaceae	Leaf	Hydrodistillation with water displayed moderate activity against the <i>M. tuberculosis</i> H37Rv with MIC of 125 μ g/ml	[233]
<i>Piper sarmentosum</i>	Piperaceae	Leaf	Extracts with petroleum ether, chloroform, and methanol, against <i>M. tuberculosis</i> H37Rv with MIC of 25, 25, and 12.5 μ g/ml	[235]
<i>Pisonia borinquena</i>	Nyctaginaceae	Leaf	95% ethanol extracts against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with 85% inhibition at 100 μ g/ml	[216]
<i>Pittosporum tenuifolium</i>	Pittosporaceae	Leaf	Ethanol extracts against <i>M. smegmatis</i> with IC ₅₀ of 0.78 mg/ml	[225]
<i>Pluchea indica</i>	Asteraceae	Flower and leaf	80% methanol extract against <i>M. tuberculosis</i> H37Rv ATCC 25618 with an MIC of 800 μ g/ml	[191]
<i>Plumbago zeylanica</i>	Plumbaginaceae	Root	Ethanol extract against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 31.25 μ g/ml	[196]
<i>Psychotria zombamontana</i>	Rubiaceae	Leaf	Acetone extract decreased 256-fold of MIC in combination with rifampicin against <i>M. aurum</i> and reduction of the MICs of the anti-TB drug ranged from 2-fold to 4-fold, 2-fold to 64-fold, and 2-fold to 64-fold for <i>M. smegmatis</i> , <i>M. aurum</i> , and <i>M. tuberculosis</i> , respectively	[207]
<i>Pterocarpus osun</i>	Fabaceae (Leguminosae)	Stem	Chloroform extract against <i>M. tuberculosis</i> H37Rv and <i>M. bovis</i> BCG with MIC of 1225 μ g/ml and 1100 μ g/ml, respectively, by MABA	[67, 236]
<i>Pterolobium stellatum</i>	Fabaceae	Root	Chloroform extracts of roots were the most active on <i>M. tuberculosis</i> H37Rv (MIC 39 μ g/ml) and AOZ8W-4 (MDR-TB clinical isolate) (MIC 0.078 mg/ml)	[229]
<i>Rhynchosia precatória</i>	Fabaceae	Root	Extracts by n-hexane, dichloromethane, ethyl acetate, and methanol individually against <i>M. tuberculosis</i> H37Rv with MIC of 15.6-125 μ g/ml	[212]
<i>Ricinus communis</i>	Euphorbiaceae	Seed	Hexane extracts against <i>M. tuberculosis</i> H37Rv sensitive strain with MIC of 2.5 mg/ml by MABA	[203]

TABLE 2: Continued.

Plant species	Plant family	Part used	Extracts	References
<i>Rosmarinus officinalis</i>	Lamiaceae	Leaf	Ethanol extracts against <i>M. tuberculosis</i> H37Ra with MIC of 6.25 µg/ml	[224]
<i>Satureja aintabensis</i>	Lamiaceae	Aerial parts	Extraction with petroleum ether, ethyl acetate, and methanol killed <i>M. tuberculosis</i> with MIC of 50-800 µg/ml	[237]
<i>Satyrium nepalense</i>	Orchidaceae	Flower	Hexane extracts against <i>M. tuberculosis</i> H37Rv TMC-102 with MIC of 15.7 µg/ml	[238]
<i>Schinus molle</i>	Anacardiaceae	Fruit	Methanol extract against a drug-resistant strain of <i>M. tuberculosis</i> CIBIN/UMF15:99 with MIC of 125 µg/ml	[202]
<i>Securidaca longepedunculata</i>	Polygalaceae	Root	Hexane extracts against <i>M. bovis</i> BCG, <i>M. tuberculosis</i> H37Ra, and H37Rv with 62.5 µg/ml	[226]
<i>Solanum torvum</i>	Solanaceae	Leaf	80% ethanol crude extracts against <i>M. tuberculosis</i> H37Ra (ATCC 25177 TM) with MIC of 156.3 µg/ml	[184]
<i>Sphaeranthus indicus</i>	Asteraceae	Floral head	Ethanol extract against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 31.25 µg/ml	[196]
<i>Sterculia setigera</i>	Sterculiaceae	Leaf	Hexane, dichloromethane, and ethyl acetate extracts against <i>M. tuberculosis</i> H37Rv with MICs of 84 µg/ml, 62 µg/ml, and 128 µg/ml, respectively	[239]
<i>Swinglea glutinosa</i>	Rutaceae	Fruit peel	Aqueous extracts for essential oils against <i>M. tuberculosis</i> H37Rv (ATCC 27294) with MIC of 100 µg/ml	[182]
<i>Tabernaemontana elegans</i>	Apocynaceae	Root	Ethyl acetate extracts against <i>M. tuberculosis</i> H37Rv with MIC of 15.6 µg/ml	[226]
<i>Tabernaemontana coronaria</i>	Apocynaceae	Leaf	Hexane partition from methanol extracts of MIC of 100 µg/ml against <i>M. tuberculosis</i> H37Rv	[206]
<i>Terminalia phanerophlebia</i>	Combretaceae	Leaf, root, twig	80% ethanol against <i>M. tuberculosis</i> H37Ra (ATCC 25177) with 0.30 and 0.78 mg/ml, respectively	[215]
<i>Terminalia sericea</i>	Combretaceae	Stem bark	Acetone extracts against <i>M. tuberculosis</i> H37Ra with MIC of 25 µg/ml	[197]
<i>Thymus sibthorpii</i>	Lamiaceae	Aerial parts	Extracts with petroleum ether, ethyl acetate, and methanol against <i>M. tuberculosis</i> with MIC of 50–800 µg/ml	[237]
<i>Trachyspermum copticum</i>	Apiaceae	Aerial parts	Hydrodistillation extracts against <i>M. kansasii</i> and MDR-TB with MICs of 78 µg/ml	[230]
<i>Urtica dioica</i>	Urticaceae	Leaf	Hexane extracts against <i>M. smegmatis</i> with MIC of 250 µg/ml	[201]
<i>Uvaria rufa</i>	Annonaceae	Leaf	Lead acetate-treated crude chloroform extracts against <i>M. tuberculosis</i> H37Rv with MIC of 8 µg/ml	[136]
<i>Vetiveria zizanioides</i>	Poaceae	Root	Ethanol extract and hexane fraction 500 µg/ml or 50 µg/ml against <i>M. tuberculosis</i> H37Rv and H37Ra	[240]
<i>Vismia baccifera</i>	Clusiaceae	Leaf	Dichloromethane-methanol extracts (1:1) inhibit the <i>M. tuberculosis</i> H37Rv at 50 µg/ml with 70.3 ± 0.5%	[188]
<i>Xylopi aethiopica</i>	Annonaceae	Fruit, bark	Methanol extracts against <i>M. tuberculosis</i> H37Ra and H37Rv with MIC of 512 µg/ml	[210]
<i>Zanthoxylum capense</i>	Rutaceae	Root	Dichloromethane extracts against <i>M. bovis</i> BCG, <i>M. tuberculosis</i> H37Ra, and H37Rv with MICs of 31.2 µg/ml	[226]
<i>Zingiber officinale</i>	Zingiberaceae	Rhizome	Ethanol extract against <i>M. tuberculosis</i> H37Ra with MIC of 2500 µg/ml by MABA	[184]

^aMinimum inhibitory concentration (MIC); ^bmicroplate alamar blue assay (MABA); ^chalf maximal inhibitory concentration (C₅₀).

extracting solvents, while chloroform, dichloromethane, acetone, and ethyl acetate were used to a lesser extent. Of course, aqueous extract method was also popularly used, which involved the process of soaking, boiling, or/and hydrodistilling. Although the plant part used for study does not determine the extraction method, as a general rule, low molecule organic solvents are recommended when there is no reference.

To date, no specific cut-off value has been established as a reference to analyze the anti-TB activity results of the plant extracts, and many different methods are available to evaluate the activity. As of date, only a few anti-TB plant extracts in Table 2 have been tested in preclinical or clinical trials. It is encouraging that more and more promising crude extracts are now paving a way for the clinical test for their therapeutic applications. This section can provide a new perspective

TABLE 3: Important anti-TB traditional medicinal plants in literature by the systemic survey on the prescribed formula.

Species number	Family number	Main families	Country or region	References
13	10	Asteraceae (3), Chrysobalanaceae, Araliaceae, Acanthaceae, Chrysobalanaceae, Cucurbitaceae, Fabaceae, Lamiaceae, Melastomataceae, Phyllanthaceae, Polygonaceae	Burundian	[241]
9	8	Apocynaceae, Verbenaceae, Rubiaceae, Goodeniaceae, Agavaceae, Moraceae, Myrtaceae, Zingiberaceae	Manus Province, Papua New Guinea	[242]
184	77	Fabaceae (21), Asteraceae (12), Malvaceae (11)	Bapedi (South Africa)	[243]
30	21	Alliaceae (3), Rutaceae (3), Apiaceae (2), Caryophyllaceae (2), Asteraceae (2), Lamiaceae (2), Myrtaceae (2), Solanaceae (2)	Nkonkobe municipality, Eastern Cape Province (South Africa)	[244]
21	12	Asteraceae, Fabaceae, Geraniaceae	Mangaung metro, Thabo Mofutsanyana, and Lejweleputswa in South Africa	[245]
25	14	Fabaceae (5), Euphorbiaceae (3), Asteraceae (3), Lamiaceae (12%)	Bas-Congo Province, Democratic Republic of Congo	[208]
15	13	Amaryllidaceae (3), Xanthorrhoeaceae (2), Arecaceae (2), Solanaceae (2), Meliaceae, Acanthaceae, Poaceae, Phyllanthaceae, Melastomataceae, Poaceae, Cyperaceae, Zingiberaceae, Amaranthaceae, Asteraceae	Greater Accra and eastern communities in Ghana	[246]
95	48	Loranthaceae (6), Caesalpiniaceae (5), Papilionaceae (5), Poaceae (4), Mimosaceae (4), Scrophulariaceae (4), Anacardiaceae (3), Combretaceae (3), Liliaceae (3), and Solanaceae (3)	Niger state, Nigeria	[247]
66	35	Rutaceae (7), Euphorbiaceae (5), Rubiaceae (4), Anacardiaceae (3), Fabaceae (3), Verbenaceae (3), Arecaceae (2), Annonaceae (2), Solanaceae (2), Moraceae (2), Rhamnaceae (2)	Lao PDR	[248]
23	20	Arecaceae (2), Aristolochiaceae (2), Rubiaceae (2)	Malaysia	[249]
181	67	Asteraceae (31), Fabaceae (14), Lamiaceae (9), Euphorbiaceae (7), Celastraceae (5)	South Africa	[250]
62	38	Asteraceae (12), Aristolochiaceae (3), Compositae (3), Rosaceae (3), Juglandaceae (2), Zygophyllaceae (2), Verbenaceae (2), Rutaceae (2), Papaveraceae (2), Fabaceae (2)	Mexico	[10]
88	36	Lamiaceae (9), Asteraceae (7), Papilionaceae (4), Acanthaceae (3), Caesalpiniaceae (3), Capparaceae (3), Euphorbiaceae (3), Mimosaceae (3)	Districts of Kamuli, Kisoro, and Nakapiripirit in Uganda	[251]
90	44	Fabaceae (13), Asteraceae (7), Moraceae (5), Rutaceae (4)	Districts of Mpigi and Butambala, Uganda	[252]
35	22	Fabaceae (5), Rutaceae (4), Apocynaceae (3), Menispermaceae (3), Malvaceae (3)	Madhya Pradesh, India	[196]
132	45	Annonaceae (14), Zingiberaceae (12), Rutaceae (10), Annonaceae (10), Asteraceae (8), Euphorbiaceae (8), Fabaceae (7)	Southeast Asian	[253]
10	8	Fabaceae (3), Canellaceae (1), Rubiaceae (1), Anacardiaceae (1), Rutaceae (1), Myrtaceae (1), Merlucciidae (1), Guttiferae (1)	Lake Victoria Basin (Uganda, Kenya, and Tanzania)	[254]
14	8	Euphorbiaceae (4), Verbenaceae (3), Rutaceae (2)	Lake Victoria region and the Samburu community	[255]
2	2	Achillea millefolium (1), Dryopteris stewartii (1)	Kel village, Neelum Valley, Azad Kashmir, Pakistan	[256]
4	3	Amaryllidaceae (1), Lauraceae (1), Amaranthaceae (1), Asteraceae (1)	Sulaymaniyah Province, Kurdistan, Iraq	[257]
22	18	Liliaceae (3), Euphorbiaceae (2), Verbenaceae (2)	India	[258]
6	6	Vitaceae (1), Poaceae (1), Pinaceae (1), Musaceae (1), Rosaceae (1), Leguminosae (1)	Arabian Peninsula	[259]
2	2	Asteraceae (1), Dryopteridaceae (1)	Pakistan	[251]
70	44	Arecaceae (4), Euphorbiaceae (4), Fabaceae (3), Piperaceae (3), Rutaceae (3)	Malaysia	[191]

TABLE 3: Continued.

Species number	Family number	Main families	Country or region	References
56	26	Ranunculaceae (8), Liliaceae (6), Asteraceae (5), Umbellifera (4), Campanulaceae (3), Rubiaceae (4), Rosaceae (2), Scrophulariaceae (2), Labiatae (2), Fabaceae (2)	China	[260]
82	30	Combretaceae (13), Fabaceae (12), Moraceae (6), Rubiaceae (5), Polygalaceae (5)	Nigeria	[261]

in expanding the anti-TB plant species for the development of anti-TB medicine in the future.

5. Medicinal Plants Only Found in Formula Prescribed by the Traditional Healers

Traditional healers continually serve the public health in most of the countries. Some ethnomedical information has been published based on many plant species in anti-TB formulas documented by different traditional healing systems, ranging from the poor documented oral African medicine to the well-documented Ayurveda or Chinese medicine. These reports inspired the scientists to find more effective compounds for tuberculosis. The investigations of medicinal plants using frontier technologies are now being reconsidered to be a feasible approach for discovering novel bioactive compounds and crude extracts, in order to solve the wide spreaded TB problems. Table 3 shows the main species or families of the traditional plant medicines and their botanical details in the published papers by the systemic survey of the prescribed formula, but very few studies reveal the working compounds or active crude extracts.

The anti-TB formulas were investigated in many countries or regions, such as China, India, Mexico, South Africa, Pakistan, Iraq, Malaysia, Congo, Lao PDR, Nigeria, Nigeria, Burundi, Papua New Guinea, Lake Victoria Basin (Uganda, Kenya, and Tanzania), Arabian Peninsula, Southeast Asian, and Manus Province. Most anti-TB formulas were found during the ethnopharmacological investigation of the indigenous plants. To strengthen the anti-TB purpose, the present review summarized the anti-TB plant in the formula from the ethnopharmacological publications, with an emphasis on their classification (Table 3). Although three reports in Table 3 did not show the botanical family of the anti-TB medical plants in detail, it still can be speculated that the family *Fabaceae* is the most represented species, followed by *Asteraceae*, *Euphorbiaceae*, and *Liliaceae* families.

We need to be aware that the plant medicine used by the traditional healers is based on their according ethnomedical traditional medical theories. In comparison with the western system of medicine, the traditional plant medicines showed certain drawbacks. An important issue of all was that the active components of the herbal drugs prescribed were unknown. The activity of the traditional herbal drugs prescribed by the traditional healers can be greatly affected by the difference in the processing methods, variation in the potency due to difference in plant species and subspecies, varying geographical location of growth, nonuniform quality

control standards, etc. Furthermore, for a given plant, a specific place, part, method, and time for collection can significantly affect the therapeutic efficacy [16, 40]. Hence, the plant medicine used by the traditional healers needs a critical evaluation to find the active components.

6. Need for Future Research

Although this review presents a big list of plants with effective anti-TB activities from different traditional medicine systems, there is a need of better therapeutic drug monitoring systems and high throughput *in vitro* assays for the routine screening to identify potentially serious and clinically significant herb-drug interactions [262]. Furthermore, there is a lack of *in vivo* information regarding the drug metabolism associated interactions with reference to the traditional medicines and the treatment of TB. This requires health care practitioners to ensure a clear communication with patients regarding the possible negative impacts of simultaneous use of certain TMs and prescribed drugs. It was reported that the widely used *Sutherlandia frutescens* in the treatment of TB in countries of Southern African Development Community interfered with the isoniazid therapy, but the mechanism of this interaction was not clear [263, 264]. Coadministration also resulted in the reduced bioavailability of ofloxacin [265], while piperine showed the ability to increase the bioavailability of the antituberculosis drug rifampicin [266–268].

Several issues affect the anti-TB activity of the components of the medicinal plants, such as the variation in the potency due to difference in species, absence of an integrated coding for every species used commonly in TMs, varying geographical location of growth and incorrect identification of drugs, and nonuniform quality control standards [269]. No clinic trial was reported on the crude extracts, and even the pure compounds from the medical plants still need to be elucidated for their constituent characterization and the mechanism of action. Till date, not many compounds have been originated from plants for further modification for use in clinic. We hope that this review will help to find a possible way to get better anti-TB results by making a combination of the compounds originated from plants based on the different TB-killing mechanisms.

Abbreviations

TB: Tuberculosis
TM: Traditional medicines

WHO: World Health Organization
 DTM: Department of Traditional Medicine
 ICTM: International Classification of Traditional Medicine
 ICD-11: International Classification of Diseases—11
 TCM: Traditional Chinese Medicine
 TAM: Traditional African medicine
 XDR: Emergence of extensively drug-resistant
 TDR: Total drug-resistant
 MIC: Minimum inhibitory concentration
 MBC: Minimum bactericidal concentration
 MIA: Minimum inhibitory amount
 IC₅₀: Half maximal inhibitory concentration
 MABA: Microplate alamar blue assay.

Data Availability

All data included in this article are available from the corresponding author upon request.

Ethical Approval

Ethical approval is not applicable.

Consent

Consent was not necessary.

Conflicts of Interest

The authors declared that there are no conflicts of interest.

Authors' Contributions

ZS and YX designed and organized the review. CK drafted the manuscript. BL revised the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We appreciated Prof. Chunhua Qiao for giving the good suggestions on drawing the compound structures. This manuscript was supported by the Beijing Municipal Science & Technology Commission No. Z181100001718181, the National Natural Science Foundation (81871691), and the Joint Key Program of Beijing Municipal Education Commission and Beijing Municipal Natural Science Foundation Committee (21JG0034).

References

- [1] WHO, *Traditional medicine strategy 2014-2023*, WHO Press, Geneva, Switzerland, 2013, https://www.who.int/medicines/publications/traditional/trm_strategy14_23/en/.
- [2] WHO, "World Health Assembly Update," 2019, <https://www.who.int/news-room/detail/25-05-2019-world-health-assembly-update>.
- [3] A. T. Pezzella, "History of pulmonary tuberculosis," *Thoracic Surgery Clinics*, vol. 29, no. 1, pp. 1-17, 2019.
- [4] T. Wirth, F. Hildebrand, C. Allix-Béguec et al., "Origin, spread and demography of the *Mycobacterium tuberculosis* complex," *PLoS Pathogens*, vol. 4, no. 9, article e1000160, 2008.
- [5] I. Comas, M. Coscolla, T. Luo et al., "Out-of-Africa migration and Neolithic coexpansion of *Mycobacterium tuberculosis* with modern humans," *Nature Genetics*, vol. 45, no. 10, pp. 1176-1182, 2013.
- [6] WHO, "Global tuberculosis report 2020," 2020, <https://www.who.int/publications/i/item/9789240013131>.
- [7] World Health Organization, *Global tuberculosis report 2019*, World Health Organization, 2019, <https://apps.who.int/iris/handle/10665/329368>.
- [8] A. H. Uc-Cachón, R. Borges-Argáez, S. Said-Fernández et al., "Naphthoquinones isolated from *Diospyros anisandra* exhibit potent activity against pan-resistant first-line drugs *Mycobacterium tuberculosis* strains," *Pulmonary Pharmacology & Therapeutics*, vol. 27, no. 1, pp. 114-120, 2014.
- [9] B. J. Enquist, X. Feng, B. Boyle et al., "The commonness of rarity: global and future distribution of rarity across land plants," *Science Advances*, vol. 5, no. 11, article eaaz0414, 2019.
- [10] S. Chen, X. Pang, J. Song et al., "A renaissance in herbal medicine identification: From morphology to DNA," *Biotechnology Advances*, vol. 32, no. 7, pp. 1237-1244, 2014.
- [11] R. Gómez-Cansino, S. L. Guzmán-Gutiérrez, M. G. Campos-Lara, C. I. Espitia-Pinzón, and R. Reyes-Chilpa, "Natural compounds from Mexican medicinal plants as potential drug leads for anti-tuberculosis drugs," *Anais da Academia Brasileira de Ciências*, vol. 89, no. 1, pp. 31-43, 2017.
- [12] R. Patel, T. Jawaid, P. Gautam, and P. Dwivedi, "Herbal remedies for gastroprotective action: a review," *International Journal of Phytopharmacy*, vol. 2, no. 2, pp. 64-67, 2012.
- [13] J. L. Ríos and M. C. Recio, "Medicinal plants and antimicrobial activity," *Journal of Ethnopharmacology*, vol. 100, no. 1-2, pp. 80-84, 2005.
- [14] D. S. Fabricant and N. R. Farnsworth, "The value of plants used in traditional medicine for drug discovery," *Environmental Health Perspectives*, vol. 109, Supplement 1, pp. 69-75, 2001.
- [15] B. R. Copp and A. N. Pearce, "Natural product growth inhibitors of *Mycobacterium tuberculosis*," *Natural Product Reports*, vol. 24, no. 2, pp. 278-297, 2007.
- [16] M. Abedinzadeh, M. Gaeini, and S. Sardari, "Natural antimicrobial peptides against *Mycobacterium tuberculosis*," *Journal of Antimicrobial Chemotherapy*, vol. 70, no. 5, pp. 1285-1289, 2015.
- [17] K. C. Chinsebu, "Tuberculosis and nature's pharmacy of putative anti-tuberculosis agents," *Acta Tropica*, vol. 153, pp. 46-56, 2016.
- [18] C. Limmatvapirat, S. Sirisopanaporn, and P. Kittakoop, "Antitubercular and antiparasitic constituents of *Abrus precatorius*," *Planta Medica*, vol. 70, no. 3, pp. 276-278, 2004.
- [19] E. Hernández-García, A. García, E. Garza-González et al., "Chemical composition of *Acacia farnesiana* (L.) wild fruits and its activity against *Mycobacterium tuberculosis* and dysentery bacteria," *Journal of Ethnopharmacology*, vol. 230, pp. 74-80, 2019.
- [20] N. Joycharat, H. Greger, O. Hofer, and E. Saifah, "Flavaglines and triterpenoids from the leaves of *Aglaia forbesii*," *Phytochemistry*, vol. 69, no. 1, pp. 206-211, 2008.

- [21] V. Kuete, A. G. B. Azebaze, A. T. Mbaveng et al., "Antioxidant, antitumor and antimicrobial activities of the crude extract and compounds of the root bark of *Allanblackia floribunda*," *Pharmaceutical Biology*, vol. 49, no. 1, pp. 57–65, 2011.
- [22] G. O'Donnell and S. Gibbons, "Antibacterial activity of two canthin-6-one alkaloids from *Allium neapolitanum*," *Phytotherapy Research*, vol. 21, no. 7, pp. 653–657, 2007.
- [23] R. Muniyan and J. Gurunathan, "Lauric acid and myristic acid from *Allium sativum* inhibit the growth of *Mycobacterium tuberculosis* H37Ra: in silico analysis reveals possible binding to protein kinase B," *Pharmaceutical Biology*, vol. 54, no. 12, pp. 2814–2821, 2016.
- [24] L. N. F. Trevizan, K. F. Nascimento, J. A. Santos et al., "Anti-inflammatory, antioxidant and anti-*Mycobacterium tuberculosis* activity of viridiflorol: The major constituent of *Allophylus edulis* (A. St.-Hil., A. Juss. & Cambess.) Radlk.," *Journal of Ethnopharmacology*, vol. 192, pp. 510–515, 2016.
- [25] H. Li, D. Webster, J. A. Johnson, and C. A. Gray, "Anti-mycobacterial triterpenes from the Canadian medicinal plant *Alnus incana*," *Journal of Ethnopharmacology*, vol. 165, pp. 148–151, 2015.
- [26] B. Gröblacher, O. Kunert, and F. Bucar, "Compounds of *Alpinia katsumadai* as potential efflux inhibitors in *Mycobacterium smegmatis*," *Bioorganic & Medicinal Chemistry*, vol. 20, no. 8, pp. 2701–2706, 2012.
- [27] I. Rivero-Cruz, L. Acevedo, J. A. Guerrero et al., "Antimycobacterial agents from selected Mexican medicinal plants," *The Journal of Pharmacy and Pharmacology*, vol. 57, no. 9, pp. 1117–1126, 2005.
- [28] N. Rastogi, J. Abaul, K. S. Goh, A. Devallois, E. Philogène, and P. Bourgeois, "Antimycobacterial activity of chemically defined natural substances from the Caribbean flora in Guadeloupe," *FEMS Immunology and Medical Microbiology*, vol. 20, no. 4, pp. 267–273, 1998.
- [29] J. Zhou, R. Xu, X. Z. du, X. D. Zhou, and Q. Li, "Saxifragifolin D attenuates phagosome maturation arrest in *Mycobacterium tuberculosis*-infected macrophages via an AMPK and VPS34-dependent pathway," *AMB Express*, vol. 7, no. 1, p. 11, 2017.
- [30] S. Deng, Y. Wang, T. Inui et al., "Anti-TB polyynes from the roots of *Angelica sinensis*," *Phytotherapy Research*, vol. 22, no. 7, pp. 878–882, 2008.
- [31] R. Lekphrom, S. Kanokmedhakul, and K. Kanokmedhakul, "Bioactive diterpenes from the aerial parts of *Anisochilus harmandii*," *Planta Medica*, vol. 76, no. 7, pp. 726–728, 2010.
- [32] H. Li, T. O'Neill, D. Webster, J. A. Johnson, and C. A. Gray, "Anti-mycobacterial diynes from the Canadian medicinal plant *Aralia nudicaulis*," *Journal of Ethnopharmacology*, vol. 140, no. 1, pp. 141–144, 2012.
- [33] Y. F. Guan, X. Song, M. H. Qiu et al., "Bioassay-guided isolation and structural modification of the anti-TB resorcinols from *Ardisia gigantifolia*," *Chemical Biology & Drug Design*, vol. 88, no. 2, pp. 293–301, 2016.
- [34] P. V. Habbu, K. M. Mahadevan, R. A. Shastry, and H. Manjunatha, "Antimicrobial activity of flavanoid sulphates and other fractions of *Argyrea speciosa* (Burm.f) Boj," *Indian Journal of Experimental Biology*, vol. 47, no. 2, pp. 121–128, 2009.
- [35] C. H. Jiang, M. L. Gan, T. T. An, and Z. C. Yang, "Bioassay-guided isolation of a *Mycobacterium tuberculosis* biofilm inhibitor from *Arisaema sinii* Krause," *Microbial Pathogenesis*, vol. 126, pp. 351–356, 2019.
- [36] V. M. Navarro-García, J. Luna-Herrera, M. G. Rojas-Briebesca, P. Álvarez-Fitz, and M. Y. Ríos, "Antibacterial activity of *Aristolochia brevipes* against multidrug-resistant *Mycobacterium tuberculosis*," *Molecules*, vol. 16, no. 9, pp. 7357–7364, 2011.
- [37] R. León-Díaz, M. Meckes, S. Said-Fernández et al., "Antimycobacterial neolignans isolated from *Aristolochia taliscana*," *Memórias do Instituto Oswaldo Cruz*, vol. 105, no. 1, pp. 45–51, 2010.
- [38] A. Sharma, R. C. Flores-Vallejo, A. Cardoso-Taketa, and M. L. Villarreal, "Antibacterial activities of medicinal plants used in Mexican traditional medicine," *Journal of Ethnopharmacology*, vol. 208, pp. 264–329, 2017.
- [39] M. Figueroa, I. Rivero-Cruz, B. Rivero-Cruz, R. Bye, A. Navarrete, and R. Mata, "Constituents, biological activities and quality control parameters of the crude extract and essential oil from *Arracacia toluensis* var. *multifida*," *Journal of Ethnopharmacology*, vol. 113, no. 1, pp. 125–131, 2007.
- [40] M. A. Jyoti, K. W. Nam, W. S. Jang et al., "Antimycobacterial activity of methanolic plant extract of *Artemisia capillaris* containing ursolic acid and hydroquinone against *Mycobacterium tuberculosis*," *Journal of Infection and Chemotherapy*, vol. 22, no. 4, pp. 200–208, 2016.
- [41] A. Puntumchai, P. Kittakoop, S. Rajviroongit, S. Vimuttipong, K. Likhitwitayawuid, and Y. Thebtaranonth, "Lakoochins A and B, new antimycobacterial stilbene derivatives from *Artocarpus lakoocha*," *Journal of Natural Products*, vol. 67, no. 3, pp. 485–486, 2004.
- [42] G. M. Molina-Salinas, J. Bórquez, A. Ardiles et al., "Antituberculosis activity of natural and semisynthetic azorellane and mulinane diterpenoids," *Fitoterapia*, vol. 81, no. 1, pp. 50–54, 2010.
- [43] G. M. Molina-Salinas, J. Bórquez, S. Said-Fernández et al., "Antituberculosis activity of alkylated mulinane diterpenoids," *Fitoterapia*, vol. 81, no. 3, pp. 219–222, 2010.
- [44] S. Boonphong, P. Puangsombat, A. Baramée, C. Mahidol, S. Ruchirawat, and P. Kittakoop, "Bioactive compounds from *Bauhinia purpurea* possessing antimalarial, antimycobacterial, antifungal, anti-inflammatory, and cytotoxic activities," *Journal of Natural Products*, vol. 70, no. 5, pp. 795–801, 2007.
- [45] P. Kittakoop, S. Nopichai, N. Thongon, P. Charoenchai, and Y. Thebtaranonth, "Bauhinoxepins A and B: new antimycobacterial Dibenzo[b,f]oxepins from *Bauhinia saccocalyx*," *Helvetica Chimica Acta*, vol. 87, no. 1, pp. 175–179, 2004.
- [46] P. S. Yang, M. J. Cheng, C. F. Peng, J. J. Chen, and I. S. Chen, "Endiandric acid analogues from the roots of *Beilschmiedia erythrophloia*," *Journal of Natural Products*, vol. 72, no. 1, pp. 53–58, 2009.
- [47] J. J. Chen, E. T. Chou, C. F. Peng, I. S. Chen, S. Z. Yang, and H. Y. Huang, "Novel epoxyfuranoid lignans and antitubercular constituents from the leaves of *Beilschmiedia tsangii*," *Planta Medica*, vol. 73, no. 6, pp. 567–571, 2007.
- [48] J. C. Aponte, Y. Estevez, R. Gilman et al., "Anti-infective and cytotoxic compounds present in *Blepharodon nitidum*," *Planta Medica*, vol. 74, no. 4, pp. 407–410, 2008.
- [49] M. R. Camacho-Corona, J. M. J. Favela-Hernández, O. González-Santiago et al., "Evaluation of some plant-derived secondary metabolites against sensitive and

- multidrug-resistant mycobacterium tuberculosis,” *Journal of the Mexican Chemical Society*, vol. 53, no. 2, pp. 71–75, 2009.
- [50] N. Promsawan, P. Kittakoop, S. Boonphong, and P. Nongkunsarn, “Antitubercular cassane furanoditerpenoids from the roots of *Caesalpinia pulcherrima*,” *Planta Medica*, vol. 69, no. 8, pp. 776–777, 2003.
- [51] H. Seo, S. Kim, H. A. Mahmud et al., “In vitro antitubercular activity of 3-deoxysappanchalcone isolated from the heartwood of *Caesalpinia sappan* Linn,” *Phytotherapy Research*, vol. 31, no. 10, pp. 1600–1606, 2017.
- [52] R. Encarnación-Dimayuga, J. Agúndez-Espinoza, A. García, G. Delgado, G. Molina-Salinas, and S. Said-Fernández, “Two new cassane-type diterpenes from *Calliandra californica* with antituberculosis and cytotoxic activities,” *Planta Medica*, vol. 72, no. 8, pp. 757–761, 2006.
- [53] J. J. Chen, H. M. Wu, C. F. Peng, I. S. Chen, and S. D. Chu, “seco-Abietane diterpenoids, a phenylethanoid derivative, and antitubercular constituents from *Callicarpa pilosissima*,” *Journal of Natural Products*, vol. 72, no. 2, pp. 223–228, 2009.
- [54] Z. Q. Xu, W. W. Barrow, W. J. Suling et al., “Anti-HIV natural product (+)-calanolide A is active against both drug-susceptible and drug-resistant strains of *Mycobacterium tuberculosis*,” *Bioorganic & Medicinal Chemistry*, vol. 12, no. 5, pp. 1199–1207, 2004.
- [55] P. Zheng, S. Somersan-Karakaya, S. Lu et al., “Synthetic calanolides with bactericidal activity against replicating and non-replicating *Mycobacterium tuberculosis*,” *Journal of Medicinal Chemistry*, vol. 57, no. 9, pp. 3755–3772, 2014.
- [56] N. Vongvanich, P. Kittakoop, P. Charoenchai, S. Intamas, K. Sriklung, and Y. Thebtaranonth, “Antiplasmodial, antimycobacterial, and Cytotoxic principles from *Camchaya calcarea*,” *Planta Medica*, vol. 72, no. 15, pp. 1427–1430, 2006.
- [57] D. Torres-Romero, I. A. Jiménez, R. Rojas, R. H. Gilman, M. López, and I. L. Bazzocchi, “Dihydro- β -agarofuran sesquiterpenes isolated from *Celastrus vulcanicola* as potential anti-*Mycobacterium tuberculosis* multidrug-resistant agents,” *Bioorganic & Medicinal Chemistry*, vol. 19, no. 7, pp. 2182–2189, 2011.
- [58] A. Jimenez-Arellanes, M. Meckes, R. Ramirez, J. Torres, and J. Luna-Herrera, “Activity against multidrug-resistant *Mycobacterium tuberculosis* in Mexican plants used to treat respiratory diseases,” *Phytotherapy Research*, vol. 17, no. 8, pp. 903–908, 2003.
- [59] A. Jiménez-Arellanes, J. Luna-Herrera, J. Cornejo-Garrido et al., “Ursolic and oleanolic acids as antimicrobial and immunomodulatory compounds for tuberculosis treatment,” *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, p. 258, 2013.
- [60] T. Akihisa, S. G. Franzblau, M. Ukiya et al., “Antitubercular activity of triterpenoids from asteraceae flowers,” *Biological & Pharmaceutical Bulletin*, vol. 28, no. 1, pp. 158–160, 2005.
- [61] A. Mehta, G. Srivastva, S. Kachhwaha, M. Sharma, and S. L. Kothari, “Antimycobacterial activity of *Citrullus colocynthis* (L.) Schrad. against drug sensitive and drug resistant *Mycobacterium tuberculosis* and MOTT clinical isolates,” *Journal of Ethnopharmacology*, vol. 149, no. 1, pp. 195–200, 2013.
- [62] N. E. Sandoval-Montemayor, A. García, E. Elizondo-Treviño, E. Garza-González, L. Alvarez, and M. del Rayo Camacho-Corona, “Chemical composition of hexane extract of *Citrus aurantifolia* and anti-*Mycobacterium tuberculosis* activity of some of its constituents,” *Molecules*, vol. 17, no. 9, pp. 11173–11184, 2012.
- [63] A. Sunthitikawinsakul, N. Kongkathip, B. Kongkathip et al., “Coumarins and carbazoles from *Clausena excavata* exhibited antimycobacterial and antifungal activities,” *Planta Medica*, vol. 69, no. 2, pp. 155–157, 2003.
- [64] R. Rojas, L. Caviedes, J. C. Aponte et al., “Aegicerin, the first oleanane triterpene with wide-ranging antimycobacterial activity, isolated from *Clavija procera*,” *Journal of Natural Products*, vol. 69, no. 5, pp. 845–846, 2006.
- [65] P. Tuntiwachwuttikul, Y. Pootaeng-on, P. Pansa, T. Srisanpang, and W. C. Taylor, “Sulfur-containing compounds from *Clinacanthus siamensis*,” *Chemical & Pharmaceutical Bulletin (Tokyo)*, vol. 51, no. 12, pp. 1423–1425, 2003.
- [66] M. Z. Pérez-González, G. A. Gutiérrez-Rebolledo, L. Yépez-Mulia, I. S. Rojas-Tomé, J. Luna-Herrera, and M. A. Jiménez-Arellanes, “Antiprotozoal, antimycobacterial, and anti-inflammatory evaluation of *Cnidioscolus chayamansa* (Mc Vaugh) extract and the isolated compounds,” *Biomedicine & Pharmacotherapy*, vol. 89, pp. 89–97, 2017.
- [67] N. N. Ibekwe, J. B. Nvau, P. O. Oladosu et al., “Some Nigerian anti-tuberculosis ethnomedicines: a preliminary efficacy assessment,” *Journal of Ethnopharmacology*, vol. 155, no. 1, pp. 524–532, 2014.
- [68] K. Asres, F. Bucar, S. Edelsbrunner, T. Kartnig, G. Höger, and W. Thiel, “Investigations on antimycobacterial activity of some Ethiopian medicinal plants,” *Phytotherapy Research*, vol. 15, no. 4, pp. 323–326, 2001.
- [69] S. Dettrakul, S. Surerum, S. Rajviroongit, and P. Kittakoop, “Biomimetic transformation and biological activities of globiferin, a terpenoid benzoquinone from *Cordia globifera*,” *Journal of Natural Products*, vol. 72, no. 5, pp. 861–865, 2009.
- [70] J. Thongtan, P. Kittakoop, N. Ruangrunsi, J. Saenboonrueng, and Y. Thebtaranonth, “New antimycobacterial and antimalarial 8,9-Secokaurane diterpenes from *Crotonkongensis*,” *Journal of Natural Products*, vol. 66, no. 6, pp. 868–870, 2003.
- [71] D. K. Agrawal, D. Saikia, R. Tiwari et al., “Demethoxycurcumin and its semisynthetic analogues as antitubercular agents,” *Planta Medica*, vol. 74, no. 15, pp. 1828–1831, 2008.
- [72] P. R. Baldwin, A. Z. Reeves, K. R. Powell et al., “Monocarboxyl analogs of curcumin inhibit growth of antibiotic sensitive and resistant strains of *Mycobacterium tuberculosis*,” *European Journal of Medicinal Chemistry*, vol. 92, pp. 693–699, 2015.
- [73] V. O. Fadipe, N. I. Mongalo, A. R. Opoku, P. M. Dikhoba, and T. J. Makhafole, “Isolation of anti-mycobacterial compounds from *Curtisia dentata* (Burm.f.) C.A.Sm (Curtisiaceae),” *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 306, 2017.
- [74] K. W. Nam, W. S. Jang, M. A. Jyoti, S. Kim, B. E. Lee, and H. Y. Song, “In vitro activity of (-)-deoxypergularinine, on its own and in combination with anti-tubercular drugs, against resistant strains of *Mycobacterium tuberculosis*,” *Phytomedicine*, vol. 23, no. 5, pp. 578–582, 2016.
- [75] S. Koysomboon, I. van Altena, S. Kato, and K. Chantrapromma, “Antimycobacterial flavonoids from *Derris indica*,” *Phytochemistry*, vol. 67, no. 10, pp. 1034–1040, 2006.
- [76] P. C. Pan, M. J. Cheng, C. F. Peng, H. Y. Huang, J. J. Chen, and I. S. Chen, “Secondary metabolites from the roots of *Litsea hypophaea* and their antitubercular activity,” *Journal of Natural Products*, vol. 73, no. 5, pp. 890–896, 2010.

- [77] P. Nareeboon, W. Kraus, U. Beifuss, J. Conrad, I. Klaiber, and S. Sutthivaiyakit, "Novel 24-nor-, 24-nor-2,3- seco-, and 3,24-dinor-2,4-seco-ursane triterpenes from *Diospyros decandra*: evidences for ring A biosynthetic transformations," *Tetrahedron*, vol. 62, no. 23, pp. 5519–5526, 2006.
- [78] D. Dey, R. Ray, and B. Hazra, "Antitubercular and antibacterial activity of quinonoid natural products against multi-drug resistant clinical isolates," *Phytotherapy Research*, vol. 28, no. 7, pp. 1014–1021, 2014.
- [79] B. Hazra, M. Dassarma, and U. Sanyal, "Separation methods of quinonoid constituents of plants used in Oriental traditional medicines," *Journal of Chromatography. B, Analytical Technologies in the Biomedical and Life Sciences*, vol. 812, no. 1-2, pp. 259–275, 2008.
- [80] J. N. Evina, D. S. N. Bikobo, A. A. A. Zintchem et al., "In vitro antitubercular activity of extract and constituents from the stem bark of *Disthemonanthus benthamianus*," *Revista Brasileira de Farmacognosia*, vol. 27, no. 6, pp. 739–743, 2017.
- [81] R. J. Case, Y. Wang, S. G. Franzblau et al., "Advanced applications of counter-current chromatography in the isolation of anti-tuberculosis constituents from *Dracaena angustifolia*," *Journal of Chromatography*, vol. 1151, no. 1-2, pp. 169–174, 2007.
- [82] Y. C. Chien, C. H. Lin, M. Y. Chiang et al., "Secondary metabolites from the root of *Ehretia longiflora* and their biological activities," *Phytochemistry*, vol. 80, pp. 50–57, 2012.
- [83] W.-Y. Lin, C. F. Peng, I. L. Tsai, J. J. Chen, M. J. Cheng, and I. S. Chen, "Antitubercular constituents from the roots of *Engelhardia roxburghiana*," *Planta Medica*, vol. 71, no. 2, pp. 171–175, 2005.
- [84] C. C. Wu, C. F. Peng, I. L. Tsai, M. H. Abd el-Razek, H. S. Huang, and I. S. Chen, "Secondary metabolites from the roots of *Engelhardia roxburghiana* and their antitubercular activities," *Phytochemistry*, vol. 68, no. 9, pp. 1338–1343, 2007.
- [85] S. Sutthivaiyakit, W. Mongkolvisut, S. Prabpai, and P. Kongsaree, "Diterpenes, sesquiterpenes, and a sesquiterpene-coumarin conjugate from *Jatropha integririma*," *Journal of Natural Products*, vol. 72, no. 11, pp. 2024–2027, 2009.
- [86] A. K. Waffo, G. A. Azebaze, A. E. Nkengfack et al., "Indicanines B and C, two isoflavonoid derivatives from the root bark of *Erythrina indica*," *Phytochemistry*, vol. 53, no. 8, pp. 981–985, 2000.
- [87] T. O. Lawal, B. A. Adeniyi, A. O. Adegoke, S. G. Franzblau, and G. B. Mahady, "In vitro susceptibility of *Mycobacterium tuberculosis* to extracts of *Eucalyptus camaldulensis* and *Eucalyptus torelliana* and isolated compounds," *Pharmaceutical Biology*, vol. 50, no. 1, pp. 92–98, 2012.
- [88] N. B. Bapela, N. Lall, P. B. Fourie, S. G. Franzblau, and C. E. J. van Rensburg, "Activity of 7-methyljuglone in combination with antituberculous drugs against *Mycobacterium tuberculosis*," *Phytomedicine*, vol. 13, no. 9-10, pp. 630–635, 2006.
- [89] Z. Yu, Y. Wei, X. Tian et al., "Diterpenoids from the roots of *Euphorbia ebracteolata* and their anti-tuberculosis effects," *Bioorganic Chemistry*, vol. 77, pp. 471–477, 2018.
- [90] Y. Wei, C. Wang, Z. Cheng et al., "Heterodimeric diterpenoids isolated from euphorbia ebracteolata roots and their inhibitory effects on α -glucosidase," *Journal of Natural Products*, vol. 80, no. 12, pp. 3218–3223, 2017.
- [91] N. Duarte, M. J. U. Ferreira, M. Martins, M. Viveiros, and L. Amaral, "Antibacterial activity of ergosterol peroxide against *Mycobacterium tuberculosis*: dependence upon system and medium employed," *Phytotherapy Research*, vol. 21, no. 7, pp. 601–604, 2007.
- [92] M. Koch, T. Bugni, C. Pond et al., "Antimycobacterial activity of *Exocarpos latifolius* due to exocarpic acid," *Planta Medica*, vol. 75, no. 12, pp. 1326–1330, 2010.
- [93] C. C. Chiang, M. J. Cheng, C. F. Peng, H. Y. Huang, and I. S. Chen, "A novel dimeric coumarin analog and antimycobacterial constituents from *Fatoua pilosa*," *Chemistry & Biodiversity*, vol. 7, no. 7, pp. 1728–1736, 2010.
- [94] Z. Z. Ibraheim, W. M. Abdel-Mageed, H. Dai, H. Guo, L. Zhang, and M. Jaspars, "Antimicrobial antioxidant daucane sesquiterpenes from *Ferula hermonis* Boiss," *Phytotherapy Research*, vol. 26, no. 4, pp. 579–586, 2012.
- [95] A. A. Kaikabo and J. N. Eloff, "Antibacterial activity of two biflavonoids from *Garcinia livingstonei* leaves against *Mycobacterium smegmatis*," *Journal of Ethnopharmacology*, vol. 138, no. 1, pp. 253–255, 2011.
- [96] Y. M. Lin, M. T. Flavin, C. S. Cassidy, A. Mar, and F. C. Chen, "Biflavonoids as novel antituberculosis agents," *Bioorganic & Medicinal Chemistry Letters*, vol. 11, no. 16, pp. 2101–2104, 2001.
- [97] H. Fouotsa, A. T. Mbaveng, C. D. Mbazona et al., "Antibacterial constituents of three Cameroonian medicinal plants: *Garcinia nobilis*, *Orcia suaveolens* and *Balsamocitrus camerunensis*," *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, p. 81, 2013.
- [98] R. S. Lekphrom, S. Kanokmedhakul, and K. Kanokmedhakul, "Bioactive styryllactones and alkaloid from flowers of *Goniotalamus laoticus*," *Journal of Ethnopharmacology*, vol. 125, no. 1, pp. 47–50, 2009.
- [99] P. Tuntiwachwuttikul, P. Phansa, Y. Pootaeng-on, and W. C. Taylor, "Chromones from the branches of *Harrisonia perforata*," *Chemical & Pharmaceutical Bulletin (Tokyo)*, vol. 54, no. 1, pp. 44–47, 2006.
- [100] N. Lall, A. A. Hussein, and J. J. M. Meyer, "Antiviral and antituberculous activity of *Helichrysum melanacme* constituents," *Fitoterapia*, vol. 77, no. 3, pp. 230–232, 2006.
- [101] T. O'Neill, J. A. Johnson, D. Webster, and C. A. Gray, "The Canadian medicinal plant *Heracleum maximum* contains antimycobacterial diynes and furanocoumarins," *Journal of Ethnopharmacology*, vol. 147, no. 1, pp. 232–237, 2013.
- [102] M. Stavri, R. Schneider, G. O'Donnell, D. Lechner, F. Bucar, and S. Gibbons, "The antimycobacterial components of hops (*Humulus lupulus*) and their dereplication," *Phytotherapy Research*, vol. 18, no. 9, pp. 774–776, 2004.
- [103] J. F. Wang, H. Q. Dai, Y. L. Wei et al., "Antituberculosis agents and an inhibitor of the para-aminobenzoic acid biosynthetic pathway from *Hydnocarpus anthelmintica* seeds," *Chemistry & Biodiversity*, vol. 7, no. 8, pp. 2046–2053, 2010.
- [104] K. Osman, D. Evangelopoulos, C. Basavannacharya et al., "An antibacterial from *Hypericum acmosepalum* inhibits ATP-dependent MurE ligase from *Mycobacterium tuberculosis*," *International Journal of Antimicrobial Agents*, vol. 39, no. 2, pp. 124–129, 2012.
- [105] D. Thangadurai, M. B. Viswanathan, and N. Ramesh, "Indigoferabietone, a novel abietane diterpenoid from *Indigofera longeracemosa* with potential antituberculous and antibacterial activity," *Die Pharmazie*, vol. 57, no. 10, pp. 714–715, 2002.

- [106] G. A. Wächter, S. Valcic, S. G. Franzblau, E. Suarez, and B. N. Timmermann, "Antitubercular activity of triterpenoids from *Lippia turbinata*," *Journal of Natural Products*, vol. 64, no. 1, pp. 37–41, 2001.
- [107] C. D. Carpenter, T. O'Neill, N. Picot et al., "Anti-mycobacterial natural products from the Canadian medicinal plant *Juniperus communis*," *Journal of Ethnopharmacology*, vol. 143, no. 2, pp. 695–700, 2012.
- [108] J. S. Mossa, F. S. el-Ferally, and I. Muhammad, "Antimycobacterial constituents from *Juniperus procera*, *Ferula communis* and *Plumbago zeylanica* and their in vitro synergistic activity with isonicotinic acid hydrazide," *Phytotherapy Research*, vol. 18, no. 11, pp. 934–937, 2004.
- [109] D. Lakshmanan, J. Werngren, L. Jose et al., "Ethyl *p*-methoxycinnamate isolated from a traditional anti-tuberculosis medicinal herb inhibits drug resistant strains of *Mycobacterium tuberculosis* in vitro," *Fitoterapia*, vol. 82, no. 5, pp. 757–761, 2011.
- [110] C. Yenjai, K. Prasanphen, S. Daodee, V. Wongpanich, and P. Kittakoop, "Bioactive flavonoids from *Kaempferia parviflora*," *Fitoterapia*, vol. 75, no. 1, pp. 89–92, 2004.
- [111] A. Jiménez-Arellanes, M. Meckes, J. Torres, and J. Luna-Herrera, "Antimycobacterial triterpenoids from *Lantana hispida* (Verbenaceae)," *Journal of Ethnopharmacology*, vol. 111, no. 2, pp. 202–205, 2007.
- [112] J. D. Guzman, D. Evangelopoulos, A. Gupta, J. M. Prieto, S. Gibbons, and S. Bhakta, "Antimycobacterials from lovage root (*Ligusticum officinale* Koch)," *Phytotherapy Research*, vol. 27, no. 7, pp. 993–998, 2013.
- [113] A. Suksamrarn, P. Poomsing, N. Aroonrerk, T. Punjanon, S. Suksamrarn, and S. Kongkun, "Antimycobacterial and antioxidant flavones from *Limnophila geoffrayi*," *Archives of Pharmacol Research*, vol. 26, no. 10, pp. 816–820, 2003.
- [114] J. L. Nkoti, D. S. Ngono Bikobo, A. Abouem A Zintchem et al., "Antitubercular evaluation of root extract and isolated phytochemicals from *Lophira lanceolata* against two resistant strains of *Mycobacterium tuberculosis*," *Pharmaceutical Biology*, vol. 56, no. 1, pp. 318–324, 2018.
- [115] A. Aguinado, V. M. Dalangin-Mallari, A. P. G. Macabeo et al., "Quinoline alkaloids from *Lunasia amara* inhibit *Mycobacterium tuberculosis* H₃₇Rv in vitro," *International Journal of Antimicrobial Agents*, vol. 29, no. 6, pp. 744–746, 2007.
- [116] B. G. Elkington, K. Sydara, A. Newsome et al., "New finding of an anti-TB compound in the genus *Marsypopetalum* (Annonaceae) from a traditional herbal remedy of Laos," *Journal of Ethnopharmacology*, vol. 151, no. 2, pp. 903–911, 2014.
- [117] C. Ma, R. J. Case, Y. Wang et al., "Anti-tuberculosis constituents from the stem bark of *Micromelum hirsutum*," *Planta Medica*, vol. 71, no. 3, pp. 261–267, 2005.
- [118] J. J. Chen, T. H. Chou, C. F. Peng, I. S. Chen, and S. Z. Yang, "Antitubercular dihydroagarofuranoid sesquiterpenes from the roots of *Microtropis fokienensis*," *Journal of Natural Products*, vol. 70, no. 2, pp. 202–205, 2007.
- [119] T. H. Chou, I. S. Chen, C. F. Peng, P. J. Sung, and J. J. Chen, "A new dihydroagarofuranoid sesquiterpene and antituberculosis constituents from the root of *Microtropis japonica*," *Chemistry & Biodiversity*, vol. 5, no. 7, pp. 1412–1418, 2008.
- [120] J. J. Chen, C. S. Yang, C. F. Peng, I. S. Chen, and C. L. Miaw, "Dihydroagarofuranoid sesquiterpenes, a lignan derivative, a benzenoid, and antitubercular constituents from the stem of *Microtropis japonica*," *Journal of Natural Products*, vol. 71, no. 6, pp. 1016–1021, 2008.
- [121] J. P. Saludes, M. J. Garson, S. G. Franzblau, and A. M. Aguinado, "Antitubercular constituents from the hexane fraction of *Morinda citrifolia* Linn. (Rubiaceae)," *Phytotherapy Research*, vol. 16, no. 7, pp. 683–685, 2002.
- [122] G. Mudur, "Indian health ministry challenges report of totally drug resistant tuberculosis," *BMJ*, vol. 344, article e702, 2012.
- [123] S. Zanetti, S. Cannas, P. Molicotti et al., "Evaluation of the Antimicrobial Properties of the Essential Oil of *Myrtus communis* L. against Clinical Strains of *Mycobacterium* spp.," *Interdisciplinary Perspectives on Infectious Diseases*, vol. 2010, Article ID 931530, 3 pages, 2010.
- [124] H. A. Mahmud, H. Seo, S. Kim et al., "Thymoquinone (TQ) inhibits the replication of intracellular *Mycobacterium tuberculosis* in macrophages and modulates nitric oxide production," *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 279, 2017.
- [125] B. S. Siddiqui, H. A. Bhatti, S. Begum, and S. Perwaiz, "Evaluation of the antimycobacterium activity of the constituents from *Ocimum basilicum* against *Mycobacterium tuberculosis*," *Journal of Ethnopharmacology*, vol. 144, no. 1, pp. 220–222, 2012.
- [126] R. S. Sundaram, M. Ramanathan, R. Rajesh, B. Satheesh, and D. Saravanan, "LC-MS quantification of rosmarinic acid and ursolic acid in theocimum sanctumlinn. leaf extract (Holy Basil, Tulsi)," *Journal of Liquid Chromatography & Related Technologies*, vol. 35, no. 5, pp. 634–650, 2012.
- [127] L. C. Pabon and L. E. Cuca, "Aporphine alkaloids from *Ocotea macrophylla* (Lauraceae)," *Quim Nova*, vol. 33, no. 4, pp. 875–879, 2010.
- [128] T. Inui, Y. Wang, D. Nikolic, D. C. Smith, S. G. Franzblau, and G. F. Pauli, "Sesquiterpenes from *Oplopanax horridus*," *Journal of Natural Products*, vol. 73, no. 4, pp. 563–567, 2010.
- [129] S. Sutthivaiyakit, O. Thongnak, T. Lhinhatrakool et al., "Cytotoxic and antimycobacterial prenylated flavonoids from the roots of *Eriosema chinense*," *Journal of Natural Products*, vol. 72, no. 6, pp. 1092–1096, 2009.
- [130] A. H. Januário, E. R. Filho, R. C. L. R. Pietro, S. Kashima, D. N. Sato, and S. C. França, "Antimycobacterial physalins from *Physalis angulata* L. (Solanaceae)," *Phytotherapy Research*, vol. 16, no. 5, pp. 445–448, 2002.
- [131] T. Rukachaisirikul, P. Siri wattanakit, K. Sukcharoenphol et al., "Chemical constituents and bioactivity of *Piper sarmentosum*," *Journal of Ethnopharmacology*, vol. 93, no. 2–3, pp. 173–176, 2004.
- [132] R. B. L. Scodro, C. T. A. Pires, V. S. Carrara et al., "Anti-tuberculosis neolignans from *Piper regnellii*," *Phytomedicine*, vol. 20, no. 7, pp. 600–604, 2013.
- [133] M. A. Lopes, K. R. C. Ferracioli, V. L. D. Siqueira et al., "In vitro interaction of eupomatenoid-5 from *Piper solmsianum* C. DC. var. *solmsianum* and anti-tuberculosis drugs," *The International Journal of Tuberculosis and Lung Disease*, vol. 18, no. 12, pp. 1513–1515, 2014.
- [134] P. Rijo, M. F. Simões, A. P. Francisco et al., "Antimycobacterial metabolites from *Plectranthus royleanone* derivatives against *Mycobacterium tuberculosis* strains," *Chemistry & Biodiversity*, vol. 7, no. 4, pp. 922–932, 2010.

- [135] J. M. Andrade, L. Custódio, A. Romagnoli et al., "Antitubercular and anti-inflammatory properties screening of natural products from *Plectranthus* species," *Future Medicinal Chemistry*, vol. 10, no. 14, pp. 1677–1691, 2018.
- [136] P. Kumar, A. Singh, U. Sharma, D. Singh, M. P. Dobhal, and S. Singh, "Anti-mycobacterial activity of plumericin and isoplumericin against MDR *Mycobacterium tuberculosis*," *Pulmonary Pharmacology & Therapeutics*, vol. 26, no. 3, pp. 332–335, 2013.
- [137] S. Kanokmedhakul, K. Kanokmedhakul, and R. Lekphrom, "Bioactive constituents of the roots of *Polyalthia cerasoides*," *Journal of Natural Products*, vol. 70, no. 9, pp. 1536–1538, 2007.
- [138] N. Panthama, S. Kanokmedhakul, and K. Kanokmedhakul, "Polyacetylenes from the roots of *Polyalthia debilis*," *Journal of Natural Products*, vol. 73, no. 8, pp. 1366–1369, 2010.
- [139] S. Kanokmedhakul, K. Kanokmedhakul, I. Kantikeaw, and N. Phonkerd, "2-Substituted furans from the roots of *Polyalthia evecta*," *Journal of Natural Products*, vol. 69, no. 1, pp. 68–72, 2006.
- [140] P. Kittakoop, S. Wanasith, P. Watts, J. Kramyu, M. Tanticharoen, and Y. Thebtaranonth, "Potent antiviral potamogetonyde and potamogetonol, new furanoid labdane diterpenes from *Potamogeton malaianus*," *Journal of Natural Products*, vol. 64, no. 3, pp. 385–388, 2001.
- [141] J. J. Chen, W. J. Lin, P. C. Shieh, I. S. Chen, C. F. Peng, and P. J. Sung, "A new long-chain alkene and antituberculosis constituents from the leaves of *Pourthia lucida*," *Chemistry and Biodiversity*, vol. 7, no. 3, pp. 717–721, 2010.
- [142] S. B. Lirio, A. P. G. Macabeo, E. M. Paragas et al., "Antitubercular constituents from *Premna odorata* Blanco," *Journal of Ethnopharmacology*, vol. 154, no. 2, pp. 471–474, 2014.
- [143] D. Dey, R. Ray, and B. Hazra, "Antimicrobial activity of pomegranate fruit constituents against drug-resistant *Mycobacterium tuberculosis* and β -lactamase producing *Klebsiella pneumoniae*," *Pharmaceutical Biology*, vol. 53, no. 10, pp. 1474–1480, 2015.
- [144] N. B. Truong, C. V. Pham, H. T. M. Doan et al., "Antituberculosis cycloartane triterpenoids from *Radermachera boniana*," *Journal of Natural Products*, vol. 74, no. 5, pp. 1318–1322, 2011.
- [145] K. Z. Deng, Y. Xiong, B. Zhou, Y. M. Guan, and Y. M. Luo, "Chemical constituents from the roots of *Ranunculus ternatus* and their inhibitory effects on *Mycobacterium tuberculosis*," *Molecules*, vol. 18, no. 10, pp. 11859–11865, 2013.
- [146] H. X. Liang, H. Q. Dai, H. A. Fu et al., "Bioactive compounds from *Rumex* plants," *Phytochemistry Letters*, vol. 3, no. 4, pp. 181–184, 2010.
- [147] A. A. Hussein, J. J. M. Meyer, M. L. Jimeno, and B. Rodríguez, "Bioactive diterpenes from *Orthosiphon labiatus* and *Salvia africana-lutea*," *Journal of Natural Products*, vol. 70, no. 2, pp. 293–295, 2007.
- [148] C. Chen, W. He, P. Huang et al., "Exploring anti-TB leads from natural products library originated from marine microbes and medicinal plants," *Antonie Van Leeuwenhoek*, vol. 102, no. 3, pp. 447–461, 2012.
- [149] D. Yang and H. Luo, "Modification of diterpenoid quinones from *Salvia miltiorrhiza*," *Journal of China Pharmaceutical University*, vol. 29, pp. 255–258, 1998.
- [150] P. Chumkaew, C. Karalai, C. Ponglimanont, and K. Chantrapromma, "Antimycobacterial activity of phorbol esters from the fruits of *Sapium indicum*," *Journal of Natural Products*, vol. 66, no. 4, pp. 540–543, 2003.
- [151] T. C. Hung, K. B. Chen, W. Y. Lee, and C. Y. C. Chen, "The Inhibition of Folylpolylglutamate Synthetase (folC) in the Prevention of Drug Resistance in *Mycobacterium tuberculosis* by Traditional Chinese Medicine," *BioMed Research International*, vol. 2014, Article ID 635152, 14 pages, 2014.
- [152] A. Suksamrarn, M. Buaprom, S. Udtip, N. Nuntawong, R. Haritakun, and S. Kanokmedhakul, "Antimycobacterial and antiplasmodial unsaturated carboxylic acid from the twigs of *Scleropyrum wallichianum*," *Chemical & Pharmaceutical Bulletin (Tokyo)*, vol. 53, no. 10, pp. 1327–1329, 2005.
- [153] C. Balachandran, V. Duraipandiyar, N. A. al-Dhabi et al., "Antimicrobial and antimycobacterial activities of methyl caffeate isolated from *Solanum torvum* Swartz. fruit," *Indian Journal of Microbiology*, vol. 52, no. 4, pp. 676–681, 2012.
- [154] J. M. Hwang, T. Oh, T. Kaneko et al., "Design, synthesis, and structure–activity relationship studies of tryptanthrins as antitubercular agents," *Journal of Natural Products*, vol. 76, no. 3, pp. 354–367, 2013.
- [155] L. A. Mitscher and W. R. Baker, "A search for novel chemotherapy against tuberculosis amongst natural products," *Pure and Applied Chemistry*, vol. 70, no. 2, pp. 365–371, 1998.
- [156] A. Mann, J. O. Amupitan, A. O. Oyewale et al., "Evaluation of in vitro antimycobacterial activity of Nigerian plants used for treatment of respiratory diseases," *African Journal of Biotechnology*, vol. 7, no. 11, pp. 1630–1636, 2008.
- [157] A. Mann, K. Ibrahim, A. O. Oyewale, J. O. Amupitan, M. O. Fatope, and J. I. Okogun, "Antimycobacterial Friedelane-terpenoid from the root bark of *Terminalia avicennioides*," *American Journal of Chemistry*, vol. 1, no. 2, pp. 52–55, 2011.
- [158] E. Y. A. Salih, R. Julkunen-Tiitto, A. M. Lampi et al., "*Terminalia laxiflora* and *Terminalia brownii* contain a broad spectrum of antimycobacterial compounds including ellagitannins, ellagic acid derivatives, triterpenes, fatty acids and fatty alcohols," *Journal of Ethnopharmacology*, vol. 227, pp. 82–96, 2018.
- [159] C. T. Higuchi, M. Sannomiya, F. R. Pavan et al., "Byrsonima fagifolia Niedenzu Apolar compounds with antitubercular activity," *Evidence-based Complementary and Alternative Medicine*, vol. 2011, Article ID 128349, 5 pages, 2011.
- [160] V. Kuete, T. K. Tabopda, B. Ngameni, F. Nana, T. E. Tshikalange, and B. T. Ngadjui, "Antimycobacterial, antibacterial and antifungal activities of *Terminalia superba* (Combretaceae)," *South African Journal of Botany*, vol. 76, no. 1, pp. 125–131, 2010.
- [161] M. C. Fomogne-Fodjo, D. T. Ndinteh, D. K. Olivier, P. Kempgens, S. van Vuuren, and R. W. M. Krause, "Secondary metabolites from *Tetracera potatoria* stem bark with antimycobacterial activity," *Journal of Ethnopharmacology*, vol. 195, pp. 238–245, 2017.
- [162] V. P. Baldin, R. B. L. Scodro, M. A. Lopes-Ortiz et al., "Antimycobacterial tuberculosis activity of essential oil and 6,7-dehydrororoleanone isolated from leaves of *Tetradenia riparia* (Hochst.) Codd (Lamiaceae)," *Phytomedicine*, vol. 47, pp. 34–39, 2018.
- [163] M. T. Gutierrez-Lugo, Y. Wang, S. G. Franzblau, E. Suarez, and B. N. Timmermann, "Antitubercular sterols from *Thalia multiflora* Horkel ex Koernicke," *Phytotherapy Research*, vol. 19, no. 10, pp. 876–880, 2005.

- [164] S. Sureram, S. P. D. Senadeera, P. Hongmanee, C. Mahidol, S. Ruchirawat, and P. Kittakoop, "Antimycobacterial activity of bisbenzylisoquinoline alkaloids from *Tiliacora triandra* against multidrug-resistant isolates of *Mycobacterium tuberculosis*," *Bioorganic & Medicinal Chemistry Letters*, vol. 22, no. 8, pp. 2902–2905, 2012.
- [165] J. Zhao, D. Evangelopoulos, S. Bhakta, A. I. Gray, and V. Seidel, "Antitubercular activity of *Arctium lappa* and *Tussilago farfara* extracts and constituents," *Journal of Ethnopharmacology*, vol. 155, no. 1, pp. 796–800, 2014.
- [166] S. Basu, A. Ghosh, and B. Hazra, "Evaluation of the antibacterial activity of *Ventilago madraspatana* Gaertn., *Rubia cordifolia* Linn. and *Lantana camara* Linn.: isolation of emodin and physcion as active antibacterial agents," *Phytotherapy Research*, vol. 19, no. 10, pp. 888–894, 2005.
- [167] X. Luo, D. Pires, J. A. Aínsa et al., "Zanthoxylum capense constituents with antimycobacterial activity against *Mycobacterium tuberculosis* in vitro and ex vivo within human macrophages," *Journal of Ethnopharmacology*, vol. 146, no. 1, pp. 417–422, 2013.
- [168] L. Bunalema, G. W. Fotso, P. Waako, J. Tabuti, and S. O. Yeboah, "Potential of *Zanthoxylum leprieurii* as a source of active compounds against drug resistant *Mycobacterium tuberculosis*," *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 89, 2017.
- [169] S. Kim, H. Seo, H. A. Mahmud et al., "In vitro activity of colinin isolated from the leaves of *Zanthoxylum schinifolium* against multidrug- and extensively drug-resistant *Mycobacterium tuberculosis*," *Phytomedicine*, vol. 46, pp. 104–110, 2018.
- [170] H. Y. Huang, T. Ishikawa, C. F. Peng, I. L. Tsai, and I. S. Chen, "Constituents of the root wood of *Zanthoxylum wutaiense* with antitubercular activity," *Journal of Natural Products*, vol. 71, no. 7, pp. 1146–1151, 2008.
- [171] P. Tangyuenyongwatana and W. Gritsanapan, "Biological evaluations of fatty acid esters originated during storage of Prasaplay, a Thai traditional medicine," *Natural Product Research*, vol. 21, no. 11, pp. 990–997, 2007.
- [172] S. Suksamrarn, P. Panseeta, S. Kunchanawatta, T. Distaporn, S. Ruktasing, and A. Suksamrarn, "Ceanothane- and lupanetype triterpenes with antiplasmodial and antimycobacterial activities from *Ziziphus cambodiana*," *Chemical & Pharmaceutical Bulletin (Tokyo)*, vol. 54, no. 4, pp. 535–537, 2006.
- [173] M. Singh, M. Kaur, and O. Silakari, "Flavones: an important scaffold for medicinal chemistry," *European Journal of Medicinal Chemistry*, vol. 84, pp. 206–239, 2014.
- [174] C. L. Cantrell, S. Franzblau, and N. Fischer, "Antimycobacterial plant terpenoids," *Planta Medica*, vol. 67, no. 8, pp. 685–694, 2001.
- [175] A. Ulubelen, G. Topcu, and C. B. Johansson, "Norditerpenoids and diterpenoids from *Salviainmulticaulis* with Antituberculous Activity," *Journal of Natural Products*, vol. 60, no. 12, pp. 1275–1280, 1997.
- [176] J. C. Wu, T. C. Warren, J. Adams et al., "A novel, dipyrrodoiazepinone inhibitor of HIV-1 reverse transcriptase acts through a nonsubstrate binding site," *Biochemistry*, vol. 30, no. 8, pp. 2022–2026, 1991.
- [177] K. R. Gustafson, H. R. Bokesch, R. W. Fuller et al., "Calanone, a novel coumarin from *Calophyllum teysmannii*," *Tetrahedron Letters*, vol. 35, no. 32, pp. 5821–5824, 1994.
- [178] R. Mu, C. Kong, W. Yu et al., "Nitrooxidoreductase Rv2466c-dependent fluorescent probe for *Mycobacterium tuberculosis* Diagnosis and Drug susceptibility testing," *ACS Infectious Diseases*, vol. 5, no. 6, pp. 949–961, 2019.
- [179] K. Tawde, "Evaluation of selected Indian traditional folk medicinal plants against *Mycobacterium tuberculosis* with antioxidant and cytotoxicity study," *Asian Pacific Journal of Tropical Medicine*, vol. 2, pp. 685–691, 2012.
- [180] S. Y. Pan, S. F. Zhou, S. H. Gao et al., "New Perspectives on How to Discover Drugs from Herbal Medicines: CAM's Outstanding Contribution to Modern Therapeutics," *Evidence-based Complementary and Alternative Medicine*, vol. 2013, Article ID 627375, 25 pages, 2013.
- [181] R. Gupta, B. Thakur, P. Singh et al., "Anti-tuberculosis activity of selected medicinal plants against multi-drug resistant *Mycobacterium tuberculosis* isolates," *The Indian Journal of Medical Research*, vol. 131, pp. 809–813, 2010.
- [182] J. G. Bueno-Sánchez, J. R. Martínez-Morales, E. E. Stashenko, and W. Ribón, "Anti-tubercular activity of eleven aromatic and medicinal plants occurring in Colombia," *Biomédica*, vol. 29, no. 1, pp. 51–60, 2009.
- [183] D. Webster, T. D. G. Lee, J. Moore et al., "Antimycobacterial screening of traditional medicinal plants using the microplate resazurin assay," *Canadian Journal of Microbiology*, vol. 56, no. 6, pp. 487–494, 2010.
- [184] J. M. Nguta, R. Appiah-Opong, A. K. Nyarko et al., "Antimycobacterial and cytotoxic activity of selected medicinal plant extracts," *Ethnopharmacology*, vol. 182, pp. 10–15, 2016.
- [185] J. B. Nvau and P. Oladosu, "Antimycobacterial evaluation of some medicinal plants used in plateau State of Nigeria for the treatment of tuberculosis," *Agriculture and Biology Journal of North America*, vol. 2, no. 9, pp. 1270–1272, 2011.
- [186] M. Billo, P. Cabalion, J. Waikedre et al., "Screening of some new Caledonian and Vanuatu medicinal plants for antimycobacterial activity," *Journal of Ethnopharmacology*, vol. 96, no. 1-2, pp. 195–200, 2005.
- [187] R. E. Robles-Zepeda, E. W. Coronado-Aceves, C. A. Velázquez-Contreras, E. Ruiz-Bustos, M. Navarro-Navarro, and A. Garibay-Escobar, "In vitro anti-mycobacterial activity of nine medicinal plants used by ethnic groups in Sonora, Mexico," *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, p. 329, 2013.
- [188] R. Gómez-Cansino, C. I. Espitia-Pinzón, M. G. Campos-Lara et al., "Antimycobacterial and HIV-1 reverse transcriptase activity of Julianaceae and Clusiaceae plant species from Mexico," *Evidence-based Complementary and Alternative Medicine*, vol. 2015, Article ID 183036, 8 pages, 2015.
- [189] A. Prabu, S. Hassan, Prabuseenivasan, A. S. Shainaba, L. E. Hanna, and V. Kumar, "Andrographolide: A potent antituberculosis compound that targets Aminoglycoside 2'-N-acetyltransferase in *Mycobacterium tuberculosis*," *Journal of Molecular Graphics & Modelling*, vol. 61, pp. 133–140, 2015.
- [190] P. Bhattar, P. Gupta, P. Daswani, P. Tetali, and T. Birdi, "Antimycobacterial efficacy of andrographis paniculata leaf extracts under intracellular and hypoxic conditions," *Journal of Evidence-Based Complementary & Alternative Medicine*, vol. 20, no. 1, pp. 3–8, 2015.
- [191] S. Mohamad, N. M. Zin, H. A. Wahab et al., "Antituberculosis potential of some ethnobotanically selected Malaysian plants," *Journal of Ethnopharmacology*, vol. 133, no. 3, pp. 1021–1026, 2011.
- [192] N. P. Komape, V. P. Bagla, P. Kabongo-Kayoka, and P. Masoko, "Anti-mycobacteria potential and synergistic

- effects of combined crude extracts of selected medicinal plants used by Bapedi traditional healers to treat tuberculosis related symptoms in Limpopo Province, South Africa," *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 128, 2017.
- [193] D. E. Cruz-Vega, M. J. Verde-Star, N. Salinas-González et al., "Antimycobacterial activity of *Juglans regia*, *Juglans mollis*, *Carya illinoensis* and *Bocconia frutescens*," *Phytotherapy Research*, vol. 22, no. 4, pp. 557–559, 2008.
- [194] S. K. Naik, S. Mohanty, A. Padhi, R. Pati, and A. Sonawane, "Evaluation of antibacterial and cytotoxic activity of *Artemisia nilagirica* and *Murraya koenigii* leaf extracts against mycobacteria and macrophages," *BMC Complementary and Alternative Medicine*, vol. 14, no. 1, p. 87, 2014.
- [195] C. N. Nkenfou, I. K. Mawabo, A. Notedji et al., "In vitro antimycobacterial activity of six Cameroonian medicinal plants using microplate alamarBlue assay," *Intl J Mycobacteriol.*, vol. 4, no. 4, pp. 306–311, 2015.
- [196] V. K. Gupta, A. Kaushik, D. S. Chauhan, R. K. Ahirwar, S. Sharma, and D. Bisht, "Anti-mycobacterial activity of some medicinal plants used traditionally by tribes from Madhya Pradesh, India for treating tuberculosis related symptoms," *Journal of Ethnopharmacology*, vol. 227, pp. 113–120, 2018.
- [197] E. Green, A. Samie, C. L. Obi, P. O. Bessong, and R. N. Ndip, "Inhibitory properties of selected South African medicinal plants against *Mycobacterium tuberculosis*," *Journal of Ethnopharmacology*, vol. 130, no. 1, pp. 151–157, 2010.
- [198] A. Y. Gordien, A. I. Gray, K. Ingleby, S. G. Franzblau, and V. Seidel, "Activity of Scottish plant, lichen and fungal endophyte extracts against *Mycobacterium aurum* and *Mycobacterium tuberculosis*," *Phytotherapy Research*, vol. 24, no. 5, pp. 692–698, 2010.
- [199] P. Arulmozhi, S. Vijayakumar, and T. Kumar, "Phytochemical analysis and antimicrobial activity of some medicinal plants against selected pathogenic microorganisms," *Microbial Pathogenesis*, vol. 123, pp. 219–226, 2018.
- [200] P. Arulmozhi, S. Vijayakumar, P. K. Praseetha, and S. Jayanthi, "Extraction methods and computational approaches for evaluation of antimicrobial compounds from *Capparis zeylanica* L.," *Analytical Biochemistry*, vol. 572, pp. 33–44, 2019.
- [201] R. Singh, S. Hussain, R. Verma, and P. Sharma, "Anti-mycobacterial screening of five Indian medicinal plants and partial purification of active extracts of *Cassia sophera* and *Urtica dioica*," *Asian Pacific Journal of Tropical Medicine*, vol. 6, no. 5, pp. 366–371, 2013.
- [202] G. M. Molina-Salinas, A. Pérez-López, P. Becerril-Montes, R. Salazar-Aranda, S. Said-Fernández, and N. W. Torres, "Evaluation of the flora of northern Mexico for in vitro antimicrobial and antituberculosis activity," *Journal of Ethnopharmacology*, vol. 109, no. 3, pp. 435–441, 2007.
- [203] S. Ullah, S. Hussain, S. N. Khan, M. Khurram, I. Khan, and M. A. Khan, "The medicinal plants in the control of tuberculosis: laboratory study on medicinal plants from the Northern Area of Pakistan," *International Journal of Mycobacteriology*, vol. 6, no. 1, pp. 102–105, 2017.
- [204] M. R. Camacho-Corona, M. A. Ramírez-Cabrera, O. G. Santiago, E. Garza-González, I. P. Palacios, and J. Luna-Herrera, "Activity against drug resistant-tuberculosis strains of plants used in Mexican traditional medicine to treat tuberculosis and other respiratory diseases," *Phytotherapy Research*, vol. 22, no. 1, pp. 82–85, 2008.
- [205] E. A. Diop, E. F. Queiroz, S. Kicka et al., "Survey on medicinal plants traditionally used in Senegal for the treatment of tuberculosis (TB) and assessment of their antimycobacterial activity," *Journal of Ethnopharmacology*, vol. 216, pp. 71–78, 2018.
- [206] S. Mohamad, N. N. Ismail, T. Parumasivam, P. Ibrahim, H. Osman, and H. A. Wahab, "Antituberculosis activity, phytochemical identification of *Costus speciosus* (J. Koenig) Sm., *Cymbopogon citratus* (DC. Ex Nees) Stapf., and *Tabernaemontana coronaria* (L.) Willd. and their effects on the growth kinetics and cellular integrity of *Mycobacterium tuberculosis* H37Rv," *BMC Complementary and Alternative Medicine*, vol. 18, no. 1, p. 5, 2018.
- [207] A. O. Aro, J. P. Dzoyem, J. N. Eloff, and L. J. McGaw, "Extracts of six Rubiaceae species combined with rifampicin have good in vitro synergistic antimycobacterial activity and good anti-inflammatory and antioxidant activities," *BMC Complementary and Alternative Medicine*, vol. 16, no. 1, p. 385, 2016.
- [208] K. N. Ngbolua, G. N. Bongo, C. M. Ashande et al., "Ethnobotanical survey and ecological study of plants resources used in folk medicine to treat symptoms of tuberculosis in Kinshasa City, Democratic Republic of the Congo," *Journal of Drug Discovery, Development and Delivery*, vol. 1, no. 4, pp. 1–6, 2014.
- [209] F. C. M. Lopes, M. C. P. Placeres, C. M. Jordão Junior et al., "Immunological and microbiological activity of *Davilla elliptica* St. Hill. (Dilleniaceae) against *Mycobacterium tuberculosis*," *Memórias do Instituto Oswaldo Cruz*, vol. 102, no. 6, pp. 769–772, 2007.
- [210] E. M. Tekwu, T. Askun, V. Kuete et al., "Antibacterial activity of selected Cameroonian dietary spices ethno-medically used against strains of *Mycobacterium tuberculosis*," *Journal of Ethnopharmacology*, vol. 142, no. 2, pp. 374–382, 2012.
- [211] L. Bunalema, C. Kirimuhuzya, J. R. Tabuti et al., "The efficacy of the crude root bark extracts of *Erythrina abyssinica* on rifampicin resistant *Mycobacterium tuberculosis*," *African Health Sciences*, vol. 11, no. 4, pp. 587–593, 2011.
- [212] E. W. Coronado-Aceves, J. J. Sánchez-Escalante, J. López-Cervantes et al., "Antimycobacterial activity of medicinal plants used by the Mayo people of Sonora, Mexico," *Journal of Ethnopharmacology*, vol. 190, pp. 106–115, 2016.
- [213] T. Rajasekar, S. Anbarasu, R. Manikkam, J. Joseph, and V. Kumar, "Inhibitory activity of *Euphorbia hirta* (Tawatawa) extracts against *Mycobacterium tuberculosis* and other non mycobacterial pathogens," *Der Pharma Chemica*, vol. 7, pp. 213–216, 2015.
- [214] L. R. Barrows, E. Powan, C. D. Pond, and T. Matainaho, "Anti-TB activity of *Evodia elleryana* bark extract," *Fitoterapia*, vol. 78, no. 3, pp. 250–252, 2007.
- [215] B. Madikizela, A. R. Ndhlala, J. F. Finnie, and J. van Staden, "Antimycobacterial, anti-inflammatory and genotoxicity evaluation of plants used for the treatment of tuberculosis and related symptoms in South Africa," *Journal of Ethnopharmacology*, vol. 153, no. 2, pp. 386–391, 2014.
- [216] M. D. Antoun, Z. Ramos, J. Vazques et al., "Evaluation of the flora of Puerto Rico for in vitro antiplasmodial and antimycobacterial activities," *Phytotherapy Research*, vol. 15, no. 7, pp. 638–642, 2001.
- [217] G. M. Molina-Salinas, M. C. Ramos-Guerra, J. Vargas-Villareal, B. D. Mata-Cárdenas, P. Becerril-Montes, and S. Said-Fernández, "Bactericidal Activity of Organic Extracts from

- Flourensia cernua* DC against Strains of *Mycobacterium tuberculosis*," *Archives of Medical Research*, vol. 37, no. 1, pp. 45–49, 2006.
- [218] D. Khelifi, M. Hamdi, A. E. Hayouni et al., "Global chemical composition and antioxidant and anti-tuberculosis activities of various extracts of *Globularia alypum* L. (Globulariaceae) leaves," *Molecules*, vol. 16, no. 12, pp. 10592–10603, 2011.
- [219] V. K. Gupta, A. Fatima, U. Faridi et al., "Antimicrobial potential of *Glycyrrhiza glabra* roots," *Journal of Ethnopharmacology*, vol. 116, no. 2, pp. 377–380, 2008.
- [220] R. Gomes-Flores, C. Arzate-Quintana, R. Quintanilla-Licea, P. Tamez-Guerra, Monreal-Cuevase, and C. Rodriguez-Padilla, "Antimicrobial activity of *Persea americana* Mill (Lauraceae) (avocado) and *Gymnosperma glutinosum* (Spreng.) Less (Asteraceae) leaf extracts and active fractions against *Mycobacterium tuberculosis*," *American-Eurasian Journal of Scientific Research (AEJSR)*, vol. 3, pp. 188–194, 2008.
- [221] B. Madikizela and L. J. McGaw, "Scientific rationale for traditional use of plants to treat tuberculosis in the eastern region of the OR Tambo district, South Africa," *Journal of Ethnopharmacology*, vol. 224, pp. 250–260, 2018.
- [222] N. K. Sethiya, N. M. Ahmed, R. M. Shekh, V. Kumar, P. Kumar Singh, and V. Kumar, "Ethnomedicinal, phytochemical and pharmacological updates on *Hygrophila auriculata* (Schum.) Hiene: an overview," *Journal of Integrative Medicine*, vol. 16, no. 5, pp. 299–311, 2018.
- [223] M. H. Kabir, N. Hasan, M. Rahman et al., "A survey of medicinal plants used by the Deb barma clan of the Tripura tribe of Moulvibazar district, Bangladesh," *Journal of Ethnobiology and Ethnomedicine*, vol. 10, no. 1, p. 19, 2014.
- [224] N. Abuzeid, S. Kalsum, R. J. Koshy et al., "Antimycobacterial activity of selected medicinal plants traditionally used in Sudan to treat infectious diseases," *Journal of Ethnopharmacology*, vol. 157, pp. 134–139, 2014.
- [225] E. A. Earl, M. Altaf, R. V. Murikoli, S. Swift, and R. O'Toole, "Native New Zealand plants with inhibitory activity towards *Mycobacterium tuberculosis*," *BMC Complementary and Alternative Medicine*, vol. 10, no. 1, p. 25, 2010.
- [226] X. Luo, D. Pires, J. A. Ainsa et al., "Antimycobacterial evaluation and preliminary phytochemical investigation of selected medicinal plants traditionally used in Mozambique," *Journal of Ethnopharmacology*, vol. 137, no. 1, pp. 114–120, 2011.
- [227] V. K. Gupta, C. Shukla, G. R. S. Bisht, D. Saikia, S. Kumar, and R. L. Thakur, "Detection of anti-tuberculosis activity in some folklore plants by radiometric BACTEC assay," *Letters in Applied Microbiology*, vol. 52, no. 1, pp. 33–40, 2011.
- [228] T. Mmushi, P. Masoko, L. Mde, M. Mokgotho, L. Mampuru, and R. Howard, "Antimycobacterial evaluation of fifteen medicinal plants in South Africa," *African Journal of Traditional, Complementary, and Alternative Medicines*, vol. 7, no. 1, pp. 34–39, 2009.
- [229] W. Kahaliw, A. Aseffa, M. Abebe, M. Teferi, and E. Engdawork, "Evaluation of the antimycobacterial activity of crude extracts and solvent fractions of selected Ethiopian medicinal plants," *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 143, 2017.
- [230] J. Kardan-Yamchi, M. Mahboubi, H. Kazemian, G. Hamzelou, and M. M. Feizabadi, "The chemical composition and anti-mycobacterial activities of *Trachyspermum copticum* and *Pelargonium graveolens* essential oils," *Recent Patents on Anti-Infective Drug Discovery*, vol. 15, no. 1, pp. 68–74, 2020.
- [231] H. Kolodziej, O. Kayser, O. A. Radtke, A. F. Kiderlen, and E. Koch, "Pharmacological profile of extracts of *Pelargonium sidoides* and their constituents," *Phytomedicine*, vol. 10, Supplement 4, pp. 18–24, 2003.
- [232] A. Jiménez-Arellanes, J. Luna-Herrera, R. Ruiz-Nicolás, J. Cornejo-Garrido, A. Tapia, and L. Yépez-Mulia, "Antiprotazoal and antimycobacterial activities of *Persea americana* seeds," *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, p. 109, 2013.
- [233] K. Z. Bernuci, C. Iwanaga, C. Fernandez-Andrade et al., "Evaluation of chemical composition and antileishmanial and antituberculosis activities of essential oils of *Piper* species," *Molecules*, vol. 21, no. 12, p. 1698, 2016.
- [234] L. E. Diaz, D. R. Munoz, R. E. Prieto et al., "Antioxidant, anti-tubercular and cytotoxic activities of *Piper imperiale*," *Molecules*, vol. 17, no. 4, pp. 4142–4157, 2012.
- [235] K. Hussain, Z. Ismail, A. Sadikun, and P. Ibrahim, "Analysis of proteins, polysaccharides, glycosaponins contents of *Piper sarmentosum* Roxb. and anti-TB evaluation for bio-enhancing/interaction effects of leaf extract with isoniazid (INH)," *Natural Product Radiance*, vol. 7, pp. 402–408, 2008.
- [236] G. C. Ebi and S. I. Ofoefule, "Antimicrobial activity of *Pterocarpus osun* stems," *Fitoterapia*, vol. 71, no. 4, pp. 433–435, 2000.
- [237] T. Askun, E. M. Tekwu, F. Satil, S. Modanlioglu, and H. Aydeniz, "Preliminary antimycobacterial study on selected Turkish plants (Lamiaceae) against *Mycobacterium tuberculosis* and search for some phenolic constituents," *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, p. 365, 2013.
- [238] M. Bhatnagar, N. Sarkar, N. Gandharv, O. Apang, S. Singh, and S. Ghosal, "Evaluation of antimycobacterial, leishmanicidal and antibacterial activity of three medicinal orchids of Arunachal Pradesh, India," *BMC Complementary and Alternative Medicine*, vol. 17, no. 1, p. 379, 2017.
- [239] I. T. Babalola, E. A. Adelakun, Y. Wang, and F. O. Shode, "Anti-TB activity of *Sterculia setigera* Del., leaves (Sterculiaceae)," *J Pharmacogn Phytochem.*, vol. 1, pp. 17–23, 2012.
- [240] D. Saikia, S. Parveen, V. K. Gupta, and S. Luqman, "Antituberculosis activity of Indian grass KHUS (*Vetiveria zizanioides* L. Nash)," *Complementary Therapies in Medicine*, vol. 20, no. 6, pp. 434–436, 2012.
- [241] J. Ngezahayo, F. Havyarimana, L. Hari, C. Stévigny, and P. Duez, "Medicinal plants used by Burundian traditional healers for the treatment of microbial diseases," *Journal of Ethnopharmacology*, vol. 173, pp. 338–351, 2015.
- [242] R. J. Case, S. G. Franzblau, Y. Wang, S. H. Cho, D. D. Soejarto, and G. F. Pauli, "Ethnopharmacological evaluation of the informant consensus model on anti-tuberculosis claims among the Manus," *Journal of Ethnopharmacology*, vol. 106, no. 1, pp. 82–89, 2006.
- [243] S. S. Semenya and A. Maroyi, "Ethnobotanical survey of plants used by Bapedi traditional healers to treat tuberculosis and its opportunistic infections in the Limpopo Province, South Africa," *South African Journal of Botany*, vol. 122, pp. 401–421, 2019.
- [244] I. O. Lawal, D. S. Grierson, and A. J. Afolayan, "Phytotherapeutic information on plants used for the treatment of tuberculosis in Eastern Cape Province, South Africa," *Evidence-*

- based *Complementary and Alternative Medicine*, vol. 2014, Article ID 735423, 11 pages, 2014.
- [245] K. V. Phungula, A. Marston, M. Khamane, P. C. Zietsman, and S. L. Bonnet, "Ethnobotanical survey of plants used for the treatment of tuberculosis by traditional health healers in the Free State Province, South Africa," *Planta Medica*, vol. 80, no. 16, p. LP75, 2014.
- [246] J. N. Nguta, R. Appiah-Opong, A. K. Nyarko, D. Yeboah-Manu, and P. G. A. Addo, "Medicinal plants used to treat TB in Ghana," *International Journal of Mycobacteriology*, vol. 4, no. 2, pp. 116–123, 2015.
- [247] A. Mann, J. O. Amupitan, A. O. Oyewale, J. I. Okogun, and K. Ibrahim, "An ethnobotanical survey of indigenous flora for treating tuberculosis and other respiratory diseases in Niger State, Nigeria," *Journal of Phytomedicine and Therapeutics*, vol. 12, no. 1, pp. 1–12, 2008.
- [248] B. G. Elkington, B. Southavong, K. Sydara et al., "Biological evaluation of plants of Laos used in the treatment of tuberculosis in Lao traditional medicine," *Pharmaceutical Biology*, vol. 47, no. 1, pp. 26–33, 2009.
- [249] S. F. Sabran, M. Mohamed, and M. F. Abu Bakar, "Ethnomedical knowledge of plants used for the treatment of tuberculosis in Johor, Malaysia," *Evidence-based Complementary and Alternative Medicine*, vol. 2016, Article ID 2850845, 12 pages, 2016.
- [250] L. J. McGaw, N. Lall, J. J. M. Meyer, and J. N. Eloff, "The potential of South African plants against *Mycobacterium* infections," *Journal of Ethnopharmacology*, vol. 119, no. 3, pp. 482–500, 2008.
- [251] J. R. Tabuti, C. B. Kukunda, and P. J. Waako, "Medicinal plants used by traditional medicine practitioners in the treatment of tuberculosis and related ailments in Uganda," *Journal of Ethnopharmacology*, vol. 127, no. 1, pp. 130–136, 2010.
- [252] L. Bunalema, S. Obakiro, J. R. S. Tabuti, and P. Waako, "Knowledge on plants used traditionally in the treatment of tuberculosis in Uganda," *Journal of Ethnopharmacology*, vol. 151, no. 2, pp. 999–1004, 2014.
- [253] S. B. Sanusi, M. F. Abu Bakar, M. Mohamed, S. F. Sabran, and M. M. Mainasara, "Southeast Asian medicinal plants as a potential source of antituberculosis agent," *Evidence-based Complementary and Alternative Medicine*, vol. 2017, Article ID 7185649, 39 pages, 2017.
- [254] A. O. J., O. P., B. T. J., O. N., M. J. J., and K. C., "Indigenous knowledge of communities around Lake Victoria Basin regarding treatment and management of tuberculosis using medicinal plants," *International Journal of Medicine and Medical Sciences*, vol. 6, no. 1, pp. 16–23, 2014.
- [255] M. A. R. Mariita, *Efficacy of medicinal plants used by communities around Lake Victoria Region and the Samburu against Mycobacteria, selected bacteria and Candida albicans Kenya*, Kenyatta University, 2006.
- [256] K. S. Ahmad, A. Hamid, F. Nawaz et al., "Ethnopharmacological studies of indigenous plants in Kel village, Neelum Valley, Azad Kashmir, Pakistan," *Journal of Ethnobiology and Ethnomedicine*, vol. 13, no. 1, p. 68, 2017.
- [257] H. M. Ahmed, "Ethnopharmacobotanical study on the medicinal plants used by herbalists in Sulaymaniyah Province, Kurdistan, Iraq," *Journal of Ethnobiology and Ethnomedicine*, vol. 12, no. 1, p. 8, 2016.
- [258] V. Arya, "A review on anti-tubercular plants," *International Journal of PharmTech Research*, vol. 3, no. 2, pp. 872–880, 2011.
- [259] A. S. Saganuwan, "Some medicinal plants of Arabian Peninsula," *Medicinal Plants Research*, vol. 4, pp. 766–788, 2010.
- [260] Y. Xu, G. M. Dai, S. H. Tang, H. J. Yang, and Z. G. Sun, "Changes of Anti-tuberculosis Herbs formula during past three decades in contrast to ancient ones," *Chinese Journal of Integrative Medicine*, vol. 27, no. 5, pp. 388–393, 2020.
- [261] S. D. Tavhare and K. Nishteswar, "Collection practices of medicinal plants-Vedic, Ayurvedic and modern perspectives," *International Journal of Pharmacy and Biological Sciences*, vol. 5, pp. 54–61, 2014.
- [262] C. Gouws and J. H. Hamman, "Recent developments in our understanding of the implications of traditional African medicine on drug metabolism," *Expert Opinion on Drug Metabolism & Toxicology*, vol. 14, no. 2, pp. 161–168, 2018.
- [263] W. R. Folk, A. Smith, H. Song et al., "Does concurrent use of some botanicals interfere with treatment of tuberculosis?," *Neuromolecular Medicine*, vol. 18, no. 3, pp. 483–486, 2016.
- [264] D. Wilson, K. Goggin, K. Williams et al., "Consumption of *Sutherlandia frutescens* by HIV-seropositive South African adults: an adaptive double-blind randomized placebo controlled trial," *PLoS One*, vol. 10, no. 7, article e0128522, 2015.
- [265] C. O. Esimone, "Drug-drug and herb-drug interactions-a comment," *Jorind*, vol. 9, no. 1, pp. 47–59, 2011.
- [266] R. K. Zutshi, R. Singh, U. Zutshi, R. K. Johri, and C. K. Atal, "Influence of piperine on rifampicin blood levels in patients of pulmonary tuberculosis," *The Journal of the Association of Physicians of India*, vol. 33, no. 3, pp. 223–224, 1985.
- [267] D. V. Tatiraju, V. B. Bagade, P. J. Karambelkar, V. M. Jadhav, and V. Kadam, "Natural bioenhancers: an overview," *Journal of Pharmacognosy and Phytochemistry*, vol. 2, no. 3, pp. 55–60, 2013.
- [268] M. Kumar-Sarangi, B. Chandra-Joshi, and B. Ritchie, "Natural bioenhancers in drug delivery: an overview," *Puerto Rico Health Sciences Journal*, vol. 37, no. 1, pp. 12–18, 2018.
- [269] Y. S. Jaiswal and L. L. Williams, "A glimpse of Ayurveda - The forgotten history and principles of Indian traditional medicine," *Journal of Traditional and Complementary Medicine*, vol. 7, no. 1, pp. 50–53, 2017.