

## Research Article

# Postoperative Bone Marrow Lesions (BMLs) Are Associated with Pain Severity in Patients Undergoing Open Wedge High Tibial Osteotomy (OWHTO)

Bo Zhu, Tong-Fu Wang, De-Sheng Chen , Jia-Wang Zhu , Zeng-Liang Wang, Jian-Gang Cao, and Jun-Wei Zhao

Department of Sports Medicine and Arthroscopy, Tianjin Hospital of Tianjin University, Tianjin, China

Correspondence should be addressed to De-Sheng Chen; 392849261@qq.com

Received 19 March 2021; Revised 27 June 2021; Accepted 30 June 2021; Published 8 July 2021

Academic Editor: Fabiano Bini

Copyright © 2021 Bo Zhu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of the study was to investigate the relationship between postoperative bone marrow lesions (BMLs) and pain severity in patients undergoing open wedge high tibial osteotomy (OWHTO). We reviewed the patients undergoing OWHTO between April 2018 and April 2020. The demographic and clinical data of patients were collected. Clinically, VAS and Knee injury and Osteoarthritis Outcome Score (KOOS) were used to assess pain level and functional outcomes of patients. The MRI Osteoarthritis Knee Score (MOAKS) was used to assess the total BMLs size in medial tibiofemoral (MTF), lateral tibiofemoral (LTF), and patellofemoral (PF) joints. 98 patients were enrolled in the study, including 57 male and 41 female patients. The VAS scores improved significantly from  $6.1 \pm 0.8$  to  $1.5 \pm 0.9$  ( $p < 0.001$ ), and all subscales of KOOS improved significantly after surgery ( $p < 0.001$ ). There were no significant differences between the pre- and postoperative total BML size of PF and LTF joints ( $p > 0.05$ ). We observed significant improvements in the total BML size of MTF joint ( $p < 0.001$ ). The VAS scores and KOOS pain scores improved better in patients without postoperative MTF joint BMLs ( $p < 0.001$ ). Postoperative MTF joint BMLs were correlated with postoperative VAS ( $p < 0.001$ ) and KOOS pain ( $p < 0.001$ ). Our study demonstrates that MTF joint BMLs improved significantly after OWHTO. We confirmed that the presence of postoperative MTF joint BMLs are strongly associated with pain severity. The greater the improvement in postoperative MTF joint BMLs, the less pain. Our findings provide valuable understandings of OWHTO in the treatment of knee osteoarthritis (KOA) and potential future directions for KOA treatment approaches.

## 1. Introduction

Knee osteoarthritis (KOA) is a common chronic disease that causes pain, stiffness, and functional disability [1–4], among which, pain is the major cause leading individuals to seek medical care [5–8]. Many studies have demonstrated that knee pain could be relieved by open wedge high tibial osteotomy (OWHTO) in case of malalignment [9–12].

Despite the clinical efficacy of OWHTO for pain relief has been demonstrated, the underlying mechanisms and corresponding histological changes remain unclear. Many studies have shown that bone marrow lesions (BMLs) are strongly associated with pain in KOA [5, 13–15]. In a study conducted by Felson et al., 401 patients with KOA were

investigated, and 272 of 351 (77.5%) patients with painful knees had BMLs, comparing with 15 of 50 (30%) patients without painful knees had BMLs [13]. Moreover, a few studies have demonstrated that BMLs of the knee were significantly improved after OWHTO [16, 17]. These findings prompted further analysis of the relevance between BMLs and pain states in patients undergoing OWHTO.

So far, only a few studies have analyzed the BML changes and their effect on the prognosis for patients undergoing OWHTO [16–18]. Yang et al. reviewed 105 patients who underwent OWHTO, and they did not find any correlation between preoperative bone marrow edema (BME) severity and postoperative outcomes [19]. To our knowledge, no studies have investigated the relationship




FIGURE 1: The BMLs in medial tibiofemoral joint improved significantly after surgery. The red arrow indicates the pre- and postoperative zone of BMLs, and the yellow arrow indicates the zone after hardware removal. BMLs: bone marrow lesions.

between postoperative BMLs and pain severity in patients undergoing OWHTO.

This study is aimed at evaluating the relationship between postoperative BMLs and pain severity in patients undergoing OWHTO.

## 2. Materials and Methods

**2.1. Patient Selection.** We retrospectively reviewed the patients undergoing OWHTO between April 2018 and April 2020. The study was approved by the institutional ethics committee, and all patients signed consent forms. The inclusion criteria were (1) symptomatic medial KOA, (2) first OWHTO on the affected side, and (3) complete clinical data. The exclusion criteria were (1) revision surgery and (2) simultaneous bilateral OWHTO.

**2.2. Perioperative Management.** All patients received standard medical care, and all procedures were performed by the same surgical team. The osteotomy was performed under fluoroscopy, and the osteotomy site was fixed with the Tomo-fix plate system (Depuy Synthes, Eimattstrasse, Switzerland). Postoperative antibiotic prophylaxis (Intravenous cefuroxime, 1.5 g, *q8h for one day*) and analgesia were administered in all patients. For the prevention of venous thromboembolism, low molecule weight heparin (4250 IU, *qd*) was injected subcutaneously for 7 days, and rivaroxaban (10 mg *qd*) was administered during the subsequent 7 days. One year after hardware implantation, the hardware was removed, and magnetic resonance imaging (MRI) was performed (Figure 1).

**2.3. MR Imaging and Analysis.** MR images were analyzed by two expert musculoskeletal radiologists who were blind to any clinical information. The intraclass correlation coefficient (ICC) was calculated to assess interobserver variability. The MRI Osteoarthritis Knee Score (MOAKS) was used to assess the total BML size in medial tibiofemoral (MTF), lateral tibiofemoral (LTF), and patellofemoral (PF) joints. The criteria of the score are as follows: 0 = no BMLs

in subregion; 1 = size of BMLs/subregional volume < 33%; 2 = size of BMLs/subregional volume < 66%; 3 = size of BMLs/subregional volume > 66%. The maximum possible BMLs size was 15 in the MTF joint, 15 in the LTF joint, and 12 in the PF joint (Figure 2).

**2.4. Data Collection.** The demographic and clinical data of patients were collected, including gender, age, height, weight, body mass index (BMI), smoking, degree of correction, and size of osteotomy gap. Clinically, VAS and KOOS were used to assess pain level and functional outcomes of patients. MOAKS was used to assess the total BML size in MTF, LTF, and PF joints.

**2.5. Statistical Analysis.** The measurement data were expressed as mean value  $\pm$  standard deviation (SD); the pre- and postoperative parameters were compared using paired *t*-tests and Wilcoxon test. An independent sample *t*-test was used to compare pain scores in patients with or without BMLs. The Pearson correlation test was used for correlation analysis. The SPSS 22.0 software was used for statistical analysis, and  $p < 0.05$  was considered statistically significant.

## 3. Results

**3.1. General Results.** 98 patients were enrolled in the study, including 57 male and 41 female patients. The mean age of the patients was  $65.1 \pm 5.7$  years, and the mean BMI was  $28.1 \pm 3.4$ . The characteristics of the patients are shown in Table 1.

**3.2. Results of the VAS, KOOS, and MOAKS.** As shown in Table 2, the VAS scores improved significantly from  $6.1 \pm 0.8$  to  $1.5 \pm 0.9$  ( $p < 0.001$ ), and all subscales of KOOS improved significantly after surgery (Figure 3). There were significant differences between smoker and nonsmoker regarding preoperative VAS scores ( $p = 0.040$ ). There were no significant differences between smoker and nonsmoker regarding postoperative VAS scores and KOOS pain scores ( $p > 0.05$ ).




FIGURE 2: The LTF joint included five subregions: central (C) lateral femoral condyle, posterior (P) lateral femoral condyle, anterior subregion of tibia, central subregion of tibia, and posterior subregion of tibia. The maximum possible score in LTF joint was 15 and similarly for the MTF joint. The PF joint included four subregions: medial (M) portion of the patella, lateral (L) portion of the patella, medial trochlea (T) portion of femur, and lateral trochlea (T) portion of femur. The maximum possible BML size in PF joint was 12. LTF joint = lateral tibiofemoral joint, MTF joint = medial tibiofemoral joint, PF joint = patellofemoral joint.

TABLE 1: Patient characteristics.

Characteristics (n = 98)	
Gender (male/female)	57/41
Mean age (years)	65.1 ± 5.7
BMI (kg/m <sup>2</sup> )	28.1 ± 3.4
Smoking (%)	26 (26.5)
Alcohol consumption (%)	31 (31.6)
Hypertension (%)	45 (45.9)
Diabetes mellitus (%)	19 (19.4)
Degree of correction (°)	10.5 ± 1.6
Osteotomy gap (mm)	11.3 ± 2.8

BMI: body mass index.

ICC for all parameters were more than 0.8 in MRI measurements. As shown in Table 3, there was no significant difference between the pre- and postoperative total BML size in PF and LTF joints. We observed significant improvements in the total BML size in MTF joint (Figure 4). There were no significant differences between smoker and nonsmoker regarding the total BML size in PF joint, LTF joint, and MTF joint ( $p > 0.05$ ).

**3.3. The Relationship between Postoperative BMLs and Pain Severity.** All patients had preoperative MTF joint BMLs; in contrast, 13 patients had no postoperative MTF joint BMLs. The VAS scores were  $1.7 \pm 0.8$  in patients with postoperative MTF joint BMLs and  $0.2 \pm 0.4$  in patients without postoperative MTF joint BMLs. The KOOS pain scores were  $76.9 \pm 9.0$  in patients with postoperative MTF joint BMLs and

$87.8 \pm 5.4$  in patients without postoperative MTF joint BMLs. The independent sample *t*-test showed that the VAS scores and KOOS pain scores improved better in patients without postoperative MTF joint BMLs (Figure 5).

Pearson correlations showed that postoperative MTF joint BMLs are correlated with postoperative VAS ( $r = 0.945$ ,  $p < 0.001$ ) and postoperative KOOS pain ( $r = -0.472$ ,  $p < 0.001$ ).

#### 4. Discussion

KOA is a degenerative joint disease, which causes pain and decreased physical function [1–4]. Knee pain is the major cause leading individuals to seek medical care [5–8]. The goal of KOA management is to relieve pain, improve knee function, and change the disease process. Although there are no approved drugs for changing KOA process, many interventions are available to address pain and function [1, 20, 21].

Many studies have demonstrated that knee pain could be relieved by OWHTO [9–12]. A systematic review found that clinical scores improved significantly after open or closed wedge high tibial osteotomy, including VAS, the American Knee Society Score, Hospital for Special Surgery Knee Score, and Lysholm score [22]. Identification of the mechanism of pain relief by OWHTO is important, an understanding of the mechanism may be helpful for targeted anti-KOA therapy and individualized therapy.

Some studies have shown that BMLs were more commonly observed in painful knees with OA than non-painful knees [13, 14]. Meanwhile, increased BMLs were strongly associated with new onset frequent knee pain in nonpainful knees [23]. Zhang et al. investigated 651

TABLE 2: Preoperative and postoperative VAS and KOOS scores.

	Preoperative	Postoperative	<i>p</i> value
VAS	6.1 ± 0.8	1.5 ± 0.9	<0.001
KOOS pain	46.5 ± 8.2	78.3 ± 9.4	<0.001
KOOS symptoms	51.2 ± 9.1	81.5 ± 7.8	<0.001
KOOS ADL	52.4 ± 10.9	78.8 ± 8.2	<0.001
KOOS SR	30.3 ± 9.1	64.2 ± 7.2	<0.001
KOOS QoL	39.4 ± 9.2	60.9 ± 8.4	<0.001

KOOS: Knee injury and Osteoarthritis Outcome Score; ADL: activities of daily living; SR: sport and recreation; QoL: quality of life.




FIGURE 3: VAS scores and all subscales of KOOS improved significantly after surgery. KOOS: Knee injury and Osteoarthritis Outcome Score; ADL: activities of daily living; SR: sport and recreation; QoL: quality of life.

TABLE 3: Preoperative and postoperative MOAKS.

	Preoperative	Postoperative	<i>p</i> value
PF joint	1.3 ± 0.8	1.3 ± 0.7	0.566
LTF joint	1.8 ± 0.8	1.9 ± 0.7	0.083
MTF joint	7.9 ± 2.2	3.4 ± 2.0	<0.001

MOAKS: MRI Osteoarthritis Knee Score; PF joint: patellofemoral joint; MTF joint: medial tibiofemoral joint; LTF joint: lateral tibiofemoral joint.

painful knees and demonstrated that improved BMLs were strongly associated with pain relief in KOA [24]. These and other studies have implied that BMLs are the result of mechanical overload and thus inducing pain in KOA [5, 25–27].

The primary purpose of OWHTO was to reduce the mechanical load of medial compartment of the knee. Interestingly, a recent study implied that reducing mechanical

load can decrease BMLs and relieve knee pain [28]. Our study demonstrates that MTF joint BMLs improved significantly after OWHTO, the greater the improvement in postoperative MTF joint BMLs, the less pain.

This study has several limitations: first, it is a small retrospective study. Second, the research is a single-center study; a further larger, multicenter research was needed to confirm our findings.




FIGURE 4: The total BML size in MTF joint improved significantly after surgery. BMLs: bone marrow lesions; PF joint: patellofemoral joint; MTF joint: medial tibiofemoral joint; LTF joint: lateral tibiofemoral joint.




FIGURE 5: The VAS scores and KOOS pain scores improved better in patients without postoperative MTF joint BMLs. KOOS: Knee injury and Osteoarthritis Outcome Score; BMLs: bone marrow lesions.

### 5. Conclusions

We confirmed that the presence of postoperative MTF joint BMLs are strongly associated with pain severity. The greater the improvement in postoperative MTF joint BMLs, the less pain. Our findings provide valuable information of OWHTO in the treatment of knee osteoarthritis (KOA) and potential future directions for KOA treatment approaches.

### Data Availability

The datasets generated and analyzed during the present study are available from the corresponding author on reasonable request.

### Disclosure

This manuscript has been released as a preprint.



## Conflicts of Interest

The authors have no conflicts of interest related to the submission of this manuscript.

## Authors' Contributions

Bo Zhu and Tong-Fu Wang contributed equally to this work and should be considered as equal to co-first authors.

## References

- [1] N. Goris, "Knee osteoarthritis," *Journal of Physiotherapy*, vol. 63, no. 3, p. 188, 2017.
- [2] E. M. Roos and N. K. Arden, "Strategies for the prevention of knee osteoarthritis," *Nature reviews Rheumatology*, vol. 12, no. 2, pp. 92–101, 2016.
- [3] C. Fingleton, K. Smart, N. Moloney, B. M. Fullen, and C. Doody, "Pain sensitization in people with knee osteoarthritis: a systematic review and meta-analysis," *Osteoarthritis and Cartilage*, vol. 23, no. 7, pp. 1043–1056, 2015.
- [4] M. de Rooij, M. van der Leeden, M. W. Heymans et al., "Prognosis of pain and physical functioning in patients with knee osteoarthritis: a systematic review and meta-analysis," *Arthritis Care & Research*, vol. 68, no. 4, pp. 481–492, 2016.
- [5] T. W. O'Neill and D. T. Felson, "Mechanisms of osteoarthritis (OA) pain," *Current Osteoporosis Reports*, vol. 16, no. 5, pp. 611–616, 2018.
- [6] T. Neogi, "The epidemiology and impact of pain in osteoarthritis," *Osteoarthritis and Cartilage*, vol. 21, no. 9, pp. 1145–1153, 2013.
- [7] A. Salma and D. Paul, "The natural history of disability and its determinants in adults with lower limb musculoskeletal pain," *The Journal of Rheumatology*, vol. 36, no. 3, pp. 583–591, 2009.
- [8] N. M. Hadler, "Knee pain is the malady—not osteoarthritis," *Annals of Internal Medicine*, vol. 116, no. 7, pp. 598–599, 1992.
- [9] P. Niemeyer, H. Schmal, O. Hauschild, J. von Heyden, N. P. Südkamp, and W. Köstler, "Open-Wedge Osteotomy Using an Internal Plate Fixator in Patients With Medial-Compartment Gonarthrosis and Varus Malalignment: 3-Year Results With Regard to Preoperative Arthroscopic and Radiographic Findings," *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, vol. 26, no. 12, pp. 1607–1616, 2010.
- [10] S. Schröter, C. E. Gonsler, L. Konstantinidis, P. Helwig, and D. Albrecht, "High complication rate after biplanar open wedge high tibial osteotomy stabilized with a new spacer plate (position HTO plate) without bone substitute," *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, vol. 27, no. 5, pp. 644–652, 2011.
- [11] N. R. Howells, L. Salmon, A. Waller, J. Scanelli, and L. A. Pinczewski, "The outcome at ten years of lateral closing-wedge high tibial osteotomy: determinants of survival and functional outcome," *The Bone & Joint Journal*, vol. 96-B, no. 11, pp. 1491–1497, 2014.
- [12] L. Kohn, M. Sauerschnig, S. Iskandar et al., "Age does not influence the clinical outcome after high tibial osteotomy," *Knee surgery, Sports Traumatology, Arthroscopy: official journal of the ESSKA*, vol. 21, no. 1, pp. 146–151, 2013.
- [13] D. T. Felson, C. E. Chaisson, C. L. Hill et al., "The association of bone marrow lesions with pain in knee osteoarthritis," *Annals of Internal Medicine*, vol. 134, no. 7, pp. 541–549, 2001.
- [14] A. J. Barr, T. Campbell, D. Hopkinson, S. R. Kingsbury, M. A. Bowes, and P. G. Conaghan, "A systematic review of the relationship between subchondral bone features, pain and structural pathology in peripheral joint osteoarthritis," *Arthritis Research & Therapy*, vol. 17, no. 1, p. 228, 2015.
- [15] M. L. Davies-Tuck, A. E. Wluka, Y. Wang, D. R. English, G. G. Giles, and F. Cicuttini, "The natural history of bone marrow lesions in community-based adults with no clinical knee osteoarthritis," *Annals of the Rheumatic Diseases*, vol. 68, no. 6, pp. 904–908, 2009.
- [16] H. Wada, K. Aso, and M. Ikeuchi, "MRI findings change after medial opening-wedge high tibial osteotomy for medial knee osteoarthritis," *Osteoarthritis and Cartilage*, vol. 28, pp. S381–S382, 2020.
- [17] J. M. Schulz, T. B. Birmingham, W. P. Maksymowych et al., "Reliability and sensitivity to change of bone marrow lesion scores using the knee inflammation MRI scoring system (KIMRISS) before and after high tibial osteotomy," *Osteoarthritis and Cartilage*, vol. 28, pp. S291–S291, 2020.
- [18] M. S. Kim, I. J. Koh, S. Sohn, H. S. Sung, and Y. in, "Degree of preoperative subchondral bone marrow lesion is associated with postoperative outcome after medial opening wedge high tibial osteotomy," *American Journal of Sports Medicine*, vol. 47, no. 10, pp. 2454–2463, 2019.
- [19] H.-Y. Yang, S.-J. Kang, W.-K. Kwak, E.-K. Song, and J.-K. Seon, "The influence of preoperative tibial bone marrow edema on outcomes after medial opening-wedge high tibial osteotomy," *The Journal of Bone and Joint Surgery American*, vol. 102, no. 23, pp. 2068–2076, 2020.
- [20] R. R. Bannuru, M. C. Osani, E. E. Vaysbrot et al., "OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis," *Osteoarthritis and Cartilage*, vol. 27, no. 11, pp. 1578–1589, 2019.
- [21] O. Bruyère, G. Honvo, N. Veronese et al., "An updated algorithm recommendation for the management of knee osteoarthritis from the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO)," *Seminars in Arthritis and Rheumatism*, vol. 49, no. 3, pp. 337–350, 2019.
- [22] X. Cheng, F. Liu, F. Xiong, Y. Huang, and A. C. Paulus, "Radiographic changes and clinical outcomes after open and closed wedge high tibial osteotomy: a systematic review and meta-analysis," *Journal of Orthopaedic Surgery and Research*, vol. 14, no. 1, p. 179, 2019.
- [23] D. T. Felson, J. Niu, A. Guermazi et al., "Correlation of the development of knee pain with enlarging bone marrow lesions on magnetic resonance imaging," *Arthritis and Rheumatism*, vol. 56, no. 9, pp. 2986–2992, 2007.
- [24] Y. Zhang, M. Nevitt, J. Niu et al., "Fluctuation of knee pain and changes in bone marrow lesions, effusions, and synovitis on magnetic resonance imaging," *Arthritis and Rheumatism*, vol. 63, no. 3, pp. 691–699, 2011.
- [25] F. Pan, J. Tian, D. Aitken, F. Cicuttini, and G. Jones, "Predictors of pain severity trajectory in older adults: a 10.7-year follow-up study," *Osteoarthritis and Cartilage*, vol. 26, no. 12, pp. 1619–1626, 2018.
- [26] J. Wang, B. Antony, Z. Zhu et al., "Association of patellar bone marrow lesions with knee pain, patellar cartilage defect and patellar cartilage volume loss in older adults: a cohort study," *Osteoarthritis and Cartilage*, vol. 23, no. 8, pp. 1330–1336, 2015.

- [27] E. Kon, M. Ronga, G. Filardo et al., "Bone marrow lesions and subchondral bone pathology of the knee," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 24, no. 6, pp. 1797–1814, 2016.
- [28] M. J. Callaghan, M. J. Parkes, C. E. Hutchinson et al., "A randomised trial of a brace for patellofemoral osteoarthritis targeting knee pain and bone marrow lesions," *Annals of the Rheumatic Diseases*, vol. 74, no. 6, pp. 1164–1170, 2015.