

## Research Article

# Effect of Cerium Oxide on Kidney and Lung Tissue in Rats with Testicular Torsion/Detorsion

Aycan Ozdemirkan <sup>1</sup>, Ali Can Kurtipek <sup>2</sup>, Aysegül Kucuk <sup>3</sup>, Cagri Ozdemir <sup>1</sup>,  
Suleyman Yesil <sup>4</sup>, Saban Cem Sezen <sup>5</sup>, Mustafa Kavutcu <sup>6</sup>, and Mustafa Arslan <sup>1,7</sup>

<sup>1</sup>Faculty of Medicine, Department of Anesthesiology and Reanimation, Gazi University, Ankara, Turkey

<sup>2</sup>Department of Internal Medicine, Ankara City Hospital Health Sciences University, Ankara, Turkey

<sup>3</sup>Faculty of Medicine, Department of Physiology, Kütahya Health Sciences University, Kütahya, Turkey

<sup>4</sup>Faculty of Medicine, Department of Urology, Gazi University, Ankara, Turkey

<sup>5</sup>Faculty of Medicine, Department of Histology and Embryology, Kırıkkale University, Kırıkkale, Turkey

<sup>6</sup>Faculty of Medicine, Department of Medical Biochemistry, Gazi University, Ankara, Turkey

<sup>7</sup>Life Sciences Application and Research Center, Gazi University, Ankara, Turkey

Correspondence should be addressed to Mustafa Arslan; mustarlan@gmail.com

Received 16 December 2021; Revised 23 February 2022; Accepted 2 March 2022; Published 22 March 2022

Academic Editor: Ahmet Özer Sehirli

Copyright © 2022 Aycan Ozdemirkan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Introduction.** Testicular torsion is a surgical emergency that results in testicular ischemia as a result of rotation of the spermatic cord around itself. Oxidative damage occurs in the testis and distant organs with the overproduction of free radicals and overexpression of proinflammatory cytokines by reperfusion after surgery. In this study, we aimed to investigate the effects of cerium oxide (CeO<sub>2</sub>), an antioxidant nanoparticle, on lung and kidney tissues in testicular torsion/detorsion (T/D) in rats. **Materials and Methods.** After ethics committee approval, 24 rats were equally (randomly) divided into 4 groups. Left inguinoscrotal incision was performed in the control (C) group. In group CeO<sub>2</sub>, 0.5 mg/kg CeO<sub>2</sub> was given intraperitoneally 30 minutes before inguinoscrotal incision. In group T/D, unilateral testicular T/D was achieved by performing an inguinoscrotal incision and rotating the left testis 720° clockwise, remaining ischemic for 120 minutes, followed by 120 minutes of reperfusion. In group CeO<sub>2</sub>-T/D, 0.5 mg/kg CeO<sub>2</sub> was given intraperitoneally 30 minutes before testicular T/D. At the end of the experiment, lung and kidney tissues were removed for histopathological and biochemical examinations. **Results.** Glomerular vacuolization (GV), tubular dilatation (TD), tubular cell degeneration and necrosis (TCDN), leukocyte infiltration (LI), and tubular cell spillage (TCS) in renal tissue were significantly different between groups ( $p = 0.012$ ,  $p = 0.049$ ,  $p < 0.003$ ,  $p = 0.046$ , and  $p = 0.049$ , respectively). GV and TCDN were significantly decreased in group CeO<sub>2</sub>-T/D compared to group T/D ( $p = 0.042$  and  $p = 0.029$ , respectively). Lung tissue neutrophil infiltration, alveolar thickening, and total lung injury score (TLIS) were significantly different between groups ( $p = 0.006$ ,  $p = 0.001$ , and  $p = 0.002$ , respectively). Neutrophil infiltration and TLIS were significantly decreased in group CeO<sub>2</sub>-T/D compared to group T/D ( $p = 0.013$  and  $p = 0.033$ , respectively). Lung and kidney tissue oxidative stress parameters were significantly different between groups ( $p < 0.05$ ). Renal tissue glutathione-S-transferase (GST), catalase (CAT), and paraoxonase (PON) activities were significantly higher, and malondialdehyde (MDA) levels were significantly lower in group CeO<sub>2</sub>-T/D than in group T/D ( $p = 0.049$ ,  $p = 0.012$ ,  $p < 0.001$ , and  $p = 0.004$ , respectively). GST and PON activities were higher, and MDA levels were lower in group CeO<sub>2</sub>-T/D than in group T/D in the lung tissue ( $p = 0.002$ ,  $p < 0.001$ , and  $p = 0.008$ , respectively). **Discussion.** In our study, cerium oxide was shown to reduce histopathological and oxidative damage in the lung and kidney tissue in a rat testis torsion/detorsion model.

## 1. Introduction

Testicular torsion is a surgical emergency that accounts for 26% of acute testicular pain causes [1]. Testicular torsion results in ischemia by interruption of venous and arterial blood flow in the scrotum as a result of the rotation of the spermatic cord around itself. The incidence of testicular torsion peaks in men aged 12 and 18 years but can be seen in any age group [2]. Its incidence has been reported as 3.5 per 100000. Testicular torsion is mostly idiopathic and can occur with trauma in 20% of cases [3]. In testicular torsion, damage to the testis may occur depending on the degree and duration of rotation [4, 5]. When surgical detorsion in the treatment of testicular torsion is performed within 4-6 hours, the testis can be saved in the vast majority of cases [2]. However, with the detorsion, reperfusion will occur in the ischemic testicular tissue. At the end of the ischemia-reperfusion process, free radical production will increase.

Overproduction of free radicals causes irreversible DNA damage, changes in enzyme activities, damage to proteins, and many damages in the organism, such as the formation of new immunological structures [6]. It has also been shown to cause germ cell apoptosis [1]. Free radicals cause lipid peroxidation, especially by acting on fatty acids in the cell membrane. Lipid peroxidation is a very harmful chemical chain reaction that alters the membrane lipid structure and indirectly produces reactive aldehydes and damages the structure and functions of other cell components. Once this reaction starts autocatalytically, it continues as a chain and if not prevented, it destroys the cell membrane, breaks down the organelles, and causes the release of lysosomal enzymes and autolysis [6]. For this reason, testicular detorsion should be considered as a typical ischemia-reperfusion (I/R) injury. In addition, reactive oxygen radicals and neutrophil infiltration that occur during reperfusion play a role in the pathogenesis of distant organ injuries [7]. There are several agents used experimentally to reduce I/R injury after testicular torsion/detorsion (T/D) such as taurine [8–11], modafinil [12], sildenafil [13], and glutathione [14]. These treatments are aimed at eliminating or reducing the oxidative damage that occurs in I/R injury in the testis.

Many pathologies with associated oxidative damage respond to compounds that can scavenge reactive oxygen species (ROS). One of these compounds is cerium oxide ( $\text{CeO}_2$ ), an antioxidant nanoparticle that mimics the activities of catalase and superoxide dismutase enzymes and has scavenging properties of radical oxygen species; its efficacy has been demonstrated experimentally in many pathologies with oxidative damage [15–21]. Based on the findings of experimental studies, it is known that it reduces oxidative damage in I/R models and plays a protective role in distant organ damage [22–24]. Although there is little information in the literature about the effects of cerium on testicular tissue, it is not known whether it has a protective effect on distant organs in testicular ischemia-reperfusion injury [25–27]. The aim of this study is to show the effects of cerium oxide on kidney and lung tissues in testicular T/D in rats.

## 2. Materials and Methods

**2.1. Animals and Experimental Protocol.** This study was approved by the Gazi University Ethics Committee (Ethics number: G.U.ET-21-060). All experiments were performed according to standards of Guide for the Care and Use of Laboratory Animals in the Gazi University Animal Laboratory. A total of 24 male Wistar albino rats (4 rats in each cage) weighing between 250 and 300 grams were used. Rats were kept in a temperature-controlled ( $21 \pm 1^\circ\text{C}$ ) and humidity-controlled (45–55%) room, which was maintained on a 12/12 reversed light cycle. Animals were fed with a standard pellet and allowed to drink water ad libitum. Animals were equally (randomly) divided into four equal groups ( $n = 6$  each): (1) control (C), (2) cerium oxide ( $\text{CeO}_2$ ), (3) torsion/detorsion (T/D), and (4) cerium oxide-T/D ( $\text{CeO}_2$ -T/D). Initially, all rats were anesthetized with 50 mg/kg intraperitoneal (ip) ketamine hydrochloride (VetaKetam, Vetagro) and ip 10 mg/kg xylazine hydrochloride (Rompun %2, BAYER).

Surgical procedures for testis torsion/detorsion are as follows: during the surgical procedures, rats were placed on a heating pad in order to maintain the body temperature. After the skin was shaved and washed with an antiseptic solution, left inguinoscrotal incision was performed using a sterile scalpel and forceps. Following exposing the left testis, unilateral testicular torsion was created by rotating the left testis  $720^\circ$  in a clockwise direction and fixing it within the hemiscrotum using a 4/0 atraumatic silk suture. Ischemia was observed with a dark purple color that appeared in the testicular tissue. Then, the skin was closed with 5/0 silk sutures and the testis was kept in torsion for 120 minutes. After 120 minutes, the incision was reopened by removing the sutures. The spermatic cord was detorsed; the testis was fixed within the hemiscrotum using a 4/0 atraumatic silk suture. Reperfusion was observed with the reappearance of pinkness in the testicular tissue. Heparin sodium (500 IU/kg, Nevparin, Mustafa Nevzat) was administered through the tail vein for the maintenance of reperfusion after occlusion. Then, the skin was closed with 5/0 silk sutures. The reperfusion phase was maintained for 120 minutes.

- (1) Control group (C): rats were only subjected to left inguinoscrotal incision
- (2) Cerium oxide group ( $\text{CeO}_2$ ): a left inguinoscrotal incision was performed. Cerium oxide was given (0.5 mg/kg, ip) 30 minutes before the incision. Cerium oxide was given at the predetermined dose [16]
- (3) Torsion-detorsion group (T/D): testis torsion/detorsion was performed surgically as described
- (4) Cerium oxide-torsion-detorsion group ( $\text{CeO}_2$ -T/D): testis torsion/detorsion was performed surgically as described. Cerium oxide was given (0.5 mg/kg ip) 30 minutes before the ischemia period

$\text{CeO}_2$  aqueous nanoparticle dispersion (<5 nm particle size, 20 wt% in  $\text{H}_2\text{O}$ , 99.5% trace metals) was obtained from Sigma-Aldrich® (St Louis, MO, USA).

At the end of the experiments, rats were sacrificed under anesthesia. The lung and kidney tissues were excised for biochemical and histopathological analysis.

**2.2. Histopathological Analysis.** Lung tissue samples were removed and fixed in 10% neutral formalin solution. Then, the lungs were examined with light microscopy by the same pathologist, who was blinded to the study. A total of 10 random areas were evaluated with 200–400 times magnified microscopy in hematoxylin and eosin- (H&E-) stained sections. Stained slides were examined under a light microscope. Neutrophil infiltration and alveolar thickness are measured in each specimen for exposing the degree of lung injury area. Each parameter was scored as any (0 point), only a little (1 point), medium amount (2 points), or severe (3 points). The two scores were added and noted as total lung injury score (TLIS).

After routine fixation process, kidney specimens were embedded in paraffin blocks; then, tissue sections of  $5\ \mu$  were mounted on slides for staining with hematoxylin and eosin (H&E). Histopathological evaluation under light microscopy was performed, and findings were scored using a scoring system by Bostan et al. [28]. Glomerular vacuolization (GV), tubular dilatation (TD), vascular vacuolization and hypertrophy (VVH), tubular cell degeneration and necrosis (TCDN), Bowman space dilatation (BSD), tubular hyaline cylinder (THC), leucocyte infiltration (LI), and tubular cell spillage (TCS) were scored using a scoring system: 0: no change; +1: minimal change; +2: medium; and +3: severe.

**2.3. Biochemical Analysis.** The lung and kidney tissues were first washed with cold NaCl solution (0.154M) to discard blood contamination and then homogenized in Diac 900 (Heidolph Instruments GmbH & Co KG, Schwabach, Germany) at 1000 U for about 3 min. After centrifugation at  $10,000 \times g$  for about 60 min, the upper clear layer was taken.

For the measurement of malondialdehyde (MDA) levels, thiobarbituric acid (TBA) reactive substances assay was performed by the method described by Van Ye et al. [29]. The reaction with TBA at 80–90°C was used to determine the MDA level, as MDA or similar substances react with TBA and produce a pink pigment that has an absorption of maximum 532 nm. To ensure protein precipitation, the sample in room temperature was mixed with cold 20% (*w/v*) trichloroacetic acid and the precipitate was then centrifuged for 10 min at 3000 rpm at room temperature to form a pellet. An aliquot of the supernatant was then placed into an equal volume of 0.6% (*w/v*) TBA in a boiling water bath for 30 min. Following cooling, sample and blank absorbance was read at 532 nm and the results were expressed as nanomole per milligram of protein, based on a graph where 1,1,3,3-tetramethoxypropane had been used as our MDA standard.

The catalase (CAT) activity is based on the measurement of absorbance decrease due to  $H_2O_2$  consumption at 240 nm as described by the Aebi H method [30].

Glutathione-s-transferase (GST) enzyme activity was measured using the method described by Habig et al. [31]. The GST activity method is based on the measurement of

absorbance increase at 340 nm due to the reduction of dinitrophenyl glutathione (DNPG). The results were expressed in international unit per milligram of protein.

Paraoxonase- (PON-) 1 activity was measured as the rate of hydrolysis of paraoxon by monitoring the increase of absorbance at 405 nm and at 25°C described by Eckerson et al.'s method [32].

**2.4. Statistical Analysis.** Statistical Package for the Social Sciences (SPSS, Chicago, IL, USA) 20.0 for Windows was used. Each categorical variable was analyzed by the Kolmogorov-Smirnov test. Biochemical and histopathological parameters were tested by using the Kruskal-Wallis test, Bonferroni correction test, and Mann-Whitney *U* test. A statistical value of less than 0.05 was considered significant. All values were expressed as mean  $\pm$  standard error (mean  $\pm$  SE).

### 3. Results

GV, TD, TCDN, LI, and TCS in the renal tissue were significantly different between groups ( $p = 0.012$ ,  $p = 0.049$ ,  $p < 0.003$ ,  $p = 0.046$ , and  $p = 0.049$ , respectively). VVH, BSD, and THC were similar among groups ( $p = 0.107$ ,  $p = 0.434$ , and  $p = 0.053$ , respectively). All renal histopathological findings, except VVH, BSD, and THC, were higher in group T/D compared to groups C and  $CeO_2$  ( $p > 0.05$ ). GV and TCDN were significantly decreased in group  $CeO_2$ -T/D compared to group T/D ( $p = 0.042$  and  $p = 0.029$ , respectively) (Table 1, Figures 1–4).

According to neutrophil infiltration/aggregation in lung tissue, there were significant differences between the groups ( $p = 0.006$ ). Neutrophil infiltration/aggregation was significantly higher in the T/D group compared to groups C and  $CeO_2$  ( $p = 0.001$  and  $p = 0.004$ , respectively) and was lower in group  $CeO_2$ -T/D compared to group T/D ( $p = 0.013$ ). The alveolar wall thickness was significantly higher in group T/D group than in group C and  $CeO_2$  ( $p < 0.001$  and  $p = 0.001$ , respectively). Also, TLISs were significantly higher in group T/D compared to groups C and  $CeO_2$  ( $p < 0.001$  and  $p = 0.002$ , respectively). TLISs were significantly lower in group  $CeO_2$ -T/D compared to group T/D ( $p = 0.033$ ) (Table 2 and Figures 5–8).

There were significant differences between groups when compared in terms of lung tissue GST, CAT, and PON enzyme activities and MDA levels ( $p = 0.002$ ,  $p = 0.024$ ,  $p = 0.008$ , and  $p < 0.001$ ). GST enzyme activity was significantly lower in group T/D compared to groups C and  $CeO_2$  ( $0.74 \pm 0.14$  vs.  $1.70 \pm 0.22$  IU/mg protein,  $p < 0.001$ , and  $0.74 \pm 0.14$  vs.  $1.42 \pm 0.13$  IU/mg protein,  $p = 0.004$ , respectively). It was significantly higher in group  $CeO_2$ -T/D when compared to group T/D ( $1.37 \pm 0.10$  vs.  $0.74 \pm 0.14$  IU/mg protein,  $p = 0.007$ ). CAT enzyme activity was significantly lower in group T/D compared to groups C and  $CeO_2$  ( $526.40 \pm 37.65$  vs.  $2038.02 \pm 575.14$  IU/mg protein,  $p = 0.004$ , and  $526.40 \pm 37.65$  vs.  $1610.35 \pm 229.98$  IU/mg protein,  $p = 0.028$ , respectively). Groups  $CeO_2$ -T/D and C were similar in terms of CAT enzyme activity ( $1476.70 \pm 184.00$  vs.  $2038.02 \pm 575.14$  IU/mg protein,  $p = 0.051$ ). MDA levels were significantly lower in group T/D compared to groups

TABLE 1: Histopathological findings in renal tissue (mean  $\pm$  SE).

	Group C (n = 6)	Group CeO <sub>2</sub> (n = 6)	Group T/D (n = 6)	Group CeO <sub>2</sub> -T/D (n = 6)	<i>p</i> **
Glomerular vacuolization (GV)	0.17 $\pm$ 0.17	0.33 $\pm$ 0.21	1.33 $\pm$ 0.33* <sup>†</sup>	0.67 $\pm$ 0.21 <sup>‡</sup>	0.012
Tubular dilatation (TD)	0.33 $\pm$ 0.21	0.50 $\pm$ 0.22	1.17 $\pm$ 0.17* <sup>†</sup>	0.67 $\pm$ 0.21	0.049
Vascular vacuolization and hypertrophy (VVH)	0.33 $\pm$ 0.21	0.50 $\pm$ 0.22	1.00 $\pm$ 0.00	0.50 $\pm$ 0.22	0.107
Tubular cell degeneration and necrosis (TCDN)	0.17 $\pm$ 0.17	0.33 $\pm$ 0.21	1.33 $\pm$ 0.21* <sup>†</sup>	0.67 $\pm$ 0.21 <sup>‡</sup>	0.003
Bowman space dilatation (BSD)	0.33 $\pm$ 0.21	0.33 $\pm$ 0.21	0.83 $\pm$ 0.31	0.50 $\pm$ 0.22	0.434
Tubular hyaline cylinder (THC)	0.50 $\pm$ 0.22	0.33 $\pm$ 0.21	1.00 $\pm$ 0.00	0.83 $\pm$ 0.17	0.053
Lymphocyte infiltration (LI)	0.33 $\pm$ 0.21	0.33 $\pm$ 0.21	1.33 $\pm$ 0.21* <sup>†</sup>	0.83 $\pm$ 0.31	0.046
Tubular cell spillage (TCS)	0.33 $\pm$ 0.21	0.50 $\pm$ 0.22	1.17 $\pm$ 0.17* <sup>†</sup>	0.67 $\pm$ 0.21	0.049

*p*\*\* : significance level with the Kruskal-Wallis test  $p < 0.05$ . \*  $p < 0.05$ : compared to group C. <sup>†</sup> $p < 0.05$ : compared to group CeO<sub>2</sub>. <sup>‡</sup> $p < 0.05$ : compared to group T/D.




FIGURE 1: Renal tissue in the control group. RC: renal cortex; pt: proximal tubule; dt: distal tubule; g: glomerular; arrowhead: bowman space; arrow: dilated tubule; a: artery (H&E  $\times$ 100).




FIGURE 2: Renal tissue in the cerium oxide group. pt: proximal tubule; dt: distal tubule; g: glomerular; arrow: dilated tubule (H&E  $\times$ 100).

C and CeO<sub>2</sub> (11.97  $\pm$  0.62 vs. 7.97  $\pm$  0.49 nmol/mg protein,  $p < 0.001$ , and 11.97  $\pm$  0.62 vs. 8.85  $\pm$  0.63 nmol/mg protein,  $p = 0.001$ , respectively). MDA levels were also significantly lower in group CeO<sub>2</sub>-T/D when compared to group T/D (7.98  $\pm$  0.58 vs. 11.97  $\pm$  0.62 nmol/mg protein,  $p < 0.001$ ). PON enzyme activity was significantly lower in group T/D compared to groups C and CeO<sub>2</sub> (2.18  $\pm$  0.43 vs. 9.09  $\pm$  2.10 IU/mg protein,  $p = 0.002$  and 2.18  $\pm$  0.43 vs. 8.60  $\pm$  1.46 IU/mg protein,  $p = 0.004$ , respectively). It was significantly higher in group CeO<sub>2</sub>-T/D when compared to group T/D (6.62  $\pm$  1.00 vs. 2.18  $\pm$  0.43 IU/mg protein,  $p = 0.034$ ) (Figure 9).

Renal tissue GST, CAT, and PON enzyme activities and MDA levels were significantly different between groups ( $p = 0.049$ ,  $p = 0.012$ ,  $p = 0.004$ , and  $p < 0.001$ , respectively). GST enzyme activity was significantly lower in group T/D compared to groups C and CeO<sub>2</sub> (1.70  $\pm$  0.14 vs. 2.73  $\pm$  0.36 IU/mg protein,  $p = 0.009$ , and 1.70  $\pm$  0.14 vs. 2.46  $\pm$  0.27 IU/mg protein,  $p = 0.047$ , respectively). It was significantly higher in group CeO<sub>2</sub>-T/D when compared to group T/D (2.50  $\pm$  0.18 vs. 1.70  $\pm$  0.14 IU/mg protein,  $p = 0.037$ ). CAT enzyme activity was significantly lower in group T/D compared to groups C and CeO<sub>2</sub> (1976.54  $\pm$  237.91 vs. 4715.22  $\pm$  701.68 IU/mg protein,  $p = 0.002$ , and 1976.54  $\pm$  237.91 vs. 3861.30  $\pm$  588.71 IU/mg protein,  $p = 0.022$ , respectively). It was significantly higher in group CeO<sub>2</sub>-T/

D when compared to group T/D (3992.94  $\pm$  503.78 vs. 1976.54  $\pm$  237.91 IU/mg protein,  $p = 0.015$ ). MDA levels were significantly higher in group T/D compared to groups C and CeO<sub>2</sub> (9.21  $\pm$  0.47 vs. 5.44  $\pm$  0.24 nmol/mg protein,  $p < 0.001$ , and 9.21  $\pm$  0.47 vs. 6.53  $\pm$  0.73 nmol/mg protein,  $p = 0.001$ , respectively). It was significantly lower in group CeO<sub>2</sub>-T/D when compared to group T/D (6.35  $\pm$  0.19 vs. 9.21  $\pm$  0.47 nmol/mg protein,  $p < 0.001$ ). PON enzyme activity was significantly lower in group T/D compared to groups C and CeO<sub>2</sub> (3.36  $\pm$  0.49 vs. 8.28  $\pm$  1.19 IU/mg protein,  $p = 0.001$ , and 3.36  $\pm$  0.49 vs. 7.68  $\pm$  0.82 IU/mg protein,  $p = 0.003$ , respectively). It was significantly higher in group CeO<sub>2</sub>-T/D when compared to group T/D (6.19  $\pm$  0.91 vs. 3.36  $\pm$  0.49 IU/mg protein,  $p = 0.036$ ) (Figure 10).

#### 4. Discussion

In this study, we evaluated the effects of CeO<sub>2</sub> on lung and renal tissue following testicular T/D in rats. While there are few data on the effects of CeO<sub>2</sub> in testicular I/R injury in the literature, the effects of CeO<sub>2</sub> on distant organ injury in this I/R model are unknown. In order to investigate the effects of CeO<sub>2</sub> in distant organ damage, we examined both histopathological and oxidative damage parameters in the lung and kidney in I/R injury. At the end of our study, it was shown that the use of CeO<sub>2</sub> in the testicular T/D model




FIGURE 3: Renal tissue in the testicular torsion/detorsion group. pt: proximal tubule; dt: distal tubule; vc: vascular congestion; conj: capillary congestion; g: glomerulus; arrow: dilated tubule; m: macula densa; a: artery; \*: degenerated glomerulus; inf: inflammation (H&E  $\times 100$ ).




FIGURE 4: Renal tissue in the testis torsion/detorsion treated with cerium oxide group. pt: proximal tubule; dt: distal tubule; g: glomerulus; a: artery; arrow: dilated tubule; double arrow: macula densa; inf: inflammation (H&E  $\times 100$ ).

significantly reduced both histopathological damage and oxidative damage in the lung and kidney.

Oxidative stress is a pathological process that occurs with the disruption of the balance between the production and accumulation of reactive oxygen radicals in cells and tissues [33]. Reactive oxygen radicals are the most important free radicals formed from oxygen in biological systems. ROS create lipid peroxidation by acting on fatty acids in the cell membrane. Lipid peroxidation causes cell apoptosis and necrosis. MDA is one of the secondary products formed during lipid peroxidation and is the most commonly used oxidative stress marker. Oxidative stress due to increased ROS during ischemia-reperfusion injury causes changes in mitochondrial oxidative phosphorylation, ATP depletion, increase in intracellular calcium, and increase in proteases and phosphatases [34]. Among the mechanisms against this oxidative damage, glutathione S-transferase plays an important role. It is responsible for the inactivation of electrophilic components and toxic substrates [35]. Catalase and paraoxonase are antioxidant enzymes. Paraoxonase is an ester hydrolase enzyme. It is named paraoxonase because it can hydrolyze paraoxon, the toxic metabolite of parathion, an organophosphate pesticide. In recent years, it has gained popularity due to its possible antioxidant effects [36]. Cata-

lase is one of the first antioxidant defense mechanisms in the cell. It provides detoxification of hydrogen peroxide, which passes from mitochondria to the cytosol during oxidative damage [37].

Ischemia-reperfusion injury in the testicular tissue is the main pathophysiology of testicular torsion/detorsion. This damage causes an inflammatory response that includes an increase in reactive oxygen radicals and cytokines. In addition, testicular torsion/detorsion causes an increase in oxidative stress markers and a decrease in antioxidant enzyme levels [38, 39]. In our study, we investigated the effects of testicular torsion and cerium oxide on oxidative stress by measuring MDA, a marker of oxidative stress, and the activities of GST, CAT, and PON against oxidative damage. Shookoobi et al., in their study in rats, showed that testicular T/D increased lipid peroxidation, thus increasing serum MDA levels [40]. Similarly, in the study of Moghimian et al., it was shown that testicular T/D increased MDA levels and decreased MDA levels with the administration of antioxidant vitamin C [5]. The results of our study were also compatible with the results of these studies. It has been shown that MDA levels increase in lung and renal tissues with testicular T/D and decrease with the administration of cerium oxide.

The effects of testicular T/D have been shown by many studies. In the study of Moghimian et al., in which they investigated the effects of *Syzygium aromaticum* in a rat testis T/D model, it was shown that systemic antioxidant enzyme levels were decreased (superoxide dismutase and glutathione peroxidase) in T/D, and enzyme levels were improved after *Syzygium aromaticum* was given [1]. In the study by Ameli et al., in which they investigated the effects of tadalafil and verapamil on oxidative stress in rat testis T/D, it was shown that the levels of systemic antioxidant enzymes were decreased (superoxide dismutase and glutathione peroxidase), and the levels were increased with tadalafil and verapamil [2]. In the testicular T/D model performed by Yuvanc et al. in rats, it was shown that PON levels decreased and following antioxidant administration PON levels increased [41]. In these studies, serum antioxidant levels were examined. In our study, we evaluated the antioxidant enzyme levels in lung and kidney tissues. As a result, we showed that antioxidant enzyme levels were low in testis T/D in lung and kidney tissues, but antioxidant enzyme levels increased in tissues with cerium oxide administration despite T/D. In the study of Akdemir and Tanyeli, oxidative damage and antioxidant enzyme levels in the testis and lung tissues were evaluated after testicular T/D. Similar to our study, it has been shown that oxidative damage in the lung tissue increased and antioxidant enzyme levels were low after testicular T/D [42]. Ischemia-reperfusion injury causes damage to distant organs as well as local damage [43]. There is information in the literature that testicular T/D is associated with lung injury [42]. However, apart from this study, we could not find any other publication in the literature showing the relationship between testicular T/D and distant organ damage. In our study, we demonstrated that after testicular ischemia-reperfusion injury, ischemia-reperfusion injury occurs in both lung and kidney tissues.

TABLE 2: Histopathological findings in lung tissue (mean  $\pm$  SE).

	Group C ( $n = 6$ )	Group CeO <sub>2</sub> ( $n = 6$ )	Group T/D ( $n = 6$ )	Group CeO <sub>2</sub> -T/D ( $n = 6$ )	$p^{**}$
Neutrophil infiltration/aggregation	0.33 $\pm$ 0.21	0.50 $\pm$ 0.22	1.50 $\pm$ 0.22* <sup>†</sup>	0.67 $\pm$ 0.21 <sup>‡</sup>	0.006
Alveolar wall thickening	0.33 $\pm$ 0.21	0.50 $\pm$ 0.22	1.50 $\pm$ 0.22* <sup>†</sup>	1.00 $\pm$ 0.00*	0.001
Total lung injury score	0.67 $\pm$ 0.42	1.00 $\pm$ 0.44	2.83 $\pm$ 0.30* <sup>†</sup>	1.67 $\pm$ 0.21* <sup>‡</sup>	0.002

$p^{**}$ : significance level with the Kruskal-Wallis test  $p < 0.05$ . \*  $p < 0.05$ : compared to group C. <sup>†</sup> $p < 0.05$ : compared to group CeO<sub>2</sub>. <sup>‡</sup> $p < 0.05$ : compared to group T/D.




FIGURE 5: Normal-structural lung tissue parenchyma in the control group. a: alveolus; sa: saccus alveolaris; da: ductus alveolaris; tb: terminal bronchiole; pa: pulmonary artery; ↓: pulmonary vascular thickening (H&E  $\times 40$ ).




FIGURE 7: Severe neutrophilic infiltration and increased alveolar wall thickness in the testis torsion/detorsion group. a: alveolus; rb: respiratory bronchiole; inf: inflammation; ↓: pulmonary vascular thickening; ↓↓: (septum) thickening; conj: capillary congestion; vc: vascular congestion (H&E  $\times 40$ ).




FIGURE 6: Mild neutrophilic infiltration and increased alveolar wall thickness in the cerium oxide group. a: alveolus; pa: pulmonary artery; rb: respiratory bronchiole; sa: saccus alveolaris; ↓↓: (septum) thickening (H&E  $\times 40$ ).




FIGURE 8: Mild neutrophilic infiltration and increased alveolar wall thickness in the testis torsion/detorsion treated with cerium oxide group. a: alveolus; da: ductus alveolaris; pa: pulmonary artery; rb: respiratory bronchiole; inf: inflammation; ↓: (septum) thickening; conj: capillary congestion; vc: vascular congestion (H&E  $\times 40$ ).

Studies have shown that the duration and degree of testicular torsion are related to the severity of ischemic injury [40, 44]. Turner et al. showed that spermatogenesis was permanently impaired within 1 hour of testicular torsion of 720° in rats, and germ cell apoptosis, neutrophil infiltration, and oxidative damage occurred after detorsion. Therefore, they stated that testicular T/D can be considered as a typical I/R injury [45]. The prognosis of testicular torsion is related to the duration of the torsion, and the resulting oxidative damage leads to different degrees of damage. Bilommi et al. studied the efficacy of glutathione in testicular T/D using a 3-hour reperfusion model after 4 hours of ischemia [14]. In the study of Arya et al., 3 hours of ischemia followed by 3 hours of reperfusion was established [46]. In both studies, injury in the testicular tissue was examined and it was

reported that the testicular tissue injury was revealed in the testicular torsion/detorsion models they used. In studies where shorter ischemia-reperfusion times were used, it was also shown that oxidative damage to the testicles and lungs occurred in a 2-hour reperfusion model after 2-hour ischemia used in studies in which Hirst and colleagues studied the effects of myricetin on rat testicular torsion/detorsion [47]. In the study in which Abbasoğlu and colleagues looked at the effects of taurine and carnosine on testicular torsion/detorsion in rats, ischemia and reperfusion times were determined as 2 hours each. As a result, it has been shown that histopathological and oxidative damage to testicular tissue occurred with these periods [9]. We used the 120-minute




FIGURE 9: Lung tissue oxidative stress parameters. GST: glutathione S-transferase; CAT: catalase; MDA: malondialdehyde; PON: paraoxonase. \* $p < 0.05$ : compared to group C. † $p < 0.05$ : compared to group CeO<sub>2</sub>. ‡ $p < 0.05$ : compared to group T/D.




FIGURE 10: Renal tissue oxidative stress parameters. \* $p < 0.05$ : compared to group C. † $p < 0.05$ : compared to group CeO<sub>2</sub>. ‡ $p < 0.05$ : compared to group T/D.

ischemia and then the 120-minute reperfusion model for the testicular torsion/detorsion model. However, we did not study the damage on testicular tissue. However, the histopathological changes and the changes in the oxidative stress parameters of the lung and kidneys have revealed that the model we used caused I/R damage in distant organs.

The reduction of oxidative damage caused by ischemia-reperfusion injury has become the target of many drug treatments. Cerium oxide is a nanoparticle that has been shown to be useful for use in medical and nonmedical fields [48,

49]. Cerium oxide exists in nature in two basic forms; cerium IV oxide (CeO<sub>2</sub>) and cerium III oxide (Ce<sub>2</sub>O<sub>3</sub>). CeO<sub>2</sub> exists in a more stable phase at room temperature and atmospheric conditions. CeO<sub>2</sub> nanoparticles are known to exist in two oxidative states; +4 and +3 oxidation states. This dual oxidation state of the nanoparticle means that these nanoparticles have oxygen vacancies. This dual oxidative state enables the emergence of antioxidant properties [18, 50, 51]. Its antioxidant properties have been proven by many different models of I/R injuries. Among

these, lower extremity, hepatic myocardial, intestinal, and spinal cord I/R injuries are involved [15–19, 23, 52, 53]. The effects of cerium oxide on testicular I/R injury are examined in studies. In these studies, cerium oxide acted as an antioxidant and reduced oxidative and histopathological damage to testicular tissue after reperfusion [46–54]. However, in our study, the effects of cerium oxide on testicular tissue were not examined and it was shown to reduce histopathological and oxidative damage in the lung and kidney tissue.

While there are many experimental animal studies and cell culture studies on cerium oxide today, there is no clinical use of this nanoparticle yet. These studies provide information about the use of cerium oxide for diagnosis and treatment in different pathophysiological conditions [50–55]. In this direction, as in other studies, we hope that the result of our study may be a study that can shed light on other experimental studies and clinical use.

The limitations of our study are that it will not be possible to adapt it to the clinical era since it is an animal study. Although the rat testicle has differences from the human testicle, rats are often used for testicular T/D models. This is due to the fact that lesions in the testicular torsion studies are comparable to those in the human testicles. Another limitation is that the experiment was performed without power analysis. The reason we did not perform a power analysis is because the ethics committee considers these numbers appropriate for the welfare of animals. We have reported the results of the study with 0.5 mg/kg ip dose we used for cerium oxide. The fact that we did not study the changes in the degree of oxidative damage at different doses may be another limitation of the study.

As a result of our study, it was shown that cerium oxide can reduce the oxidative and histopathological injury in the lung and kidney after testicular ischemia reperfusion. Reducing the oxidative damage associated with I/R has become the goal of drug studies conducted in this area. In many studies, many molecules that can inhibit oxidative stress have also been studied. It may be thought that cerium oxide may also be one of these potential inhibitors in testicular I/R injury. With this result, our study is the first to investigate the effects of cerium oxide on distant organ damage in a testicular I/R model. For this reason, considering the limitations of the study, contributions to the literature can be made by supporting the results with future studies to be carried out with new methods.

## Data Availability

The data that support the findings of this study are available from the corresponding author (Arslan, M), upon reasonable request (Mustafa Arslan mustarslan@gmail.com).

## Conflicts of Interest

The authors declare no conflict of interests.

## References

- [1] M. Moghimian, S. H. Abtahi-Evari, M. Shokoohi, M. Amiri, and M. Soltani, "Effect of *Syzygium aromaticum* (clove) extract on seminiferous tubules and oxidative stress after testicular torsion in adult rats," *Physiology and Pharmacology*, vol. 21, no. 4, pp. 343–350, 2017.
- [2] M. Ameli, M. S. Hashemi, M. Moghimian, and M. Shokoohi, "Protective effect of tadalafil and verapamil on testicular function and oxidative stress after torsion/detorsion in adult male rat," *Andrologia*, vol. 50, no. 8, article e13068, 2018.
- [3] H. Türk, O. Çelik, C. S. İşoğlu, H. Tarhan, Y. Ö. İlbey, and Y. Ö. İlbey, "Testicular Torsion In Adult," *The Journal of Tepecik Education and Research Hospital*, vol. 24, no. 1, pp. 73–76, 2014.
- [4] A. E. Sessions, R. Rabinowitz, W. C. Hulbert, M. M. Goldstein, and R. A. Mevorach, "Testicular torsion: direction, degree, duration and disinformation," *The Journal of Urology*, vol. 169, no. 2, pp. 663–665, 2003.
- [5] M. Moghimian, M. Soltani, H. Abtahi, J. Adabi, and N. Jajarmy, "Protective effect of tunica albuginea incision with tunica vaginalis flap coverage on tissue damage and oxidative stress following testicular torsion: Role of duration of ischemia," *Journal of Pediatric Urology*, vol. 12, no. 6, pp. 390.e1–390.e6, 2016.
- [6] E. Tabakoğlu and R. Durgut, "Veteriner hekimlikte oksidatif stres ve bazı önemli hastalıklarda oksidatif stresin etkileri," *AVKAE Dergisi*, vol. 3, no. 1, pp. 69–75, 2013.
- [7] S. Shimizu, P. Tsounapi, F. Dimitriadis et al., "Testicular torsion-detorsion and potential therapeutic treatments: a possible role for ischemic postconditioning," *International Journal of Urology*, vol. 23, no. 6, pp. 454–463, 2016.
- [8] S.-M. Wei, Z.-Z. Yan, and J. Zhou, "Taurine reduces testicular ischemia/reperfusion-induced neutrophil recruitment to testis probably by downregulation of pro-inflammatory cytokines and E-selectin," *Urology*, vol. 72, no. 2, pp. 464–465, 2008.
- [9] L. Abbasoğlu, E. B. Kalaz, M. Soluk-Tekkeşin, V. Olgaç, S. Dođru-Abbasođlu, and M. Uysal, "Beneficial effects of taurine and carnosine in experimental ischemia/reperfusion injury in testis," *Pediatric Surgery International*, vol. 28, no. 11, pp. 1125–1131, 2012.
- [10] T. R. Aydos, M. M. Bařar, O. Kul et al., "Effects of ozone therapy and taurine on ischemia/reperfusion-induced testicular injury in a rat testicular torsion model," *Turkish Journal of Medical Sciences*, vol. 44, no. 5, pp. 749–755, 2014.
- [11] S.-M. Wei, Z.-Z. Yan, and J. Zhou, "Beneficial effect of taurine on testicular ischemia-reperfusion injury in rats," *Urology*, vol. 70, no. 6, pp. 1237–1242, 2007.
- [12] H. Yousefi-Manesh, S. Shirooie, S. Hemati et al., "Protective effects of modafinil administration on testicular torsion/detorsion damage in rats," *Experimental and Molecular Pathology*, vol. 111, article 104305, 2019.
- [13] A. Beheshtian, A. H. Salmasi, S. Payabvash et al., "Protective effects of sildenafil administration on testicular torsion/detorsion damage in rats," *World Journal of Urology*, vol. 26, no. 2, pp. 197–202, 2008.
- [14] R. Bilommi, B. A. Nawas, D. D. Kusmayadi, R. Diposarosa, A. Chairul, and B. S. Hernowo, "The effects of glutathione on malondialdehyde expression and seminiferous tubule damage in experimental testicular torsion-detorsion in Wistar rats," *Journal of Pediatric Urology*, vol. 9, no. 6, pp. 1059–1063, 2013.



- [15] T. Tatar, Y. Polat, F. M. Comu et al., "Effect of cerium oxide on erythrocyte deformability in rat lower extremity ischemia reperfusion injury," *Bratislava Medical Journal-Bratislavské Lekárske Listy*, vol. 119, no. 7, pp. 441–443, 2018.
- [16] A. Tuncay, V. Sivgin, A. Ozdemirkan et al., "The effect of cerium oxide on lung tissue in lower extremity ischemia reperfusion injury in sevoflurane administered rats," *International Journal of Nanomedicine*, vol. Volume 15, pp. 7481–7489, 2020.
- [17] F. Pagliari, C. Mandoli, G. Forte et al., "Cerium oxide nanoparticles protect cardiac progenitor cells from oxidative stress," *ACS Nano*, vol. 6, no. 5, pp. 3767–3775, 2012.
- [18] K. A. Amin, M. S. Hassan, E. S. Awad, and K. S. Hashem, "The protective effects of cerium oxide nanoparticles against hepatic oxidative damage induced by monocrotaline," *International Journal of Nanomedicine*, vol. 6, pp. 143–149, 2011.
- [19] E. O. Gubernatorova, X. Liu, A. Othman et al., "Europium-doped cerium oxide nanoparticles limit reactive oxygen species formation and ameliorate intestinal ischemia-reperfusion injury," *Advanced Healthcare Materials*, vol. 6, no. 14, article 1700176, 2017.
- [20] A. A. Bhargava, N. Sathy, S. Singh, and M. Das, "Cerium oxide nanoparticles protect rodent lungs from hypobaric hypoxia-induced oxidative stress and inflammation," *International Journal of Nanomedicine*, vol. 8, pp. 4507–4520, 2013.
- [21] S. M. Hirst, A. Karakoti, S. Singh et al., "Bio-distribution and *in vivo* antioxidant effects of cerium oxide nanoparticles in mice," *Environmental Toxicology*, vol. 28, no. 2, pp. 107–118, 2013.
- [22] C. Xu, X. Qu, and X. Qu, "Cerium oxide nanoparticle: a remarkably versatile rare earth nanomaterial for biological applications," *NPG Asia Materials*, vol. 6, no. 3, p. e90, 2014.
- [23] N. D. P. K. Manne, R. Arvapalli, V. A. Graffeo et al., "Prophylactic treatment with cerium oxide nanoparticles attenuate hepatic ischemia reperfusion injury in Sprague Dawley rats," *Cellular Physiology and Biochemistry*, vol. 42, no. 5, pp. 1837–1846, 2017.
- [24] C. J. Wingard, D. M. Walters, B. L. Cathey et al., "Mast cells contribute to altered vascular reactivity and ischemia-reperfusion injury following cerium oxide nanoparticle instillation," *Nanotoxicology*, vol. 5, no. 4, pp. 531–545, 2011.
- [25] H. Moridi, S. A. Hosseini, H. Shateri et al., "Protective effect of cerium oxide nanoparticle on sperm quality and oxidative damage in malathion-induced testicular toxicity in rats: an experimental study," *International Journal of Reproductive Biomedicine*, vol. 16, no. 4, pp. 261–266, 2018.
- [26] N. D. Nosenko, N. M. Zholobak, L. I. Polyakova et al., "Morphofunctional state of reproductive system of ageing male rats in case of using nanocerium," *Fiziologicheskii Zhurnal*, vol. 60, no. 1, pp. 11–17, 2014.
- [27] N. M. Kobylak, T. M. Falalyeyeva, O. G. Kuryk et al., "Antioxidative effects of cerium dioxide nanoparticles ameliorate age-related male infertility: optimistic results in rats and the review of clinical clues for integrative concept of men health and fertility," *The EPMA Journal*, vol. 6, no. 1, p. 12, 2015.
- [28] H. Bostan, Y. Kalkan, Y. Tomak et al., "Reversal of Rocuronium-Induced Neuromuscular Block with Sugammadex and Resulting Histopathological Effects in Rat Kidneys," *Renal Failure*, vol. 33, no. 10, pp. 1019–1024, 2011.
- [29] T. M. Van Ye, A. M. Roza, G. M. Pieper, J. Henderson Jr., C. P. Johnson, and M. B. Adams, "Inhibition of intestinal lipid peroxidation does not minimize morphologic damage," *Journal of Surgical Research*, vol. 55, no. 5, pp. 553–558, 1993.
- [30] H. Aebi, "B. Isolation, purification, characterization, and assay of antioxygenic enzymes [13] Catalase *in vitro*," *Methods in Enzymology*, vol. 105, pp. 121–126, 1984.
- [31] W. H. Habig, M. J. Pabst, and W. B. Jakoby, "Glutathione S-transferases. The first enzymatic step in mercapturic acid formation," *Journal of Biological Chemistry*, vol. 249, no. 22, pp. 7130–7139, 1974.
- [32] H. W. Eckerson, C. M. Wyte, and B. N. La Du, "The human serum paraoxonase/arylesterase polymorphism," *American Journal of Human Genetics*, vol. 35, pp. 1126–1138, 1983.
- [33] G. Pizzino, N. Irrera, M. Cucinotta et al., "Oxidative Stress: Harms and Benefits for Human Health," *Oxidative Medicine and Cellular Longevity*, vol. 2017, Article ID 8416763, 2017.
- [34] M. Nisari, A. R. Yay, T. Ertekin et al., "Evaluation of protective effects of melatonin on free radical metabolism in rat kidney during ischemia-reperfusion," *Ukrainian Journal of Nephrology and Dialysis*, vol. 4, no. 64, pp. 20–29, 2019, <https://ukrjnd.com.ua/index.php/journal/article/view/369>.
- [35] E. Röth, N. Marczin, B. Balatonyi et al., "Effect of a glutathione S-transferase inhibitor on oxidative stress and ischemia-reperfusion-induced apoptotic signalling of cultured cardiomyocytes," *Experimental and Clinical Cardiology*, vol. 16, no. 3, pp. 92–96, 2011.
- [36] T. M. van Himbergen, L. J. H. van Tits, M. Roest, and A. F. H. Stalenhoef, "The story of PON1: how an organophosphate-hydrolysing enzyme is becoming a player in cardiovascular medicine," *The Netherlands Journal of Medicine*, vol. 64, no. 2, pp. 34–38, 2006.
- [37] R. Radi, J. F. Turrens, L. Y. Chang, K. M. Bush, J. D. Crapo, and B. A. Freeman, "Detection of catalase in rat heart mitochondria," *Journal of Biological Chemistry*, vol. 266, no. 32, pp. 22028–22034, 1991.
- [38] E. Yuluğ, S. Türedi, E. Karagüzel, O. Kutlu, A. Menteşe, and A. Alver, "The short term effects of resveratrol on ischemia-reperfusion injury in rat testis," *Journal of Pediatric Surgery*, vol. 49, no. 3, pp. 484–489, 2014.
- [39] M. Ayan, U. Tas, E. Sogut et al., "Protective effect of thymoquinone against testicular torsion induced oxidative injury," *Andrologia*, vol. 48, no. 2, pp. 143–151, 2016.
- [40] M. Shokoohi, E. O. Madarek, A. Khaki et al., "Investigating the Effects of Onion Juice on Male Fertility Factors and Pregnancy Rate After Testicular Torsion/Detorsion by Intrauterine Insemination Method," *International Journal of Women's Health and Reproduction Sciences*, vol. 6, no. 4, pp. 499–505, 2018, <http://ijwhr.net/text.php?id=386>.
- [41] E. Yuvanc, D. Tuglu, T. Ozan et al., "Investigation of the antioxidant effects of pheniramine maleate and nebivolol on testicular damage in rats with experimentally induced testis torsion," *Acta Cirúrgica Brasileira*, vol. 33, no. 2, pp. 125–133, 2018.
- [42] F. Akdemir and A. Tanyeli, "The effect of Fraxin against lung and testis damage induced by testicular torsion/detorsion in rats," *Annals of Medical Research*, vol. 27, no. 10, article 2769, 2020, <https://www.ejmanager.com/fulltextpdf.php?mno=95498>.
- [43] D. L. Carden and D. N. Granger, "Pathophysiology of ischaemia-reperfusion injury," *The Journal of Pathology*, vol. 190, no. 3, pp. 255–266, 2000.

- [44] C. Yang, B. Song, J. Tan, X. Liu, and G. Wei, "Testicular torsion in children: a 20-year retrospective study in a single institution," *The Scientific World Journal*, vol. 11, Article ID 575908, 2011.
- [45] T. T. Turner, H. J. Bang, and J. L. Lysiak, "The molecular pathology of experimental testicular torsion suggests adjunct therapy to surgical repair," *The Journal of Urology*, vol. 172, 6 Part 2, pp. 2574–2578, 2004.
- [46] S. S. Ashraf-Talesh and A. Amniattalab, "Histopathologic Evaluation of Intraperitoneal Administration of Cerium Oxide Nanoparticles on Ischemia Reperfusion Injury in Rat Testicular Torsion and Detorsion Model," *Iranian Journal of Veterinary Surgery*, vol. 16, no. 2, pp. 84–90, 2021.
- [47] D. Öztürk, A. Tanyeli, D. Güzel, M. C. Güler, E. Eraslan, and H. Baylan, "Mirisetinin testiküler iskemi reperfüzyon ile indüklenen testis ve akciğer hasarına karşı etkileri," *Sakarya Medical Journal*, vol. 11, no. 1, pp. 109–114, 2021.
- [48] B. Nelson, M. Johnson, M. Walker, K. Riley, and C. Sims, "Antioxidant Cerium Oxide Nanoparticles in Biology and Medicine," *Antioxidants*, vol. 5, no. 2, p. 15, 2016.
- [49] B. Stephen Inbaraj and B. H. Chen, "An overview on recent *in vivo* biological application of cerium oxide nanoparticles," *Asian Journal of Pharmaceutical Sciences*, vol. 15, no. 5, pp. 558–575, 2020.
- [50] I. Celardo, J. Z. Pedersen, E. Traversa, and L. Ghibelli, "Pharmacological potential of cerium oxide nanoparticles," *Nano-scale*, vol. 3, no. 4, p. 1411, 2011.
- [51] G. Casals, M. Perramón, E. Casals et al., "Cerium Oxide Nanoparticles: A New Therapeutic Tool in Liver Diseases," *Antioxidants*, vol. 10, no. 5, p. 660, 2021.
- [52] T. Zhao, W. Wu, L. Sui et al., "Reactive oxygen species-based nanomaterials for the treatment of myocardial ischemia reperfusion injuries," *Bioactive Materials*, vol. 7, pp. 47–72, 2022.
- [53] L. Dong, X. Kang, Q. Ma et al., "Novel Approach for Efficient Recovery for Spinal Cord Injury Repair via Biofabricated Nano-Cerium Oxide Loaded PCL With Resveratrol to Improve *in Vitro* Biocompatibility and Autorecovery Abilities," *Dose-Response*, vol. 18, no. 3, article 155932582093351, 2020.
- [54] A. Mousavi, A. Gharzi, M. Gholami, F. Beyranvand, and M. Takesh, "The therapeutic effect of cerium oxide nanoparticle on ischaemia/reperfusion injury in rat testis," *Andrologia*, vol. 53, no. 11, article e14231, 2021.
- [55] N. Feng, Y. Liu, X. Dai, Y. Wang, Q. Guo, and Q. Li, "Advanced applications of cerium oxide based nanozymes in cancer," *RSC Advances*, vol. 12, pp. 1486–1493, 2022.