









Research Article

Determination of Therapeutic and Safety Effects of *Zygophyllum coccineum* Extract in Induced Inflammation in Rats

Mohammed Yosri ¹, Mahmoud M. Elaasser ¹, Marwa M. Abdel-Aziz ¹,
Marwa M. Hassan,² Abdulmohsen Hussen Alqhtani ³, Naif Al-Gabri ^{4,5},
Ahmed B. A. Ali ⁶, A. Pokoo-Aikins ⁷, and Basma H. Amin ¹

¹The Regional Center for Mycology and Biotechnology, Al Azhar University, Nasr City, Cairo, Egypt

²Anatomy and Embryology Department, Faculty of Medicine, Helwan University, Egypt

³Animal Production Department, Food and Agriculture Sciences College, King Saud University, Riyadh, Saudi Arabia

⁴Department of Pathology, Faculty of Veterinary Medicine, Thamar University, Yemen

⁵Laboratory of Pathology, Salam Veterinary Group, Buraydah, Al Qassim 51911, Saudi Arabia

⁶Department of Animal and Veterinary Science, Clemson University, SC, USA

⁷US National Poultry Research Center, Toxicology & Mycotoxin Research Unit, USDA, ARS, Athens, GA 30605, USA

Correspondence should be addressed to Mohammed Yosri; mohammed.yosri@yahoo.com
and Abdulmohsen Hussen Alqhtani; ahalqhtani@ksu.edu.sa

Received 1 April 2022; Revised 28 May 2022; Accepted 14 June 2022; Published 18 July 2022

Academic Editor: Wen-Wu Li

Copyright © 2022 Mohammed Yosri et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. *Z. coccineum* is a facultative plant with many medicinal applications. This study examined the anti-inflammatory activity of *Zygophyllum coccineum* (*Z. coccineum*) in an arthritis animal model. **Materials and Methods.** Seventy-Six Wistar Albino rats of either sex randomly divided into six groups (12/each). The inflammation model was done using Complete Freund's Adjuvant in albino rats. The anti-inflammatory activities of the extract were estimated at different dose levels (15.6, 31, and 60 mg/kg) as well as upon using methotrexate (MTX) as a standard drug (0.3 mg/kg). Paw volume and arthritis index scores have been tested in all examined animals' treatments. Histological examination of joints was also performed. Flow cytometric studies were done to isolated osteoclasts. Cytokines assay as well as biochemical testing was done in the examined samples. **Results.** *In vitro* studies reported an IC_{50} of 15.6 μ g/ml for *Z. coccineum* extract in lipoxigenase inhibition assay (L.O.X.). Moreover, it could be noticed that isorhamnetin-3-O-glucoside, tribuloside, and 7-acetoxy-4-methyl coumarin were the most common compounds in *Z. coccineum* extract separated using L.C.–ESI–TOF–M.S. (liquid chromatography-electrospray ionization ion-trap time-of-flight mass spectrometry). Microscopic examinations of synovial tissue and hind limb muscles revealed the effect of different doses of *Z. coccineum* extract on restoring chondrocytes and muscles structures. Osteoclast size and apoptotic rate examinations revealed the protective effect of *Z. coccineum* extract on osteoclast. The results upon induction of animals and upon treatment using of MTX significantly increased apoptotic rate of osteoclast compared to control, while using of 15.6 μ g/ml. for *Z. coccineum* extract lead to recover regular apoptotic rate demonstrating the protective effect of the extract. *Z. coccineum* extract regulated the secretion of proinflammatory and anti-inflammatory cytokines. Biochemical tests indicated the safety of *Z. coccineum* extract on kidney and liver functions. **Conclusion.** *Z. coccineum* extract has efficient and safe anti-inflammatory potential in an induced rat model.

1. Introduction

Rheumatoid arthritis (R.A.) is a form of arthritis that leads to pain, swelling, and distortion of joints functions. It is a

long-lived disorder developed by combined genetic, epigenetic, and environmental circumstances. Many immune and nonimmune cells, as well as inflammatory factors, have essential functions in the process of inflammation and joint

destruction [1]. The synovium considers as potential tissue in R.A. [2], and the formation of a pannus leads to the destruction of cartilage and bone. Chondrocytes are affected by many autoantibodies and secreted extracellular components [3]. Osteoclasts are multinucleated cells, and they act as bone modulator cells. They are generated in the bone marrow through the stimulatory effects of macrophage colony-stimulating factor (M-CSF) and receptor activator of nuclear factor κ B ligand (R.A.N.K.L.) [4–6]. Cytokines are pivotal in both humoral and adaptive responses in developing rheumatoid arthritis [7]. The equilibrium in pro and anti-inflammatory cytokines using either natural or synthetic factors is innovative in overcoming rheumatoid arthritis pathology [8].

The available nonsteroidal anti-inflammatory drugs to treat rheumatoid arthritis have many drawbacks, including abdominal discomfort in many internal body organs affecting their functions. Alternative therapy is a modern path that has been used to combat many human diseases. It has been reported that 50–80% of rheumatoid arthritis patients use conventional herbal therapies [9–11]. Different natural extracts have been used with various beneficial activities including regulating liver damage [12–15], various types of cancer cells [16, 17], diabetes [18], and COVID-19 [19], through regulating various signaling pathways [20–23].

Z. coccineum is a succulent shrub from the family Zygophyllaceae. It can grow in different kinds of soil and can tolerate high salt levels inland. It is distributed in many African, Asian, and Mediterranean areas as well as Australia [24–26]. The aerial parts contained bioactive molecules with promising antifungal and insecticidal actions [27, 28]. It has been used in the removal of high content of heavy metals, including copper and lead, in water [29]. *Z. coccineum* extract has been tested against induced diabetic rats, showing promising activity to control high blood glucose with minimal burden in body organs, including liver and kidney [30]. *Z. coccineum* aqueous extract was reported to control elevated blood pressure and heart rate in induced rats [31]. In this study, the anti-inflammatory properties of *Z. coccineum* extract have been investigated *in vitro* and in induced rats, illustrating its therapeutic effects with a noticeable safety profile.

2. Materials and Methods

2.1. Preparation of Extract. *Z. coccineum* was collected in 2020 from the deltaic coast of Egypt and identified by the institutional plant taxonomist (Prof. Abbas Elgamry) at the Botany and Microbiology Department, Faculty of Science, Al Azhar University according to (Voucher No. ID: 0044). 500 gm whole plant was air-dried, crushed with blinder extracted from aq.-ethanol 70%, dried, and were stored at -20°C for further analysis [32].

2.2. Determination of Phytochemicals. Separation was achieved on a $5\ \mu\text{m}$ C18 column ($50 \times 2.0\ \text{mm}$ internal diameter; Bohus, Sweden; 45°C) using Agilent L.C.–ESI–TOF–M.S. (Agilent Technologies, Palo Alto, CA, U.S.A.) System. The LC mobile phases consisted of (A) 5 mM NH_4HCO_2 in 1% CH_3OH (pH 3.0) and (B) 5 mM

NH_4HCO_2 in 1% CH_3OH (pH 8.0). Gradient elution was done at 0.2 ml/min and 0–20 min, 10% B; 21–35 min, 90% B; 35.01–60 min, 10% B; and then 90% B to the end of the run. The sample was injected for LC-MS analysis in both positive and negative modes. The LC-MS analysis was also performed before the sample injection for blank and quality control samples for confidence in the experiment. Infusion experiments were carried out to optimize the negative ion ESI-MS/MS parameters for maximal generation of deprotonated molecules and effective generation of characteristic fragment ions for all analyses. The MS-DIAL V. 3.70 open-source software was used for the identification and calculation of the relative percentage of compounds [32, 33]. MS-DIAL 3.70 software was used for small molecule analysis of the separated sample. According to the acquisition mode, ReSpect positive (2737 records) or ReSpect negative (1573 records) databases were used as reference databases [34].

2.3. In Vitro Lipoxigenase (L.O.X.) Inhibition Assay. *Z. coccineum* extract and the reference compound (Ibuprofen) were tested in order to investigate the anti-inflammatory response by inhibiting the L.O.X. enzyme (lipoxigenase enzyme) from Glycine max (type I-B) according to Granica et al. [35] with slight modifications. Briefly, in 96-well plates, $100\ \mu\text{l}$ of soybean L.O.X. solution (1000 U/ml in borate buffer solution, pH 9) and $200\ \mu\text{l}$ of borate buffer were mixed together with varying concentrations of the sample to a final concentration range of 0.98–125 $\mu\text{g}/\text{ml}$ at 25°C for 15 min. Samples were preincubated with $100\ \mu\text{l}$ of linoleic acid (substrate) to start the reaction. The inhibitory activity was determined by monitoring the absorbance's increase at 234 nm using a microplate reader (BioTek, U.S.A.).

2.4. Animals and Treatments. Male Wister albino rats (10 weeks old) weighing 130–150 g were purchased from the animal unit of Faculty of Science, Al-Azhar University, left for acclimation for ten days, and split into six groups (twelve rats each). The first group was observed as negative control (N.C.) infused subcutaneously with saline and 10% Tween-80, twice per week for two weeks. The other groups were interjected subcutaneously twice a week for two weeks at the base of the tail with $100\ \mu\text{L}$ C.F.A. (Complete Freund's Adjuvant) (Sigma-Aldrich, U.S.A.) to convene arthritis model [36]. The second group presents the positive control (P.C.) that was kept untreated. Three estimated doses (15.6, 31, and 60 mg/kg) of *Z. coccineum* extract were administrated subcutaneously in the third, fourth, and fifth groups, respectively. Lastly, antirheumatic standard drug MTX (Methotrexate) (Orion Pharma, Espoo, Finland) was injected. Signs of arthritis were raised after 14 days, and then *Z. coccineum* extract was used three times per week for 2 weeks at a dose of 15.6, 31, and 60 mg/kg extract B.W. MTX was used subcutaneously twice per week at a dose of 0.3 mg/Kg B.W. [37] as shown in supplement (1). The animal studies were approved by the ethical committee in the Regional Center for Mycology and Biotechnology (No. RCMB26062020).

2.5. Assessing Swelling Scoring. Severity scores and paw volume were assessed daily to evaluate inflammation. The

severity score test is graded on a scale of 0–5, where a total score of 5 points indicates severe inflammation deficits and a score of 0 indicates normal performance; 4 points indicates severe injury, 3 indicates mean to moderate injury, and 1–2 indicate mild injury. Evaluation was performed after induction by technicians who were blinded to the experiments [38]. Paw volumes have been measured using a digital plethysmometer (B.V.K., India) [39].

2.6. Histopathology Studies. Animals were euthanized by cervical dislocation. Tibiofemoral joint and hind limb muscles were harvested, for tibiofemoral joint samples only that were decalcified using 5% nitric acid for 10 days. There are both types of samples (joints and muscles). Sections of samples were cut at 5 μm on a rotary microtome, processed, mounted on slides, and stained (Leica Autostainer XL) with hematoxylin and eosin, to assess tissue morphology and imaged at $\times 20$ (Zeiss microscope and imaging system, Carl Zeiss Inc., Germany) [40].

2.7. Transmission Electron Microscopy. To test ultrastructure changes of decalcified tibiofemoral joint and hind limb muscles, transmission electron microscopy was used. Samples chemically fixed in an aqueous solution of glutaraldehyde (2.5%) for two days at 4°C, followed by postfixation with a 1% osmium tetroxide solution for four hours at 4°C followed by sectioned using ultra-microtome (Leica, Germany) and analyzed using 1010 T.E.M. microscopy (J.E.O.L., Japan) in the Regional Center for Mycology and Biotechnology [41].

2.8. Cell Culture Studies. Bone marrow-derived monocytes/macrophages were detached from the tibias of male rats by rub the bone-marrow hole with D.M.E.M. enriched by 10% fetal bovine serum, 5% l-glutamine, 100 U ml⁻¹ penicillin, and 100 μg ml⁻¹ streptomycin. The cells were bred for six h to split nonadherent and adherent cells into six well tissue culture plates. Nonadherent cells were cultured in 6-well tissue culture plates at 2×10^5 cells/well in the presence of 10 ng/ml rh M-CSF (Invitrogen, U.S.A.) for 3 days to collect macrophage-like osteoclast precursor cells. After 3 days, the nonadherent cells were collected, and preosteoclasts were cultivated in the existence of 10 ng/ml M-CSF, 50 ng/ml R.A.N.K.L., and diverse levels of sodium butyrate for 4 days to form osteoclasts. On day 2, the medium was replaced with fresh medium consisting of M-CSF, R.A.N.K.L., and sodium butyrate. Photos were captured using inverted microscopy Zeiss microscope and imaging system (Carl Zeiss Inc., Germany) [42].

2.9. Flow Cytometry Analysis. Cultured osteoclasts in 6-well plate were used for this test. Osteoclasts were split using trypsin in 0.25% pancreatin and washed using phosphate-buffered saline. The death rate was assessed by an Annexin V-FITC and propidium iodide (P.I.) staining kit (B.D. Bioscience, U.S.A.), and cells were suspended using buffer which has Annexin V-FITC and/or P.I. stock solution and kept away from light at room temperature for 15 min. The cells were examined by flow cytometry (B.D. Bioscience, U.S.A.) [43, 44].

2.10. ELISA Testing. Blood samples were collected from rat's eyes in heparinized Eppendorf tubes before sacrificing animals and centrifuged at 5000 rpm for 15 min at 5°C. Then serum was collected and kept at -80°C till analysis. Serum levels of IFN- γ , IL-1 β , IgG1a, IgG2a, IL4, IL-6, and IL-17 were detected by kits (Abcam, U.S.A.) using the manufacturer's explained steps.

2.11. Biochemical Tests. The levels of A.L.T., creatinine, and C-reactive protein in serum were determined by the standard protocol using the biochemical diagnostic kits (Diamond, UK) [45].

2.12. Statistical Analysis. Calculations of statistical variance were done by GraphPad Prism 5 software (San Diego, CA). Either two groups were represented in the test, an unpaired two-sided Student's *t*-test was done; for relations of more than two groups, one-way analysis (ANOVA) was used followed by Turkey's post-hoc test to detect significance between groups where $P < 0.05$ is expressed significant.

3. Results

3.1. LC-ESI-TOF-M.S. Analysis. The aq.-ethanolic extract of the *Z. coccineum* was chromatographically separated using L.C.-M.S. resulting in generation of characteristic fragment ions that were identified by the system software, and the relative percentages of the separated compounds were determined. However, it could be noticed that isorhamnetin-3-O-glucoside, tribuloside, 7-acetoxy-4-methyl coumarin, luteolin, and zygophyloside-S were the most common compounds in *Z. coccineum* extract as shown in Table 1 and Figure 1. Moreover, in general, the separated compounds were included into nine phytochemical groups, though fourteen compounds belonged to flavonoids that represented 40.89% of the total LC-MS chromatogram contents. The terpenoid compounds (including saponins) were in second place, representing 36.73% of the total contents detected by 12 compounds. Interestingly, eight quinovic acid-based triterpenoid saponins were detected with a relative percentage value of 16.74% of the total plant extract's constituents that were characteristic of this plant species. Furthermore, five phenolic compounds, including coumarins, represent 16.35% of the total LC-MS chromatogram contents. Also, only two compounds, i.e., cinnamaldehyde and syringaldehyde, were separated as aldehydes (1.94%). The lipid percentage was 1.82% of the total contents, and it contained two sterols as well as one fatty acid. Alkaloids (0.55%), alcohols (0.34%), organic acids (0.24), and anthocyanins (0.07%) were also presented as minor contents.

3.2. Effect of *Z. coccineum* Extract on Lipoxxygenase Enzymatic Activity. *Z. coccineum* extract showed promising inhibition of lipoxxygenase action. The IC₅₀ value for 15.6 \pm 2.1 μg /ml, whereas for ibuprofen as reference compound, it was estimated as IC₅₀ value of 1.4 \pm 1.2 μg /ml as shown in Figure 2.

3.3. Effect of *Z. coccineum* Extract on Paw Volume and Arthritis Score. Paw volume and arthritis score were

TABLE 1: Identified compounds in the aqueous-ethanolic extract of *Z. coccineum* by L.C.–ESI–TOF–M.S (detected in positive and negative mode of ESI).

No.	R.T. min.	Characteristic mass fragments	Molecular weight (m/z)	Compound name	Chemical formula	% ¹
1	12.69	43, 68, 88	88.06	Pyruvic acid	C ₃ H ₄ O ₃	0.09
2	15.34	105, 133	133.10	Cinnamaldehyde	C ₉ H ₈ O	0.21
3	16.51	43, 71, 89	134.08	Malic acid	C ₄ H ₆ O ₅	0.15
4	18.37	93,137	137.02	p-Hydroxybenzoic acid	C ₇ H ₆ O ₃	0.11
5	24.88	129,146	146.16	Spermidine	C ₇ H ₁₉ N ₃	0.16
6	25.71	120, 138, 147, 154	154.16	4-Hydroxy-3-methoxybenzyl alcohol	C ₈ H ₁₀ O ₃	0.34
7	26.32	119, 163	163.04	p-Coumaric acid	C ₉ H ₈ O ₃	0.53
8	26.86	43, 172	172.26	Capric acid	C ₁₀ H ₂₀ O ₂	0.91
9	27.29	135, 150, 161, 179	179.05	Caffeic acid	C ₉ H ₈ O ₄	1.35
10	30.67	129, 180	180.17	1,7-Dimethylxanthine	C ₇ H ₈ N ₄ O ₂	0.39
11	31.78	123, 140, 168, 183	183.09	Syringaldehyde	C ₉ H ₁₀ O ₄	1.73
12	33.19	161, 219	219.15	7-Acetoxy-4-methyl coumarin	C ₁₂ H ₁₀ O ₄	11.07
13	33.24	179, 223	223.13	Sinapic acid	C ₁₁ H ₁₂ O ₅	3.29
14	33.76	153, 219, 263	263.13	Abscisic acid	C ₁₅ H ₂₀ O ₄	2.14
15	34.12	204, 251, 252, 267	267.07	Formononetin	C ₁₆ H ₁₂ O ₄	1.56
16	34.95	119, 121, 153, 225, 271	271.06	Apigenin	C ₁₅ H ₁₀ O ₅	5.19
17	35.77	137, 153, 241, 269, 287	287.05	Luteolin	C ₁₅ H ₁₀ O ₆	10.21
18	36.21	147, 273	273.14	Naringenin	C ₁₅ H ₁₂ O ₅	0.37
19	37.04	153, 286, 301	301.07	Kaempferide	C ₁₆ H ₁₂ O ₆	0.22
20	38.13	153, 165, 229, 247, 285, 303	303.04	Quercetin	C ₁₅ H ₁₀ O ₇	0.37
21	38.41	285, 303	303.09	Taxifolin	C ₁₅ H ₁₂ O ₇	0.43
22	39.72	121, 153, 229, 302, 317	317.06	Isorhamnetin	C ₁₆ H ₁₂ O ₇	0.35
23	40.38	414	414.70	β -Sitosterol	C ₂₁ H ₂₄ O ₉	0.42
24	42.65	165, 241 287, 433	433.11	Kaempferol-3-O- β -L-rhamnoside	C ₂₁ H ₂₀ O ₁₀	0.19
25	45.73	43, 456	456.23	Ursolic acid	C ₃₀ H ₄₈ O ₃	0.36
26	51.28	302, 317,479	479.12	Isorhamnetin-3-O-glucoside	C ₂₂ H ₂₂ O ₁₂	21.14
27	52.17	229, 247, 303, 465	465.14	Hyperoside (quercetin-3-O-galactoside)	C ₂₁ H ₂₀ O ₁₂	0.38
28	52.86	414, 576	576.85	β -Sitosterolglucoside	C ₃₅ H ₆₀ O ₆	0.49
29	54.79	287, 433,579	579.14	Kaempferol3,7-di-O- β -L-rhamnoside	C ₂₇ H ₃₀ O ₁₄	0.09
30	57.18	287, 449, 595	595.16	Cyanidin-3-O-rutinoside	C ₂₇ H ₃₁ O ₁₅	0.07
31	57.93	153, 229, 303, 465, 611	611.16	Rutin	C ₂₇ H ₃₀ O ₁₆	0.12
32	59.04	300, 315, 623	623.28	Isorhamnetin-3-O-rutinoside	C ₂₈ H ₃₂ O ₁₆	0.27
33	61.58	602, 647	647.37	3-O- $[\beta$ -D-Glucopyranosyl] quinovic acid	C ₃₆ H ₅₆ O ₁₀	0.14
34	61.87	285, 593	593.15	Tribuloside	C ₃₀ H ₂₆ O ₁₃	17.35
35	64.72	485, 631, 763	763.02	3-O- $[\alpha$ -L-Arabinopyranosyl-(1 \rightarrow 2)- β -D-quinovopyranosyl] quinovic acid	C ₄₀ H ₆₂ O ₁₄	0.43
36	65.83	254, 587, 749, 793	793.44	3-O- $[\beta$ -D-Quinovopyranosyl] quinovic acid-28- β -D-glucopyranosyl ester	C ₄₂ H ₆₆ O ₁₄	0.28
37	67.71	254, 587, 749, 803	803.41	Zygophylloside S (3-O- $[\alpha$ -L-arabinopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl] quinovic acid)	C ₄₁ H ₆₄ O ₁₄	6.53
38	69.28	603, 647, 809	809.43	Zygophyloside-K (3-O- $[\beta$ -D-glucopyranosyl] quinovic acid-28-O- β -D-glucopyranosyl ester)	C ₄₂ H ₆₆ O ₁₅	3.48
39	70.23	587, 667, 711, 873	873.40	Zygophyloside-F	C ₄₂ H ₆₆ O ₁₇	2.32

TABLE 1: Continued.

No.	R.T. min.	Characteristic mass fragments	Molecular weight (m/z)	Compound name	Chemical formula	% ¹
40	71.65	97, 727, 845,889	889.39	Zygophyloside-G	C ₄₂ H ₆₆ O ₁₈	1.19
41	72.89	603, 647, 809, 891	891.24	3-O-[α-L-Arabinopyranosyl-(1 → 2)-β-D-quinovopyranosyl] quinovic acid-28-O-β-D-glucopyranosyl ester	C ₄₂ H ₆₆ O ₁₅	1.17
42	73.27	602, 893	893.13	3-O-[β-D-(2-O-Sulphonyl)-quinovopyranosyl] quinovic acid	C ₄₂ H ₆₆ O ₁₄ S	1.34

¹Relative percentages of compounds in *Z. coccineum* extract were calculated based on the total peak area in the chromatogram.




FIGURE 1: Total ion chromatograph (TIC) of separated compounds in the aqueous-ethanolic extract of *Z. coccineum* detected by L.C.–ESI-TOF–M.S. Each separated peak represents a chemical constituent at specified retention time.




FIGURE 2: Graphical representation of in vitro anti-inflammatory assay of *Z. coccineum* extract. Results are expressed as a mean ± SEM ($n = 3$), where ($IC_{50} = 15.6 \pm 2.1 \mu\text{g/ml}$ for *Z. coccineum* extract while, $IC_{50} = 1.4 \pm 1.2 \mu\text{g/ml}$ for standard).

measured daily after the induction of animals. The paw volume and arthritis score were dramatically increased in the induced rats versus negative control. The paw volume and arthritis score significantly decreased ($P < 0.05$) in the induced animals after receiving 15.6 and 31 mg/kg of *Z. coc-*

cineum extract, similar to obtained results for induced animals and treated with standard drug, while treatment using concentration of 60 mg/kg of *Z. coccineum* extract showed a slight improvement in both tested parameters, indicating that 15 mg/kg of the extract could be significantly improving harmful symptoms of arthritis in the current experimental model, as shown in Figure 3.

3.4. Histology and T.E.M. Observations. At the end of the experiment (30 days), rats in the negative control group exhibited normal chondrocytes, with a regular sized nucleus and paw cartilage with a flat rough surface as shown in Figure 4(a), A. In the induced rats' extensive inflammatory cells and condensed membrane, deformed nuclei of chondrocytes and clear fat droplets with a score (5) were shown in Figure 4(b), B. The examined results of induced rats upon receiving 15 mg/kg of *Z. coccineum* extract indicated significantly recovering of chondrocyte structure and nucleus size compared to the experimental group as an optimum dose for healing as shown in Figure 4(c), C, while administration of 31 mg/kg of *Z. coccineum* extract for induced rats led to minimal inflammation, as shown through a decreased number of inflammatory cells, thickening, and mild cartilage destruction and slightly altered smaller chondrocytes structure with a score (2-3) as shown in Figure 4(d), D.




FIGURE 3: Effect of 15.6, 31, and 60 mg/kg of *Z. coccineum* extract on (a) arthritis index score and (b) paw volume in arthritic rats. The induced elevation in paw volume and arthritis index score was inhibited by 15.6 and 31 $\mu\text{g/ml/kg}$ of *Z. coccineum* extract. The treatment with 60 mg/kg of *Z. coccineum* extract displayed minimal diminution in both paw volume and arthritis index score. Values are represented as mean \pm S.E.M. of triplicate values ($n = 3$), where $P < 0.05$.

Furthermore, treating infected animals with 60 mg/kg of *Z. coccineum* extract led to moderate signs of inflammation, reflecting the ineffectiveness of the used dose to overcome the tissue alterations with a score (3) as shown in Figures 4(e) and 4(f). All of these examined results were also compared to the positive control group showing regular tissue structure with mild pathological changes in mitochondria, endoplasmic reticulum, and Golgi bodies, as shown in Figure 4(f), F.

To investigate the role of various doses of *Z. coccineum* extract on the morphological alterations in hind limb and muscle, tissues were collected and stained with hematoxylin and eosin and processed for transmission electron microscopy examinations. In the first negative control group, classical muscle structure could be seen with regular fibers and nuclear structure with homogenous distribution, as shown in Figure 5(a), A. There was an increase in inflammation and myofiber degradation in the positive control group along with a decrease in muscle mass and fiber size (Figure 5(b), B). However, by using 15.6 mg/kg of *Z. coccineum* extract for the induced rats, muscle bundles recovered their regular structure as numerous fibers with normal

nuclei distribution could be observed as shown in Figure 5(c), C. Furthermore, upon administration of 31 and 60 mg/kg of *Z. coccineum* extract, the examined muscles appeared with a significantly decreased level of cell infiltration and huge irregular fibers with numerous centrally located nuclei as shown in Figures 5(d), D and 5(e), E. In comparison to MTX as a standard drug in induced animals, muscles appeared with slight necrosis and slight aggregations to nuclei, as shown in Figure 5(f), F.

3.5. *Z. coccineum* Extract Modulated Osteoclast Activity. To test the effect of the *Z. coccineum* on osteoclasts, isolated osteoclast cells from different groups of animals were cultured for 7 days and examined using an inverted microscope. A significant decrease ($P < 0.05$) could be seen in size of osteoclast cells isolated from induced rats compared to cells isolated from the negative control group of animals, while cultured osteoclast cells from animals treated with 15.6 mg/kg of *Z. coccineum* extract resulted in recovering of osteoclast cells its normal structure compared to the negative control group. Furthermore, a slight change could be seen in osteoclast size in isolated cultured cells from




FIGURE 4: Effects of synovial tissue histopathology of rats with antigen-induced arthritis (magnifications, $\times 20$ and $\times 8000$) using H&E staining and transmission electron microscope. (a, A) Control group with regular chondrocytes (green arrow) with normal nucleus, (b, B) model group with elongated chondrocytes (green arrow), elongated nucleus and many inflammatory cells infiltrated the synovium, (f, F) induced group and receiving MTX as the standard drug group, and cartilaginous tissue could be clearly detected, while (c, C; d, D; e, E) tissues of induced rats showed less inflammation, less lymphocyte accumulation, and regained normal structure of chondrocytes (green arrow) with less cartilage damage decreased upon receiving 15.6, 31, and 60 mg/kg of *Z. coccineum* extract, respectively.




FIGURE 5: Histology of hind limb muscles, where five groups (a, A) are as follows: control group showing regular muscle structure with elongated nuclei (white arrow). (b, B) Infected with C.F.A. to induce A.I.A. showing injured muscles with severe infiltration and aggregated lymphocytes, macrophages, and plasma cells (red arrow). (c, C; e, E) Induced animals plus 15.6, 31, and 60 mg/kg of *Z. coccineum* extract. (f, F) Induced animals plus treatment with standard drug showing necrosis (green arrow) and slight nuclei aggregation; muscles obtained after animal sacrifice and were stained with H&E and photos captured at a magnification of $\times 20$ and prepared for a transmission electron microscope and images taken at a magnification of $\times 8000$.

animals treated with 31 and 60 mg/kg *Z. coccineum* extract as well as MTX as shown in Figure 6(a). This *ex vivo* result indicated that 15.6 mg/kg of *Z. coccineum* could enhance the propagation of mature osteoclasts in arthritic rats.

To detect apoptotic rate for osteoclasts, Annexin V-FITC kit was used to stain cells. It could be noticed that induction using C.F.A dramatically increased apoptotic rate of osteoclasts ($P < 0.05$). However, using of 15.6, 31, and 60 mg/kg of *Z. coccineum* extract resulted in recovering of osteoclast cells its normal apoptotic rate compared to the negative control group where using of 15.6 mg/kg made the optimal recovery. Furthermore, administration of MTX to induced animals increased apoptotic rate of osteoclasts ($P < 0.05$) relative to negative control group as shown in Figure 6(b).

3.6. *Z. coccineum* Extract Regulated the Expression of Cytokines. ELISA was done to evaluate proinflammatory factors, including $IFN-\gamma$, $IL-1\beta$, $IL-6$, $IL-17$, and anti-inflammatory factors as $IL-4$ as well as $IgG1a$ and $IgG2a$ in rat serum. The cytokines levels in serum concentrations of $IFN-\gamma$, $IL-1\beta$, $IL-6$, $IL-17$, and $IgG1a$ and $IgG2a$ in A.I.A. (antigen-induced arthritis) rats were significantly higher than in control ($P < 0.05$, 0.001). $IL-4$ declined in serum in A.I.A. rat's ($P < 0.05$). Treatment with 15.6, 31, and 60 mg/kg of *Z. coccineum* extract significantly decreased inflammation by downregulation of $IL-1\beta$, $IL-6$, $IFN-\gamma$, and $IL-17$ and upregulation of $IL-4$ in the serum ($P < 0.05$, 0.001). A dose of 15.6 mg/kg *Z. coccineum* extract shows anti-inflammatory action as those of animals treated with MTX as a standard





FIGURE 6: Continued.




FIGURE 6: (a) Microscopic examination of osteoclasts in medium enriched with recombinant R.A.N.K.L. Images are representative of cell appearance after the 7 days in culture using phase-contrast microscopy (A) control cells and (B) cells cultured from induced rat. (C)–(E) Cells cultured from induced rat and treated with 15.6, 31, and 60 mg/kg *Z. coccineum* extract, respectively, and (F) cells cultured from induced rat and treated with standard drug where (A) control cells and (B) cells are cultured from induced rat. (C, D, E) Cells cultured from induced rat and treated with 15.6, 31, and 60 mg/kg *Z. coccineum* extract, respectively, and (F) cells cultured from induced rat and treated with standard drug. (G) A bar chart depicts difference in osteoclast size between control group and various treatment; there is a dramatic decrease in osteoclast size in cells cultured cells upon from induced rats ($P < 0.05$), and cells recover their size upon different treatments (15.6, 31, 60 mg/kg *Z. coccineum* extract and MTX). (b) Annexin V-FITC and propidium iodide were used to detect apoptosis in osteoclasts and analyzed by flow cytometer: (A) control cells and (B) cells cultured from induced rat. (C)–(E) Cells cultured from induced rat and treated with 15.6, 31, and 60 mg/kg *Z. coccineum* extract, respectively, and (F) Cells cultured from induced rat and treated with standard drug. (G) A bar chart depicts difference in osteoclast apoptotic rates between control group and various treatment; there is a dramatic increase in osteoclast apoptotic rates in cells cultured cells upon from induced rats and upon using MTX ($P < 0.05$), and cells recover their regular apoptotic rate different treatments (15.6, 31 and 60 mg/kg *Z. coccineum* extract).




FIGURE 7: Continued.




FIGURE 7: Various concentrations (15.6, 31, and 60 mg/kg) of *Z. coccineum* extract of the levels of cytokines in A.I.A. rats versus control and upon treatment using the standard drug. The level of (a) IFN- γ , (b) IL-1 β , (c) IgG1a, (d) IgG2a, (e) IL-4, (f) IL-6, and (g) IL-17 in rat serum was detected by ELISA. Results are presented as means \pm SEM, $n = 3$. * $P < 0.05$, *** $P < 0.01$.

drug as shown in Figure 7. In the current model, regulations of cytokine levels likely lead to the decreased inflammation observed.

3.7. Effect of *Z. coccineum* Extract on Various Biochemical Investigations in Rats. The data showed that A.S.T. and creatinine levels were significantly elevated in the group of rats that used MTX as a standard drug for treatment compared to other tested groups ($P < 0.05$), which indicated harmful effects in either kidney or liver functions of this group of tested animals. Furthermore, the induced group of animals using C.F.A. showed a dramatic increase in C-reactive protein level compared to other tested groups ($P < 0.05$). Moreover, upon using 15.6, 31, and 60 mg/kg *Z. coccineum* extract, the C-reactive protein level reached to regular level

with slight differences between groups in induced rats, indicating the antiarthritic activity of extract as shown in Figure 8.

4. Discussion

Due to the restrictions surrounding human samples from arthritis patients, animal models are a viable alternative to test new therapeutics. [46–48]. Plants contain many compounds that could be used separately or combined in therapeutic applications, including anti-inflammatory functions [49, 50]. In the present study, we test the impact of *Z. coccineum* extract against inflammation symptoms in the antigen-induced arthritis rat model.




FIGURE 8: Effect of different concentration (15.6, 31, and 60 mg/kg) of *Z. coccineum* extract on serum liver and kidney functions as well as C-reactive protein in A.I.A. rats versus control and upon treatment using the standard drug. The level of (a) creatinine, (b) A.L.T., (c) C-reactive protein in rat serum was detected by biochemical kits. Results are presented as means \pm S.E.M. ** $P < 0.05$.

In the current study, the separated compounds were included into nine phytochemical groups; flavonoids (40.89%), terpenoid compounds (including saponins 36.73%), phenolic (including coumarins 16.35%), aldehydes (1.94%), lipids (sterols and fatty acid 1.82%, alkaloids (0.55%), alcohols (0.34%), organic acids (0.24), and anthocyanins (0.07%) of the total LC-MS chromatogram contents. However, it could be noticed that isorhamnetin-3-O-glucoside, tribuloside, and 7-acetoxy-4-methyl coumarin were the most common compounds in *Z. coccineum* extract.

Many previous reports recounted the presence of triterpene-saponins, phenolics, and flavonoids in *Z. coccineum* extracts [51, 52]. It has also been reported that *Z. coccineum* possesses potent antioxidant activity and has traditionally been used for hypertension, diabetes, and rheumatoid fever [53].

In this study, the chemotaxonomic marker of the genus *Zygophyllum*, i.e., isorhamnetin-3-O-rutinoside was identified in the *Z. coccineum* ethanolic extract. Tribuloside (also known as kaempferol-3-O-(p-coumaroyl)-glucoside) was

detected in the current study as the major flavonoid glycoside constituent as reported by other groups [54–56]. However, the presence of flavonoids and phenolic contents with different structural formulae in *Z. coccineum* extract as obtained by LCMS separation in this study suggested varying roles of the antioxidant activities that was consistent with previously reported by El-Shora et al. [57]. Previously secondary metabolites such as flavonoids derivatives were evaluated in Zygophyllaceae [58]. A recent study on phytochemical screening of Zygophyllaceae family proved its potential as antioxidants and might have a role *in vivo* studies as potent therapeutic agents validating their ethnopharmacological usages [59]. Due to their antioxidant capabilities, flavonoids, and phenolic constituents in the plant, suppressing the dangerous oxidative stress and protecting the body from the detrimental free radicals' effects by reducing reactive oxygen species may be essential for biological activities. In accordance with Onguéné et al. [60] who screened the antimicrobial, anti-inflammatory, and antimalarial effects of alkaloids, terpenoids, and coumarin isolated from African plants. Moreover, Matsuda et al. [61] *in vitro* investigated the role of flavonoids stilbene from medicinal plants against Basophilic Leukemia cells. Furthermore, Xu et al. [62] reported the presence of polyphenols and flavonoids from medical plants with anti-inflammatory applications.

Interestingly, eight quinovic acid-based triterpenoid saponins were detected with a relative percentage value of 16.74% of the total plant extract's constituents that were characteristic of this plant species. Moreover, the previous report of Mohammed et al. [32] identified five quinovic acid-based triterpenoid saponins with a 27.12% relative percentage of the total *Z. coccineum* extract's constituents. Also, these types of saponins (quinovic acid-based triterpenoid saponins) were reported from the genus *Zygophyllum* and might be considered a chemical marker for the genus [63].

This study evaluated the effect of using different doses of *Z. coccineum* extract on arthritis score and paw volume to downregulated inflammation signs compared to normal and treatment using the standard drug. Gautam et al. [64] reported the role of bioactive ingredients in butanol fraction of *Punica granatum* in enhancing biophysical arthritic symptoms. Besides, Manan et al. [65] examined the beneficial roles of phenolics and flavonoids derived from *Alternanthera bettzickiana* in regulating inflammatory symptoms. Furthermore, Singh et al. [66] explained the oxidative role of phytoconstituents in enhancing arthritic symptoms. In the present report, *Z. coccineum* extract showed promising *in vitro* anti-inflammatory action in accordance with Shahzad et al. [67], who reported antiviral and anti-inflammatory effects of herbs, including *Zygophyllum* growing in Asia and Africa. Besides, Elbadry et al. [68] recorded *in vitro* antimicrobial and larvicidal activity of Egyptian *Zygophyllum* extract.

The anti-inflammatory action of the Zygophyllaceae family can be attributed to their wide range biological activities [69]. Different species of Zygophyllaceae (*Tribulus longipetalus* and *T. terrestris*) were tested for albumin denaturation assay, and they exhibited significant albumin

inhibition and representing significant anti-inflammatory activity [70]. Also, *Tribulus terrestris* extract ameliorated mice macrophages and triggered excessive release of NO and inflammatory cytokines including interleukin-1 beta, interleukin-6, and tumor necrosis factor-alpha [59]. Consequently, we concluded that *Z. coccineum* extract offered an anti-inflammatory effect related to downregulation of NF- κ B and mRNA expression inhibition of inflammatory factors including TNF- α , IL-1 β , and IL-6. The inhibition of Akt/M.A.P.K. is the crucial mechanism suggested the anti-inflammatory activities of natural products [71, 72]. Previously, good anti-inflammatory activity results were documented by Mnafgui et al., [73] for *Z. album* by measuring serum level of C-reactive protein and tumor necrosis factor. Also, the *Z. macropodium* ethanolic extract displayed significant inhibitory activity on increased vascular permeability in mice induced writhing by acetic acid compared to normal control [74].

Previous investigations have also revealed that osteoclasts are cells that have essential roles in bone remodeling [75]. In the current study using of 15.6 mg/kg of *Z. coccineum* extract in induced animals which led to recover osteoclast its regular structure and apoptotic rate, microscopic examinations of joints and muscles revealed that restored chondrocytes and muscles structures restored in induced rats upon using 15.6 mg/kg of *Z. coccineum* extract. Furthermore, Bourebaba et al. [76] *in vitro* investigated the role of *Cladophora glomerata* extract in chondrogenic enhancement. Besides, Saud et al. [77] explained the role of bioactive constituents in controlling gene expression in muscle cells and chondrocytes. Different dose levels of 15.6 mg/kg, 31 mg/kg, and 60 mg/kg *Z. coccineum* extract were tested, and various variations were reported in the current report. In same line with, Amalraj et al. [78] who reported that either low or high doses of curcumin has enhanced signs of inflammation relative to placebo. However, Antiarthritic activity of 50,100, and 150 mg/kg doses of *Berberis orthobotrys* has been tested in the arthritis animal model where the highest action was seen at 150 mg/kg [79].

Similarly, Lee et al. [69] reported that a compound isolated from a Zygophyllaceae plant *Tribulus terrestris*, known as tribulusamide D, inhibited LPS-induced inflammatory response in RAW264.7 cell model via suppression of NF- κ B and P38 signaling cascades. Besides, Ito et al. [80] explained the role of ginger in the differentiation of osteoclasts, while Kim et al. [81] reported the role of *Leonurus sibiricus* in gene expression regulating osteoclast differentiation. In the present study, *Z. coccineum* extract regulated the secretion of pro- and anti-inflammatory cytokines, shifting cytokine profile into protective form. In accordance with Byun et al.; Kulyar et al., [82, 83], who reported the role of medicinal plants in reinforcing immune synapse regulating many diseases. Methotrexate has been used in the treatment of many autoimmune diseases. It has many adverse effects on body organs [84]. Thus, using alternative therapies from natural products has become a significant need in many categories. In the current study, we noticed the beneficial effects of *Z. coccineum* extract in the arthritic model with minimal burden on kidney and liver functions.

5. Conclusions

Rheumatoid arthritis (R.A.) is an autoimmune disorder with inflammation in synovium and impacts numerous people all over the world, and there is an emerging trend in using natural products in treatments as safe alternatives. In this study, we have *in vitro* and *in vivo* investigated the anti-inflammatory impact of *Z. coccineum* extract reporting its role in enhancing outer inflammation signs as well as in chondrocytes, osteoclasts, and inflammatory cytokines in a dose-dependent effect with a notable safety effect on the internal body functions as shown in supplement (2) for proposed flow chart of the mechanism. *Z. coccineum* extract could be used as an effective natural source for the treatment of rheumatoid arthritis with high safety profile after more confirmation and verification of results in future studies.

Data Availability

All data that support the findings of this study are available from the corresponding authors upon reasonable request.

Conflicts of Interest

The authors declare that they have no competing interests.

Acknowledgments

“This work was supported by grants from the Academy of Scientific Research and Technology (grant number #1956) and Jessor#1956. This work was supported by Researchers Supporting Project number (RSP2022R439), King Saud University, Riyadh, Saudi Arabia.”

Supplementary Materials

Supplement (1): various groups of animals and different treatments using *Z. coccineum* extract in the A.I.A model. Supplement (2): flow diagram illustrating proposed various cells playing some roles in treatment of A.I.A after using *Z. coccineum* extract. (*Supplementary Materials*)

References

- [1] Q. Fang, C. Zhou, and K. S. Nandakumar, “Molecular and cellular pathways contributing to joint damage in rheumatoid arthritis,” *Mediators of Inflammation*, vol. 2020, Article ID 3830212, 20 pages, 2020.
- [2] C. Orr, E. Vieira-Sousa, D. L. Boyle et al., “Synovial tissue research: a state-of-the-art review,” *Nature Reviews Rheumatology*, vol. 13, no. 8, pp. 463–475, 2017.
- [3] G. R. Burmester, E. Feist, and T. Dörner, “Emerging cell and cytokine targets in rheumatoid arthritis,” *Nature Reviews Rheumatology*, vol. 10, no. 2, pp. 77–88, 2014.
- [4] S. E. Wilson, R. R. Mohan, M. Netto et al., “RANK, RANKL, OPG, and M-CSF expression in stromal cells during corneal wound healing,” *Investigative Ophthalmology & Visual Science*, vol. 45, no. 7, pp. 2201–2211, 2004.
- [5] N. Zhao, H. Tsuda, T. Murofushi et al., “Chaetocin inhibits RANKL-induced osteoclast differentiation through reduction of Blimp1 in Raw264.7 cells,” *Life Sciences*, vol. 143, no. 143, pp. 1–7, 2015.
- [6] L. Kong, W. Smith, and D. Hao, “Overview of RAW264.7 for osteoclastogenesis study: phenotype and stimuli,” *Journal of Cellular and Molecular Medicine*, vol. 23, no. 5, pp. 3077–3087, 2019.
- [7] I. B. McInnes, C. D. Buckley, and J. D. Isaacs, “Cytokines in rheumatoid arthritis – shaping the immunological landscape,” *Nature Reviews Rheumatology*, vol. 12, no. 1, pp. 63–68, 2016.
- [8] J. Alam, I. Jantan, and S. N. A. Bukhari, “Rheumatoid arthritis: recent advances on its etiology, role of cytokines and pharmacotherapy,” *Biomedicine & Pharmacotherapy*, vol. 92, pp. 615–633, 2017.
- [9] M. H. Farzaei, F. Farzaei, M. Abdollahi, Z. Abbasabadi, A. H. Abdolghaffari, and B. Mehraban, “A mechanistic review on medicinal plants used for rheumatoid arthritis in traditional Persian medicine,” *The Journal of Pharmacy and Pharmacology*, vol. 68, no. 10, pp. 1233–1248, 2016.
- [10] X. Z. Li and S. N. Zhang, “Herbal compounds for rheumatoid arthritis: literatures review and cheminformatics prediction,” *Phytotherapy Research*, vol. 34, no. 1, pp. 51–66, 2020.
- [11] M. Akram, M. Daniyal, S. Sultana et al., “Traditional and modern management strategies for rheumatoid arthritis,” *Clinica Chimica Acta*, vol. 512, pp. 142–155, 2021.
- [12] A. A. Hamza, G. H. Heeba, S. Hamza, A. Abdalla, and A. Amin, “Standardized extract of ginger ameliorates liver cancer by reducing proliferation and inducing apoptosis through inhibition oxidative stress/ inflammation pathway,” *Biomedicine & Pharmacotherapy*, vol. 134, article 111102, 2021.
- [13] S. M. El-Dakhly, A. A. A. Salama, S. O. M. Hassanin, N. N. Yassen, A. A. Hamza, and A. Amin, “Aescin and diosmin each alone or in low dose- combination ameliorate liver damage induced by carbon tetrachloride in rats,” *BMC Research Notes*, vol. 13, no. 1, p. 259, 2020.
- [14] A. A. Hamza, F. M. Lashin, M. Gamel, S. O. Hassanin, Y. Abdalla, and A. Amin, “Hawthorn herbal preparation from *Crataegus oxyacantha* attenuates in vivo carbon tetrachloride-induced hepatic fibrosis via modulating oxidative stress and inflammation,” *Antioxidants (Basel)*, vol. 9, no. 12, p. 1173, 2020.
- [15] A. A. Hamza, M. G. Mohamed, F. M. Lashin, and A. Amin, “Dandelion prevents liver fibrosis, inflammatory response, and oxidative stress in rats,” *The Journal of Basic and Applied Zoology*, vol. 81, no. 1, p. 43, 2020.
- [16] C. Murali, P. Mudgil, C. Y. Gan et al., “Camel whey protein hydrolysates induced G2/M cellcycle arrest in human colorectal carcinoma,” *Scientific Reports*, vol. 11, no. 1, p. 7062, 2021.
- [17] A. Amin, A. Farrukh, C. Murali et al., “Saffron and its major Ingredients' effect on colon cancer cells with mismatch repair deficiency and microsatellite instability,” *Molecules*, vol. 26, no. 13, p. 3855, 2021.
- [18] A. Amin, M. Lotfy, D. Mahmoud-Ghoneim et al., “Pancreas-protective effects of chlorella in STZ-induced diabetic animal model: insights into the mechanism,” *Journal of Diabetes Mellitus*, vol. 1, no. 3, pp. 36–45, 2011.
- [19] C. Mu, Y. Sheng, Q. Wang, A. Amin, X. Li, and Y. Xie, “Potential compound from herbal food of *Rhizoma Polygonati* for treatment of COVID-19 analyzed by network pharmacology: viral and cancer signaling mechanisms,” *Journal of Functional Foods*, vol. 77, article 104149, 2021.

- [20] B. Al-Dabbagh, I. A. Elhaty, C. Murali, A. Al Madhoon, and A. Amin, "Salvadora persica (Miswak): antioxidant and promising antiangiogenic insights," *American Journal of Plant Sciences*, vol. 9, no. 6, pp. 1228–1244, 2018.
- [21] Y. Xie, C. Mu, B. Kazybay et al., "Network pharmacology and experimental investigation of Rhizoma polygonati extract targeted kinase with herbzyme activity for potent drug delivery," *Drug Delivery*, vol. 28, no. 1, pp. 2187–2197, 2021.
- [22] A. Abdalla, C. Murali, and A. Amin, "Safranal inhibits angiogenesis via targeting HIF-1 α /VEGF machinery: in vitro and ex vivo insights," *Frontiers in Oncology*, vol. 11, article 789172, 2021.
- [23] A. A. Hamza, S. O. Hassanin, S. Hamza, A. Abdalla, and A. Amin, "Polyphenolic-enriched olive leaf extract attenuated doxorubicin-induced cardiotoxicity in rats via suppression of oxidative stress and inflammation," *The Journal of Basic and Applied Zoology*, vol. 82, no. 1, p. 54, 2021.
- [24] I. Hammad and S. H. Qari, "Genetic diversity among Zygophyllum (Zygophyllaceae) populations based on RAPD analysis," *Genetics and Molecular Research*, vol. 9, no. 4, pp. 2412–2420, 2010.
- [25] H. M. Hammada, N. M. Ghazy, F. M. Harraz, M. M. Radwan, M. A. ElSohly, and I. I. Abdallah, "Chemical constituents from Tribulus terrestris and screening of their antioxidant activity," *Phytochemistry*, vol. 92, pp. 153–159, 2013.
- [26] Y. A. El-Amier, H. M. El-Shora, and M. Hesham, "Ecological study on Zygophyllum coccineum L. in coastal and inland desert of Egypt," *Journal of Agriculture and Ecology Research*, vol. 6, no. 4, pp. 1–17, 2016.
- [27] E. Amin, S. S. El-Hawary, M. M. Fathy et al., "Triterpenoidal saponins: bioactive secondary metabolites from Zygophyllum coccineum," *Planta Medica*, vol. 77, no. 5, pp. 488–491, 2011.
- [28] E. Mohamed, A. M. M. A. Kasem, A. A. Gobouri, A. Elkesh, and E. Azab, "Influence of maternal habitat on Salinity tolerance of Zygophyllum coccineum with Regard to Seed germination and growth Parameters," *Plants (Basel)*, vol. 9, no. 11, p. 1504, 2020.
- [29] A. N. Amro and M. K. Abhary, "Removal of lead and copper ions from water using powdered Zygophyllum coccineum biomass," *International Journal of Phytoremediation*, vol. 21, no. 14, pp. 1457–1462, 2019.
- [30] H. A. Mansour, A. S. Newairy, M. I. Yousef, and S. A. Sheweita, "Biochemical study on the effects of some Egyptian herbs in alloxan-induced diabetic rats," *Toxicology*, vol. 170, no. 3, pp. 221–228, 2002.
- [31] S. Gibbons and M. A. Oriowo, "Antihypertensive effect of an aqueous extract of Zygophyllum coccineum L. in rats," *Phytotherapy Research*, vol. 15, no. 5, pp. 452–455, 2001.
- [32] H. A. Mohammed, R. A. Khan, A. A. Abdel-Hafez et al., "Phytochemical Profiling, In Vitro and In Silico Anti-Microbial and Anti-Cancer Activity Evaluations and Staph Gyrase B and h-TOP-II β Receptor-Docking Studies of Major Constituents of Zygophyllum coccineum L. Aqueous-Ethanol Extract and Its Subsequent Fractions: An Approach to Validate Traditional Phytomedicinal Knowledge," *Molecules*, vol. 26, no. 3, p. 577, 2021.
- [33] L. Li, S. Liang, F. Du, and C. Li, "Simultaneous quantification of multiple licorice flavonoids in rat plasma," *Journal of the American Society for Mass Spectrometry*, vol. 18, no. 4, pp. 778–782, 2007.
- [34] H. Tsugawa, T. Cajka, T. Kind et al., "MS-DIAL: data-independent MS/MS deconvolution for comprehensive metabolome analysis," *Nature Methods*, vol. 12, no. 6, pp. 523–526, 2015.
- [35] S. Granica, M. E. Czerwińska, J. P. Piwowarski, M. Ziąja, and A. K. Kiss, "Chemical composition, antioxidative and anti-inflammatory activity of extracts prepared from aerial parts of Oenothera biennis L. and Oenothera paradoxa Hudziak obtained after seeds cultivation," *Journal of Agricultural and Food Chemistry*, vol. 61, no. 4, pp. 801–810, 2013.
- [36] A. Bendele, "Animal models of rheumatoid arthritis," *Journal of Musculoskeletal & Neuronal Interactions*, vol. 1, no. 4, pp. 377–385, 2001.
- [37] H. Suzuki, N. Hirano, C. Watanabe, and Y. Tarumoto, "Carbon tetrachloride does not induce micronucleus in either mouse bone marrow or peripheral blood," *Mutation Research*, vol. 394, no. 1-3, pp. 77–80, 1997.
- [38] C. Mossiat, D. Laroche, C. Prati, T. Pozzo, C. Demougeot, and C. Marie, "Association between arthritis score at the onset of the disease and long-term locomotor outcome in adjuvant-induced arthritis in rats," *Arthritis Research & Therapy*, vol. 17, no. 1, p. 184, 2015.
- [39] Z. Zhang, A. Chinnathambi, S. A. Alharbi, and L. Bai, "Copper oxide nanoparticles from Rabdosia rubescens attenuates the complete Freund's adjuvant (CFA) induced rheumatoid arthritis in rats via suppressing the inflammatory proteins COX-2/PGE2," *Arabian Journal of Chemistry*, vol. 13, no. 6, pp. 5639–5650, 2020.
- [40] K. H. Collins, H. A. Paul, R. A. Reimer, R. A. Seerattan, D. A. Hart, and W. Herzog, "Relationship between inflammation, the gut microbiota, and metabolic osteoarthritis development: studies in a rat model," *Osteoarthritis and Cartilage*, vol. 23, no. 11, pp. 1989–1998, 2015.
- [41] X. Cong, X. M. Zhang, Y. Zhang et al., "Disruption of endothelial barrier function is linked with hyposecretion and lymphocytic infiltration in salivary glands of Sjögren's syndrome," *Biochimica et Biophysica Acta - Molecular Basis of Disease*, vol. 1864, no. 10, pp. 3154–3163, 2018.
- [42] T. Sun, Z. Li, X. Zhong et al., "Strontium inhibits osteoclastogenesis by enhancing LRP6 and β -catenin-mediated OPG targeted by miR-181d-5p," *Journal of Cell Communication and Signaling*, vol. 13, no. 1, pp. 85–97, 2019.
- [43] W. Liu, C. Xu, H. Zhao et al., "Osteoprotegerin induces apoptosis of osteoclasts and osteoclast precursor cells via the Fas/Fas ligand pathway," *PLoS One*, vol. 10, no. 11, article e0142519, 2015.
- [44] K. Yuan, J. Mei, D. Shao et al., "Cerium oxide nanoparticles regulate osteoclast differentiation bidirectionally by modulating the cellular production of reactive oxygen species," *International Journal of Nanomedicine*, vol. 15, pp. 6355–6372, 2020.
- [45] Y. Zhao, Y. Sun, G. Wang, S. Ge, and H. Liu, "Dendrobium Officinale polysaccharides protect against MNNG-induced PLGC in rats via activating the NRF2 and antioxidant enzymes HO-1 and NQO-1," *Oxidative Medicine and Cellular Longevity*, vol. 2019, Article ID 9310245, 11 pages, 2019.
- [46] W. B. van den Berg, L. A. Joosten, and P. L. van Lent, "Murine antigen-induced arthritis," *Methods in Molecular Medicine*, vol. 136, pp. 243–253, 2007.
- [47] N. Choudhary, L. K. Bhatt, and K. S. Prabhavalkar, "Experimental animal models for rheumatoid arthritis," *Immunopharmacology and Immunotoxicology*, vol. 40, no. 3, pp. 193–200, 2018.

- [48] B. Grötsch, A. Bozec, and G. Schett, "In vivo models of rheumatoid arthritis," *Methods in Molecular Biology*, vol. 1914, pp. 269–280, 2019.
- [49] F. Maione, R. Russo, H. Khan, and N. Mascolo, "Medicinal plants with anti-inflammatory activities," *Natural Product Research*, vol. 30, no. 12, pp. 1343–1352, 2016.
- [50] V. Giorgi, D. Marotto, A. Batticciotto, F. Atzeni, S. Bongiovanni, and P. Sarzi-Puttini, "Cannabis and autoimmunity: possible mechanisms of action," *ImmunoTargets and Therapy*, vol. 10, pp. 261–271, 2021.
- [51] E. Amin, S. S. El-Hawary, M. M. Fathy, R. Mohammed, Z. Ali, and I. A. Khan, "Zygophylloside S, a New Triterpenoid Saponin from the Aerial Parts of *Zygophyllum coccineum* L.," *Planta Medica*, vol. 76, 2010.
- [52] E. Amin, Y. H. Wang, B. Avula et al., "Simultaneous determination of bioactive secondary metabolites involved in the antioxidant, anti-inflammatory and anticancer activities of the edible halophyte *Zygophyllum album* Desf.," *Food Chemistry*, vol. 139, pp. 1073–1080, 2013.
- [53] K. Pöllmann, S. Gagel, M. H. A. Elgamil, K. H. Shaker, and K. Seifert, "Triterpenoid saponins from the roots of *Zygophyllum* species," *Phytochemistry*, vol. 40, pp. 1233–1236, 1995.
- [54] H. A. Hassanean and E. K. Desoky, "An acylated isorhamnetin glucoside from *Zygophyllum simplex*," *Phytochemistry*, vol. 31, no. 9, pp. 3293–3294, 1992.
- [55] S. R. Hussein, M. M. Marzouk, L. Ibrahim, S. A. Kawashty, and N. A. M. Saleh, "Flavonoids of *Zygophyllum album* L.f. and *Zygophyllum simplex* L. (*Zygophyllaceae*)," *Biochemical Systematics and Ecology*, vol. 39, no. 4-6, pp. 778–780, 2011.
- [56] W. M. Ksouri, F. Medini, K. Mkadmini et al., "LC-ESI-TOF-MS identification of saponins and a flavonoid from aerial parts of *Zygophyllum coccineum* L.," *Journal of AOAC International*, vol. 95, pp. 757–762, 2012.
- [57] H. M. El-Shora, Y. A. El-Amier, and M. H. Awad, "Comparative phytochemical studies on *Zygophyllum coccineum* L. from different habitats, Egypt," *British Journal of Applied Science & Technology*, vol. 15, pp. 1–9, 2016.
- [58] J. H. Salem, I. Chevalot, C. Harscoat-Schiavo, C. Paris, M. Fick, and C. Humeau, "Biological activities of flavonoids from *Nitraria retusa* (Forssk.) Asch. and their acylated derivatives," *Food Chemistry*, vol. 124, no. 2, pp. 486–494, 2011.
- [59] W. R. Zhao, W. T. Shi, J. Zhang et al., "Tribulus terrestris L. extract protects against lipopolysaccharide-induced inflammation in RAW 264.7 macrophage and zebrafish via inhibition of Akt/MAPKs and NF- κ B/iNOS-NO signaling pathways," *Evidence-based Complementary and Alternative Medicine*, vol. 2021, Article ID 6628561, 11 pages, 2021.
- [60] B. D. Bekono, F. Ntie-Kang, P. A. Onguéné et al., "The potential of anti-malarial compounds derived from African medicinal plants: a review of pharmacological evaluations from 2013 to 2019," *Malaria Journal*, vol. 19, no. 1, p. 183, 2020.
- [61] H. Matsuda, S. Nakamura, and M. Yoshikawa, "Degranulation inhibitors from medicinal plants in antigen-stimulated rat basophilic leukemia (RBL-2H3) cells," *Chemical and Pharmaceutical Bulletin*, vol. 64, no. 2, pp. 96–103, 2016.
- [62] D. P. Xu, Y. Li, X. Meng et al., "Natural antioxidants in foods and medicinal plants: extraction, assessment and resources," *International Journal of Molecular Sciences*, vol. 18, no. 1, p. 96, 2017.
- [63] K. Pöllmann, K. Schaller, U. Schweizer, M. H. A. Elgamil, K. H. Shaker, and K. Seifert, "Triterpenoid saponins from *Zygophyllum decumbens*," *Phytochemistry*, vol. 48, no. 5, pp. 875–880, 1998.
- [64] R. K. Gautam, S. Sharma, K. Sharma, and G. Gupta, "Evaluation of antiarthritic activity of butanol fraction of *Punica granatum* Linn. rind extract against Freund's complete adjuvant-induced arthritis in rats," *Journal of Environmental Pathology, Toxicology and Oncology*, vol. 37, no. 1, pp. 53–62, 2018.
- [65] M. Manan, U. Saleem, M. S. H. Akash et al., "Antiarthritic potential of comprehensively standardized extract of *Alternanthera bettzickiana*," *A.C.S. Omega*, vol. 5, no. 31, pp. 19478–19496, 2020.
- [66] S. Singh, T. G. Singh, K. Mahajan, and S. Dhiman, "Medicinal plants used against various inflammatory biomarkers for the management of rheumatoid arthritis," *The Journal of Pharmacy and Pharmacology*, vol. 72, no. 10, pp. 1306–1327, 2020.
- [67] F. Shahzad, D. Anderson, and M. Najafzadeh, "The antiviral, anti-inflammatory effects of natural medicinal herbs and mushrooms and SARS-CoV-2 infection," *Nutrients*, vol. 12, no. 9, p. 2573, 2020.
- [68] M. A. Elbadry, M. M. Elaasser, H. H. Elshiekh, and M. M. Sheriff, "Evaluation of Antimicrobial, Cytotoxic and Larvicidal Activity of *Zygophyllum Coccineum* North Sinai, Egypt," *Medicinal and Aromatic Plants*, vol. 4, no. 5, p. 214, 2015.
- [69] H. H. Lee, E. K. Ahn, S. S. Hong, and J. S. Oh, "Anti-inflammatory effect of tribulusamide D isolated from *Tribulus terrestris* in lipopolysaccharide-stimulated RAW264.7 macrophages," *Molecular Medicine Reports*, vol. 16, no. 4, pp. 4421–4428, 2017.
- [70] M. S. Mohammed, M. F. Alajmi, P. Alam, H. S. Khalid, A. M. Mahmoud, and W. J. Ahmed, "Chromatographic finger print analysis of anti-inflammatory active extract fractions of aerial parts of *Tribulus terrestris* by HPTLC technique," *Asian Pacific journal of tropical biomedicine*, vol. 4, no. 3, pp. 203–208, 2014.
- [71] M. Tanaka, Y. Kishimoto, M. Sasaki et al., "Terminalia bellirica (Gaertn.) Roxb. extract and gallic acid attenuate LPS-induced inflammation and oxidative stress via MAPK/NF- κ B and Akt/AMPK/Nrf2 pathways," *Oxidative Medicine and Cellular Longevity*, vol. 2018, Article ID 9364364, 15 pages, 2018.
- [72] Y. Nie, Z. Wang, G. Chai et al., "Dehydrocostus lactone suppresses LPS-induced acute lung injury and macrophage activation through NF- κ B signaling pathway mediated by p38 MAPK and Akt," *Molecules*, vol. 24, no. 8, p. 1510, 2019.
- [73] K. Mnafigui, M. Kchaou, H. Ben Salah et al., "Essential oil of *Zygophyllum album* inhibits key-digestive enzymes related to diabetes and hypertension and attenuates symptoms of diarrhea in alloxan-induced diabetic rats," *Pharmaceutical Biology*, vol. 54, no. 8, pp. 1326–1333, 2016.
- [74] Y. Xiao-rong, Z. Xiang-feng, Z. Xue-mei, and G. Hong-yan, "Analgesic and anti-inflammatory activities and mechanisms of 70% ethanol extract of *Zygophyllum macropodium* in animals," *Chinese Herbal Medicines*, vol. 10, no. 1, pp. 59–65, 2018.
- [75] M. Jia, Y. Nie, D. P. Cao et al., "Potential antiosteoporotic agents from plants: a comprehensive review," *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 364604, 28 pages, 2012.
- [76] L. Bourebaba, I. Michalak, M. Baouche, K. Kucharczyk, and K. Marycz, "Cladophora glomerata methanolic extract promotes chondrogenic gene expression and cartilage phenotype differentiation in equine adipose-derived mesenchymal

- stromal stem cells affected by metabolic syndrome,” *Stem Cell Research & Therapy*, vol. 10, no. 1, p. 392, 2019.
- [77] B. Saud, R. Malla, and K. Shrestha, “A review on the effect of plant extract on mesenchymal stem cell proliferation and differentiation,” *Stem Cells International*, vol. 2019, Article ID 7513404, 13 pages, 2019.
- [78] A. Amalraj, K. Varma, J. Jacob et al., “A novel highly bioavailable curcumin formulation improves symptoms and diagnostic indicators in rheumatoid arthritis patients: a randomized, double-blind, placebo-controlled, two-dose, three-arm, and parallel-group study,” *Journal of Medicinal Food*, vol. 20, no. 10, pp. 1022–1030, 2017.
- [79] A. M. Uttra and U. H. Hasan, “Anti-arthritis activity of aqueous-methanolic extract and various fractions of *Berberis orthobotrys* Bien ex aitch,” *BMC Complementary and Alternative Medicine*, vol. 17, p. 371, 2017.
- [80] S. Ito, A. Ohmi, A. Sakamiya et al., “Ginger hexane extract suppresses RANKL-induced osteoclast differentiation,” *Bioscience, Biotechnology, and Biochemistry*, vol. 80, no. 4, pp. 779–785, 2016.
- [81] J. H. Kim, M. Kim, H. S. Jung, and Y. Sohn, “*Leonurus sibiricus* L. ethanol extract promotes osteoblast differentiation and inhibits osteoclast formation,” *International Journal of Molecular Medicine*, vol. 44, no. 3, pp. 913–926, 2019.
- [82] J. H. Byun, C. W. Choi, M. J. Jang, S. H. Lim, H. J. Han, and S. Y. Choung, “Anti-osteoarthritic mechanisms of *chrysanthemum zawadskii* var. *latilobum* in MIA-induced osteoarthritic rats and interleukin-1 β -induced SW1353 human chondrocytes,” *Medicina (Kaunas, Lithuania)*, vol. 56, no. 12, p. 685, 2020.
- [83] M. F. Kulyar, R. Li, K. Mehmood, M. Waqas, K. Li, and J. Li, “Potential influence of *Nagella sativa* (Black cumin) in reinforcing immune system: a hope to decelerate the COVID-19 pandemic,” *Phytomedicine*, vol. 85, article 153277, 2021.
- [84] D. P. Misra, A. Y. Gasparyan, and O. Zimba, “Benefits and adverse effects of hydroxychloroquine, methotrexate and colchicine: searching for repurposable drug candidates,” *Rheumatology International*, vol. 40, no. 11, pp. 1741–1751, 2020.