Research Article

CZYH Alleviates β -Amyloid-Induced Cognitive Impairment and Inflammation Response via Modulation of JNK and NF- κ B Pathway in Rats

Yuanyuan Deng,^{1,2} Lianzhi Ye,^{1,2} Cheng Yu,^{1,2} Caixia Yin,^{1,2} Jingshan Shi,^{1,2} and Qihai Gong^{1,2}

¹Key Laboratory of Basic Pharmacology of Ministry of Education, Department of Pharmacology, Zunyi Medical University, Zunyi, Guizhou 563000, China

²Joint International Research Laboratory of Ethnomedicine of Ministry of Education, Zunyi Medical University, Zunyi, Guizhou 563000, China

Correspondence should be addressed to Qihai Gong; zmcgqh@qq.com

Received 19 May 2019; Revised 17 August 2019; Accepted 9 September 2019; Published 4 November 2019

Academic Editor: Luigi Ferini-Strambi

Copyright © 2019 Yuanyuan Deng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cu-Zhi-Yi-Hao (CZYH), an empirical formula of traditional Chinese medicine (TCM), has been used for amnesia treatment in clinical practice. However, its underlying pharmacological mechanism has not been fully illuminated. The current study was designed to investigate the neuroprotective effect of CZYH on a β -amyloid 25-35- ($A\beta_{25-35}$ -) induced learning and memory deficit rat model. CZYH (200, 400, or 800 mg/kg), donepezil (1.0 mg/kg), or distilled water was given to $A\beta_{25-35}$ -stimulated animals for 17 days consecutively. The Morris water maze test revealed that CZYH (400 or 800 mg/kg) administration improved the $A\beta_{25-35}$ -induced cognitive impairments in rats, and Nissl staining demonstrated that CZYH mitigated the $A\beta$ -caused neuron loss. In addition, CZYH treatment markedly inhibited the activation of microglia as evidenced by a decreased level of IBA-1 and increased YM-1/2 protein expression. The protein expression levels of TNF- α , IL-1 β , and COX-2 were also repressed by CZYH. Besides, CZYH treatment alleviated $A\beta$ -induced I κ B- α degradation and NF- κ B p65 phosphorylation, as well as reduced the JNK phosphorylation level. In conclusion, the present study suggests that CZYH could improve learning and memory abilities and relieve neuron loss in $A\beta_{25-35}$ -induced rats, at least partly through inhibition of the neuroinflammatory response via inhibiting the JNK-dependent NF- κ B activation, indicating that CZYH might be a promising formula for the treatment of AD.

1. Introduction

Alzheimer's disease (AD) is the most prevalent form of neurodegenerative disease [1], with memory loss, cognitive impairment, and behavior disorders as the main clinical manifestations [2], which lead a person unable to execute fundamental bodily functions such as walking and eventually to be bed-bound, seriously decreasing the life quality of patients and increasing the social burden. There are about 44 million people worldwide suffering from AD or a related form of dementia. As the population ages, the number of AD patients will be more than triple to 152 million by 2050 as estimated by epidemiology research.

Thus, AD is becoming the most feared disease among elderly people [3].

The pathogenesis of AD is complicated and not fully clarified yet. $A\beta$ hypothesis is predominant in AD research, which suggests that the abnormal release and following accumulation of $A\beta$ are driving causes in the pathogenesis of AD [4]. $A\beta$ is a small peptide generated when the membrane-associated amyloid precursor protein (APP) is cleaved by β -secretase (BACE-1) and γ -secretase. Meanwhile, $A\beta$ could be degraded by enzymes, such as the insulin-degrading enzyme (IDE) and neprilysin (NEP) [5, 6]. The level of $A\beta$ peptides abnormally increase when the equilibrium between production and clearance is altered,

and then the aggregation of $A\beta$ triggers a series of toxic cascade events, including microglial activation, tau phosphorylation, and neuronal degeneration and loss, which lead to memory deficits eventually [7]. Recent evidence indicate AD is also an inflammatory process, as excessive release of $A\beta$ spontaneously aggregates into soluble oligomers, activating the microglia, accompanied by the liberation of inflammatory factors and corresponding inflammation response. These elevated inflammatory factors hamper the resident microglia to remove the $A\beta$ -formed senile plaques, which in turn amplify the inflammation response and $A\beta$ toxicity, speeding up the AD process [8, 9].

Of the more than one hundred drugs that have been tested in the past 2-3 decades, only four have clinical uses; these are cholinesterase inhibitors, including donepezil, rivastigmine, galantamine, and an N-methyl-D-aspartate receptor antagonist memantine. They can alleviate the symptoms of dementia in some patients, but only for some people, and they also bring multiple side effects at the same time due to their single target [10]. Thus, it is imperative to have a combined strategy which takes the body as a whole instead of just one factor.

Traditional Chinese medicine (TCM), based on the idea that everything in our body is connected and considers the overall condition to manage diseases, has been postulated as a promising strategy for AD treatment in Asian countries as it has advantages to keep balance in the human body and then reduce side effects. The Cu-Zhi-Yi-Hao (CZYH) decoction is an empirical formula of TCM in amnesia treatment, composed of eight Chinese herbal medicines including Epimedium, Pueraria, Rehmannia, male Antheraea silkmoths, Salt psyllium, Goji (Wolfberry), Fu Ling (*Poria*), and Coix seed (Table 1). In this decoction, Epimedium and Pueraria are the main components. Epimedium is the leaf part from *Epimedium brevicornum Maxim* and has been used to treat sexual dysfunction, cardiovascular diseases, and dementia for many years. Increasing evidence demonstrated that the two main active compounds of Epimedium, icariin and icarisiide II, improve cognition in APP/PS1 transgenic mouse models as well as different cognitive deficit rat models induced by streptozocin, $A\beta$, ibotenic acid, etc. [11–14]. Pueraria itself or its extract has neurotherapeutic effects on cognitive disabled animals [15], Puerarin mitigates $A\beta$ -induced cognitive decline via inhibition of apoptosis response [16], and the ethanol extract of Pueraria acts against trimethyltin-generated learning and memory impairments by repression of acetylcholinesterase (AChE) [17]. A clinical trial of a TCM formula which is composed of Epimedium and another five herbs has displayed a promising therapeutic effect in mild AD patients, and the improving scores were even better than donepezil [18], drawing more attention to use the TCM formula to treat cognitive deficits.

Therefore, in the current study, the effect of the formula CZYH was assessed on an $A\beta_{25-35}$ -induced learning and memory impairment rat model. The possible underlying mechanisms were also investigated.

TABLE 1: Composition of Cu-Zhi-Yi-Hao.

Common name	Latin name	Quantity (g)
Epimedium	<i>Epimedium brevicornum Maxim</i>	10
Pueraria	<i>Pueraria edulis Pampan</i>	10
Rehmannia	<i>Rehmanniae Radix</i>	8
Male Antheraea silkmoths	<i>Bombycidae</i>	8
Salt psyllium	<i>Semen Plantaginis</i>	4
Goji	<i>Lycii Fructus</i>	6
Fu Ling	<i>Poria</i>	8
Coix seed	<i>Semen Coicis</i>	8

2. Materials and Methods

2.1. Reagents. Donepezil hydrochloride was bought from Salvage Pharmaceutical Company (Guizhou, China), ground, and dissolved in double-distilled water before use. $A\beta_{25-35}$ was purchased from Sigma-Aldrich (St. Louis, MO, USA) and aggregated by incubation in sterilized normal saline (2 $\mu\text{g}/\mu\text{L}$) at 37°C for 5 days before injection [11].

2.2. Preparation of CZYH Decoction. CZYH is composed of eight constituents (as seen in Table 1), including Epimedium 10 g, Pueraria 10 g, Rehmannia 8 g, male Antheraea silkmoths 8 g, Salt psyllium 4 g, Goji 6 g, Fu Ling 8 g, and Coix seed 8 g. This formula was provided by the Affiliated Hospital of Zunyi Medical University and identified by Professor Jianwen Yang (Zunyi Medical University, Zunyi, China). The mixture was soaked for 1 h with 500 mL distilled water and boiled with gentle heating for 2 h, the dregs were filtered with gauze and the filtrate was collected, then another 375 mL distilled water was added to the dregs and decocted for 2 h, and the above procedure was repeated once, combining the filtrate three times. The extraction was further concentrated, combined, and lyophilized as previously described to get the lyophilized powder [19]. The yield was 25.1% compared with the original crude amount.

2.3. Animals. Ninety-eight specific pathogen-free- (SPF-) grade male Sprague Dawley (SD) rats (250–300 g) were provided by the Laboratory Animal Center of the Daping Hospital (license no. SCXK2012-0011) and maintained in a SPF condition (temperature $22 \pm 1^\circ\text{C}$, humidity $58 \pm 5\%$, and 12 h light/12 h dark cycle) in the Animal Center of Key Laboratory of Basic Pharmacology of Ministry of Education, Zunyi Medical University. All animal handling procedures followed the Animal Management Rules of Zunyi Medical University (argument number [2015] 2-043).

2.4. Experimental Design. Animals were randomly allocated into seven groups: sham, sham+CZYH-H (800 mg/kg), model ($A\beta_{25-35}$), model+CZYH-L (200 mg/kg), model+CZYH-M (400 mg/kg), model+CZYH-H (800 mg/kg), and model+donepezil (1.0 mg/kg) as the positive drug group ($n = 14$ for each). The sample size was determined by previous experience [20]. The $A\beta_{25-35}$ -induced model

FIGURE 1: CZYH alleviates learning and memory impairments in A β_{25-35} -induced rats: (a) experimental design; (b) escape latency; (c) time percentage spent in target quadrant; (d) swimming speed. Data are means $\pm$ SEM, $n = 12-14$. $^{\#}P < 0.05$ and $^{##}P < 0.01$ vs. sham; $^*P < 0.05$ and $^{**}P < 0.01$ vs. A β_{25-35} .

was conducted following our previous report [11]. Briefly, animals were given a bilateral hippocampal injection of 5 μ L A β_{25-35} (2 μ g/ μ L) or the same volume NS (sham groups) after being anesthetized with 2% pentobarbital sodium (i.p.). CZYH or donepezil was administered orally daily for 17 days according to the above-described dosage regimen after the animals were woken up from the surgery (Figure 1(a)).

2.5. Morris Water Maze. The Morris water maze (MWM) task was applied to estimate the spatial learning and memory function as previously described [12]. The animals were trained and exposed to spatial acquisition trials from day 11 to day 16 after surgery to examine their ability to escape and find the hidden platform. On day 17, a spatial probe trial was carried out to observe the final memory consolidation. All the activities during the trials were automatically recorded and analyzed by TopScan Version 3.00 system.

2.6. Nissl Staining. All rats were deeply anesthetized with 2% pentobarbital sodium (intraperitoneally) after the MWM test and perfused with precold 0.01 M phosphate-buffered saline

(PBS). The brains were dissociated, and five brains from each group were fixed in 4% paraformaldehyde at 4°C overnight and subsequently embedded in paraffin. Sections (5 μ m thick, coronal) of brain tissue were stained with toluidine blue. Blue-purple-stained Nissl bodies of the CA1 and DG regions in the hippocampus were observed and counted to estimate the morphological changes [21].

2.7. Western Blotting. The hippocampus tissues were separated and then homogenized in RIPA buffer. The protein concentration of each sample was measured with the BCA method and then heat-blocked and separated as we described before [22]. Immunoblots were probed using primary antibodies against IL-1 β (1:2,000) and COX (1:1,000) (obtained from the Abcam company (USA)) and I κ B- α (1:2,000), NF- κ B p65 (1:1,000), p-NF- κ B p65 (1:1,000), Akt (1:1,000), p-Akt (1:1,000), total p38 MAPK (1:1,000), p-p38 MAPK (1:1,000), JNK (1:1,000), and p-JNK (1:1,000) (from Cell Signaling Technology (USA)), as well as internal antibodies β -actin (1:5000) and GAPDH (1:5000) (purchased from Beyotime (China)). Subsequently, the membranes were incubated with corresponding

(a)

(b)

(c)

FIGURE 2: CZYH mitigates $A\beta_{25-35}$ -induced neuronal loss in rats. (a) Nissl staining of hippocampus CA1 and DG regions (scale bar: 100 μm and 50 μm). (b, c) Quantitative analysis of Nissl bodies in (b) the CA1 hippocampal region and (c) the DG hippocampal region. Values are means $\pm$ SEM, $n = 5$. ^{##} $P < 0.01$ vs. sham; ^{*} $P < 0.05$ and ^{**} $P < 0.01$ vs. $A\beta_{25-35}$.

FIGURE 3: CZYH suppresses Aβ₂₅₋₃₅-induced proinflammatory microglial activation in the hippocampus of rats. (a) The antibody-reactive band of IBA-1, TNF-α, and YM-1/2. (b-d) Quantification of (b) IBA-1, (c) TNF-α, and (d) YM-1/2 protein expression levels. Values are means ± SEM, $n = 5$. # $P < 0.05$ and ## $P < 0.01$ vs. sham; * $P < 0.05$ and ** $P < 0.01$ vs. Aβ₂₅₋₃₅.

secondary antibodies for 90 min at room temperature. The blots were visualized with ECL detection reagents (Beyotime, China) and quantified by One-4.6.7 (Bio-Rad, USA).

2.8. Statistics. Statistical analysis was executed using Graph-Pad Prism 7.04. Values are given as means ± SEM. Repeated-measure analysis of variance (ANOVA) was used for analyzing the MWM data, while one-way ANOVA was performed for the other data; they were followed with Dunnett's test to compare differences among groups.

3. Results

3.1. CZYH Attenuates Aβ-Induced Cognitive Impairments. In the MWM test, animals displayed declined escape latency across the training period and the Aβ₂₅₋₃₅-induced rats have longer escape latency than those of the sham group in day 5 and day 6 (Figure 1(b); $P < 0.05$, $P < 0.01$, respectively), indicative of impaired spatial learning ability in the model group. However, CZYH treatment relieved the prolongation of escape latency triggered by Aβ₂₅₋₃₅ ($F(6, 85) = 3.316$, $P = 0.0056$). The percentage of time spent in the target quadrant was shorter in the Aβ₂₅₋₃₅-induced group

compared with the sham group in the spatial probe test ($F(6, 85) = 3.25$; $P = 0.0063$) ($P < 0.05$), while CZYH 400 mg/kg- or 800 mg/kg-treated rats spent more time in the target quadrant than vehicle-treated animals ($P < 0.05$, $P < 0.01$, respectively). Meanwhile, the swimming speeds of these groups did not exhibit differences (Figure 1(d)), suggesting there is no motor dysfunction between groups. Overall, these results indicated that CZYH alleviated spatial learning and memory deficits induced by Aβ₂₅₋₃₅.

3.2. CZYH Mitigates Aβ-Induced Neuronal Loss. The neuronal alterations were assessed by examining the numbers and morphology of neuronal cells in CA1 and DG regions of the hippocampus using Nissl staining. As depicted in Figure 2, highly dense pyramidal layer neurons with uniform and intact structure can be seen in sham groups, whereas lower density with atrophied and disordered neurons were found in the hippocampal CA1 and DG regions of Aβ-induced animals (CA1: $F(6, 28) = 22.78$, $P < 0.0001$; DG: $F(6, 28) = 14.12$, $P < 0.0001$) ($P < 0.01$, $P < 0.01$, respectively). However, the neurons in the CA1 and DG regions revert back to a neat and dense arrangement after

FIGURE 4: CZYH decreases the expression of inflammatory factors in the hippocampus of A β_{25-35} -induced rats. (a) The antibody-reactive band of COX-2 and IL-1 β protein expression. Quantification of (b) COX-2 and (c) IL-1 β protein expression levels. Values are means $\pm$ SEM, $n = 5$. * $P < 0.05$ and ** $P < 0.01$ vs. sham; ## $P < 0.01$ vs. A β_{25-35} .

CZYH 400 mg/kg or 800 mg/kg treatment. In general, these data suggested that CZYH treatment attenuated neuronal loss induced by A β .

3.3. CZYH Suppresses A β -Induced Microglial Activation and Subsequent Inflammatory Response. Since the inflammatory process is involved in A β toxicity, Western blotting was used to determine the activation of microglia and the expression of inflammatory factors. The expression of microglial activation marker IBA-1 was enhanced in the hippocampi of A β -treated rats compared to those of the sham group ($F(6, 28) = 4.702$, $P = 0.0020$) ($P < 0.01$, Figure 3), suggesting microglial activation and neuroinflammation. Meanwhile, the level of the proinflammatory microglial (M1) phenotype marker TNF- α was increased ($F(6, 28) = 6.462$, $P = 0.0002$) ($P < 0.01$), and the anti-inflammatory microglial (M2) marker YM-1/2 was decreased notably ($F(6, 28) = 4.225$, $P = 0.0038$) ($P < 0.05$). However, CZYH treatment reduced the expression of both IBA-1 and TNF- α and upregulated the levels of YM-1/2, pointing that treatment with CZYH represses microglial M1 activation and promotes microglia to polarize to the neuroprotective M2 phenotype. Additionally, the inflammatory factors including COX-2 and IL-1 β expression were increased in A β -

injected animals ($F(6, 28) = 19.85$, $P < 0.0001$; IL-1 β : $F(6, 28) = 5.735$, $P = 0.0005$) ($P < 0.01$, $P < 0.01$, respectively) and decreased after CZYH or donepezil treatment (Figure 4), further confirming CZYH could subside inflammatory response induced by A β .

3.4. CZYH Lessens A β -Induced I κ B- α Degradation and NF- κ B p65 Phosphorylation. As NF- κ B is a pivotal transcription factor in regulating inflammation, immune responses, and cell survival [23], the phosphorylation of NF- κ B p65 and expression of its major inhibitor protein I κ B- α were further scrutinized. As illustrated in Figure 5, the I κ B- α level was markedly declined after A β injection ($F(6, 28) = 4.338$, $P = 0.0032$) ($P < 0.01$) and was escalated following CZYH 400 mg/kg or 800 mg/kg treatment ($P < 0.01$, $P < 0.05$). Conversely, the phosphorylation level of NF- κ B p65 was increased in A β -induced rats compared to the level of the sham group ($F(6, 28) = 5.499$, $P = 0.0007$) ($P < 0.01$), and the elevated NF- κ B p65 phosphorylation was efficiently repressed by CZYH 800 mg/kg and donepezil administration.

3.5. CZYH Restrains A β -Induced JNK Phosphorylation. Mitogen-activated protein kinases (MAPKs) are a family of

FIGURE 5: CZYH represses IκB-α degradation and NF-κB p65 phosphorylation induced by Aβ₂₅₋₃₅ in the hippocampus of rats. (a) The antibody-reactive band of IκB-α and p-NF-κB p65 protein expression. (b, c) Statistics of (b) IκB-α and (c) p-NF-κB p65 levels. Values are means ± SEM, *n* = 5. ^{##}*P* < 0.01 vs. sham; ^{*}*P* < 0.05 and ^{**}*P* < 0.01 vs. Aβ₂₅₋₃₅.

kinases involved in mediating intracellular signaling including inflammation. In order to clarify the mechanisms underlying the protection effects of CZYH against Aβ-induced cognitive deficits, Western blot was utilized to examine the activation of MAPK family members: c-Jun N-terminal kinase (JNK), extracellular signal-related kinase (ERK), and p38 MAPK. All the three MAPK members' (JNK, ERK, and p38) phosphorylation levels were upregulated after Aβ stimulation (*P* < 0.01, *P* < 0.05, and *P* < 0.05, respectively), while the upraised phosphorylation level of JNK was reduced after CZYH or donepezil treatment (Figure 6). However, the elevated level of p-ERK can only be suppressed by donepezil instead of different doses of CZYH. Broadly speaking, CZYH treatment could attenuate the Aβ-induced phosphorylation of JNK, but not ERK and p38.

4. Discussion

Although AD is a multifactorial disease with a not clearly understood pathogenesis, multiple studies have shown that Aβ aggregation and deposition is the driving cause of AD. The intrahippocampal injection of a fragment of Aβ can mimic pathological and behavioral features of AD; Aβ-induced animal models have been widely used to evaluate pharmacological therapies for AD. Consistent with previous studies [11], we confirmed that a single injection of Aβ₂₅₋₃₅

into the rat's hippocampus leads to learning and memory impairments and neuronal damage in CA1 and DG subfields of the hippocampus. Importantly, the above deficits could be effectively prevented by CZYH 400 or 800 mg/kg treatment. It is worth noting that, even if 200 mg/kg CZYH administration was able to reduce the abnormally elevated escape latency for Aβ₂₅₋₃₅-induced rats in the MWM test, it has insignificant effect on improving spatial memory (Figure 1(c)) and neuron loss (Figure 2), suggesting that the effective dose of CZYH is above 200 mg/kg. Moreover, the effect of CZYH was not in a dose-dependent manner; it was speculated that the disparity between doses in the current study is too close to cause a significant difference in effect.

Neuroinflammation is linked to the mechanisms of Aβ-induced toxicity and AD pathogenesis. Excessive level of Aβ can activate microglial cells to release inflammatory cytokines, including COX-2, IL-1β, and TNF-α [24]. These elevated proinflammatory factors impede resident microglia to phagocytosis Aβ, which in turn aggravate the inflammatory response and AD development [25]. There are two phenotypes of activated microglia; one is the classic activated microglia (also known as the proinflammatory state, M1) which has been characterized by secreting inflammatory cytokines; another one is the alternative activated state (anti-inflammatory state, M2) and is regarded to be beneficial as it may delay the progression of the disease. The present

FIGURE 6: CZYH attenuates JNK phosphorylation induced by Aβ₂₅₋₃₅ in the hippocampus of rats. (a) Representative bands of p-JNK, JNK, p-ERK, ERK, p-p38, and p38. (b–d) Statistics of (b) JNK, (c) ERK, and (d) p38 phosphorylation. Values are means ± SEM, $n = 5$ for JNK and ERK and $n = 3$ for p38. ^{*} $P < 0.05$ and ^{##} $P < 0.01$ vs. sham; ^{*} $P < 0.05$ and ^{**} $P < 0.01$ vs. Aβ₂₅₋₃₅.

study revealed that the expression of M1 markers was significantly enhanced after Aβ₂₅₋₃₅ stimulation, whereas YM-1/2 expression (M2 marker) was decreased, consistent with the recent study which showed Aβ₁₋₄₂ can activate N9 microglia into the M1 state [26], and the current findings firstly reveal that Aβ₂₅₋₃₅ could induce hippocampus microglial M1 activation *in vivo*. In addition, elevated levels of COX-2, IL-1β, and TNF-α were also detected in Aβ₂₅₋₃₅-induced animals, implying that exogenous injection of Aβ₂₅₋₃₅ stimulates microglia to M1 polarization which then secrete proinflammatory factors, forming chronic neuroinflammation conditions in the brain, eventually resulting in neuronal loss and associated cognitive decline. Consistent with the studies about donepezil [6, 27], we also find that donepezil inhibited M1 activation and promotes M2 polarization in microglia and then suppressed the release of inflammatory cytokines. It is of note that similar anti-inflammatory effects have been seen in CZYH as evidenced by the reduced level of IBA-1 and the elevated level of YM-1/2, and decreased expression of

inflammatory factors was also observed in CZYH-treated groups.

NF-κB, a well-characterized transcription factor, has an important role in regulating the expressions of several proinflammatory cytokines. NF-κB was found activated in microglial cells in the brains of AD patients [28], and it has been demonstrated that NF-κB is involved in the neuroinflammatory response induced by Aβ [21, 29]. NF-κB is retained as inactive by binding with IκB proteins in physiological conditions. Upon stimulation, the IκB kinase is phosphorylated and degraded to empower NF-κB nuclear translocation, subsequently initiating the transcription of downstream genes, including inflammatory cytokines. The current study confirmed that Aβ₂₅₋₃₅ mediated the degradation of IκB-α and the subsequent NF-κB p65 phosphorylation, leading to the release of the microglia-derived proinflammatory factors. However, the activation of NF-κB p65/IκB-α could be ameliorated by CZYH treatment, which is consistent with the observations that *Epimedium* and a

prescription contain *Epimedium* modulate neuroinflammatory response via suppressing of the NF- κ B pathway [30–32].

MAPKs are a family of protein kinases that are specific to the amino acids serine and threonine. Increasing lines of evidence provided strong support that MAPK regulates the activities of several transcription factors, including NF- κ B [33, 34]. There are three major subgroups of MAPK: JNK, ERK, and p38, with distinct roles in regulating cell functions. ERK is involved in growth factor signaling, including cell proliferation, cell cycle progression, cell division, and cell differentiation [35], while the JNK and p38 signaling pathways are responsive to stress stimuli, such as cytokines, heat shock, and radiation [36, 37]. This study found that A β_{25-35} injection elicited the phosphorylation of ERK and p38, as well as JNK, in the hippocampus, adding more evidence to support and extend the notion that neuroinflammatory signals stimulated by A β in microglia via phosphorylation of MAPK initiate I κ B- α degradation following NF- κ B activation, which immediately mobilizes the transcription of inflammatory molecules. In addition, we observed that CZYH reduced the JNK kinase phosphorylation activity in the hippocampus of A β_{25-35} -induced rats but has no significant effect on the phosphorylation of p38 or ERK. However, donepezil is able to alleviate both the JNK phosphorylation and the p38 phosphorylation, which is coincident with a previous report [38]. These results suggest that CZYH mediated decrease in JNK phosphorylation and then inhibited the downstream activation of NF- κ B, accounting for the suppression of neuroinflammatory response in A β_{25-35} -induced rats.

In conclusion, this study demonstrated that CZYH attenuates cognitive impairments and neuron loss induced by A β_{25-35} , owing to its anti-inflammatory effects via the suppression of JNK phosphorylation and the blockade of the NF- κ B/I κ B- α pathway. Therefore, our findings suggest that Chinese herbal formula CZYH could be beneficial and effective for AD treatment.

Abbreviations

AD:	Alzheimer's disease
A β :	Beta-amyloid
ANOVA:	Analysis of variance
APP:	Beta-amyloid precursor protein
BACE1:	Beta-site APP-cleaving enzyme 1
BCA:	Bicinchoninic acid
COX-2:	Cyclooxygenase-2
CZYH:	Cu-Zhi-Yi-Hao
ERK:	Extracellular signal-related kinase
IDE:	Insulin-degrading enzyme
JNK:	c-Jun N-terminal kinase
MAPK:	Mitogen-activated protein kinase
MWM:	Morris water maze
NF- κ B:	Nuclear factor κ B
PVDF:	Polyvinylidene fluoride
SD:	Sprague Dawley
SEM:	Standard error of the mean
SPF:	Specific pathogen free
TCM:	Traditional Chinese medicine
TNF- α :	Tumor necrosis factor- α .

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

This study has no financial interests.

Acknowledgments

This study was supported by the Natural Science Foundation of China (Grant No. 81560585), the Jinshan Shi's Tutor Studio of Pharmacology (GZS-2016(07)), the Program for Changjiang Scholars and Innovative Research Team in University, China (Grant No. IRT17R113), and the Science and Technology Joint Fund of 2016 combined Zunyi Science and Technology Bureau with Zunyi Medical University (Grant No. [2016]29).

References

- [1] C. M. Roe, G. M. Babulal, D. M. Head et al., "Preclinical Alzheimer's disease and longitudinal driving decline," *Alzheimer's & Dementia: Translational Research & Clinical Interventions*, vol. 3, no. 1, pp. 74–82, 2017.
- [2] M. Wiesmann, V. Zerbi, D. Jansen et al., "Hypertension, cerebrovascular impairment, and cognitive decline in aged A β PP/PS1 mice," *Theranostics*, vol. 7, 5, pp. 1277–1289, 2017.
- [3] Alzheimer's Association, "2018 Alzheimer's disease facts and figures," *Alzheimer's & Dementia*, vol. 14, no. 3, pp. 367–429, 2018.
- [4] J. Cedernaes, H. B. Schiöth, and C. Benedict, "Efficacy of antibody-based therapies to treat Alzheimer's disease: just a matter of timing?," *Experimental Gerontology*, vol. 57, pp. 104–106, 2014.
- [5] B. Maloney and D. K. Lahiri, "The Alzheimer's amyloid β -peptide (A β) binds a specific DNA A β -interacting domain (A β ID) in the APP, BACE1, and APOE promoters in a sequence-specific manner: characterizing a new regulatory motif," *Gene*, vol. 488, no. 1-2, pp. 1–12, 2011.
- [6] J. S. Miners, Z. V. Helmond, P. G. Kehoe, and S. Love, "Changes with age in the activities of β -secretase and the A β -degrading enzymes neprilysin, insulin-degrading enzyme and angiotensin-converting enzyme," *Brain Pathology*, vol. 20, no. 4, pp. 794–802, 2010.
- [7] E. S. Musiek and D. M. Holtzman, "Three dimensions of the amyloid hypothesis: time, space and 'wingmen'," *Nature Neuroscience*, vol. 18, no. 6, pp. 800–806, 2015.
- [8] J. M. Rubio-Perez and J. M. Morillas-Ruiz, "A review: inflammatory process in Alzheimer's disease, role of cytokines," *The Scientific World Journal*, vol. 2012, Article ID 756357, 15 pages, 2012.
- [9] Y.-J. Lee, S. B. Han, S.-Y. Nam, K.-W. Oh, and J. T. Hong, "Inflammation and Alzheimer's disease," *Archives of Pharmacological Research*, vol. 33, no. 10, pp. 1539–1556, 2010.
- [10] P. Scheltens, K. Blennow, M. M. Breteler et al., "Alzheimer's disease," *Lancet*, vol. 388, pp. 505–517, 2016.
- [11] Y. Deng, L. Long, K. Wang et al., "Icariside II, a broad-spectrum anti-cancer agent, reverses beta-amyloid-induced cognitive impairment through reducing inflammation and

- apoptosis in rats," *Frontiers in Pharmacology*, vol. 8, p. 39, 2017.
- [12] L. Yan, Y. Deng, J. Gao et al., "Icariside II effectively reduces spatial learning and memory impairments in Alzheimer's disease model mice targeting beta-amyloid production," *Frontiers in Pharmacology*, vol. 8, p. 106, 2017.
 - [13] C. Yin, Y. Deng, J. Gao, X. Li, Y. Liu, and Q. Gong, "Icariside II, a novel phosphodiesterase-5 inhibitor, attenuates streptozotocin-induced cognitive deficits in rats," *Neuroscience*, vol. 328, pp. 69–79, 2016.
 - [14] N. Zong, F. Li, Y. Deng, J. Shi, F. Jin, and Q. Gong, "Icariin, a major constituent from *Epimedium brevicornum*, attenuates ibotenic acid-induced excitotoxicity in rat hippocampus," *Behavioural Brain Research*, vol. 313, pp. 111–119, 2016.
 - [15] K. Anukulthanakorn, I. S. Parhar, S. Jaroenporn, T. Kitahashi, G. Watanbe, and S. Malaivijitnond, "Neurotherapeutic effects of *Pueraria mirifica* extract in early-and late-stage cognitive impaired rats," *Phytotherapy Research*, vol. 30, no. 6, pp. 929–939, 2016.
 - [16] J. Li, G. Wang, J. Liu et al., "Puerarin attenuates amyloid-beta-induced cognitive impairment through suppression of apoptosis in rat hippocampus in vivo," *European Journal of Pharmacology*, vol. 649, no. 1–3, pp. 195–201, 2010.
 - [17] Y.-M. Seo, S. J. Choi, C. K. Park, M. C. Gim, and D.-H. Shin, "Synergistic effect of Korean red ginseng and *Pueraria montana* var. *lobata* against trimethyltin-induced cognitive impairment," *Food Science and Biotechnology*, vol. 27, no. 4, pp. 1193–1200, 2018.
 - [18] Y. Zhang, C. Lin, L. Zhang et al., "Cognitive improvement during treatment for mild Alzheimer's disease with a Chinese herbal formula: a randomized controlled trial," *PLoS One*, vol. 10, no. 6, p. e0130353, 2015.
 - [19] K.-K. Jia, Y.-J. Zheng, Y.-X. Zhang et al., "Banxia-houpu decoction restores glucose intolerance in CUMS rats through improvement of insulin signaling and suppression of NLRP3 inflammasome activation in liver and brain," *Journal of Ethnopharmacology*, vol. 209, pp. 219–229, 2017.
 - [20] L. Zeng, J. Gao, Y. Deng, J. Shi, and Q. Gong, "CZ2HF mitigates β -amyloid 25–35 fragment-induced learning and memory impairment through inhibition of neuroinflammation and apoptosis in rats," *International Journal of Molecular Medicine*, vol. 43, no. 1, pp. 557–566, 2019.
 - [21] H. Liu, Y. Deng, J. Gao, Y. Liu, J. Shi, and Q. Gong, "Sodium hydrosulfide attenuates beta-amyloid-induced cognitive deficits and neuroinflammation via modulation of MAPK/NF- κ B pathway in rats," *Current Alzheimer Research*, vol. 12, no. 7, pp. 673–683, 2015.
 - [22] C. Yin, Y. Deng, Y. Liu, J. Gao, L. Yan, and Q. Gong, "Icariside II ameliorates cognitive impairments induced by chronic cerebral hypoperfusion by inhibiting the amyloidogenic pathway: involvement of BDNF/TrkB/CREB signaling and up-regulation of PPAR α and PPAR γ in rats," *Frontiers in Pharmacology*, vol. 9, p. 1211, 2018.
 - [23] S. Vallabhapurapu and M. Karin, "Regulation and function of NF- κ B transcription factors in the immune system," *Annual Review of Immunology*, vol. 27, no. 1, pp. 693–733, 2009.
 - [24] M. Gold and J. El Khoury, " β -Amyloid, microglia, and the inflammasome in Alzheimer's disease," *Seminars in Immunopathology*, vol. 37, no. 6, pp. 607–611, 2015.
 - [25] H. Sarlus and M. T. Heneka, "Microglia in Alzheimer's disease," *Journal of Clinical Investigation*, vol. 127, p. 3240, 2017.
 - [26] H. Cai, Q. Liang, and G. Ge, "Gypenoside attenuates β amyloid-induced inflammation in N9 microglial cells via SOCS1 signaling," *Neural Plasticity*, vol. 2016, Article ID 6362707, 10 pages, 2016.
 - [27] T. Chen, R. Hou, S. Xu, and C. Wu, "Donepezil regulates 1-methyl-4-phenylpyridinium-induced microglial polarization in Parkinson's disease," *ACS Chemical Neuroscience*, vol. 6, no. 10, pp. 1708–1714, 2015.
 - [28] Z.-M. Shi, Y.-W. Han, X.-H. Han et al., "Upstream regulators and downstream effectors of NF- κ B in Alzheimer's disease," *Journal of the Neurological Sciences*, vol. 366, pp. 127–134, 2016.
 - [29] J. Lai, M. Hu, H. Wang et al., "Montelukast targeting the cysteinyl leukotriene receptor 1 ameliorates A β_{1-42} -induced memory impairment and neuroinflammatory and apoptotic responses in mice," *Neuropharmacology*, vol. 79, pp. 707–714, 2014.
 - [30] Z. Wenliang, L. Rongheng, W. Shumei, M. Fangzheng, and J. Ping, "Effect of Chinese traditional herb *Epimedium grandiflorum* C. Morren and its extract icariin on osteoarthritis via suppressing NF- κ B pathway," *Indian Journal of Experimental Biology (IJEB)*, vol. 54, no. 4, pp. 313–321, 2013.
 - [31] L.-L. Yin, L.-L. Lin, L. Zhang, and L. Li, "Epimedium flavonoids ameliorate experimental autoimmune encephalomyelitis in rats by modulating neuroinflammatory and neurotrophic responses," *Neuropharmacology*, vol. 63, no. 5, pp. 851–862, 2012.
 - [32] K.-W. Zeng, T. Zhang, H. Fu, G.-X. Liu, and X.-M. Wang, "Modified Wu-Zi-Yan-Zong prescription, a traditional Chinese polyherbal formula, suppresses lipopolysaccharide-induced neuroinflammatory processes in rat astrocytes via NF- κ B and JNK/p38 MAPK signaling pathways," *Phytomedicine*, vol. 19, no. 2, pp. 122–129, 2012.
 - [33] H. Lv, Z. Yu, Y. Zheng et al., "Isovitexin exerts anti-inflammatory and anti-oxidant activities on lipopolysaccharide-induced acute lung injury by inhibiting MAPK and NF- κ B and activating HO-1/Nrf2 pathways," *International Journal of Biological Sciences*, vol. 12, no. 1, pp. 72–86, 2016.
 - [34] J. Park, J.-S. Min, B. Kim et al., "Mitochondrial ROS govern the LPS-induced pro-inflammatory response in microglia cells by regulating MAPK and NF- κ B pathways," *Neuroscience Letters*, vol. 584, pp. 191–196, 2015.
 - [35] J.-C. Chambard, R. Lefloch, J. Pouyssegur, and P. Lenormand, "ERK implication in cell cycle regulation," *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, vol. 1773, no. 8, pp. 1299–1310, 2007.
 - [36] S. J. Harper and P. LoGrasso, "Signalling for survival and death in neurones: the role of stress-activated kinases, JNK and p38," *Cellular Signalling*, vol. 13, no. 5, pp. 299–310, 2001.
 - [37] E. K. Kim and E.-J. Choi, "Pathological roles of MAPK signaling pathways in human diseases," *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, vol. 1802, no. 4, pp. 396–405, 2010.
 - [38] L. He, Y. Deng, J. Gao, L. Zeng, and Q. Gong, "Icariside II ameliorates ibotenic acid-induced cognitive impairment and apoptotic response via modulation of MAPK pathway in rats," *Phytomedicine*, vol. 41, pp. 74–81, 2018.