

## Research Article

# (3, 2)-Fuzzy Sets and Their Applications to Topology and Optimal Choices

Hariwan Z. Ibrahim,<sup>1</sup> Tareq M. Al-shami ,<sup>2</sup> and O. G. Elbarbary<sup>3</sup>

<sup>1</sup>Department of Mathematics, Faculty of Education, University of Zakho, Zakho, Iraq

<sup>2</sup>Department of Mathematics, Sana'a University, Sana'a, Yemen

<sup>3</sup>Department of Mathematics, Faculty of Science, Tanta University, Tanta, Egypt

Correspondence should be addressed to Tareq M. Al-shami; tareqalshami83@gmail.com

Received 15 September 2021; Accepted 11 October 2021; Published 27 October 2021

Academic Editor: Ahmed Mostafa Khalil

Copyright © 2021 Hariwan Z. Ibrahim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of this paper is to define the concept of (3, 2)-fuzzy sets and discuss their relationship with other kinds of fuzzy sets. We describe some of the basic set operations on (3, 2)-fuzzy sets. (3, 2)-Fuzzy sets can deal with more uncertain situations than Pythagorean and intuitionistic fuzzy sets because of their larger range of describing the membership grades. Furthermore, we familiarize the notion of (3, 2)-fuzzy topological space and discuss the master properties of (3, 2)-fuzzy continuous maps. Then, we introduce the concept of (3, 2)-fuzzy points and study some types of separation axioms in (3, 2)-fuzzy topological space. Moreover, we establish the idea of relation in (3, 2)-fuzzy set and present some properties. Ultimately, on the basis of academic performance, the decision-making approach of student placement is presented via the proposed (3, 2)-fuzzy relation to ascertain the suitability of colleges to applicants.

## 1. Introduction

The concept of fuzzy sets was proposed by Zadeh [1]. The theory of fuzzy sets has several applications in real-life situations, and many scholars have researched fuzzy set theory. After the introduction of the concept of fuzzy sets, several research studies were conducted on the generalizations of fuzzy sets. The integration between fuzzy sets and some uncertainty approaches such as soft sets and rough sets has been discussed in [2–4].

The idea of intuitionistic fuzzy sets suggested by Atanassov [5] is one of the extensions of fuzzy sets with better applicability. Applications of intuitionistic fuzzy sets appear in various fields, including medical diagnosis, optimization problems, and multicriteria decision making [6–8]. Yager [9] offered a new fuzzy set called a Pythagorean fuzzy set, which is the generalization of intuitionistic fuzzy sets. Fermatean fuzzy sets were introduced by Senapati and Yager [10], and they also defined basic operations over the Fermatean fuzzy sets.

The concept of fuzzy topological spaces was introduced by Chang [11]. He studied the topological concepts like continuity and compactness via fuzzy topological spaces. Then, Lowen [12] presented a new type of fuzzy topological spaces. Çoker [13] subsequently initiated a study of intuitionistic fuzzy topological spaces. Recently, Olgun et al. [14] presented the concept of Pythagorean fuzzy topological spaces and Ibrahim [15] defined the concept of Fermatean fuzzy topological spaces.

The main purpose of this paper is to introduce the concept of (3, 2)-fuzzy sets and compare them with the other types of fuzzy sets. We introduce the set of operations for the (3, 2)-fuzzy sets and explore their main features. Following the idea of Chang, we define a topological structure via (3, 2)-fuzzy sets as an extension of fuzzy topological space, intuitionistic fuzzy topological space, and Pythagorean fuzzy topological space. We discuss the main topological concepts in (3, 2)-fuzzy topological spaces such as continuity and compactness. In addition, the concept of relation to (3, 2)-fuzzy sets is investigated. Finally, an improved version of

max-min-max composite relation for (3, 2)-fuzzy sets is proposed.

## 2. (3, 2)-Fuzzy Sets

In this section, we initiate the notion of (3, 2)-fuzzy sets and study their relationship with other kinds of fuzzy sets. Then, we furnish some operations to (3, 2)-fuzzy sets.

*Definition 1.* Let  $X$  be a universal set. Then, the (3, 2)-fuzzy set (briefly, (3, 2)-FS)  $D$  is defined by the following:

$$D = \{\langle r, \alpha_D(r), \beta_D(r) \rangle : r \in X\}, \quad (1)$$

where  $\alpha_D(r): X \rightarrow [0, 1]$  is the degree of membership and  $\beta_D(r): X \rightarrow [0, 1]$  is the degree of non-membership of  $r \in X$  to  $D$ , with the condition

$$0 \leq (\alpha_D(r))^3 + (\beta_D(r))^2 \leq 1. \quad (2)$$

The degree of indeterminacy of  $r \in X$  to  $D$  is defined by

$$\pi_D(r) = \sqrt[5]{1 - [(\alpha_D(r))^3 + (\beta_D(r))^2]}. \quad (3)$$

It is clear that  $(\alpha_D(r))^3 + (\beta_D(r))^2 + (\pi_D(r))^5 = 1$ , and  $\pi_D(r) = 0$  whenever  $(\alpha_D(r))^3 + (\beta_D(r))^2 = 1$ . In the interest of simplicity, we shall mention the symbol  $D = (\alpha_D, \beta_D)$  for the (3, 2)-FS  $D = \{\langle r, \alpha_D(r), \beta_D(r) \rangle : r \in X\}$ .

*Definition 2.* Let  $X$  be a universal set. Then, the intuitionistic fuzzy set (IFS) [5] (resp. Pythagorean fuzzy set (PFS) [9] and Fermatean fuzzy set (FFS) [10]) is defined by the following:

$$K = \{\langle r, \alpha_K(r), \beta_K(r) \rangle : r \in X\}, \quad (4)$$

with the condition  $0 \leq \alpha_K(r) + \beta_K(r) \leq 1$  (resp.  $0 \leq (\alpha_K(r))^2 + (\beta_K(r))^2 \leq 1$ ,  $0 \leq (\alpha_K(r))^3 + (\beta_K(r))^3 \leq 1$ ), where  $\alpha_K(r): X \rightarrow [0, 1]$  is the degree of membership and  $\beta_K(r): X \rightarrow [0, 1]$  is the degree of non-membership of every  $r \in X$  to  $K$ .

To illustrate the importance of (3, 2)-FS to extend the grades of membership and non-membership degrees, assume that  $\alpha_D(r) = 0.9$  and  $\beta_D(r) = 0.5$  for  $X = \{r\}$ . We obtain  $0.9 + 0.5 = 1.40 > 1$  and  $(0.9)^2 + (0.5)^2 = 1.06 > 1$  which means that  $D = (0.9, 0.5)$  neither follows the condition of IFS nor follows the condition of PFS. On the other hand,  $(0.9)^3 + (0.5)^2 = 0.979 < 1$  which means we can apply the (3, 2)-FS to control it. That is,  $D = (0.9, 0.5)$  is a (3, 2)-FS.

**Theorem 1.** *The set of (3, 2)-fuzzy membership grades is larger than the set of intuitionistic membership grades and Pythagorean membership grades.*

*Proof.* It is well known that for any two numbers  $r_1, r_2 \in [0, 1]$ , we have

$$\begin{aligned} r_1^3 &\leq r_1^2 \leq r_1, \\ r_2^2 &\leq r_2. \end{aligned} \quad (5)$$

Then, we get

$$\begin{aligned} r_1 + r_2 &\leq 1 \\ \Rightarrow r_1^2 + r_2^2 &\leq 1 \\ \Rightarrow r_1^3 + r_2^2 &\leq 1. \end{aligned} \quad (6)$$

Hence, the space of (3, 2)-fuzzy membership grades is larger than the space of intuitionistic membership grades and Pythagorean membership grades. This development can be evidently recognized in Figure 1.  $\square$

**Lemma 1.** *Let  $X = \{r_j: j = 1, \dots, k\}$  be a universal set and  $D$  be (3, 2)-FS. If  $\pi_D(r_j) = 0$ , then  $|\alpha_D(r_j)| = \sqrt[3]{|(\beta_D(r_j) - 1)(\beta_D(r_j) + 1)|}$ .*

*Proof.* Presume that  $D$  is (3, 2)-FS and  $\pi_D(r_j) = 0$  for  $r_j \in X$ ; then,

$$\begin{aligned} (\alpha_D(r_j))^3 + (\beta_D(r_j))^2 = 1 &\Rightarrow -(\alpha_D(r_j))^3 = (\beta_D(r_j))^2 - 1 \Rightarrow -(\alpha_D(r_j))^3 = (\beta_D(r_j) - 1)(\beta_D(r_j) + 1) \Rightarrow |(\alpha_D(r_j))^3| \\ &= |(\beta_D(r_j) - 1)(\beta_D(r_j) + 1)| \Rightarrow |\alpha_D(r_j)|^3 = |(\beta_D(r_j) - 1)(\beta_D(r_j) + 1)| \Rightarrow |\alpha_D(r_j)| = \sqrt[3]{|(\beta_D(r_j) - 1)(\beta_D(r_j) + 1)|}. \end{aligned} \quad (7)$$

*Example 1.* Let  $D$  be (3, 2)-FS and  $r \in X$  such that  $\beta_D(r) = 0.82$  and  $\pi_D(r) = 0$ . Then,  $|\alpha_D(r)| = \sqrt[3]{|(\beta_D(r) - 1)(\beta_D(r) + 1)|} = \sqrt[3]{|(-0.18)(1.82)|} = \sqrt[3]{0.3276}$ .

*Definition 3.* Let  $\delta$  be a positive real number ( $\delta > 0$ ). If  $D_1 = (\alpha_{D_1}, \beta_{D_1})$  and  $D_2 = (\alpha_{D_2}, \beta_{D_2})$  are two (3, 2)-FSs, then their operations are defined as follows:

- (1)  $D_1 \cap D_2 = (\min\{\alpha_{D_1}, \alpha_{D_2}\}, \max\{\beta_{D_1}, \beta_{D_2}\})$ .
- (2)  $D_1 \cup D_2 = (\max\{\alpha_{D_1}, \alpha_{D_2}\}, \min\{\beta_{D_1}, \beta_{D_2}\})$ .
- (3)  $D_1^c = (\beta_{D_1}, \alpha_{D_1})$ .

$$(4) \delta D_1 = (\sqrt[5]{1 - (1 - \alpha_{D_1}^3)^\delta}, \beta_{D_1}^\delta).$$

$$(5) D_1^\delta = (\alpha_{D_1}^\delta, \sqrt[5]{1 - (1 - \beta_{D_1}^2)^\delta}).$$

*Remark 1.* We will use supremum “sup” instead of maximum “max” and infimum “inf” instead of minimum “min” if the union and the intersection are infinite.

*Example 2.* Assume that  $D_1 = (\alpha_{D_1} = 0.9, \beta_{D_1} = 0.5)$  and  $D_2 = (\alpha_{D_2} = 0.89, \beta_{D_2} = 0.49)$  are both (3, 2)-FSs. Then,




FIGURE 1: Comparison of grade space of IFSs, PFSs, and (3, 2)-FSs.

- (1)  $D_1 \cap D_2 = (\min\{\alpha_{D_1}, \alpha_{D_2}\}, \max\{\beta_{D_1}, \beta_{D_2}\}) = (\min\{0.9, 0.89\}, \max\{0.5, 0.49\}) = (0.89, 0.5)$ .
- (2)  $D_1 \cup D_2 = (\max\{\alpha_{D_1}, \alpha_{D_2}\}, \min\{\beta_{D_1}, \beta_{D_2}\}) = (\max\{0.9, 0.89\}, \min\{0.5, 0.49\}) = (0.9, 0.49)$ .
- (3)  $D_1^c = (0.5, 0.9)$ .
- (4)  $\delta D_1 = (\sqrt[5]{1 - (1 - \alpha_{D_1}^3)^\delta}, \beta_{D_1}^\delta) = (\sqrt[5]{1 - (1 - 0.9^3)^\delta}, 0.5^4) \approx (0.99892, 0.06250)$ , for  $\delta = 4$ .
- (5)  $D_1^\delta = (\alpha_{D_1}^\delta, \sqrt[5]{1 - (1 - \beta_{D_1}^2)^\delta}) = (0.9^4, \sqrt[5]{1 - (1 - 0.5^2)^\delta}) \approx (0.65610, 0.92674)$ , for  $\delta = 4$ .

**Theorem 2.** Let  $L_1 = (\alpha_{L_1}, \beta_{L_1})$  and  $L_2 = (\alpha_{L_2}, \beta_{L_2})$  be two (3, 2)-FSs; then, the following properties hold:

$$\begin{aligned} L_1 \cap (L_2 \cap L_3) &= (\alpha_{L_1}, \beta_{L_1}) \cap (\min\{\alpha_{L_2}, \alpha_{L_3}\}, \max\{\beta_{L_2}, \beta_{L_3}\}) = (\min\{\alpha_{L_1}, \min\{\alpha_{L_2}, \alpha_{L_3}\}\}, \max\{\beta_{L_1}, \max\{\beta_{L_2}, \beta_{L_3}\}\}) \\ &= (\min\{\min\{\alpha_{L_1}, \alpha_{L_2}\}, \alpha_{L_3}\}, \max\{\max\{\beta_{L_1}, \beta_{L_2}\}, \beta_{L_3}\}) \\ &= (\min\{\alpha_{L_1}, \alpha_{L_2}\}, \max\{\beta_{L_1}, \beta_{L_2}\}) \cap (\alpha_{L_3}, \beta_{L_3}) = (L_1 \cap L_2) \cap L_3. \end{aligned} \quad (8)$$

(2) The proof is similar to (1).

- (1)  $L_1 \cap L_2 = L_2 \cap L_1$ .
- (2)  $L_1 \cup L_2 = L_2 \cup L_1$ .
- (3)  $(L_1 \cap L_2) \cup L_2 = L_2$ .
- (4)  $(L_1 \cup L_2) \cap L_2 = L_2$ .

*Proof.* From Definition 3, we can obtain

- (1)  $L_1 \cap L_2 = (\min\{\alpha_{L_1}, \alpha_{L_2}\}, \max\{\beta_{L_1}, \beta_{L_2}\}) = (\min\{\alpha_{L_2}, \alpha_{L_1}\}, \max\{\beta_{L_2}, \beta_{L_1}\}) = L_2 \cap L_1$ .
- (2) The proof is similar to (1).
- (3)  $(L_1 \cap L_2) \cup L_2 = (\min\{\alpha_{L_1}, \alpha_{L_2}\}, \max\{\beta_{L_1}, \beta_{L_2}\}) \cup (\alpha_{L_2}, \beta_{L_2}) = (\max\{\min\{\alpha_{L_1}, \alpha_{L_2}\}, \alpha_{L_2}\}, \min\{\max\{\beta_{L_1}, \beta_{L_2}\}, \beta_{L_2}\}) = (\alpha_{L_2}, \beta_{L_2}) = L_2$ .
- (4) The proof is similar to (3).  $\square$

**Theorem 3.** Let  $L_1 = (\alpha_{L_1}, \beta_{L_1})$ ,  $L_2 = (\alpha_{L_2}, \beta_{L_2})$  and  $L_3 = (\alpha_{L_3}, \beta_{L_3})$  be three (3, 2)-FSs and  $\delta > 0$ ; then,

- (1)  $L_1 \cap (L_2 \cap L_3) = (L_1 \cap L_2) \cap L_3$ .
- (2)  $L_1 \cup (L_2 \cup L_3) = (L_1 \cup L_2) \cup L_3$ .
- (3)  $\delta(L_1 \cup L_2) = \delta L_1 \cup \delta L_2$ .
- (4)  $(L_1 \cup L_2)^\delta = L_1^\delta \cup L_2^\delta$ .

*Proof.* For the three (3, 2)-FSs  $L_1, L_2$ , and  $L_3$  and  $\delta > 0$ , according to Definition 3, we can obtain

(1)

$$\begin{aligned} \delta(L_1 \cup L_2) &= \delta(\max\{\alpha_{L_1}, \alpha_{L_2}\}, \min\{\beta_{L_1}, \beta_{L_2}\}) = \left( \sqrt[5]{1 - (1 - \max\{\alpha_{L_1}^3, \alpha_{L_2}^3\})^\delta}, \min\{\beta_{L_1}^\delta, \beta_{L_2}^\delta\} \right), \\ \delta L_1 \cup \delta L_2 &= \left( \sqrt[5]{1 - (1 - \alpha_{L_1}^3)^\delta}, \beta_{L_1}^\delta \right) \cup \left( \sqrt[5]{1 - (1 - \alpha_{L_2}^3)^\delta}, \beta_{L_2}^\delta \right) \\ &= \left( \max\left\{ \sqrt[5]{1 - (1 - \alpha_{L_1}^3)^\delta}, \sqrt[5]{1 - (1 - \alpha_{L_2}^3)^\delta} \right\}, \min\{\beta_{L_1}^\delta, \beta_{L_2}^\delta\} \right) \\ &= \left( \sqrt[5]{1 - (1 - \max\{\alpha_{L_1}^3, \alpha_{L_2}^3\})^\delta}, \min\{\beta_{L_1}^\delta, \beta_{L_2}^\delta\} \right) = \delta(L_1 \cup L_2). \end{aligned} \quad (9)$$

(4) The proof is similar to (3).  $\square$

In the following result, we claim that  $L^c$  is (3, 2)-FS for any (3, 2)-FS  $L$ .

**Theorem 4.** Let  $L_1 = (\alpha_{L_1}, \beta_{L_1})$  and  $L_2 = (\alpha_{L_2}, \beta_{L_2})$  be two (3, 2)-FSs such that  $L_1^c$  and  $L_2^c$  are (3, 2)-FSs. Then,

$$(1) (L_1 \cap L_2)^c = L_1^c \cup L_2^c.$$

$$(2) (L_1 \cup L_2)^c = L_1^c \cap L_2^c.$$

*Proof.* For the two (3, 2)-FSs  $L_1$  and  $L_2$ , according to Definition 3, we can obtain

$$(1)$$

$$\begin{aligned} (L_1 \cap L_2)^c &= (\min\{\alpha_{L_1}, \alpha_{L_2}\}, \max\{\beta_{L_1}, \beta_{L_2}\})^c = (\max\{\beta_{L_1}, \beta_{L_2}\}, \\ &\min\{\alpha_{L_1}, \alpha_{L_2}\}) = (\beta_{L_1}, \alpha_{L_1}) \cup (\beta_{L_2}, \alpha_{L_2}) = L_1^c \cup L_2^c. \end{aligned} \quad (10)$$

(2) The proof is similar to (1).  $\square$

**Definition 4.** Let  $D_1 = (\alpha_{D_1}, \beta_{D_1})$  and  $D_2 = (\alpha_{D_2}, \beta_{D_2})$  be two (3, 2)-FSs; then,

- (1)  $D_1 = D_2$  if and only if  $\alpha_{D_1} = \alpha_{D_2}$  and  $\beta_{D_1} = \beta_{D_2}$ .
- (2)  $D_1 \geq D_2$  if and only if  $\alpha_{D_1} \geq \alpha_{D_2}$  and  $\beta_{D_1} \leq \beta_{D_2}$ .
- (3)  $D_2 \subset D_1$  or  $D_1^{D_2}$  if  $D_1 \geq D_2$ .

**Example 3.**

- (1) If  $D_1 = (0.9, 0.5)$  and  $D_2 = (0.9, 0.5)$  for  $X = \{x\}$ , then  $D_1 = D_2$ .
- (2) If  $D_1 = (0.9, 0.5)$  and  $D_2 = (0.81, 0.61)$  for  $X = \{x\}$ , then  $D_2 \leq D_1$  and  $D_2 \subset D_1$ .

### 3. Topology with respect to (3, 2)-Fuzzy Sets

In this section, we formulate the concept of (3, 2)-fuzzy topology on the family of (3, 2)-fuzzy sets whose complements are (3, 2)-fuzzy sets and scrutinize main properties. Then, we define (3, 2)-fuzzy continuous maps and give some

characterizations. Finally, we establish two types of (3, 2)-fuzzy separation axioms and reveal the relationships between them.

#### 3.1. (3, 2)-Fuzzy Topology

**Definition 5.** Let  $\tau$  be a family of (3, 2)-fuzzy subsets of a non-empty set  $X$ . If

- (1)  $1_X, 0_X \in \tau$  where  $1_X = (1, 0)$  and  $0_X = (0, 1)$ ,
- (2)  $D_1 \cap D_2 \in \tau$ , for any  $D_1, D_2 \in \tau$ ,
- (3)  $\cup_{i \in I} D_i \in \tau$ , for any  $\{D_i\}_{i \in I} \subset \tau$ ,

then  $\tau$  is called a (3, 2)-fuzzy topology on  $X$  and  $(X, \tau)$  is a (3, 2)-fuzzy topological space. We call  $D$  an open (3, 2)-FS if it is a member of  $\tau$  and call its complement a closed (3, 2)-FS.

**Remark 2.** We call  $\tau = \{1_X, 0_X\}$  the indiscrete (3, 2)-fuzzy topology on  $X$ . If  $\tau$  contains all (3, 2)-fuzzy subsets, then we call  $\tau$  the discrete (3, 2)-fuzzy topology on  $X$ .

**Example 4.** Let  $\tau = \{1_X, 0_X, D_1, D_2, D_3, D_4, D_5\}$  be the family of (3, 2)-fuzzy subsets of  $X = \{x_1, x_2\}$ , where

$$\begin{aligned} D_1 &= \{\langle x_1, \alpha_{D_1}(x_1) = 0.8, \beta_{D_1}(x_1) = 0.62 \rangle, \langle x_2, \alpha_{D_1}(x_2) = 0.81, \beta_{D_1}(x_2) = 0.61 \rangle\}, \\ D_2 &= \{\langle x_1, \alpha_{D_2}(x_1) = 0.83, \beta_{D_2}(x_1) = 0.53 \rangle, \langle x_2, \alpha_{D_2}(x_2) = 0.82, \beta_{D_2}(x_2) = 0.62 \rangle\}, \\ D_3 &= \{\langle x_1, \alpha_{D_3}(x_1) = 0.79, \beta_{D_3}(x_1) = 0.63 \rangle, \langle x_2, \alpha_{D_3}(x_2) = 0.8, \beta_{D_3}(x_2) = 0.63 \rangle\}, \\ D_4 &= \{\langle x_1, \alpha_{D_4}(x_1) = 0.83, \beta_{D_4}(x_1) = 0.53 \rangle, \langle x_2, \alpha_{D_4}(x_2) = 0.82, \beta_{D_4}(x_2) = 0.61 \rangle\}, \\ D_5 &= \{\langle x_1, \alpha_{D_5}(x_1) = 0.8, \beta_{D_5}(x_1) = 0.62 \rangle, \langle x_2, \alpha_{D_5}(x_2) = 0.81, \beta_{D_5}(x_2) = 0.62 \rangle\}. \end{aligned} \quad (11)$$

Hence,  $\tau$  is (3, 2)-fuzzy topology on  $X$ .

**Remark 3.** We showed that every fuzzy set  $D$  on a set  $X$  is a (3, 2)-fuzzy set having the form  $D = \{\langle r, \alpha_D(r), 1 - \alpha_D(r) \rangle : r \in X\}$ . Then, every fuzzy topological space  $(X, \tau_1)$  in the sense of Chang is obviously a (3, 2)-fuzzy

topological space in the form  $\tau = \{D : \alpha_D \in \tau_1\}$  whenever we identify a fuzzy set in  $X$  whose membership function is  $\alpha_D$  with its counterpart  $D = \{\langle r, \alpha_D(r), 1 - \alpha_D(r) \rangle : r \in X\}$ . Similarly, one can note that every intuitionistic fuzzy topology (Pythagorean fuzzy topology) is (3, 2)-fuzzy topology. The following examples explain this note.

*Example 5.* Consider  $\tau = \{1_X, 0_X, D_1, D_2\}$  as family of fuzzy subsets of  $X = \{x\}$ , where

$$\begin{aligned} 1_X &= \{\langle c, \alpha_{1_X}(x) = 1, 1 - \alpha_{1_X}(x) = \beta_{1_X}(x) = 0 \rangle\}, \\ 0_X &= \{\langle c, \alpha_{0_X}(x) = 0, 1 - \alpha_{0_X}(x) = \beta_{0_X}(x) = 1 \rangle\}, \\ D_1 &= \{\langle c, \alpha_{D_1}(x) = 0.7, 1 - \alpha_{D_1}(x) = \beta_{D_1}(x) = 0.3 \rangle\}, \\ D_2 &= \{\langle c, \alpha_{D_2}(x) = 0.2, 1 - \alpha_{D_2}(x) = \beta_{D_2}(x) = 0.8 \rangle\}. \end{aligned} \quad (12)$$

Then,  $\tau$  is fuzzy topology on  $X$ , and hence it is (3, 2)-fuzzy topology.

*Example 6.* Let  $\tau = \{1_X, 0_X, D_1, D_2\}$  be the family of (3, 2)-fuzzy subsets on  $X = \{x_1, x_2\}$  where

$$\begin{aligned} D_1 &= \{\langle x_1, \alpha_{D_1}(x_1) = 0.76, \beta_{D_1}(x_1) = 0.74 \rangle, \langle x_2, \alpha_{D_1}(x_2) = 0.6, \beta_{D_1}(x_2) = 0.83 \rangle\}, \\ D_2 &= \{\langle x_1, \alpha_{D_2}(x_1) = 0.75, \beta_{D_2}(x_1) = 0.74 \rangle, \langle x_2, \alpha_{D_2}(x_2) = 0.59, \beta_{D_2}(x_2) = 0.83 \rangle\}. \end{aligned} \quad (13)$$

Hence,  $\tau$  is (3, 2)-fuzzy topology. On the other hand,  $\tau$  is neither intuitionistic fuzzy topology nor Pythagorean fuzzy topology.

*Definition 6.* Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space and  $D = \{\langle x, \alpha_D(x), \beta_D(x) \rangle : x \in X\}$  be a (3, 2)-FS in  $X$ . Then, the (3, 2)-fuzzy interior and (3, 2)-fuzzy closure of  $D$  are, respectively, defined by

- (1)  $\text{cl}(D) = \cap \{H : H \text{ is a closed (3, 2)-FS in } X \text{ and } D \subset H\}$ .
- (2)  $\text{int}(D) = \cup \{G : G \text{ is an open (3, 2)-FS in } X \text{ and } G \subset D\}$ .

*Remark 4.* Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space and  $D$  be any (3, 2)-FS in  $X$ . Then,

- (1)  $\text{int}(D)$  is an open (3, 2)-FS.
- (2)  $\text{cl}(D)$  is a closed (3, 2)-FS.
- (3)  $\text{int}(1_X) = \text{cl}(1_X) = 1_X$  and  $\text{int}(0_X) = \text{cl}(0_X) = 0_X$ .

*Example 7.* Consider the (3, 2)-fuzzy topological space  $(X, \tau)$  in Example 4. If  $D = \{\langle c_1, 0.67, 0.81 \rangle, \langle c_2, 0.75, 0.74 \rangle\}$ , then  $\text{int}(D) = 0_X$  and  $\text{cl}(D) = 1_X$ .

**Theorem 5.** Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space and  $D_1, D_2$  be (3, 2)-FSs in  $X$ . Then, the following properties hold:

- (1)  $\text{int}(D_1) \subset D_1$  and  $D_1 \subset \text{cl}(D_1)$ .
- (2) If  $D_1 \subset D_2$ , then  $\text{int}(D_1) \subset \text{int}(D_2)$  and  $\text{cl}(D_1) \subset \text{cl}(D_2)$ .
- (3)  $D_1$  is an open (3, 2)-FS if and only if  $D_1 = \text{int}(D_1)$ .
- (4)  $D_1$  is a closed (3, 2)-FS if and only if  $D_1 = \text{cl}(D_1)$ .

*Proof.* (1) and (2) are obvious.

- (3) and (4) follow from Definition 6.  $\square$

**Corollary 1.** Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space and  $D_1, D_2$  be (3, 2)-FSs in  $X$ . Then, the following properties hold:

- (1)  $\text{int}(D_1) \cup \text{int}(D_2) \subset \text{int}(D_1 \cup D_2)$ .
- (2)  $\text{cl}(D_1 \cap D_2) \subset \text{cl}(D_1) \cap \text{cl}(D_2)$ .
- (3)  $\text{int}(D_1 \cap D_2) = \text{int}(D_1) \cap \text{int}(D_2)$ .
- (4)  $\text{cl}(D_1) \cup \text{cl}(D_2) = \text{cl}(D_1 \cup D_2)$ .

*Proof.* (1) and (2) follows from (1) of the above theorem. (3): since  $\text{int}(D_1 \cap D_2) \subset \text{int}(D_1)$  and  $\text{int}(D_1 \cap D_2) \subset \text{int}(D_2)$ , we obtain  $\text{int}(D_1 \cap D_2) \subset \text{int}(D_1) \cap \text{int}(D_2)$ . On the other hand, from the facts  $\text{int}(D_1) \subset D_1$  and  $\text{int}(D_2) \subset D_2$ , we have  $\text{int}(D_1) \cap \text{int}(D_2) \subset D_1 \cap D_2$  and  $\text{int}(D_1) \cap \text{int}(D_2) \in \tau$ ; we see that  $\text{int}(D_1) \cap \text{int}(D_2) \subset \text{int}(D_1 \cap D_2)$ , and hence  $\text{int}(D_1 \cap D_2) = \text{int}(D_1) \cap \text{int}(D_2)$ . (4) can be proved similar to (3).  $\square$

**Theorem 6.** Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space and  $D$  be (3, 2)-FS in  $X$ . Then, the following properties hold:

- (1)  $\text{cl}(D^c) = \text{int}(D)^c$ .
- (2)  $\text{int}(D^c) = \text{cl}(D)^c$ .
- (3)  $\text{cl}(D^c)^c = \text{int}(D)$ .
- (4)  $\text{int}(D^c)^c = \text{cl}(D)$ .

*Proof.* We only prove (1); the other parts can be proved similarly.

Let  $D = \{\langle x, \alpha_D(x), \beta_D(x) \rangle : x \in X\}$  and suppose that the family of open (3, 2)-fuzzy sets contained in  $D$  is indexed by the family  $\{\langle x, \alpha_{U_i}(x), \beta_{U_i}(x) \rangle : i \in J\}$ . Then,  $\text{int}(D) = \{\langle x, \vee \alpha_{U_i}(x), \wedge \beta_{U_i}(x) \rangle\}$ . Therefore,  $\text{int}(D)^c = \{\langle x, \wedge \beta_{U_i}(x), \vee \alpha_{U_i}(x) \rangle\}$ . Now,  $D^c = \{\langle x, \beta_D(x), \alpha_D(x) \rangle\}$  such that  $\alpha_{U_i} \leq \alpha_D$ ,  $\beta_{U_i} \geq \beta_D$  for each  $i \in J$ . This implies that  $\{\langle x, \beta_{U_i}(x), \alpha_{U_i}(x) \rangle : i \in J\}$  is the family of all closed (3, 2)-fuzzy sets containing  $D^c$ . That is,

$\text{cl}(D^c) = \{\langle x, \wedge \beta_{U_i}(x), \vee \alpha_{U_i}(x) \rangle\}$ . Hence,  $\text{cl}(D^c) = \text{int}(D)^c$ .  $\square$

### 3.2. (3, 2)-Fuzzy Continuous Maps

**Definition 7.** Let  $f: X \rightarrow Y$  be a map and  $A$  and  $B$  be (3, 2)-fuzzy subsets of  $X$  and  $Y$ , respectively. The functions of membership and non-membership of the image of  $A$ , denoted by  $f[A]$ , are, respectively, calculated by

$$\alpha_{f[A]}(y) := \begin{cases} \sup_{z \in f^{-1}(y)} \alpha_A(z), & \text{if } f^{-1}(y) \neq \phi, \\ 0, & \text{otherwise,} \end{cases} \quad (14)$$

$$\beta_{f[A]}(y) := \begin{cases} \inf_{z \in f^{-1}(y)} \beta_A(z), & \text{if } f^{-1}(y) \neq \phi, \\ 1, & \text{otherwise.} \end{cases}$$

The functions of membership and non-membership of preimage of  $B$ , denoted by  $f^{-1}[B]$ , are, respectively, calculated by

$$\alpha_{f^{-1}[B]}(x) := \alpha_B(f(x)),$$

$$\beta_{f^{-1}[B]}(x) := \beta_B(f(x)). \quad (15)$$

**Remark 5.** To show that  $f[A]$  and  $f^{-1}[B]$  are (3, 2)-fuzzy subsets, consider  $\gamma_A(z)^5 = \alpha_A(z)^3 + (\beta_A(z))^2$ . If  $f^{-1}(y)$  is non-empty, then we obtain

$$\begin{aligned} (\alpha_{f[A]}(y))^3 + (\beta_{f[A]}(y))^2 &= \left( \sup_{z \in f^{-1}(y)} \alpha_A(z) \right)^3 + \left( \inf_{z \in f^{-1}(y)} \beta_A(z) \right)^2 \\ &= \sup_{z \in f^{-1}(y)} (\alpha_A(z))^3 + \inf_{z \in f^{-1}(y)} (\beta_A(z))^2 \\ &= \sup_{z \in f^{-1}(y)} ((\gamma_A(z))^5 - (\beta_A(z))^2) + \inf_{z \in f^{-1}(y)} (\beta_A(z))^2 \\ &\leq \sup_{z \in f^{-1}(y)} (1 - (\beta_A(z))^2) + \inf_{z \in f^{-1}(y)} (\beta_A(z))^2 = 1. \end{aligned} \quad (16)$$

In contrast,  $f^{-1}(y) = \phi$  leads to the fact that  $(\alpha_{f[A]}(y))^3 + (\beta_{f[A]}(y))^2 = 1$ .

It is easy to prove the case of  $f^{-1}[B]$ .

**Theorem 7.** Let  $f: X \rightarrow Y$  be a map s.t.  $A$  and  $B$  are (3, 2)-fuzzy subsets of  $X$  and  $Y$ , respectively. Then, we have

- (1)  $f^{-1}[B^c] = f^{-1}[B]^c$ .
- (2)  $f[A]^c \subseteq f[A^c]$ .
- (3) If  $B_1 \subseteq B_2$ , then  $f^{-1}[B_1] \subseteq f^{-1}[B_2]$  where  $B_1$  and  $B_2$  are (3, 2)-fuzzy subsets of  $Y$ .
- (4) If  $A_1 \subseteq A_2$ , then  $f[A_1] \subseteq f[A_2]$  where  $A_1$  and  $A_2$  are (3, 2)-fuzzy subsets of  $X$ .
- (5)  $f[f^{-1}[B]] \subseteq B$ .
- (6)  $A \subseteq f^{-1}[f[A]]$ .

*Proof*

- (1) Consider  $v \in X$  and let  $B$  be a (3, 2)-fuzzy subset of  $Y$ . Then,

$$\alpha_{f^{-1}[B^c]}(v) = \alpha_{B^c}(f(v)) = \beta_B(f(v)) = \beta_{f^{-1}[B]}(v) = \alpha_{f^{-1}[B]^c}(v). \quad (17)$$

Similarly, one can have  $\beta_{f^{-1}[B^c]}(v) = \beta_{f^{-1}[B]^c}(v)$ . Therefore,  $f^{-1}[B^c] = f^{-1}[B]^c$ , as required.

- (2) For any  $w \in Y$  such that  $f^{-1}(w) \neq \phi$  and for any (3, 2)-fuzzy subset  $A$  of  $X$ , we can write

$$\begin{aligned} (\gamma_{f[A]}(w))^5 &= (\alpha_{f[A]}(w))^3 + (\beta_{f[A]}(w))^2 \\ &= \sup_{z \in f^{-1}(w)} (\alpha_A(z))^3 + \inf_{z \in f^{-1}(w)} (\beta_A(z))^2 \\ &= \sup_{z \in f^{-1}(w)} ((\gamma_A(z))^5 - (\beta_A(z))^2) + \inf_{z \in f^{-1}(w)} (\beta_A(z))^2 \\ &\leq \sup_{z \in f^{-1}(w)} ((\gamma_A(z))^5) - \inf_{z \in f^{-1}(w)} (\beta_A(z))^2 \\ &\quad + \inf_{z \in f^{-1}(w)} (\beta_A(z))^2 \\ &= \sup_{z \in f^{-1}(w)} ((\gamma_A(z))^5). \end{aligned} \quad (18)$$

Now from (18), we have

$$\begin{aligned}
\alpha_{f[A^c]}(w) &= \sup_{z \in f^{-1}(w)} \alpha_{A^c}(z) \\
&= \sup_{z \in f^{-1}(w)} \beta_A(z) \\
&= \sup_{z \in f^{-1}(w)} \sqrt{(\gamma_A(z))^5 - (\alpha_A(z))^3} \\
&\geq \sqrt{\sup_{z \in f^{-1}(w)} (\gamma_A(z))^5 - \sup_{z \in f^{-1}(w)} (\alpha_A(z))^3} \\
&\geq \sqrt{(\gamma_{f[A]}(w))^5 - (\alpha_{f[A]}(w))^3} \\
&= \beta_{f[A]}(w) \\
&= \alpha_{f[A]^c}(w).
\end{aligned} \tag{19}$$

The proof is easy when  $f^{-1}(w) = \phi$ . Following a similar technique, we obtain  $\beta_{f[A^c]}(w) \leq \beta_{f[A]}(w)$ , which means that  $f[A]^c \subseteq f[A^c]$ .

(3) Assume that  $B_1 \subseteq B_2$ . Then, for each  $v \in X$ ,  $\alpha_{f^{-1}[B_1]}(v) = \alpha_{B_1}(f(v)) \leq \alpha_{B_2}(f(v)) = \alpha_{f^{-1}[B_2]}(v)$ . Also,  $\beta_{f^{-1}[B_1]}(v) \geq \beta_{f^{-1}[B_2]}(v)$ . Hence, we obtain the desired result.

(4) Assume that  $A_1 \subseteq A_2$  and  $w \in Y$ . The proof is easy when  $f(w) = \phi$ . So, presume that  $f(w) \neq \phi$ . Then,

$$\alpha_{f[A_1]}(w) = \sup_{z \in f^{-1}(w)} \alpha_{A_1}(z) \leq \sup_{z \in f^{-1}(w)} \alpha_{A_2}(z) = \alpha_{f[A_2]}(w). \tag{20}$$

Thus,  $\alpha_{f[A_1]} \leq \alpha_{f[A_2]}$  follows. Similarly, we have  $\beta_{f[A_1]} \geq \beta_{f[A_2]}$ .

(5) For any  $w \in Y$  s.t.  $f(w) \neq \phi$ , we find that

$$\begin{aligned}
\alpha_{f[f^{-1}[B]]}(w) &= \sup_{z \in f^{-1}(w)} \alpha_{f^{-1}[B]}(z) = \sup_{z \in f^{-1}(w)} \alpha_B(f(z)) \\
&\leq \alpha_B(w).
\end{aligned} \tag{21}$$

On the other hand, we have  $\alpha_{f[f^{-1}[B]]}(w) = 0 \leq \alpha_B(w)$  when  $f(w) = \phi$ . Similarly, we have  $\beta_{f[f^{-1}[B]]}(w) = 0 \geq \beta_B(w)$ .

(6) For any  $v \in X$ , we have

$$\alpha_{f^{-1}[f[A]]}(v) = \alpha_{f[A]}(f(v)) = \sup_{z \in f^{-1}(f(v))} \alpha_A(z) \geq \alpha_A(v). \tag{22}$$

Similarly, we have  $\beta_{f^{-1}[f[A]]} \leq \beta_A$ .  $\square$

The proof of the following result is easy, and hence it is omitted.

**Theorem 8.** *Let  $X$  and  $Y$  be two non-empty sets and  $f: X \rightarrow Y$  be a map. Then, the following statements are true:*

- (1)  $f[\cup_{i \in I} A_i] = \cup_{i \in I} f[A_i]$  for any (3, 2)-fuzzy subset  $A_i$  of  $X$ .
- (2)  $f^{-1}[\cup_{i \in I} B_i] = \cup_{i \in I} f^{-1}[B_i]$  for any (3, 2)-fuzzy subset  $B_i$  of  $Y$ .
- (3)  $f[A_1 \cap A_2] \subseteq f[A_1] \cap f[A_2]$  for any two (3, 2)-fuzzy subsets  $A_1$  and  $A_2$  of  $X$ .
- (4)  $f^{-1}[\cap_{i \in I} B_i] = \cap_{i \in I} f^{-1}(B_i)$  for any (3, 2)-fuzzy subset  $B_i$  of  $Y$ .

**Definition 8.** In a (3, 2)-fuzzy topological space, consider that  $A$  and  $U$  are two (3, 2)-fuzzy subsets. We call  $U$  a neighborhood of  $A$ , briefly nbd, if there exists an open (3, 2)-fuzzy subset  $E$  such that  $A \subseteq E \subseteq U$ .

**Theorem 9.** *A (3, 2)-fuzzy subset  $A$  is open iff it contains a nbd of its each subset.*

*Proof.* The proof is easy.  $\square$

**Definition 9.** A map  $f: (X, \tau_1) \rightarrow (Y, \tau_2)$  is said to be (3, 2)-fuzzy continuous if for any (3, 2)-fuzzy subset  $A$  of  $X$  and for any nbd  $V$  of  $f[A]$  there is a nbd  $U$  of  $A$  s.t.  $f[U] \subseteq V$ .

**Theorem 10.** *The following statements are equivalent for a map  $f: (X, \tau_1) \rightarrow (Y, \tau_2)$ :*

- (1)  $f$  is (3, 2)-fuzzy continuous.
- (2) For each (3, 2)-FS  $A$  of  $X$  and each nbd  $V$  of  $f[A]$ , there is a nbd  $U$  of  $A$  s.t. for each  $B \subseteq U$ , we obtain  $f[B] \subseteq V$ .
- (3) For each (3, 2)-FS  $A$  of  $X$  and for each nbd  $V$  of  $f[A]$ , there is a nbd  $U$  of  $A$  s.t.  $U \subseteq f^{-1}[V]$ .
- (4) For each (3, 2)-FS  $A$  of  $X$  and for each nbd  $V$  of  $f[A]$ ,  $f^{-1}[V]$  is a nbd of  $A$ .

*Proof.*

(1)  $\Rightarrow$  (2): let  $f$  be a (3, 2)-fuzzy continuous map. Consider  $A$  as a (3, 2)-FS of  $X$  and  $V$  as a nbd of  $f[A]$ . Then, there is a nbd  $U$  of  $A$  s.t.  $f[U] \subseteq V$ . If  $B \subseteq U$ , we obtain  $f[B] \subseteq f[U] \subseteq V$ .

(2)  $\Rightarrow$  (3): assume  $A$  as a (3, 2)-FS of  $X$  and  $V$  as a nbd of  $f[A]$ . According to (2), there is a nbd  $U$  of  $A$  s.t. for each  $B \subseteq U$ , we find  $f[B] \subseteq V$ . Therefore,  $B \subseteq f^{-1}[f[B]] \subseteq f^{-1}[V]$ . Since  $B$  is chosen arbitrarily,  $U \subseteq f^{-1}[V]$ .

(3)  $\Rightarrow$  (4): presume  $A$  is a (3, 2)-FS of  $X$  and  $V$  is a nbd of  $f[A]$ . According to (3), there is a nbd  $U$  of  $A$  s.t.  $U \subseteq f^{-1}[V]$ . Since  $U$  is a nbd of  $A$ , there is an open (3, 2)-FS  $K$  of  $X$  s.t.  $A \subseteq K \subseteq U$ . On the other hand, we obtain  $A \subseteq K \subseteq f^{-1}[V]$  because  $U \subseteq f^{-1}[V]$ . This means that  $f^{-1}[V]$  is a nbd of  $A$ .

(4)  $\Rightarrow$  (1): suppose that  $A$  is a (3, 2)-FS of  $X$  and  $V$  is a nbd of  $f[A]$ . By hypothesis,  $f^{-1}[V]$  is a nbd of  $A$ . So, there is an open (3, 2)-FS  $K$  of  $X$  s.t.  $A \subseteq K \subseteq f^{-1}[V]$  which means  $f[K] \subseteq f[f^{-1}[V]] \subseteq V$ . Moreover,  $K$  is an

open (3, 2)-FS, so it is a nbd of  $A$ . Hence, we obtain the proof that  $f$  is (3, 2)-fuzzy continuous.  $\square$

**Theorem 11.** *A map  $f: (X, \tau_1) \longrightarrow (Y, \tau_2)$  is (3, 2)-fuzzy continuous iff  $f^{-1}[B]$  is an open (3, 2)-FS of  $X$  for each open (3, 2)-FS  $B$  of  $Y$ .*

*Proof.* Necessity: presume  $f$  as a (3, 2)-fuzzy continuous map. Consider an open (3, 2)-FS  $B$  of  $Y$  s.t.  $A \subseteq f^{-1}[B]$ . This directly gives that  $f[A] \subseteq B$ . It follows from Theorem 9 that there is a nbd  $V$  of  $f[A]$  satisfying  $V \subseteq B$ . Now,  $f$  is (3, 2)-fuzzy continuous, so by (4) of Theorem 10, we obtain that  $f^{-1}[V]$  is a nbd of  $A$ . Also, it follows from (3) of Theorem 7 that  $f^{-1}[V] \subseteq f^{-1}[B]$ . So,  $f^{-1}[B]$  is a nbd of  $A$ . Since  $A$  is an arbitrary subset of  $f^{-1}[B]$ , then by Theorem 9, the (3, 2)-FS  $f^{-1}[B]$  is open.  $\square$

**3.2.1. Sufficiency.** Presume  $A$  is a (3, 2)-FS of  $X$  and  $V$  is a nbd of  $f[A]$ . Then,  $\tau_2$  contains a (3, 2)-FS  $L$  of s.t.  $f[A] \subseteq L \subseteq V$ . By hypothesis,  $f^{-1}[L]$  is an open (3, 2)-FS. Also, we have  $A \subseteq f^{-1}[f[A]] \subseteq f^{-1}[L] \subseteq f^{-1}[V]$ . Thus,  $f^{-1}[V]$  is a nbd of  $A$  which demonstrates that  $f$  is (3, 2)-fuzzy continuous.

We build the following two examples such that the first one provides a (3, 2)-fuzzy continuous map, whereas the second one presents a fuzzy map that is not (3, 2)-fuzzy continuous.

**Example 8.** Consider  $X = \{a_1, a_2\}$  with the (3, 2)-fuzzy topology  $\tau_1 = \{1_X, 0_X, A_1\}$  and  $Y = \{b_1, b_2\}$  with the (3, 2)-fuzzy topology  $\tau_2 = \{1_Y, 0_Y, B_1\}$ , where

$$\begin{aligned} A_1 &= \{\langle a_1, 0.7, 0.78 \rangle, \langle a_2, 0.9, 0.5 \rangle\}, \\ B_1 &= \{\langle b_1, 0.9, 0.5 \rangle, \langle b_2, 0.7, 0.78 \rangle\}. \end{aligned} \quad (23)$$

Let  $f: X \longrightarrow Y$  be defined as follows:

$$f(x) = \begin{cases} b_2, & \text{if } x = a_1, \\ b_1, & \text{if } x = a_2. \end{cases} \quad (24)$$

Since  $1_Y, 0_Y$ , and  $B_1$  are open (3, 2)-fuzzy subsets of  $Y$ , then

$$\begin{aligned} f^{-1}[1_Y] &= \{\langle a_1, 1, 0 \rangle, \langle a_2, 1, 0 \rangle\}, \\ f^{-1}[0_Y] &= \{\langle a_1, 0, 1 \rangle, \langle a_2, 0, 1 \rangle\}, \\ f^{-1}[B_1] &= \{\langle a_1, 0.7, 0.78 \rangle, \langle a_2, 0.9, 0.5 \rangle\} \end{aligned} \quad (25)$$

are open (3, 2)-fuzzy subsets of  $X$ . Thus,  $f$  is (3, 2)-fuzzy continuous.

**Example 9.** Consider  $X = \{a_1, a_2\}$  with the (3, 2)-fuzzy topology  $\tau_1 = \{1_X, 0_X\}$  and  $Y = \{b_1, b_2\}$  with the (3, 2)-fuzzy topology  $\tau_2 = \{1_Y, 0_Y, B_1\}$ , where  $B_1 = \{\langle b_1, 0.82, 0.62 \rangle, \langle b_2, 0.52, 0.90 \rangle\}$ .

Let  $f: X \longrightarrow Y$  be defined as follows:

$$f(x) = \begin{cases} b_1, & \text{if } x = a_1, \\ b_2, & \text{if } x = a_2. \end{cases} \quad (26)$$

Since  $B_1$  is an open (3, 2)-fuzzy subset of  $Y$ , but  $f^{-1}[B_1] = \{\langle a_1, 0.82, 0.62 \rangle, \langle a_2, 0.52, 0.90 \rangle\}$  is not an open (3, 2)-fuzzy subset of  $X$ ,  $f$  is not (3, 2)-fuzzy continuous.

**Theorem 12.** *The following are equivalent to each other:*

- (1)  $f: (X, \tau_1) \longrightarrow (Y, \tau_2)$  is (3, 2)-fuzzy continuous.
- (2) For each closed (3, 2)-fuzzy subset  $B$  of  $Y$  we have that  $f^{-1}[B]$  is a closed (3, 2)-fuzzy subset of  $X$ .
- (3)  $cl(f^{-1}[B]) \subseteq f^{-1}[cl(B)]$  for each (3, 2)-fuzzy set in  $Y$ .
- (4)  $f^{-1}[int(B)] \subseteq int(f^{-1}[B])$  for each (3, 2)-fuzzy set in  $Y$ .

*Proof.* They can be easily proved using Theorems 6, 7, and 11.  $\square$

**Theorem 13.** *Let  $(Y, \tau)$  be a (3, 2)-fuzzy topological space and  $f: X \longrightarrow Y$  be a map. Then, there is a coarsest (3, 2)-fuzzy topology  $\tau_1$  over  $X$  such that  $f$  is (3, 2)-fuzzy continuous.*

*Proof.* Let us define a class of (3, 2)-fuzzy subsets  $\tau_1$  of  $X$  by

$$\tau_1 := \{f^{-1}[V] : V \in \tau\}. \quad (27)$$

We prove that  $\tau_1$  is the coarsest (3, 2)-fuzzy topology over  $X$  such that  $f$  is (3, 2)-fuzzy continuous.

- (1) We can write for any  $x \in X$  that

$$\alpha_{f^{-1}[0_Y]}(x) = \alpha_{0_Y}(f(x)) = 0 = \alpha_{0_X}(x). \quad (28)$$

Similarly, we immediately have  $\beta_{f^{-1}[0_Y]}(x) = \beta_{0_X}(x)$  for any  $x \in X$  which implies  $f^{-1}[0_Y] = 0_X$ . Now, as  $0_Y \in \tau$ , we have  $0_X = f^{-1}[0_Y] \in \tau_1$ . In a similar manner, it is easy to see that  $1_X = f^{-1}[1_Y] \in \tau_1$ .

- (2) Assume that  $D_1, D_2 \in \tau_1$ . Then, for  $i = 1, 2$ , there exists  $B_i \in \tau$  such that  $f^{-1}[B_i] = D_i$  which implies  $\alpha_{f^{-1}[B_i]} = \alpha_{D_i}$  and  $\beta_{f^{-1}[B_i]} = \beta_{D_i}$ . Thus, we obtain for any  $x \in X$  that

$$\begin{aligned} \alpha_{D_1 \cap D_2}(x) &= \min\{\{\alpha_{D_1}(x), \alpha_{D_2}(x)\}\} = \min\{\alpha_{f^{-1}[B_1]}(x), \alpha_{f^{-1}[B_2]}(x)\} = \min\{\alpha_{B_1}(f(x)), \alpha_{B_2}(f(x))\} \\ &= \alpha_{B_1 \cap B_2}(f(x)) = \alpha_{f^{-1}[B_1 \cap B_2]}(x). \end{aligned} \quad (29)$$



Similarly, it is not difficult to see that  $\beta_{D_1 \cap D_2} = \beta_{f^{-1}[B_1 \cap B_2]}$ . Hence, we get  $D_1 \cap D_2 \in \tau_1$ .

- (3) Assume that  $\{D_i\}_{i \in I}$  is an arbitrary subfamily of  $\tau_1$ . Then, for any  $i \in I$ , there exists  $B_i \in \tau_1$  such that  $f^{-1}[B_i] = D_i$  which implies  $\alpha_{f^{-1}[B_i]} = \alpha_{D_i}$  and  $\beta_{f^{-1}[B_i]} = \beta_{D_i}$ . Therefore, one can get for any  $x \in X$  that

$$\begin{aligned} \alpha_{\cup_{i \in I} D_i}(x) &= \sup_{i \in I} \alpha_{D_i}(x) \\ &= \sup_{i \in I} \alpha_{f^{-1}[B_i]}(x) \\ &= \sup_{i \in I} (f(x)) \\ &= \alpha_{\cup_{i \in I} B_i}(f(x)) \\ &= \alpha_{f^{-1}[\cup_{i \in I} B_i]}(x). \end{aligned} \quad (30)$$

On the other hand, it is easy to see that  $\beta_{\cup_{i \in I} D_i} = \beta_{f^{-1}[\cup_{i \in I} B_i]}$ . Thus, we have  $\cup_{i \in I} D_i \in \tau_1$ .

From Theorem 11, the (3, 2)-fuzzy continuity of  $f$  is trivial. Now, we prove that  $\tau_1$  is the coarsest (3, 2)-fuzzy topology over  $X$  such that  $f$  is (3, 2)-fuzzy continuous. Let  $\tau_2 \subseteq \tau_1$  be a (3, 2)-fuzzy topology over  $X$  such that  $f$  is (3, 2)-fuzzy continuous. If  $B \in \tau_1$ , then there is  $V \in \tau_2$  such that  $f^{-1}[V] = B$ . Since  $f$  is (3, 2)-fuzzy continuous with respect to  $\tau_2$ , we have  $B = f^{-1}[V] \in \tau_2$ . Hence,  $\tau_2 = \tau_1$ , as required.  $\square$

**3.3. (3, 2)-Fuzzy Separation Axioms.** Separation axioms are one of the most important and popular notions in topological studies. They have been studied and applied to model some real-life issues in soft setting as explained in [16, 17].

**Definition 10.** Let  $X \neq \emptyset$  and  $x \in X$  be a fixed element in  $X$ . Suppose that  $r_1 \in (0, 1]$  and  $r_2 \in [0, 1)$  are two fixed real numbers such that  $r_1^3 + r_2^2 \leq 1$ . Then, a (3, 2)-fuzzy point  $p_{(r_1, r_2)}^x = \{\langle x, \alpha_p(x), \beta_p(x) \rangle\}$  is defined to be a (3, 2)-fuzzy set of  $X$  as follows.

$$p_{(r_1, r_2)}^x(y) := \begin{cases} (r_1, r_2), & \text{if } y = x, \\ (0, 1), & \text{otherwise,} \end{cases} \quad (31)$$

for  $y \in X$ . In this case,  $x$  is called the support of  $p_{(r_1, r_2)}^x$ . A (3, 2)-fuzzy point  $p_{(r_1, r_2)}^x$  is said to belong to a (3, 2)-fuzzy set  $D = \{\langle x, \alpha_D(x), \beta_D(x) \rangle\}$  of  $X$  denoted by  $p_{(r_1, r_2)}^x \in D$  if  $r_1 \leq \alpha_D(x)$  and  $r_2 \geq \beta_D(x)$ . Two (3, 2)-fuzzy points are said to be distinct if their supports are distinct.

**Remark 6.** Let  $D_1 = \{\langle x, \alpha_{D_1}(x), \beta_{D_1}(x) \rangle\}$  and  $D_2 = \{\langle x, \alpha_{D_2}(x), \beta_{D_2}(x) \rangle\}$  be two (3, 2)-fuzzy sets of  $X$ . Then,  $D_1 \subseteq D_2$  if and only if  $p_{(r_1, r_2)}^x \in D_1$  implies  $p_{(r_1, r_2)}^x \in D_2$  for any (3, 2)-fuzzy point  $p_{(r_1, r_2)}^x$  in  $X$ .

**Definition 11.** Let  $r_1, r_3 \in (0, 1]$ ,  $r_2, r_4 \in [0, 1)$ , and  $x, y \in X$ . A (3, 2)-fuzzy topological space  $(X, \tau)$  is said to be

- (1)  $T_0$  if for each pair of distinct (3, 2)-fuzzy points  $p_{(r_1, r_2)}^x, p_{(r_3, r_4)}^y$  in  $X$ , there exist two open (3, 2)-fuzzy sets  $L$  and  $K$  such that

$$\begin{aligned} L &= \{\langle x, 1, 0 \rangle, \langle y, 0, 1 \rangle\}, \\ \text{or } K &= \{\langle x, 0, 1 \rangle, \langle y, 1, 0 \rangle\}. \end{aligned} \quad (32)$$

- (2)  $T_1$  if for each pair of distinct (3, 2)-fuzzy points  $p_{(r_1, r_2)}^x, p_{(r_3, r_4)}^y$  in  $X$ , there exist two open (3, 2)-fuzzy sets  $L$  and  $K$  such that

$$\begin{aligned} L &= \{\langle x, 1, 0 \rangle, \langle y, 0, 1 \rangle\}, \\ K &= \{\langle x, 0, 1 \rangle, \langle y, 1, 0 \rangle\}. \end{aligned} \quad (33)$$

**Proposition 1.** Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space. If  $(X, \tau)$  is  $T_1$ , then  $(X, \tau)$  is  $T_0$ .

*Proof.* The proof is straightforward from Definition 11.  $\square$

Here is an example which shows that the converse of above proposition is not true in general.

**Example 10.** Consider  $X = \{c_1, c_2\}$  with the (3, 2)-fuzzy topology  $\tau = \{1_X, 0_X, D\}$ , where  $D = \{\langle c_1, 1, 0 \rangle, \langle c_2, 0, 1 \rangle\}$ . Then,  $(X, \tau)$  is  $T_0$  but not  $T_1$  because there does not exist an open (3, 2)-fuzzy set  $K$  such that  $K = \{\langle x, 0, 1 \rangle, \langle y, 1, 0 \rangle\}$ .

**Theorem 14.** Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space,  $r_1, r_3 \in (0, 1]$ , and  $r_2, r_4 \in [0, 1)$ . If  $(X, \tau)$  is  $T_0$ , then for each pair of distinct (3, 2)-fuzzy points  $p_{(r_1, r_2)}^x, p_{(r_3, r_4)}^y$  of  $X$ ,  $cl(p_{(r_1, r_2)}^x) \neq cl(p_{(r_3, r_4)}^y)$ .

*Proof.* Let  $(X, \tau)$  be  $T_0$  and  $p_{(r_1, r_2)}^x, p_{(r_3, r_4)}^y$  be any two distinct (3, 2)-fuzzy points of  $X$ . Then, there exist two open (3, 2)-fuzzy sets  $L$  and  $K$  such that

$$\begin{aligned} L &= \{\langle x, 1, 0 \rangle, \langle y, 0, 1 \rangle\}, \\ \text{or } K &= \{\langle x, 0, 1 \rangle, \langle y, 1, 0 \rangle\}. \end{aligned} \quad (34)$$

Let  $L = \{\langle x, 1, 0 \rangle, \langle y, 0, 1 \rangle\}$  exist. Then,  $L^c = \{\langle x, 0, 1 \rangle, \langle y, 1, 0 \rangle\}$  is a closed (3, 2)-fuzzy set which does not contain  $p_{(r_1, r_2)}^x$  but contains  $p_{(r_3, r_4)}^y$ . Since  $cl(p_{(r_3, r_4)}^y)$  is the smallest closed (3, 2)-fuzzy set containing  $p_{(r_3, r_4)}^y$ , then  $cl(p_{(r_3, r_4)}^y) \subseteq L^c$ , and therefore  $p_{(r_1, r_2)}^x \notin cl(p_{(r_3, r_4)}^y)$ . Consequently,  $cl(p_{(r_1, r_2)}^x) \neq cl(p_{(r_3, r_4)}^y)$ .  $\square$

**Theorem 15.** Let  $(X, \tau)$  be a (3, 2)-fuzzy topological space. If  $p_{(1, 0)}^x$  is closed (3, 2)-fuzzy set for every  $x \in X$ , then,  $(X, \tau)$  is  $T_1$ .

*Proof.* Suppose  $p_{(1, 0)}^x$  is a closed (3, 2)-fuzzy set for every  $x \in X$ . Let  $p_{(r_1, r_2)}^x, p_{(r_3, r_4)}^y$  be any two distinct (3, 2)-fuzzy points of  $X$ ; then,  $x \neq y$  implies that  $p_{(1, 0)}^x c$  and  $p_{(1, 0)}^y c$  are two open (3, 2)-fuzzy sets such that

$$\begin{aligned} p_{(1, 0)}^y c &= \{\langle x, 1, 0 \rangle, \langle y, 0, 1 \rangle\}, \\ p_{(1, 0)}^x c &= \{\langle x, 0, 1 \rangle, \langle y, 1, 0 \rangle\}. \end{aligned} \quad (35)$$

Thus,  $(X, \tau)$  is  $T_1$ .  $\square$

#### 4. (3, 2)-Fuzzy Relations

A relation is a mathematical description of a situation where certain elements of sets are related to one another in some way. The system of fuzzy relation equations was first studied by Sanchez [18–21], who used it in medical research. Biswas [22] defined the method of intuitionistic medical diagnosis which involves intuitionistic fuzzy relations. Kumar et al. [23] used the applications of intuitionistic fuzzy set theory in diagnosis of various types of diseases. The notion of max-min-max composite relation for Pythagorean fuzzy sets was studied by Ejegwa [24], and the approach was improved and applied to medical diagnosis.

In this section, we introduce the notions of max-min-max composite relation and improved composite relation for (3, 2)-FSs. Moreover, we provide a numerical example to elaborate on how we can apply the composite relations to obtain the optimal choices.

*Definition 12.* Let  $X$  and  $Y$  be two (crisp) sets. The (3, 2)-fuzzy relation  $R$  (briefly, (3, 2)-FR) from  $X$  to  $Y$  is a (3, 2)-FS of  $X \times Y$  characterized by the degree of membership function  $\alpha_R$  and degree of non-membership function  $\beta_R$ . The (3, 2)-FR  $R$  from  $X$  to  $Y$  will be denoted by  $R(X \longrightarrow Y)$ . If  $D$  is a (3, 2)-FS of  $X$ , then

- (1) The max-min-max composition of the (3, 2)-FR  $R(X \longrightarrow Y)$  with  $D$  is a (3, 2)-FS  $C$  of  $Y$  denoted by  $C = R \circ D$  and is defined by

$$\begin{aligned}\alpha_{RoD}(n) &= \bigvee_m [\alpha_D(m) \wedge \alpha_R(m, n)], \\ \beta_{RoD}(n) &= \bigwedge_m [\beta_D(m) \vee \beta_R(m, n)],\end{aligned}\quad (36)$$

for all  $n \in Y$ .

- (2) The improved composite relation of  $R(X \longrightarrow Y)$  with  $D$  is a (3, 2)-FS  $C$  of  $Y$  denoted by  $C = R \circ D$ , such that

$$\begin{aligned}\alpha_{RoD}(n) &= \bigvee_m \left[ \frac{\alpha_D(m) + \alpha_R(m, n)}{2} \right], \\ \beta_{RoD}(n) &= \bigwedge_m \left[ \frac{\beta_D(m) + \beta_R(m, n)}{2} \right],\end{aligned}\quad (37)$$

for all  $n \in Y$ .

*Definition 13.* Let  $Q(X \longrightarrow Y)$  and  $R(Y \longrightarrow Z)$  be two (3, 2)-FRs. Then, for all  $(m, r) \in X \times Z$  and  $n \in Y$ ,

- (1) The max-min-max composition  $R \circ Q$  is the (3, 2)-fuzzy relation from  $X$  to  $Z$  defined by

$$\begin{aligned}\alpha_{RoQ}(m, r) &= \bigvee_n [\alpha_Q(m, n) \wedge \alpha_R(n, r)], \\ \beta_{RoQ}(m, r) &= \bigwedge_n [\beta_Q(m, n) \vee \beta_R(n, r)].\end{aligned}\quad (38)$$

- (2) The improved composite relation  $R \circ Q$  is the (3, 2)-fuzzy relation from  $X$  to  $Z$  such that

$$\begin{aligned}\alpha_{RoQ}(m, r) &= \bigvee_n \left[ \frac{\alpha_Q(m, n) + \alpha_R(n, r)}{2} \right], \\ \beta_{RoQ}(m, r) &= \bigwedge_n \left[ \frac{\beta_Q(m, n) + \beta_R(n, r)}{2} \right].\end{aligned}\quad (39)$$

*Remark 7.* The improved composite and max-min-max composite relations for (3, 2)-fuzzy sets are calculated by the following:

$$S_R = \alpha_{RoQ} - \beta_{RoQ} \cdot \pi_{RoQ}. \quad (40)$$

*Example 11.* Let  $D_1$  and  $D_2$  be two (3, 2)-fuzzy sets for  $X = \{x_1, x_2, x_3, x_4\}$ . Assume that

$$\begin{aligned}D_1 &= \{\langle x_1, 0.8, 0.61 \rangle, \langle x_2, 0.5, 0.87 \rangle, \langle x_3, 0.85, 0.55 \rangle, \langle x_4, 0.8, 0.69 \rangle\}, \\ D_2 &= \{\langle x_1, 0.7, 0.79 \rangle, \langle x_2, 0.78, 0.73 \rangle, \langle x_3, 0.6, 0.85 \rangle, \langle x_4, 0.89, 0.54 \rangle\}.\end{aligned}\quad (41)$$

By using Definitions 12 (1) and 13 (1), respectively, we find the max-min-max composite relation with application to  $D_1$  and  $D_2$  as follows:

$$\begin{aligned}\alpha_C(d_{1i}, d_{2k}) &= \bigvee_{x_j} [0.7, 0.5, 0.6, 0.8] = 0.8, \\ \beta_C(d_{1i}, d_{2k}) &= \bigwedge_{x_j} [0.79, 0.87, 0.85, 0.69] = 0.69.\end{aligned}\quad (42)$$

It is obvious that the minimum value of the membership values of the elements (that is,  $x_1, x_2, x_3, x_4$ ) in  $D_1$  and  $D_2$ , respectively, is 0.7, 0.5, 0.6, and 0.8. Also, the maximum value of the non-membership values of the elements (that is,  $x_1, x_2, x_3, x_4$ ) in  $D_1$  and  $D_2$ , respectively, is 0.79, 0.87, 0.85, and 0.69. From Remark 7, we can get

$$S_R = 0.8 - (0.69) \cdot (\sqrt{[5]} 0.0119) \approx 0.52. \quad (43)$$

Again, by using Definitions 12 (2) and 13 (2), respectively, we find the improved composite relation with application to  $D_1$  and  $D_2$  as follows:

$$\begin{aligned}\alpha_C(d_{1i}, d_{2k}) &= \bigwedge_{x_j} [0.75, 0.64, 0.725, 0.845] = 0.845, \\ \beta_C(d_{1i}, d_{2k}) &= \bigwedge_{x_j} [0.7, 0.8, 0.7, 0.615] = 0.615.\end{aligned}\quad (44)$$

From Remark 7, we can get

$$S_R = 0.845 - (0.615) \cdot (\sqrt{5}) \cdot 0.018423875 \approx 0.57. \quad (45)$$

Hence, from (43) and (45), we obtain that the improved composite relation produces better relation with greater relational value when compared to max-min-max composite relation.

## 5. Application of (3, 2)-Fuzzy Sets

We localize the idea of (3, 2)-FR as follows.

Let  $S = \{r_1, \dots, r_l\}$  be a finite set of subjects related to the colleges,  $C = \{b_1, \dots, b_m\}$  be a finite set of colleges, and  $A = \{t_1, \dots, t_n\}$  be a finite set of students. Suppose that we have two (3, 2)-FRs,  $U(A \rightarrow S)$  and  $R(S \rightarrow C)$ , such that

$$\begin{aligned}U &= \{ \langle (t, r), \alpha_U(t, r), \beta_U(t, r) \rangle \mid (t, r) \in A \times S \}, \\ R &= \{ \langle (r, b), \alpha_R(r, b), \beta_R(r, b) \rangle \mid (r, b) \in S \times C \}.\end{aligned}\quad (46)$$

where

$\alpha_U(t, r)$  denotes the degree to which the student ( $t$ ) passes the related subject requirement ( $r$ ).

$\beta_U(t, r)$  denotes the degree to which the student ( $t$ ) does not pass the related subject requirement ( $r$ ).

$\alpha_R(r, b)$  denotes the degree to which the related subject requirement ( $r$ ) determines the college ( $b$ ).

$\beta_R(r, b)$  denotes the degree to which the related subject requirement ( $r$ ) does not determine the college ( $b$ ).

$T = RoU$  is the composition of  $R$  and  $U$ . This describes the state in which the students,  $t_i$ , with respect to the related subject requirement,  $r_j$ , fit the colleges,  $b_k$ . Thus,

$$\begin{aligned}\alpha_T(t_i, b_k) &= \bigvee_{r_j \in S} [\alpha_U(t_i, r_j) \wedge \alpha_R(r_j, b_k)], \\ \beta_T(t_i, b_k) &= \bigwedge_{r_j \in S} [\beta_U(t_i, r_j) \vee \beta_R(r_j, b_k)],\end{aligned}\quad (47)$$

$\forall t_i \in A$  and  $b_k \in C$ , where  $i, j$ , and  $k$  take values from  $1, \dots, n$ .

The values of  $\alpha_{RoU}(t_i, b_k)$  and  $\beta_{RoU}(t_i, b_k)$  of the composition  $T = R \circ U$  are as follows (Table 1).

If the value of  $T$  is given by the following:

$$T = \alpha_T(t_i, b_k) - \beta_T(t_i, b_k) \cdot \pi_T(t_i, b_k), \quad (48)$$

then the student placement can be achieved.


**5.1. Application Example.** By using a hypothetical case with quasi-real data, we apply this method. Let  $A = \{t_1, t_2, t_3, t_4, t_5\}$  be the set of students for the colleges;  $S = \{\text{English Lang., Mathematics, Biology, Physics, Chemistry, Computer Sci.}\}$  be the set of related subject requirement to the set of colleges; and  $C = \{\text{College of Engineering (E), College of Medicine (M), College of Agricultural Engineering Sciences (AE), College of Sport Sciences (Sp), College of Science (S)}\}$  be the set of colleges the students are vying for (Algorithm 1).

From Table 4 and based on suitability of the students to the list of colleges, this decision making is made:

- (1)  $t_1$  and  $t_2$  are suitable to study at College of Agricultural Engineering Sciences.
- (2)  $t_3$  is suitable to study at College of Agricultural Engineering Sciences, College of Sport Sciences, and College of Science.
- (3)  $t_4$  is suitable to study at College of Medicine.
- (4)  $t_5$  is suitable to study at College of Agricultural Engineering Sciences and College of Science.

## 6. Discussion

The main idea of this work is to introduce a new type of fuzzy set called (3, 2)-FS. We illustrated that this type produces membership grades larger than intuitionistic and Pythagorean fuzzy sets which are already defined in the literature. However, Fermatean fuzzy sets give a larger space of membership grades than (3, 2)-FS. Figure 2 illustrates the relationships between these types of fuzzy sets.



We summarize the relationships in terms of the space of membership and non-membership grades in the following figure.

Regarding topological structure, we illustrated that every fuzzy topology in the sense of Chang (intuitionistic fuzzy topology and Pythagorean fuzzy topology) is a (3, 2)-fuzzy topology. In contrast, every (3, 2)-fuzzy topological space is a Fermatean fuzzy topological space because every (3, 2)-fuzzy subset of a set can be considered as a Fermatean fuzzy subset. The next example elaborates that Fermatean fuzzy topological space need not be a (3, 2)-fuzzy topological space.

**Example 12.** Let  $X = \{x_1, x_2\}$ . Consider the following family of Fermatean fuzzy subsets  $\tau = \{1_X, 0_X, D_1, D_2\}$ , where

TABLE 1: The composition  $R \circ U$ .

$R \circ U$	$E$	$M$	AE	Sp	$S$
$t_1$	(0.81, 0.60)	(0.81, 0.60)	(0.81, 0.61)	(0.81, 0.60)	(0.81, 0.60)
$t_2$	(0.82, 0.59)	(0.82, 0.60)	(0.82, 0.61)	(0.82, 0.59)	(0.82, 0.60)
$t_3$	(0.82, 0.60)	(0.82, 0.60)	(0.82, 0.61)	(0.82, 0.61)	(0.82, 0.61)
$t_4$	(0.82, 0.60)	(0.83, 0.60)	(0.82, 0.61)	(0.82, 0.61)	(0.82, 0.60)
$t_5$	(0.83, 0.59)	(0.83, 0.59)	(0.83, 0.60)	(0.83, 0.59)	(0.83, 0.60)

TABLE 2: The (3, 2)-fuzzy relation  $U(A \rightarrow S)$ .

$U(A \rightarrow S)$	Mathematics	Computer Sci.	English Lang.	Biology	Physics	Chemistry
$t_1$	(0.81, 0.61)	(0.80, 0.62)	(0.81, 0.61)	(0.80, 0.61)	(0.71, 0.71)	(0.81, 0.60)
$t_2$	(0.80, 0.61)	(0.81, 0.61)	(0.80, 0.61)	(0.62, 0.80)	(0.82, 0.60)	(0.82, 0.59)
$t_3$	(0.82, 0.61)	(0.82, 0.60)	(0.82, 0.61)	(0.80, 0.62)	(0.62, 0.80)	(0.81, 0.61)
$t_4$	(0.81, 0.62)	(0.83, 0.60)	(0.81, 0.61)	(0.81, 0.61)	(0.80, 0.61)	(0.82, 0.60)
$t_5$	(0.83, 0.59)	(0.82, 0.60)	(0.83, 0.60)	(0.82, 0.59)	(0.81, 0.59)	(0.83, 0.59)

TABLE 3: The (3, 2)-fuzzy relation  $R(S \rightarrow C)$ .

$R(S \rightarrow C)$	$E$	$M$	AE	Sp	$S$
Mathematics	(0.83, 0.59)	(0.84, 0.59)	(0.80, 0.62)	(0.82, 0.61)	(0.83, 0.60)
Computer Sci.	(0.82, 0.60)	(0.83, 0.59)	(0.80, 0.61)	(0.80, 0.62)	(0.80, 0.61)
English Lang.	(0.84, 0.59)	(0.83, 0.60)	(0.84, 0.59)	(0.83, 0.60)	(0.84, 0.59)
Biology	(0.81, 0.61)	(0.80, 0.609)	(0.80, 0.62)	(0.81, 0.61)	(0.81, 0.60)
Physics	(0.83, 0.60)	(0.82, 0.60)	(0.82, 0.61)	(0.82, 0.60)	(0.82, 0.60)
Chemistry	(0.83, 0.59)	(0.83, 0.60)	(0.82, 0.61)	(0.84, 0.59)	(0.83, 0.60)

TABLE 4: Greatest value given by  $T = \alpha_T(t_i, b_k) - \beta_T(t_i, b_k) \cdot \pi_T(t_i, b_k)$ .

$T$	$E$	$M$	AE	Sp	$S$
$t_1$	0.425	0.425	0.434	0.425	0.425
$t_2$	0.447	0.450	0.455	0.447	0.450
$t_3$	0.450	0.450	0.455	0.455	0.455
$t_4$	0.450	0.479	0.455	0.455	0.450
$t_5$	0.474	0.474	0.479	0.474	0.479

*Step 1.* The (3, 2)-fuzzy relation  $U(A \rightarrow S)$  and the (3, 2)-fuzzy relation  $R(S \rightarrow C)$  are given as in Tables 2 and 3, respectively. These data in (3, 2)- $F$  values are assumably obtained after students finished from preparatory school.

*Step 2.* Compute the composition  $R \circ U$  as in Table 1.

*Step 3.* Calculate  $T = \alpha_T(t_i, b_k) - \beta_T(t_i, b_k) \cdot \pi_T(t_i, b_k)$  as in Table 4.

*Step 4.* We present the decision making from Table 4. The greatest value of relation between students and colleges is taken for decisions.

ALGORITHM 1: Determination of the optimal college for students.

$$\begin{aligned}
 D_1 &= \{ \langle x_1, \alpha_{D_1}(x_1) = 0.75, \beta_{D_1}(x_1) = 0.81 \rangle, \langle x_2, \alpha_{D_1}(x_2) = 0.85, \beta_{D_1}(x_2) = 0.7 \rangle \}, \\
 D_2 &= \{ \langle x_1, \alpha_{D_2}(x_1) = 0.76, \beta_{D_2}(x_1) = 0.81 \rangle, \langle x_2, \alpha_{D_2}(x_2) = 0.86, \beta_{D_2}(x_2) = 0.7 \rangle \}.
 \end{aligned} \tag{49}$$




FIGURE 2: Comparison of grade space of IFs, PFs, FFs, and (3, 2)-FSs.

Observe that  $(X, \tau)$  is a Fermatean fuzzy topological space, but  $(X, \tau)$  is not a (3, 2)-fuzzy topological space.

## 7. Conclusions

In this paper, we have introduced a new generalized intuitionistic fuzzy set called (3, 2)-fuzzy sets and studied their relationship with intuitionistic fuzzy, Pythagorean fuzzy, and Fermatean fuzzy sets. In addition, some operators on (3, 2)-fuzzy sets are defined and their relationships have been proved. The notions of (3, 2)-fuzzy topology, (3, 2)-fuzzy neighborhood, and (3, 2)-fuzzy continuous mapping were studied. Furthermore, we introduced the concept of (3, 2)-fuzzy points and studied separation axioms in (3, 2)-fuzzy topological space. We also introduced the concept of relation to (3, 2)-fuzzy sets, called (3, 2)-FR. Moreover, based on academic performance, the application of (3, 2)-FSs was explored on student placement using the proposed composition relation.

In future work, more applications of (3, 2)-fuzzy sets may be studied; also, (3, 2)-fuzzy soft sets may be studied. In addition, we will try to introduce the compactness and connectedness in (3, 2)-fuzzy topological spaces. The motivation and objectives of this extended work are given step by step in this paper.

## Data Availability

No data were used to support this study.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## References

- [1] L. A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, no. 3, pp. 338–353, 1965.
- [2] B. Ahmad and A. Kharal, "On fuzzy soft sets," *Advances in Fuzzy Systems*, vol. 2009, Article ID 586507, 6 pages, 2009.
- [3] M. Atef, M. I. Ali, and T. M. Al-shami, "Fuzzy soft covering-based multi-granulation fuzzy rough sets and their applications," *Computational and Applied Mathematics*, vol. 40, no. 4, p. 115, 2021.
- [4] N. Çağman, S. Enginoğlu, and F. Çitak, "Fuzzy soft set theory and its application," *Iranian Journal of Fuzzy Systems*, vol. 8, no. 3, pp. 137–147, 2011.
- [5] K. T. Atanassov, "Intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol. 20, no. 1, pp. 87–96, 1986.
- [6] H. Garg and S. Singh, "A novel triangular interval type-2 intuitionistic fuzzy set and their aggregation operators," *Iranian Journal of Fuzzy Systems*, vol. 15, pp. 69–93, 2018.
- [7] H. Garg and K. Kumar, "An advanced study on the similarity measures of intuitionistic fuzzy sets based on the set pair analysis theory and their application in decision making," *Soft Computing*, vol. 22, no. 15, pp. 4959–4970, 2018.
- [8] H. Garg and K. Kumar, "Distance measures for connection number sets based on set pair analysis and its applications to decision-making process," *Applied Intelligence*, vol. 48, no. 10, pp. 3346–3359, 2018.
- [9] R. R. Yager, "Pythagorean fuzzy subsets," in *Proceedings of the 2013 joint IFSA world congress and NAFIPS annual meeting (IFSA/NAFIPS)*, pp. 57–61, IEEE, Edmonton, Canada, 2013.
- [10] T. Senapati and R. R. Yager, "Fermatean fuzzy sets," *Journal of Ambient Intelligence and Humanized Computing*, vol. 11, no. 2, pp. 663–674, 2020.
- [11] C. L. Chang, "Fuzzy topological spaces," *Journal of Mathematical Analysis and Applications*, vol. 24, no. 1, pp. 182–190, 1968.
- [12] R. Lowen, "Fuzzy topological spaces and fuzzy compactness," *Journal of Mathematical Analysis and Applications*, vol. 56, no. 3, pp. 621–633, 1976.
- [13] D. Çoker, "An introduction to intuitionistic fuzzy topological spaces," *Fuzzy Sets and Systems*, vol. 88, no. 1, pp. 81–89, 1997.
- [14] M. Olgun, M. Ünver, and Ş. Yardımcı, "Pythagorean fuzzy topological spaces," *Complex & Intelligent Systems*, vol. 5, no. 2, pp. 177–183, 2019.
- [15] H. Z. Ibrahim, "Fermatean fuzzy topological spaces," *Journal of Applied Mathematics and Informatics*, 2022, In press.
- [16] T. M. Al-shami, "On soft separation axioms and their applications on decision-making problem," *Mathematical Problems in Engineering*, vol. 2021, Article ID 8876978, 12 pages, 2021.
- [17] T. M. Al-shami and M. E. El-Shafei, "Partial belong relation on soft separation axioms and decision-making problem, two birds with one stone," *Soft computing*, vol. 24, no. 7, pp. 5377–5387, 2020.
- [18] E. Sanchez, "Resolution of composite fuzzy relation equations," *Information and Control*, vol. 30, no. 1, pp. 38–48, 1976.
- [19] E. Sanchez, "Solutions in composite fuzzy relation equations. Application to medical diagnosis in Brouwerian logic," in *Fuzzy Automata and Decision Processes*, M. M. Gupta, G. N. Saridis, and B. R. Gaines, Eds., pp. 221–234, Elsevier/North Holland, New York, NY, USA, 1977.
- [20] E. Sanchez, "Resolution of eigen fuzzy sets equations," *Fuzzy Sets and Systems*, vol. 1, no. 1, pp. 69–74, 1978.
- [21] E. Sanchez, "Compositions of fuzzy relations," in *Advances in Fuzzy Set Theory and Applications*, M. M. Gupta, R. K. Ragade, and R. R. Yager, Eds., pp. 421–433, North-Holland, Amsterdam, Netherlands, 1979.
- [22] R. Biswas, "Intuitionistic fuzzy relations," *Bull. Sous. Ens. Flous. Appl. (BUSEFAL)*, vol. 70, pp. 22–29, 1997.

- [23] S. Kumar, R. Biswas, and A. R. Ranjan, "An application of intuitionistic fuzzy sets in medical diagnosis," *Fuzzy Sets and Systems*, vol. 117, pp. 209–213, 2001.
- [24] P. A. Ejegwa, "Improved composite relation for pythagorean fuzzy sets and its application to medical diagnosis," *Granular Computing*, vol. 5, no. 2, pp. 277–286, 2020.