

N-acetyltransferase 2 slow acetylator genotype associated with adverse effects of sulfasalazine in the treatment of inflammatory bowel disease

Min Chen PhD¹, Bing Xia MD PhD^{1,2,3}, Bixiao Chen MD²,
Qiusha Guo PhD², Jin Li MD PhD¹, Mei Ye MD PhD³, Zhengguo Hu MD³

M Chen, B Xia, B Chen, et al. *N*-acetyltransferase 2 slow acetylator genotype associated with adverse effects of sulfasalazine in the treatment of inflammatory bowel disease. *Can J Gastroenterol* 2007;21(3):155-158.

AIM: *N*-acetyltransferase 2 (NAT2) is an important enzyme catalyzing *N*-acetylation of sulfasalazine (SASP). The aim of the present study was to investigate associations of the genotypes of NAT2 with inflammatory bowel disease (IBD), and with adverse effects of SASP, which is used as the first-line treatment of IBD.

PATIENTS AND METHODS: The wildtype allele (NAT2*4) and three variant alleles (NAT2*5B, NAT2*6A and NAT2*7B) of the NAT2 gene were determined in 101 patients with IBD (84 patients with ulcerative colitis and 17 patients with Crohn's disease) and 109 healthy controls by the polymerase chain reaction-restriction fragment length polymorphism method. Sixty-eight patients with IBD treated with SASP were followed, and their adverse reactions were recorded.

RESULTS: Eleven patients (16%) experienced adverse effects from SASP, including nine cases of sulfapyridine (SP) dose-related adverse effects and two cases of hypersensitivity (skin rash). Patients with the slow acetylator genotypes without the NAT2*4 allele experienced adverse effects more frequently (36%) than those with the fast acetylator genotypes with at least one NAT2*4 allele (11%), but the results were not significantly different (OR of 0.26, 95% CI 0.065 to 1.004; P=0.051). However, those with the slow acetylator genotypes experienced more SP dose-related adverse effects than those with the fast acetylator genotypes (36% versus 8%, OR of 0.17, 95% CI 0.039 to 0.749; P=0.019).

CONCLUSIONS: The NAT2 gene polymorphism was not associated with susceptibility to IBD in Chinese populations, but the NAT2 slow acetylator genotypes were significantly associated with SP dose-related adverse effects of SASP in the treatment of IBD.

Key Words: Adverse effects; Genetic polymorphism; Inflammatory bowel disease; *N*-acetyltransferase 2; Sulfasalazine

The human *N*-acetyltransferase 2 (NAT2) gene is located on human chromosome 8p22, and is highly polymorphic. To date, one allele coding for fast acetylator genotype (wild-type NAT2*4) and several mutated alleles coding for impaired acetylator genotype have been identified (1). Polymorphism of the NAT2 enzyme causes interindividual variations of the metabolism of drugs, arylamine and potential carcinogens (2). Studies on the polymorphism of NAT2 to evaluate genetic predisposition for development of certain diseases or to individualize drug therapy are now underway.

Inflammatory bowel disease (IBD), including ulcerative colitis (UC) and Crohn's disease (CD), is an entity of chronic intestinal

Génotype acétylateur lent de la *N*-acétyltransférase 2 associé aux effets indésirables de la sulfasalazine dans le traitement des maladies inflammatoires de l'intestin

OBJECTIF : La *N*-acétyltransférase 2 (NAT2) est une importante enzyme qui catalyse la *N*-acétylation de la sulfasalazine. Le but de la présente étude est d'analyser les liens entre les génotypes de la NAT2, les maladies inflammatoires de l'intestin (MII) et les effets indésirables de la SASP, utilisée en traitement de première intention dans les MII.

PATIENTS ET MÉTHODES : L'allèle sauvage (NAT2*4) et ses trois variantes (NAT2*5B, NAT2*6A et NAT2*7B) du gène NAT2 ont été identifiés chez 101 patients souffrant de MII (84 patients atteints de colite ulcéreuse et 17 atteints de la maladie de Crohn) et chez 109 sujets témoins en bonne santé au moyen de la méthode d'analyse du polymorphisme de la longueur des fragments de restriction des produits de la réaction en chaîne de la polymérase. Soixante-huit patients atteints de MII et traités par sulfasalazine ont été suivis et on a pris note de leurs réactions indésirables.

RÉSULTATS : Onze patients (16 %) ont manifesté des réactions indésirables à la sulfasalazine, dont neuf cas d'effets indésirables liés à la dose de sulfapyridine (SP) et deux cas d'hypersensibilité (éruptions cutanées). Les patients qui présentaient les génotypes acétylateurs lents, sans l'allèle NAT2*4, ont manifesté plus d'effets secondaires (36 %) que les porteurs des génotypes acétylateurs rapides, qui avaient au moins un allèle NAT2*4 (11 %), mais les résultats ne se sont pas révélés significativement différents (RR 0,26, IC 95 %, 0,065 à 1,004; P = 0,051). Par contre, les sujets qui présentaient les génotypes acétylateurs lents ont manifesté plus d'effets indésirables liés à la dose de SP que les porteurs de gènes acétylateurs rapides (36 %, contre 8 %, RR 0,17, IC 95 % 0,039 à 0,749; P = 0,019).

CONCLUSIONS : Le polymorphisme du gène NAT2 n'a pas été associé à une propension aux MII dans les populations chinoises, mais les génotypes acétylateurs lents du NAT2 ont été significativement associés aux effets indésirables liés à la dose de SP avec la sulfasalazine dans le traitement des MII.

inflammation with unknown etiology. Genetic susceptibility of this disease has been demonstrated by epidemiological studies, such as increased familial aggregations, an increased concordance in monozygotic twins and ethnic variations. Furthermore, environmental factors also contribute to the etiology of IBD. Xenobiotics that could initiate the development of IBD have been identified. Some of them are believed to be activating factors involved in the initiation of the disease. NAT2 is an important xenobiotic-metabolizing enzyme and is expressed in the intestinal mucosa (3). Current studies have shown an association between NAT2 gene polymorphisms and various autoimmune diseases such as systemic lupus erythematosus (4), rheumatoid arthritis (5)

¹Research Center of Digestive Diseases; ²Key Laboratory of Allergy and Immune-Related Diseases, School of Medicine, Wuhan University;

³Departments of Internal Medicine and Geriatrics, Wuhan University Zhongnan Hospital, Wuhan, People's Republic of China

Correspondence and reprints: Dr Bing Xia, Departments of Internal Medicine and Geriatrics, Wuhan University Zhongnan Hospital, Donghu Road 169, Wuhan 430071, Hubei Province, People's Republic of China. Telephone 86-27-67812985, fax 86-27-87307622, e-mail bingxia2004@yahoo.com.cn

Received for publication December 16, 2005. Accepted May 10, 2006

TABLE 1
Restriction sites and lengths of the nucleotide fragments produced by the digestion of the 535 base pair (bp) amplification products with Kpn I, Bam HI and Taq I enzymes

NAT2 alleles	Position (mutation)	Restriction enzymes	Recognition motif	Fragment pattern (bp); (bp); point alleles (wt)	Fragment pattern mutation
NAT2*5B	481 (C→T)	Kpn I	G-GTAC'C	483;52	535
NAT2*6A	590 (G→A)	Taq I	T'CG-A	205;170;160	330;205
NAT2*7B	857 (G→A)	Bam HI	G'GATC-C	428;107	535

A Adenine; C Cytosine; G Guanine; NAT2 N-acetyltransferase 2; T Thymine; wt Wildtype

and Behcet's disease (6). NAT2 gene polymorphisms may also be involved in the pathogenesis of IBD, but the association between NAT2 gene polymorphisms and IBD has not been studied yet.

Sulfasalazine (SASP) is the first-line therapy for induction and maintenance of remission in patients with UC and patients with CD, and is widely used in China. SASP is split into 5-aminosalicylic acid (5-ASA) and sulfapyridine (SP) by bacterial azoreductases in the colon. 5-ASA is effective in the treatment of IBD and inactivated to N-acetyl-5-ASA via NAT1 in the colon mucosa. SP is well absorbed from the colon, does not have beneficial effects in patients with IBD and results in adverse effects. It is inactivated to N-acetyl-SP via NAT2, subsequently hydroxylated and conjugated with glucuronide, and excreted in the urine.

It is well known that interindividual variability of therapeutic effectiveness and/or adverse reactions of drugs are often influenced by polymorphisms of metabolic enzymes. Inheritance of a slow acetylator genotype for NAT1 could potentially lead to increased efficacy. Similarly, adverse effects may occur more frequently in patients with slow acetylator NAT2 genotypes who are treated with SASP. Patients with rheumatoid arthritis with slow acetylator genotypes NAT2 have been shown to have more adverse effects than those with fast acetylator genotypes when taking SASP (7). We are interested in studying whether we can find the same results in patients with IBD when they take SASP.

In the present study, we investigated the association between NAT2 gene polymorphism and the susceptibility to IBD, as well as the association between the NAT2 genotype and the incidence of adverse effects of SASP in patients with IBD.

PATIENTS AND METHODS

Subjects

One hundred one Chinese patients with IBD (55 males and 46 females, age range 17 to 69 years, and mean age 37.8±1.38 years) were studied. The diagnosis of IBD was established according to previously published criteria (8). Of the 101 patients with IBD, 84 patients had UC and 17 patients had CD (CD is not common in China). The patients were recruited from the Departments of Internal Medicine and Geriatrics of Wuhan University Zhongnan Hospital (Wuhan, People's Republic of China) between 2000 and 2003. One hundred nine healthy physical examiners (62 men and 47 women, age range 25 to 76 years, mean age 45.82±8.73 years) made up the control group and were all affiliated with the same hospital. All subjects were from the Hubei province, were unrelated to each other and were of Chinese Han nationality. The study protocol was approved by the ethics committee of Wuhan University Zhongnan Hospital, and all subjects gave informed consent.

Determination of NAT2 genetic polymorphism

The most common mutations in the Chinese population, at positions C481T, G590A and G857A (nucleotides: adenine [A], thymine [T], cytosine [C] and guanine [G]) of the NAT2 gene, were determined by the polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method as described previously (9). According to the nomenclature of the NAT2 gene, the wildtype and three variant alleles were defined as NAT2*4, and NAT2*5B, NAT2*6A and NAT2*7B. Determination of the NAT2 genotype was performed by individuals who were blind to the clinical data of the patients.

Briefly, genomic DNA was isolated from 5 mL of venous blood by conventional proteinase K and phenol/chloroform method. The primers were as follows: NAT2(+) 5'-GCCTCAGGTGC-CTTGCATTT-3' and NAT2(-) 5'-CGTGAGGGTAGAGAG-GATAT-3'. Each PCR cycle was performed in a total volume of 50 µL (containing 30 ng of DNA, 10 mM of tris hydrochloride, 1.5 mM of magnesium chloride, 200 µM of each deoxyribonucleotide triphosphate, 20 pmol of primer and two units of Taq I polymerase). After initial denaturation at 95°C for 5 min, 30 cycles of amplification were carried out with denaturation at 95°C for 50 s, annealing at 55°C for 50 s and extension at 72°C for 50 s, followed by the final extension at 72°C for 5 min. To establish the presence or absence of each single nucleotide polymorphism, PCR products were digested with 10 U Kpn I, Bam HI or Taq I (TaKaRa Biotechnology Inc, Dalian, People's Republic of China) at 37°C for 3 h and the fragments were separated on 2% agarose gel. Restriction sites and lengths of the nucleotide fragments produced by the digestion of the 535 base pair PCR products with Kpn I, Bam HI and Taq I enzymes, are shown in Table 1.

Acetylator genotype

NAT2 acetylator genotypes were produced according to previously published data (10). Homozygotes (NAT2*4/NAT2*4) or heterozygotes (NAT2*4/NAT2*5B, NAT2*4/NAT2*6A and NAT2*4/NAT2*7B) for the dominant NAT2*4 wildtype allele were classified as fast acetylator genotypes, and homozygotes of the mutant alleles (NAT2*5B, NAT2*6A and NAT2*7B) were classified as slow acetylator genotypes.

Definition of adverse effects

All of the 101 patients with IBD were followed by telephone, mail or interview every three months. The clinical characteristics, drug histories and adverse effects of the medications of each patient were carefully recorded on the registry form of IBD patients. This form was designed and validated by the IBD Collaborative Study Group in China supervised by the Chinese Association of Gastroenterology. The definition of adverse effects was according to the criteria proposed by Edwards and Aronson (11). Namely, the adverse effects were classified as 'certain', 'probable', 'possible', 'unlikely', 'conditional' and 'unassessable' according to the definition.

Statistical analysis

The data were analyzed using SPSS (SPSS Inc, USA) statistical software. The departure of the Hardy-Weinberg equilibrium was tested using the χ^2 test. Frequency of the allele and acetylator genotypes in each group was compared by χ^2 test or Fisher's exact tests if needed. The association between NAT2 acetylator genotype and adverse effects of SASP was assessed by binary logistic regression analysis. ORs and 95% CIs were also calculated. P<0.05 was considered to be statistically significant.

TABLE 2
Distribution of *N*-acetyltransferase 2 (*NAT2*) alleles and acetylator genotypes in inflammatory bowel disease (IBD) patients and in healthy controls (con)

	Alleles				Acetylator genotypes	
	<i>NAT2</i> *4 n (%)	<i>NAT2</i> *5B n (%)	<i>NAT2</i> *6A n (%)	<i>NAT2</i> *7B n (%)	Fast n (%)	Slow n (%)
Con (n=109)	130 (59.6)	12 (5.5)	51 (23.4)	25 (11.5)	86 (78.9)	23 (21.1)
IBD (n=101)	116 (57.4)	9 (4.5)	49 (24.3)	28 (13.9)	79 (78.2)	22 (21.8)
UC (n=84)	98 (58.3)	8 (4.8)	40 (23.8)	22 (13.1)	68 (80.9)	16 (19.1)
CD (n=17)	18 (52.9)	1 (2.9)	9 (26.5)	6 (17.6)	11 (64.7)	6 (35.3)

CD Crohn's disease; UC Ulcerative colitis

RESULTS

Distribution of the *NAT2* allele and acetylator genotypes

The distribution of allele and acetylator genotypes of *NAT2* in patients with IBD and healthy controls is shown in Table 2. The allele frequencies of *NAT2**4, *NAT2**5B, *NAT2**6A and *NAT2**7B were 0.57, 0.05, 0.24 and 0.14, respectively, in patients with IBD, and the allele frequencies of *NAT2**4, *NAT2**5B, *NAT2**6A and *NAT2**7B were 0.60, 0.06, 0.23 and 0.12, respectively, in healthy controls. *NAT2* slow acetylator genotypes were detected in 21.8% of patients with IBD and 21.1% of healthy controls. The frequencies of *NAT2* alleles and slow acetylator genotypes of the IBD patients were not significantly different from those of the controls.

Association between *NAT2* acetylator genotype and adverse effects of SASP

Of the 101 patients with IBD, 68 patients (67.3%) were treated with SASP. Adverse effects of SASP occurred in 11 patients (16%). Nine patients experienced SP dose-related adverse effects (four patients had severe gastrointestinal disorders, four had headaches and one patient had renal dysfunction). The other two patients experienced hypersensitivity (skin rash).

As shown in Table 3, patients with the slow acetylator genotypes who had no *NAT2**4 allele experienced adverse effects more frequently than those with the fast acetylator genotypes who had at least one *NAT2**4 allele, but the results were not significantly different (36% versus 11%, OR of 0.26, 95% CI 0.065 to 1.004; P=0.051). When the overall adverse effects were subdivided into two groups – SP dose-related adverse effects and hypersensitive reactions – those with the slow acetylator genotypes experienced SP dose-related adverse effects more frequently than the fast acetylator genotypes (36% versus 8%, OR of 0.17, 95% CI 0.039 to 0.749; P=0.019).

DISCUSSION

The *NAT2* gene is highly polymorphic. Each mutant allele contains a combination of one or more nucleotide substitutions, which occur at the following positions: 191 (G→A [Arg64→Glu]), 282 (C→T [Tyr94]), 341 (T→C [Ile114→Thr]), 481 (C→T [Leu161]), 590 (G→A [Arg197→Gln]), 803 (A→T [Lys268→Arg]) and 857 (G→A [Gly286→Glu]). Of the 25 *NAT2* mutant alleles, three mutant alleles (*NAT2**5A, *NAT2**6A and *NAT2**7B) have been shown to be commonly associated with slow acetylator status, and can be identified by the detection of the mutations at positions 481, 590 and 857 of the coding exon of the *NAT2* gene. By detecting these three main alleles, many studies (12-14) have observed a high correlation

TABLE 3
Risk factors for adverse effects of sulfasalazine (SASP) in the treatment of inflammatory bowel disease

Risk factors	Overall adverse effects		Serum SP dose-related adverse effects	
	With adverse effects	Without adverse effects	With adverse effects	Without adverse effects
Male/female, n	33/24	5/6	33/24	3/6
Age, years (mean ± SD)	39.19±11.74	42.36±17.81	39.19±11.74	44.78±18.38
SASP dose (mean ± SD)	2.92±0.82	2.55±0.68	2.92±0.82	2.55±0.72
Treatment time, months (mean ± SD)	13.00±8.52	16.73±7.54	13.00±8.52	16.67±7.66
Fast/slow acetylator genotypes, n	48/9	6/5*	48/9	4/5†

*36% versus 11%, OR of 0.26, 95% CI 0.065 to 1.004, P=0.051; †36% versus 8%, OR of 0.17, 95% CI 0.039 to 0.749, P=0.019. SP Sulfapyridine

between the genotype and phenotype of *NAT2*. The prediction rate has been reported to be between 88% and 100%. Gao et al (15) determined these three mutant alleles by real-time fluorescence light-cycle technique, and found that the frequencies of *NAT2**4, *NAT2**6A, *NAT2**7B and *NAT2**5B among the Chinese were 60.4%, 24.1%, 13.7% and 1.8%, respectively. Our results have shown that the frequencies of *NAT2**4, *NAT2**6A, *NAT2**7B and *NAT2**5B were 59.6%, 23.4%, 11.5% and 5.5%, respectively, and were similar to the results found by Gao et al (15).

NAT2 slow acetylator genotypes have been linked with increased susceptibility to certain autoimmune diseases (4-6). It has been suggested that nonacetylated xenobiotics may accumulate in slow acetylators and are subsequently metabolized by other enzymes into reactive intermediates. These reactive intermediates may alter self-proteins presented to the immune system and stimulate T cells, which in turn initiate pathological and clinical signs of autoimmunity (16). However, our study failed to detect any association between the *NAT2* slow acetylator genotypes and susceptibility to IBD in the Chinese population, suggesting that the *NAT2* slow acetylator genotypes may not be an independent risk factor in the pathogenesis of IBD.

The incidence of adverse effects of SASP among the Western populations has ranged from 25% to 40% (14,17), but in our study, the incidence of adverse effects of SASP among the Chinese people was only 16%. This difference may be explained by the low frequency of slow acetylator genotypes in the Asian population (18). Surprisingly, 36% of those with slow acetylator genotypes who were treated with SASP had experienced adverse effects, while adverse effects occurred in only 11% of those with fast acetylator genotypes. It is suggested that those with slow acetylator genotypes are more likely to manifest SP dose-related effects than those with fast acetylator genotypes. However, these data also suggest that other factors may be involved, and prospective studies are needed to determine whether closer monitoring of those with slow acetylator genotypes will lead to a decrease in SP adverse events.

In general, the adverse effects of SASP were divided into two groups: adverse events (eg, vomiting and headache) appeared to be dependent on the serum SP concentration (greater than 50 µg/mL) (19), and hypersensitive reactions. The latter are independent of SP concentration, and are often

TABLE 4
N-acetyltransferase 2 (NAT2) allele frequencies in the Chinese population and other ethnic populations

Population (reference)	2n	NAT2*4 (%)	NAT2*5B (%)	NAT2*6A (%)	NAT2*7B (%)	NAT2*14 (%)
Chinese (present case)	218	59.6	5.50	23.4	11.5	0.0
Japanese (20)	316	64.0	1.90*	23.1	11.1	0.0
Caucasian (21)	744	25.0†	45.0†	28.0	1.30†	0.0
African (21)	256	36.0†	30.0†	22.0	2.00†	9.0†

* $P < 0.05$; † $P < 0.01$ compared with the Chinese population

manifested as skin rash and fever. In our study, we found no significant association between the NAT2 slow acetylator genotypes and the overall adverse effects of SASP. However, when we subdivided the overall adverse effects into two groups – SP-related adverse effects and hypersensitive reactions – we found a strong association between the NAT2 slow acetylator genotypes and adverse effects of SASP. Tanaka et al (7) reported a strong association between the NAT2 slow acetylator genotypes and overall adverse effects of SASP, while Kumagai et al (14) and Ricart et al (17) did not find these overall associations significant, but they found that adverse effects occurred more frequently in those with slow acetylator genotypes. In their studies, adverse effects of hypersensitive reactions were very common (61% and 52%, respectively). Although slow acetylator genotypes are also suspected to be a risk factor for development of hypersensitivity reactions (9,17), the results are controversial (13). In our study, only two patients had hypersensitivity reactions and it was difficult to evaluate the

REFERENCES

- Hein DW, Doll MA, Fretland AJ, et al. Molecular genetics and epidemiology of the NAT1 and NAT2 acetylation polymorphisms. *Cancer Epidemiol Biomarkers Prev* 2000;9:29-42.
- Hein DW. Molecular genetics and function of NAT1 and NAT2: Role in aromatic amine metabolism and carcinogenesis. *Mutat Res* 2002;506-507:65-77.
- Hickman D, Pope J, Patil SD, et al. Expression of arylamine N-acetyltransferase in human intestine. *Gut* 1998;42:402-9.
- Gunnarsson I, Kanerud L, Pettersson E, Lundberg I, Lindblad S, Ringertz B. Predisposing factors in sulphasalazine-induced systemic lupus erythematosus. *Br J Rheumatol* 1997;36:1089-94.
- Pawlik A, Ostaneck L, Brzosko I, Gawroska-Szklarz B, Brzosko M, Dabrowska-Zamojcin E. Increased genotype frequency of N-acetyltransferase 2 slow acetylation in patients with rheumatoid arthritis. *Clin Pharmacol Ther* 2002;72:319-25.
- Najim RA, Sharquie KE, Al-Janabi MH. Acetylator phenotype in Behcet's disease. *Adv Exp Med Biol* 2003;528:131-8.
- Tanaka E, Taniguchi A, Urano W, et al. Adverse effects of sulfasalazine in patients with rheumatoid arthritis are associated with diplotype configuration at the N-acetyltransferase 2 gene. *J Rheumatol* 2002;29:2492-9.
- Sanderson JD. IBD: Diagnosis and control. *Practitioner* 1997;241:120-2,127-8,130.
- Zielinska E, Niewiarowski W, Bodalski J, et al. Genotyping of the arylamine N-acetyltransferase polymorphism in the prediction of idiosyncratic reactions to trimethoprim-sulfamethoxazole in infants. *Pharm World Sci* 1998;20:123-30.
- Yokogawa K, Nakaharu T, Ishizaki J, et al. Kinetic phenotypic diagnosis of N-acetylation polymorphism in patients based on ratio of urinary metabolites of salicylazosulfapyridine. *Int J Pharm* 2001;229:183-91.
- Edwards IR, Aronson JK. Adverse drug reactions: Definitions, diagnosis, and management. *Lancet* 2000;356:1255-9.
- Deguchi T, Mashimo M, Suzuki T. Correlation between acetylator phenotypes and genotypes of polymorphic arylamine N-acetyltransferase in human liver. *J Biol Chem* 1990;265:12757-60.
- Tanigawara Y, Kita T, Aoyama N, et al. N-acetyltransferase 2 genotype-related sulfapyridine acetylation and its adverse events. *Biol Pharm Bull* 2002;25:1058-62.
- Kumagai S, Komada F, Kita T, et al. N-acetyltransferase 2 genotype-related efficacy of sulfasalazine in patients with rheumatoid arthritis. *Pharm Res* 2004;21:324-9.
- Gao JB, Huang YD, Lin JA, Zhu QC, Liang JP. [Relationship between genetic polymorphisms of N-acetyltransferase and susceptibility to hepatocellular carcinoma]. *Zhonghua Gan Zang Bing Za Zhi* 2003;11:20-2.
- von Schmiedeberg S, Goebel C, Gleichmann E, Uetrecht J. Neutrophils and drug metabolism. *Science* 1995;268:585-6. (Lett)
- Ricart E, Taylor WR, Loftus EV, et al. N-acetyltransferase 1 and 2 genotypes do not predict response or toxicity to treatment with mesalamine and sulfasalazine in patients with ulcerative colitis. *Am J Gastroenterol* 2002;97:1763-8.
- Cascorbi I, Brockmoller J, Mrozikiewicz PM, Muller A, Roots I. Arylamine N-acetyltransferase activity in man. *Drug Metab Rev* 1999;31:489-502.
- Das KM, Eastwood MA, McManus JP, Sircus W. The metabolism of salicylazosulfapyridine in ulcerative colitis. I. The relationship between metabolites and the response to treatment in inpatients. *Gut* 1973;14:631-41.
- Lin HJ, Han CY, Lin BK, Hardy S. Ethnic distribution of slow acetylator mutations in the polymorphic N-acetyltransferase (NAT2) gene. *Pharmacogenetics* 1994;4:125-34.
- Bell DA, Taylor JA, Butler MA, et al. Genotype/phenotype discordance for human arylamine N-acetyltransferase (NAT2) reveals a new slow-acetylator allele common in African-Americans. *Carcinogenesis* 1993;14:1689-92.

associations. We found that the genetic polymorphism of NAT2 showed great differences among different ethnic groups.

The comparisons of NAT2 allele frequencies between the Chinese population and other ethnic populations are shown in Table 4. The frequency of the NAT2*4 allele in the Chinese population was similar to that of the Japanese population (20), and was significantly higher than the Caucasian and African populations (21). Compared with the Caucasian (NAT2*5B, 45%; NAT2*7B, 1.3%) and African (NAT2*5B, 30%; NAT2*7B, 2.0%) populations, the Chinese exhibited a very low-frequency of NAT2*5B (5.5%) and a relatively higher frequency of NAT2*7B (11.5%). The frequency in the Chinese population was similar to that in the Japanese population. NAT2*14 only existed in the African population. It had been found that the frequency of the slow acetylator activities presented in almost 50% of Caucasians and only 10% in Asians (18). The genetic differences among the populations may cause this difference in the phenotype of the populations. This difference should be considered when drug medication is developed by internationally harmonized clinical trials and marketed in various countries.

CONCLUSION

The NAT2 genetic polymorphism was not associated with the susceptibility to IBD in Chinese patients, but the slow acetylator genotypes were significantly associated with SP dose-related adverse effects of SASP in IBD treatment.

ACKNOWLEDGEMENT: The present study was supported by a grant received by Dr Bing Xia from the Ministry of Education of China (20030486013).



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

