

Description of two new *ABCB11* mutations responsible for type 2 benign recurrent intrahepatic cholestasis in a French-Canadian family

Yannick Beauséjour MD PhD¹, Fernando Alvarez MD², Martin Beaulieu PhD³, Marc Bilodeau MD¹

Y Beauséjour, F Alvarez, M Beaulieu, M Bilodeau. Description of two new *ABCB11* mutations responsible for type 2 benign recurrent intrahepatic cholestasis in a French-Canadian family. *Can J Gastroenterol* 2011;25(6):311-314.

Benign recurrent intrahepatic cholestasis is a rare clinical entity that is caused by mutations in the canalicular transport genes. The present report describes two individuals from the same family whose symptoms were typical of the clinical characteristics of type 2 benign recurrent intrahepatic cholestasis. Sequencing of the *ABCB11* gene revealed two previously unreported mutations that predict the absence of expression of the protein. The clinical presentation of the current cases are discussed, as are the differential diagnosis and genetic characteristics of the hereditary cholestatic disorders, overemphasizing the possibility of making a definite genetic diagnosis.

Key Words: *ABCB11*; Benign recurrent cholestasis

Benign recurrent intrahepatic cholestasis (BRIC) is a rare cause of hereditary cholestatic liver disorders, and is sometimes difficult to differentiate from progressive familial intrahepatic cholestasis (PFIC). The main difference between BRIC and PFIC is their clinical and biochemical evolution. BRIC is characterized by a benign course, with recurrent episodes of cholestasis and complete normalization of liver biochemistry between each episode (1). PFIC is characterized by the progressive appearance of pruritis, growth failure, coagulopathy and, eventually, liver failure at a relatively young age (2-4). However, the boundary between these two entities can sometimes overlap, which makes precise, early diagnosis difficult (5). Therefore, they are often considered as a phenotypic continuum with variable course. In recent years, several proteins involved in biliary physiology, such as hepatocanalicular transporters, have been found to harbour mutations that were later associated with BRIC and PFIC syndromes. Therefore, based on these findings, redefinition of these disorders is currently underway.

Bile formation is the result of the concerted action of membrane transporters in hepatocytes, and is mainly driven by the secretion of bile acids. Indeed, the canalicular membrane is equipped with transporters belonging to the ATP binding cassette (ABC) superfamily – the hepatocanalicular transporters. Bile constituents such as bile salts, phosphatidylcholine and cholesterol are actively secreted by *ABCB11* (bile salt export pump [BSEP]), *ABCB4* (MDR3) and *ABCG5/8* respectively. In addition, *ATP1B1* (FIC1), which is a P-type ATPase (ATP-dependent aminophospholipid translocase), is involved in maintaining the integrity of the canalicular membrane (6,7). Therefore, it is not surprising that mutations of these transporters are associated with inherited liver diseases, albeit in an autosomal recessive pattern (8) (Table 1).

La description de deux nouvelles mutations *ABCB11* responsables d'une cholestase intrahépatique bénigne de type 2 dans une famille canadienne-française

La cholestase intrahépatique récurrente bénigne est une entité clinique rare causée par des mutations des gènes de transport canaliculaires. Le présent rapport décrit le cas de deux personnes d'une même famille dont les symptômes sont représentatifs des avec les caractéristiques cliniques de la cholestase intrahépatique récurrente bénigne de type 2. Le séquençage du gène *ABCB11* a révélé deux mutations jamais signalées auparavant, lesquelles prédisaient l'absence d'expression de la protéine. La présentation clinique de ces cas est exposée, de même que le diagnostic différentiel et les caractéristiques génétiques des troubles cholestatiques héréditaires. On insiste sur la possibilité de poser un diagnostic génétique précis.

CASE PRESENTATION

The present report describes the discovery of two previously unreported mutations in the *ABCB11* gene identified in two Caucasian sisters previously diagnosed with BRIC. The history of the clinical development of their syndromes is described.

Both sisters were diagnosed in early childhood – one in the neonatal period and the second at nine months of age. Intractable intermittent pruritis and jaundice were the dominant clinical features. Ursodeoxycholic acid was prescribed soon after diagnosis. Pruritis was controlled with rifampicin and/or cholestyramine. During the 20 to 25 years of follow-up, neither patient exhibited signs or complications of chronic liver failure; in fact, the liver always appeared normal when examined by ultrasound. Therefore, liver enlargement and splenomegaly were never detected. Of note, one of the two sisters developed acute pancreatitis at a young age, which was judged to be secondary to biliary sludge without, however, any evidence of cholelithiasis. This same patient was also diagnosed with unilateral deafness of unexplained etiology at a very young age.

In the course of the evolution of their disease, these patients developed two to three episodes of clinically significant jaundice of approximately six months to 1.5 years duration each. Interestingly, the duration and severity of each occurrence appeared to be longer and more severe. Moderate hepatocyte injury was observed each time, and was characterized by a two- to eightfold increase in the upper limit of normal (ULN) levels of serum aspartate aminotransferase and alanine aminotransferase, three times the ULN for alkaline phosphatase levels, and bilirubin levels reaching more than 200 µmol/L. However, gamma-glutamyl transpeptidase activity always remained normal. Both patients experienced high blood levels of low-density lipoprotein and low levels

¹Liver Unit, Centre hospitalier de l'Université de Montréal; ²Department of Pediatrics, Centre hospitalier universitaire Sainte-Justine; ³Department of Biochemistry, Centre hospitalier de l'Université de Montréal, Montréal, Québec

Correspondence: Dr Marc Bilodeau, Liver Unit, Centre hospitalier de l'Université de Montréal, 1058 St-Denis, Montréal, Québec H2X 3J4.

Telephone 514-890-8000 ext 35749, fax 514-412-7314, e-mail marc.bilodeau@umontreal.ca

Received for publication September 28, 2010. Accepted December 9, 2010

TABLE 1
Canalicular transporter and related mutations

Gene involved	Chromosome	Canalicular transport defect	Clinical impact	Histopathology
<i>ATP8B1</i>	18q21-q22	Aminophospholipid flippase	Cholestasis, intractable pruritis and extrahepatic symptoms (pancreatitis, deafness)	Bland cholestasis
		BRIC-I	Diarrhea	Portal-track fibrosis
		PFIC-I	Coagulopathy (PFIC-I)	
			Progression to liver failure (PFIC-I)	
<i>ABCB11</i>	2q24-31	Bile salt export pump	Cholestasis, intractable pruritis	Bland cholestasis
		BRIC-II	Lithiasis	Portal-track fibrosis
		PFIC-II	Coagulopathy (PFIC-II)	Bile duct proliferation
		ICP	Progression to liver failure (PFIC-II)	Portal inflammation and giant-cell hepatitis
			Increased risk for hepatocarcinoma	
		Severe itching during the third trimester of pregnancy (ICP)		
		Premature delivery and stillbirth (ICP)		
<i>ABCB4</i>	7q21	Phosphatidylcholine floppase	Mild pruritis	Bland cholestasis
		PFIC-III	Microlithiasis, cholesterol stones, sludge	Marked bile duct proliferation
			Progressive cholestasis	
			High gamma-glutamyl transpeptidase levels	
		Biliary cirrhosis		
<i>ABCC2</i>		Dubin-Johnson syndrome	Mild icterus	Black liver, pigment contained in lysosomes (black or blue)
		MRP-2	Vague abdominal pain	
			Weakness	
			Hepatosplenomegaly	
<i>ABCG5/8</i>		Sitosterolemia	Anemia, premature atherosclerosis	
		Cholesterol floppase		

BRIC Benign recurrent intrahepatic cholestasis; *ICP* Intrahepatic cholestasis of pregnancy; *MRP-2* Multidrug resistance protein-2; *PFIC* Progressive familial intrahepatic cholestasis

of high-density lipoprotein cholesterol in the blood during the episodes as well (total cholesterol 7.68 mmol/L and 5.68 mmol/L; low-density lipoprotein cholesterol 6.3 mmol/L and 4.0 mmol/L; and high-density lipoprotein cholesterol 0.34 mmol/L and 0.66 mmol/L). These episodes were most often triggered by mild upper respiratory tract infections; however, some occurred without a clear triggering factor. Of note, on one occasion, cholestasis appeared in the first two weeks following the introduction of minimal doses of oral contraceptives, suggesting that both events were linked. Neither of the sisters have become pregnant. Overall, these two patients were subjected to multiple liver biopsies during their childhood. Results showed evidence of canalicular cholestasis, mainly located in the centrilobular area, without any signs of biliary duct damage, proliferation or fibrosis. In between episodes, both sisters reported chronic mild pruritis.

To assess the genetic transmission of their syndrome, the phenotype of the parents of the two sisters was investigated and a family pedigree was created. Both parents were of French-Canadian descent and had been living in a small area (± 100 km²) of the Lanaudière region (Quebec) for at least the past four generations, but they were not related. They had another unaffected 28-year-old female child.

The mother (55 years of age) had no history of jaundice, cholestatic complications during her pregnancies or mutations in the hemochromatosis *HFE* gene. Her liver biochemistry was completely normal. She was only diagnosed with osteoporosis. On the other hand, nine of her 11 brothers and sisters underwent gallbladder surgery in the past.

The father (57 years of age) was diagnosed with hereditary hemochromatosis 20 years previously; four of his eight brothers and sisters were also diagnosed with the disease. The condition is currently well controlled biochemically with phlebotomy treatment. A physical examination showed mild joint deformities without darkening of the skin or liver enlargement. His liver biochemistry was normal, except for a slightly elevated total bilirubin level (less than two times ULN). Genetic screening revealed that he was homozygous for the C282Y mutation in the hemochromatosis *HFE* gene. Noteworthy, the father had no history of jaundice, pruritis or pancreatitis. However, he underwent cholecystectomy at 47 years of age for symptomatic gallstones.

Finally, he suffers from partial hereditary deafness, which required stapes prosthetic replacement. Four of his eight brothers have the same disorder.

Mutations in the *ABCB11* gene were investigated by direct sequencing of polymerase chain reaction products generated from genomic DNA extracts of peripheral blood cells drawn from one of the two sisters. A splice mutation (390 G→T [G130G]) was found on one allele, while a missense sequence variant (830 C→A [A277E]) was found on the other allele. Mutation nomenclature is based on the recommendation of the American College of Medical Genetics designating the initial ATG start codon as +1 using the reference sequences NT_005403 and NM_003742. The genetic codons in which these mutations are localized are well conserved in mammals according to DNA alignment software. The splice mutation is located at a well-identified splice site that is translated in the canalicular domain of the protein (9). Therefore, splice modifications are likely to lead to significant changes in the protein. The missense mutation is located in the second intracellular domain of the protein. Without detailed mutagenesis studies, however, it is difficult to predict the impact of this mutation on the protein. Similar missense mutations have been reported in this region of the protein (9). The other sister was found to harbour the exact two same mutations. Sequencing of *ABCB4* and *ATP8B1* genes was also performed for one of the two sisters, with no mutations identified. Therefore, it was decided that these cases be classified as type 2 BRIC (BRIC-II) (Table 1).

Finally, to evaluate the genetic phase transmission of these mutations, the *ABCB11* gene in both parents was sequenced. The mother was found to have the 390 G→T (G130G) mutation, while the father harboured the 830 C→A (A277E) mutation. Therefore, the daughters' genotype was transmitted from both parents.

DISCUSSION

Synthesized and recycled bile acids are transported from hepatocytes across canalicular membranes against a concentration gradient driven by an ATP-dependent pump. This ATP-dependent transporter is encoded by the gene *ABCB11*, which is also known as the BSEP. The

BSEP is expressed primarily in the liver, but is also expressed at lower levels in other organs such as human testis (10), brain cortex (11), placenta (12), small and large intestines (13), and kidneys (14). The functional role of this protein in nonhepatic tissues is not completely clear. Autosomal recessive mutations in this gene are responsible for rare hereditary forms of cholestatic liver disorders known as BRIC-II/PFIC-II. To date, approximately 150 mutations in the *ABCB11* gene have been identified (9,15-19). These mutations include insertions, deletions, splice or nonsense mutations of the gene leading to reduced expression of the protein in hepatocytes, thus reducing or completely inhibiting its activity. Among the two most common mutations in individuals of European descent are the E297G and D482G mutations, which collectively account for approximately 58% of all cases (9).

In the present report, we described two young sisters who shared two previously unreported mutations in *ABCB11*; however, even if these mutations are expected to inactivate the product of this gene, it would require protein expression studies to confirm it. The sisters inherited these genes from their parents in a genetic trans phase because each of the two parents is heterozygous for one of the identified mutations. As previously mentioned, BRIC is inherited in an autosomal recessive pattern, and parents of a proband are generally obligate carriers of the disease-causing mutation. Interestingly, the probability of observing, as was the case in the present study, two sisters with the same mutation in an autosomal recessive pattern is only 6%.

Clinically, both sisters presented with bouts of jaundice that lasted approximately six months to 1.5 years, and were characterized by a mixed pattern of hepatocellular injury with normal levels of gamma-glutamyl transpeptidase. As documented previously, minor viral infections have been responsible for these acute episodes. Interestingly, each new exacerbation seems to be more severe in intensity and in duration over time. However, neither of the sisters have shown evidence of liver failure to date, with serum albumin and coagulogram levels remaining close to normal. For the moment, this benign evolution seems to be consistent with the diagnosis of BRIC-II.

A strong penetrance for cholelithiasis seems to be present in this family, especially for the splice mutation variant G130G. Several mutations have been shown to increase the risk of cholelithiasis in cases of nonfamilial BRIC. Our observation is based only on a family review and not on genetic sequencing; therefore, the results should be interpreted with caution. Previous reports (20) have linked mutations in *ABCB11* and *ABCB4*, along with *ABCG8* of the canalicular *ABCG5/ABCG8* cholesterol transporter with gallstone disease. This can be easily understood considering that these transporters carry compounds that have all been implicated in the formation of gallstones. Furthermore, a coding variant in *ABCG8* has been documented

to be significantly associated with gallstones in large patient panels (21,22); however, no *ABCB11* variants have been implicated in gallstone formation outside the context of BRIC-II (23).

Others have reported a correlation between the type of mutation and disease severity. Indeed, mutations that were predicted to affect expression or function of the gene protein to a greater extent, such as frameshift, nonsense and large deletions, were detected more frequently in patients with progressive disease (3). According to this hypothesis, the new mutations described in the present report could be sufficient to cause acquired gallstone disease in heterozygous patients, and BRIC-II in homozygotes. One intriguing finding, however, is that the two homozygote cases have not presented with gallstones to date.

It was also surprising to find evidence of probable extrahepatic manifestations of the disease. Indeed, deafness is believed to be exclusively observed in BRIC-I/PFIC-I disease (24,25). In our case study, one of the two daughters has a hearing problem of unclear origin. We do not believe that the history of stapes disorder in the father's family was linked to the *ABCB11* mutations because this has not been previously documented, and because it was not the diagnosis in the affected daughter. Finally, the history of acute nonlithiatic pancreatitis in the same case is also original because this type of extrahepatic manifestation is, again, more characteristic of BRIC-I even though it has also been rarely observed in BRIC-II (4,24).

Aside from genotyping, it is also possible to perform immunohistochemical analysis of the BSEP protein on liver biopsy specimens (9,18). This may enable a more rapid diagnosis if the technique is already mastered in the local laboratory. Despite generally good agreement between the immunohistochemical evaluation of the BSEP protein and the presence of *ABCB11* mutations, discrepancies exist, particularly with certain genotypes (18). Finally, transport studies have also been performed to test the functional capacity of the mutated protein *in vitro* (18,26).

Clearly, more data need to be accumulated to better characterize the clinical spectrum of these rare disorders. This is especially important to be able to more effectively predict the clinical evolution of these patients. In particular, we acknowledge that hepatocellular carcinoma has been described in individuals harbouring *ABCB11* mutations (9,27). Furthermore, it is possible to observe some BRIC patients who eventually evolve toward liver failure and subsequently adopt a PFIC phenotype (28). Finally, the identification of such new mutations in a French-Canadian family raises the possibility that these mutations are specific to this particular population, as has been described in other areas of the world (29). With the advent of molecular genetic testing, it is now possible to identify and categorize these individuals more clearly.

REFERENCES

- Luketic VA, Shiffman ML. Benign recurrent intrahepatic cholestasis. *Clin Liv Dis* 2004;8:133-49.
- Davit-Spraul A, Gonzales E, Baussan C, Jacquemin E. Progressive familial intrahepatic cholestasis. *Orphanet J Rare Dis* 2009;4:1.
- Alissa FT, Jaffe R, Shneider BL. Update on progressive familial intrahepatic cholestasis. *J Pediatr Gastroenterol Nutr* 2008;46:241-52.
- van Mil SW, van der Woerd WL, van der Brugge G, et al. Benign recurrent intrahepatic cholestasis type 2 is caused by mutations in *ABCB11*. *Gastroenterology* 2004;127:379-84.
- van Ooteghem NA, Klomp LW, van Berge-Henegouwen GP, Houwen RH. Benign recurrent intrahepatic cholestasis progressing to progressive familial intrahepatic cholestasis: Low GGT cholestasis is a clinical continuum. *J Hepatol* 2002;36:439-43.
- Harris MJ, Le Couteur DG, Arias IM. Progressive familial intrahepatic cholestasis: Genetic disorders of biliary transporters. *J Gastroenterol Hepatol* 2005;20:807-17.
- Bull LN, van Eijk MJ, Pawlikowska L, et al. A gene encoding a P-type ATPase mutated in two forms of hereditary cholestasis. *Nat Genet* 1998;18:219-24.
- Karsen TH, Hov JR. Genetics of cholestatic liver disease in 2010. *Curr Opin Gastroenterol* 2010;26:251-8.
- Strautnieks SS, Byrne JA, Pawlikowska L, et al. Severe bile salt export pump deficiency: 82 different *ABCB11* mutations in 109 families. *Gastroenterology* 2008;134:1203-14.
- Langmann T, Mauerer R, Zahn A, et al. Real-time reverse transcription-PCR expression profiling of the complete human ATP-binding cassette transporter superfamily in various tissues. *Clin Chem* 2003;49:230-8.
- Loncar J, Popovic M, Zaja R, Smital T. Gene expression analysis of the ABC efflux transporters in rainbow trout (*Oncorhynchus mykiss*). *Comp Biochem Physiol C Toxicol Pharmacol* 2010;151:2:209-15.
- Patel P, Weerasekera N, Hitchins M, Boyd CA, Johnston DG, Williamson C. Semi quantitative expression analysis of *MDR3*, *FIC1*, *BSEP*, *OATP-A*, *OATP-C*, *OATP-D*, *OATP-E* and *NTCP* gene transcripts in 1st and 3rd trimester human placenta. *Placenta* 2003;24:39-44.
- Torok M, Gutmann H, Fricker G, Drewe J. Sister of P-glycoprotein expression in different tissues. *Biochem Pharmacol* 1999;57:833-5.
- Huls M, van den Heuvel JJ, Dijkman HB, Russel FG, Masereeuw R. ABC transporter expression profiling after ischemic reperfusion injury in mouse kidney. *Kidney Int* 2006;69:2186-93.
- Chen HL, Liu YJ, Su YN, et al. Diagnosis of *BSEP/ABCB11* mutations in Asian patients with cholestasis using denaturing high performance liquid chromatography. *J Pediatr* 2008;153:825-32.

16. Goto K, Sugiyama K, Sugiura T, et al. Bile salt export pump gene mutations in two Japanese patients with progressive familial intrahepatic cholestasis. *J Pediatr Gastroenterol Nutr* 2003;36:647-50.
 17. Chen HL, Chang PS, Hsu HC, et al. *FIC1* and *BSEP* defects in Taiwanese patients with chronic intrahepatic cholestasis with low gamma-glutamyltransferase levels. *J Pediatr* 2002;140:119-24.
 18. Noe J, Kullak-Ublick GA, Jochum W, et al. Impaired expression and function of the bile salt export pump due to three novel *ABCB11* mutations in intrahepatic cholestasis. *J Hepatol* 2005;43:536-43.
 19. Pauli-Magnus C, Stieger B, Meier Y, Kullak-Ublick GA, Meier PJ. Enterohepatic transport of bile salts and genetics of cholestasis. *J Hepatol* 2005;43:342-57.
 20. Hoblinger A, Lammert F. Genetics of biliary tract diseases: New insights into gallstone disease and biliary tract cancers. *Curr Opin Gastroenterol* 2008;24:363-71.
 21. Buch S, Schafmayer C, Volzke H, et al. A genome-wide association scan identifies the hepatic cholesterol transporter *ABCG8* as a susceptibility factor for human gallstone disease. *Nat Genet* 2007;39:995-9.
 22. Grunhage F, Acalovschi M, Tirziu S, et al. Increased gallstone risk in humans conferred by common variant of hepatic ATP-binding cassette transporter for cholesterol. *Hepatology* 2007;46:793-801.
 23. Schafmayer C, Tepel J, Franke A, et al. Investigation of the Lith1 candidate genes *ABCB11* and *LXRA* in human gallstone disease. *Hepatology* 2006;44:650-7.
 24. Pawlikowska L, Strautnieks S, Jankowska I, et al. Differences in presentation and progression between severe *FIC1* and *BSEP* deficiencies. *J Hepatol*;53:170-8.
 25. Stapelbroek JM, Peters TA, van Beurden DH, et al. *ATP8B1* is essential for maintaining normal hearing. *Proc Natl Acad Sci USA* 2009;106:9709-14.
 26. Ho RH, Leake BF, Kilkenny DM, et al. Polymorphic variants in the human bile salt export pump (*BSEP*; *ABCB11*): Functional characterization and interindividual variability. *Pharmacogenet Genomics* 2010;20:45-57.
 27. Knisely AS, Strautnieks SS, Meier Y, et al. Hepatocellular carcinoma in ten children under five years of age with bile salt export pump deficiency. *Hepatology* 2006;44:478-86.
 28. Davit-Spraul A, Fabre M, Branchereau S, et al. *ATP8B1* and *ABCB11* analysis in 62 children with normal gamma-glutamyl transferase progressive familial intrahepatic cholestasis (PFIC): Phenotypic differences between PFIC1 and PFIC2 and natural history. *Hepatology* 2010;51:1645-55.
 29. Liu LY, Wang ZL, Wang XH, et al. *ABCB11* gene mutations in Chinese children with progressive intrahepatic cholestasis and low gamma glutamyltransferase. *Liver Int* 2010;30:809-15.
-



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

