

Review Article

Mechanisms for the Invasion and Dissemination of *Salmonella*

Qiao Li 

Tongji Hospital, Tongji Medical College, Huazhong University of Sciences and Technology, Wuhan, Hubei, China

Correspondence should be addressed to Qiao Li; graceliqiao@126.com

Received 26 November 2021; Revised 15 May 2022; Accepted 30 May 2022; Published 9 June 2022

Academic Editor: Jayaraman Tharmalingam

Copyright © 2022 Qiao Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Salmonella enterica is a gastroenteric Gram-negative bacterium that can infect both humans and animals and causes millions of illnesses per year around the world. *Salmonella* infections usually occur after the consumption of contaminated food or water. Infections with *Salmonella* species can cause diseases ranging from enterocolitis to typhoid fever. *Salmonella* has developed multiple strategies to invade and establish a systemic infection in the host. Different cell types, including epithelial cells, macrophages, dendritic cells, and M cells, are important in the infection process of *Salmonella*. Dissemination throughout the body and colonization of remote organs are hallmarks of *Salmonella* infection. There are several routes for the dissemination of *Salmonella typhimurium*. This review summarizes the current understanding of the infection mechanisms of *Salmonella*. Additionally, different routes of *Salmonella* infection will be discussed. In this review, the strategies used by *Salmonella enterica* to establish persistent infection will be discussed. Understanding both the bacterial and host factors leading to the successful colonization of *Salmonella enterica* may enable the rational design of effective therapeutic strategies.

1. Introduction

Salmonella is a Gram-negative, intracellular pathogen. There are more than 2,600 serovars of *Salmonella* characterized to date that are differentiated on the basis of the lipopolysaccharide (LPS) O antigen and the flagellar H antigen [1]. There are just two species of *Salmonella*: *S. enterica* and *S. bongori* [2]. *Salmonella typhi* and *Salmonella typhimurium* are well-known members of the *S. enterica* species. *S. typhimurium* and *S. enteritidis* are predominantly associated with gastroenteritis in humans [3]. Each year, there are about 155,000 deaths due to nontyphoidal *Salmonella* (NTS) infections. *Salmonella enterica* serovar typhi infections cause a staggering 20 million infections and 200,000 deaths annually [4]. Gastroenteritis induced by *Salmonella* infections is a major cause of morbidity and mortality in children under 5 years of age [5]. Diarrhea caused by *Salmonella* species causes a global human health burden that contributes to significant annual morbidity and mortality and requires new therapeutic strategies for effective management. Almost 60% of *Salmonella* strains have developed resistance to first-line antibiotics [6]. Most patients recover from infections after treatment. However, 3–5% of patients become

chronic carriers, with chronic infection in the gall bladder [7]. Chronic carriers can intermittently shed the bacteria through their feces and urine throughout the rest of their lives [8]. The liver is also a reservoir for chronic infections with *Salmonella Typhi*; from the liver, the bacteria can be intermittently shed into the gallbladder [9].

Salmonella typhi infections can cause fever, hepatomegaly, splenomegaly, and bacteremia. In the disease process, the bacteria disseminate into the gall bladder, liver, and spleen [10]. Approximately 90% of chronic *Salmonella* carriers have gallstones [10–12], and are at significantly increased risk for gallbladder cancer (GC) [12, 13]. Del-Giorno et al. reported that persistent *Salmonella* infections can cause pancreatitis in a murine model of infection [14]. Some *Salmonella* carriers are asymptomatic. Roughly 2–5% of *Salmonella*-infected patients fail to clear the bacteria within one year [12]. Such chronic infections, especially asymptomatic infections, pose a huge socioeconomic burden, especially in South Asian and African countries, by unknowingly spreading infections to others, who may experience symptomatic infections and suffer economic costs as a result. Understanding the cellular routes of *Salmonella* invasion and dissemination in the host and the mechanisms

of *Salmonella* persistent infection may facilitate the exploration of novel treatment strategies for patients with chronic infections. Ultimately, this may help eliminate the asymptomatic carriage of *Salmonella* as a concern for public health.

Salmonella infections can result from the ingestion of contaminated foods because they can survive the low pH of the stomach [15]. Although bile in the small intestine poses a challenge for *S. typhimurium*, the PhoQ/PhoP two-component regulatory system mediates resistance to bile [16]. *Salmonella* predominantly causes inflammation of the terminal ileum and colon [17]. *S. typhimurium* can spread systemically in mice, and *S. typhimurium* infections in mice are used as an animal model for typhoid fever in humans [18]. Pretreatment of mice with streptomycin prior to *Salmonella* infection disturbs the healthy microbiota and facilitates infection of the intestinal lumen with *Salmonella typhimurium* [19]. Streptomycin-treated mice are therefore often used as animal models of *S. Typhimurium*-induced gastroenteritis [19]. Before the oral infection of *S. typhimurium* and *S. enteritidis*, approximately 20 mg of streptomycin treatment by intragastric administration in the mice will allow a high colonization level in the cecum and colon of the mice [20]. Acute microbiota depletion will reduce the colonization resistance and facilitate the infection of the bacteria. Microbiota can limit *Salmonella* colonization, and diet can affect microbiota composition. Low-fiber or high-fat diets will increase *S. typhimurium* colonization in mice [21]. Fat can promote *S. typhimurium* infection in mice by eliciting bile salts, which help fat digestion [21]. A high-fat diet will cause microbiota perturbation [21]. *E. coli* may limit *S. typhimurium* infections during diet shifts [21].

Mice with a mutation in the natural resistance-associated macrophage protein 1 gene (*Nramp1*), such as CL57/BL6 or BALB/C mice, are susceptible to *Salmonella* infection [22]. *Nramp1* is a macrophage-specific exporter, and the *Nramp1* gene codes for an ion transporter that pumps ions out of *Salmonella*-containing vacuoles (SCV) [22]. The SCV is the intracellular vacuolar niche in which *Salmonella* can replicate and achieve dormant infection. Wild type 129 \times 1/Sv mice, which possess the *Nramp1*^{+/+} allele, are used as an animal model for chronic *S. typhimurium* infection [23]. Mice with a mutation in the natural resistance-associated macrophage protein 1 gene (*Nramp1*), such as CL57/BL6 or BALB/C mice, are susceptible to *Salmonella* infection [22]. *Nramp1* is a macrophage-specific exporter, and the *Nramp1* gene codes for an ion transporter that pumps ions out of SCV [22]. The SCV is the intracellular vacuolar niche in which *Salmonella* can replicate and achieve dormant infection. Wild type 129 \times 1/Sv mice, which possess the *Nramp1*^{+/+} genotype, are used as an animal model for chronic *S. typhimurium* [23].

2. M Cells

Enteropathogenic infections start in the intestinal lumen. Dissemination through microfold or membranous (M) cells is one of the best-understood routes of *Salmonella* dissemination [24]. M cells are specialized follicle-associated epithelial (FAE) enterocytes on the surface of mucosa-

associated lymphoid tissues [25, 26]. *Salmonella typhimurium* initiates infection in mice by infecting and destroying the specialized epithelial M cells and then traveling to the mesenteric lymph nodes [24]. See Figure 1.

Salmonella directly invades M cells but can also transform follicle-associated epithelial cells into M cells to provide additional routes for intestinal invasion [27]. Indeed, Tahoun et al. found that *S. Typhimurium* can induce an epithelial-mesenchymal transition (EMT) of FAE enterocytes and transition the FAE to M cells [27]. These processes rely on the bacterial type III effector protein SopB [27]. Through the activation of NF- κ B and Wnt/b-Catenin signaling pathways, *Salmonella* induces host cell transdifferentiation through receptor activator of NF- κ B ligand (RANKL) [27]. This finding was the first report that *S. typhimurium* can transform epithelial cells into M cells using a single virulence factor.

Intestinal immunity is the first defense barrier that enteropathogens encounter during infection. Lymphotoxin signaling is important for maintaining intestinal immune balance. LT β R can also be activated by lymphotoxin (LT $\alpha\beta$) [28]. Lymphotoxin signaling promotes the differentiation of M cells from intestinal epithelial cells [29]. This signaling is involved in the regulation of intestinal inflammation, as shown by the DSS-induced colitis model [30]. Mice with knocked-out lymphotoxin signaling molecules (LT α_3 , LT $\alpha_2\beta_1$, and LT $\alpha_1\beta_2$) have abnormal lymphoid development [31]. Lymphotoxin β -receptor knockout mice lack all lymph nodes and gut-associated lymphatic tissues, including Peyer's patches (PPs) [32]. These lymph node-defective mice are a good model for the systemic dissemination of *S. typhimurium*. Infection of *Salmonella* in LT β R^{-/-} mice demonstrates that organized lymph tissues are dispensable for the systemic infection of the host [20]. As shown by a study from Barthel et al., without Peyer's patches (PPs), bacteria can still reach remote organs [20]. This phenomenon indicates the importance of dendritic cell-mediated transportation in the dissemination of *S. typhimurium* [33]. *Salmonella* exploited dendritic cells as vesicles for dissemination. Cheminay et al. showed that after infection by *Salmonella*, dendritic cells could upregulate the CCR7 receptor and migrate via the CCR7 ligands CCL19 and CCL211 [13, 33–35].

A study by Wroblewska et al. showed that lymphotoxin signaling is essential for the clearance of *Salmonella* from the intestinal lumen [36]. A lack of LT β R signaling did not impact the initiation of inflammation induced by *Salmonella*. However, the resolution of *Salmonella* infection was impaired [36]. The infectious processes in *S. typhimurium* in WT and LT β R^{-/-} mice lacking Peyer's patches (PPs) and MLN are highly similar [20].

3. Epithelial Cell

S. typhimurium can invade polarized gallbladder epithelial cells and replicate inside the epithelial cells [37]. Gallbladder epithelial cells are a reservoir for *Salmonella* colonization [37]. Long-term colonization of *Salmonella* in the




FIGURE 1: The multiple routes of *Salmonella* dissemination. *Salmonella* can be phagocytized by macrophages or dendritic cells and disseminate to the Peyer's patches (PP), mesenteric lymph nodes, and eventually the liver and spleen. *Salmonella* can also reach the circulation by the transportation of dendritic cells, e.g., CD18-expressing phagocytes. *Salmonella* can also invade the intestinal epithelial cells directly by way of a zipper or trigger mechanism. *Salmonella* can achieve persistent infection in epithelial cells and phagocytic cells. The dormant persisters can be released and induce the recurrence of the infection.

gallbladder cells can drive the premalignant transformations of the cells. *Salmonella* can invade the polarized gallbladder cells. *Salmonella* can induce the extrusion of epithelial cells, which is accompanied by caspase-1 activation-related cell death. Epithelial cells can provide a shelter for the bacteria to survive and replicate in the cytosol of the epithelial cells [38, 39]. The type III secretion system is involved in the priming of the bacteria for invasion. Cytosolic bacteria can induce the extrusion of epithelial cells and be released into the intestinal lumen [38, 39].

Unlike M cells, *Salmonella*'s invasion of epithelial cells does not rely on phagocytosis. The type III secretion system (T3SS) is the most important virulence factor for *Salmonella* species, and one is encoded on *Salmonella* pathogenicity island 1 (SPI1) and the other is encoded on *Salmonella* pathogenicity island 2 (SPI2) [40]. The type III secretion system is a molecular syringe that can translocate the effector proteins directly from the bacteria into the cytosol of cells. Effector proteins are injected into the cytoplasm of the host by a T3SS gene cluster. SPI1 is involved in the invasion process of *Salmonella* [41]. After invading host cells, *Salmonella* survives in SCVs by using elements encoded on SPI2 [42–45]. Approximately 4–6 h after the cellular invasion, bacterial replication is initiated [46].

Salmonella can induce membrane ruffling in intestinal cells to cause them to engulf the bacteria [47]. Various *S. Typhimurium* fimbrial operons contribute to bacterial attachment and invasion of epithelial cells [48]. The zipper and trigger mechanisms are two well-studied mechanisms of *Salmonella* entry into epithelial cells [49, 50]. The trigger mechanism is activated by the type III secretory system [49]. SipB/C in *Salmonella* type III secretory system assembles a pore in the epithelial cell, bacteria and epithelial cells can contact through the continuum created by the SipB/C [49]. Cytoskeletal reorganizations known as “membrane ruffles”

and “internalization” are two key elements of the trigger mechanism [51]. Bacteria are internalized in SCV following a trigger mechanism [49].

In contrast, there are only minor cytoskeletal protein rearrangements involved in the zipper mechanism [50]. Instead, the zipper mechanism is mainly mediated by interactions between bacterial ligands such as Rck and host cell surface receptors [52]. There are many outer membrane proteins that participate in the invasion process of *Salmonella typhimurium* [53]. Rck is a 17 kDa outer membrane protein (OMP), which are membrane proteins found in the outer membranes of Gram-negative bacteria. Rck is encoded by the rck gene on the large virulence plasmid [54]. They are a family of highly conserved OMPs within the Enterobacteriaceae family. This receptor binding leads to downstream signal activation mediated by the phosphorylation of tyrosine kinase. The zipper mechanism is activated by the binding of host cell receptors by the bacterial ligands. Actin polymerization and membrane extension are initiated by the activated downstream signaling.

PagN is another OMP [54] and is widely conserved in the *Salmonella* genus [55]. The PagN protein interacts with cell surface heparin sulfate proteoglycans to invade cells [53]. Binding between OmpV and the extracellular matrix components fibronectin and $\alpha1\beta1$ integrin leads to the adhesion of *Salmonella typhimurium* to intestinal epithelial cells and ultimately activates actin modulation [56]. PAMPs of *Salmonella* can be recognized by the innate immune response receptors through MyD88-dependent TLR signaling [57]. Infection with SPI1 T3SS disrupted *Salmonella* can still induce colitis in C57BL/6 mice through a mechanism that is dependent on MyD88 signaling [58]. The effectors of type III secretion systems in the invasion and dissemination of *Salmonella* are summarized in Table 1.

TABLE 1: Effectors of Type III secretion systems in the invasion and dissemination of *Salmonella*.

Type III secretion System	Function	References
SPI-1	Invasion of nonphagocytic cells, including epithelial cells	[40]
SPI-2	Survive in the phagocytic cells	[59]
SseI	Inhibition of macrophages and DCs migration	[60]
SipA	Promotion of cytoskeletal rearrangements, invasion of epithelial cells	[61, 62]
SopA	Invasion of epithelial cells	[63]
SopB	Invasion of epithelial cells	[63]
SopD	Invasion of epithelial cells	[63]
SopE2	Invasion of epithelial cells	[63]
SipB/C	Translocator to deliver the effectors into the cell	[49]

The binding of pattern recognition receptors (PRRs) with pathogen-associated molecular patterns (PAMPs), including peptidoglycan, lipopolysaccharide, flagellin, can mediate *Salmonella* invasion [64, 65]. TLR4 and TLR5 play a role in the host response to *Salmonella* [66]. In human macrophages, *Salmonella* can activate NAIP/NLRC4 and canonical NLRP3 Inflammasomes by its flagellin [67]. Caspase-1 will be activated after binding with NLRC4 and NLRP3 inflammasomes in response to *Salmonella*. *Salmonella* colonization was much higher in caspase 11 deficient mice than in wild-type mice [68]. Casp1^{-/-} and Casp1/11^{-/-} monolayers showed significantly increased intracellular bacteria, accompanied by low intestinal epithelial cells (IECs) shedding and death [68]. Caspase activation is important for limiting the intracellular replication of *Salmonella* [68].

4. Dendritic Cells

Intestinal dendritic cells are found in Peyer's patches [69], in the lamina propria [70], in the subepithelial dome [71], and under the follicle epithelium [72, 73]. The phagocytosis of *Salmonella* by dendritic cells and macrophages is mediated by the interactions between specific pathogen-associated molecular patterns (PAMP) and cellular receptors on the phagocyte surface, such as pattern recognition receptors (PRRs), which include Toll-like receptors (TLRs), NOD-like receptors (NLRs), and C-type Lectin receptors [74]. NOD-like receptors (NLRs), nucleotide-binding leucine-rich repeat-containing proteins, are intracellular innate immune receptors that belong to the pattern recognition receptors (PRRs) [75]. NLR is short for nucleotide-binding domain leucine-rich repeat. MyD88-and TRIF-dependent pathways can be regulated by NLRs [75].

Dendritic cells are exploited by *Salmonella* typhimurium as "Trojan horses" to enable systemic dissemination [76]. This strategy of manipulating host cell migration to facilitate broader dissemination is common among other pathogens such as *Mycobacterium tuberculosis*, HIV, and a range of other Gram-negative bacteria [77–80]. For example, after phagocytes are infected by HIV, the gp120 protein on the virus binds with the C-type Lectin receptor DC-Sign, initiating phagocytosis by dendritic cells that then migrate to lymph nodes and release viral particles that proceed to infect CD4+ lymphocytes [77–79]. Several Gram-negative bacteria also disseminate through antigen-presenting cells, as

demonstrated by Yang et al. [76, 81–88]. After binding to C-type lectin receptors with core LPS, bacteria are transported throughout the host by antigen-presenting cells [76, 81–88].

By exploiting migratory dendritic cells, the *Salmonella* can thus traffic from the intestinal lumen to systemic organs [34]. During active infection, the dendritic cells' expression of CCR7, a receptor for the chemokines CCL19 and CCL21, is increased [34]. This allows dendritic cells to migrate along chemotactic gradients to remote sites like the lymph nodes and spleen [34]. *Salmonella* survives inside the dendritic cells, subverts the function of dendritic cells, impairs the activation of adaptive immune responses, prevents fusion and lyso-endosomal degradation, and achieves systemic dissemination [45]. Cheminay et al. published the first example that *Salmonella* can inhibit antigen presentation by dendritic cells by altering MHC-II-dependent antigen presentation in an SPI2-dependent manner [89]. Through subversion of the antigen presentation of dendritic cells, the bacteria reduce the activation of the active immune response. Lapaque et al. demonstrated that *Salmonella* can inhibit the surface expression of MHC class II antigens on dendritic cells through ubiquitination [90].

CD103⁺CD11b⁺ DCs have been reported to transport *Salmonella* typhimurium to the mesenteric lymph nodes (MLN) after oral infection [91]. CD103⁺ dendritic cells (DCs) typically phagocytose bacteria from the small intestine and present antigens to T cells [91]. Another group of dendritic cells that can facilitate the dissemination of *Salmonella* is intestinal CD11c⁺ lamina propria cells (LPCs), which do so in a TLR5⁻dependent manner [92]; the migration of *Salmonella* typhimurium from the intestinal tract to MLN is impaired in TLR5^{-/-} mice. In TLR5^{-/-} mice, migration of bacteria by CD11c + LPCs is impaired [92, 93].

Distinct populations of dendritic cells participate in the processing and immune sampling of *Salmonella*. Specialized DC subsets in Peyer's patches (PPs), CCR6 (+) DCs, are recruited to the dome regions of Peyer's patches (PPs) to sample the bacteria and present to CD4⁺ T cells [94, 95]. CX3CR1-positive lamina propria DCs take up *S. typhimurium* by transepithelial processes [96]. Indeed, CX3CR1 deficiency leads to reduced bacterial sampling in the intestinal lumen by lamina propria DCs [96]. Further, these CX3CR1-positive DCs lacked CCR6 expression, which is different from the Peyer's patches (PPs) associated-dendritic cells [96].

S. typhimurium can be taken up by sub-epithelial DCs and can survive within murine PP dendritic cells [97]. The *S. typhimurium* strain PhoP^r has a point mutation in the phoP/Q locus [98] that attenuates its capacity to survive in macrophages but was able to persist for several weeks *in vivo* [97]. *Salmonella* can persist in the dendritic cells in the Peyer's patch. They can also be directly sampled by dendritic cells that express tight junction proteins, such as the interepithelial dendritic cells in the intestinal villi that penetrate gut epithelial monolayers by opening tight junctions and directly sampling bacteria from the mucus [99].

Infection of CD11c⁻CD18⁺ dendritic cells can lead to rapid entry into the systemic circulation. It has been reported by Vazquez-Torres et al. that *Salmonella* can achieve systemic dissemination through CD18-expressing phagocytes [100]. One hour after infection, *Salmonella* can be detected in the blood. At sites other than M cells and Peyer's patches, *Salmonella* can also disseminate from the gastrointestinal tract to the spleen. Downregulation of DC cells in the lamina propria can limit the invasion of *Salmonella* [100].

5. Macrophage

During the intracellular life of *Salmonella* in the host cells, *Salmonella* can interfere with the antigen-presenting process of the dendritic cells, for example, by interfering with the antigen presentation of bacteria on dendritic cells and inhibiting the adaptive immunity, *Salmonella* can affect the polarization of macrophages to the M2 phenotype, which will inhibit the inflammatory process and facilitate the persistent survival of *Salmonella* in the host. The manipulation of the macrophage is a strategy that *Salmonella* derived during its evolution. Uchiya et al. demonstrated that *Salmonella* can interfere with the function of macrophages to escape immune responses. Uchiya et al. reported that *Salmonella* can inhibit cytokine signaling in macrophages via the Janus kinase/signal transducer and activator of transcription (JAK/STAT) signaling pathway through SPI2 [101].

In addition to dendritic cells and M cells, *S. typhimurium* can also disseminate via inflammatory monocytes. Monocytes are recruited to the inflammatory sites where they differentiate into macrophages. Macrophages serve as a reservoir in which *Salmonella* can survive and replicate [102]. Inside the macrophage, *Salmonella* can induce micropinocytosis [103], and spacious phagosomes (SP) are formed after *Salmonella* enters the macrophage and persists in the cytoplasm [103]. A T3SS encoded by SPI2 allows survival and avoids the NADPH oxidase-dependent killing of macrophages [104].

The PhoQ/PhoP regulatory system is utilized by *S. typhimurium* to enable survival in macrophages [105]. The PhoQ/PhoP two-component system is one of the most important regulatory mechanisms for the virulence of *Salmonella*. Inside the SCV, the low PH and low Mg²⁺ environment activate the two-component PhoQ/PhoP system [106]. The gene regulating the expression of O antigen, rfb, is inhibited inside the SCV [107]. Thus, the length of O antigen

is decreased under the regulation of the two-component PhoQ/PhoP system. The protease PgtE in *Salmonella typhimurium*, a homologue for Pla in *Yersinia Pestis* and OmpT in *E. coli*, is then expressed [108]. Expression of PgtE protease dissolves the extracellular matrix and facilitates the cellular dissemination of *Salmonella in vivo*. *S. typhimurium*, when released from the macrophage, can then be phagocytosed by other cells, including other macrophages [109].

Salmonella can modify macrophage polarization during chronic infection. Macrophages can differentiate into two groups after bacterial infection; the classically activated macrophages (M1 type) or the alternatively activated macrophages (M2 type). Cytokines are the primary determinant of macrophage polarization. The M1 type is proinflammatory and activates a Th1 immune response [110]. IFN γ - and LPS-induced activation of TLR4 signaling can shift the macrophage to the M1 phenotype. In contrast, the M2 type is antiinflammatory and activates the Th2 immune response [110]. The cytokine IL-4 shifts macrophages to the M2 phenotype. Usually, macrophages will exhibit M1 polarization after sensing the stimuli from bacteria or viruses. *Salmonella* phagocytized by the macrophage can shift the macrophage polarization state. Saliba et al. reported that macrophages harboring nongrowing *Salmonella* are prone to proinflammatory M1 polarization, but macrophages harboring growing bacteria shifted to an antiinflammatory M2-like state [111]. *S. typhimurium* preferentially lives in M2 macrophages during chronic infections [110]. Thus, *Salmonella* has mechanisms to shift the differentiation of macrophages into the M2 phenotype [110]. Intracellular glucose levels are higher in M2 macrophages, contributing to their permissiveness for the intracellular replication of *Salmonella* [112].

S. typhimurium persists within splenic granulomas enriched with CD11b⁺CD11c⁺Ly6C⁺ macrophages [4, 113]. Trung et al. previously reported that *Salmonella* can manipulate granuloma macrophage polarization towards the M2 phenotype [4]. As previously discussed, *S. typhimurium* preferentially persists in M2-reprogrammed macrophages. The bacterial effector SteE contributes to the establishment of persistent infection by downregulating tumor necrosis factor (TNF) signaling [4]. The bacteria have to develop strategies to overcome the immune response and persist chronically. *S. typhimurium* can polarize the primary macrophages to M2 polarization through the *e* SPI2 T3SS effector SteE. Macrophage M2 polarization can contribute to the systemic persistence of the bacteria [113].

Studies have shown that *Salmonella* can induce host cell death during infection [114]. Monack et al. found that caspase-1 is exploited by *Salmonella* to colonize the Peyer's patches (PPs) [115]. Systemic dissemination after an oral challenge with *Salmonella* is impaired in Casp-1^{-/-} mice. This indicates that caspase-1 is important for the systemic dissemination of *Salmonella* [115]. Caspase-1 (Casp-1), an interleukin [IL]-1 β -converting enzymes, can induce apoptosis in mammalian cells. Caspase 1 can cleave the proinflammatory cytokines IL-1 β and IL-18. Mice lacking

Casp-1 (Casp-1^{-/-} mice) showed a 1,000-fold higher lethal dose (LD50) of *S. typhimurium* in the mice than wide-type mice [115]. Casp-1^{-/-} mice were colonized by lower intracellular bacteria and did not show systemic dissemination of the bacteria, reduced colonization of bacteria in the Peyer's patches (PP) and spleens [115]. It suggests that Casp-1 is necessary for the establishment of systematic infection by *S. typhimurium* in mice [38, 67, 68, 115, 116]. *Salmonella* colonization was much higher in Caspase 11 deficient mice than in wild-type mice [68]. Casp1^{-/-} and Casp1/11^{-/-} monolayers showed significantly increased intracellular bacteria, accompanied by low intestinal epithelial cells (IECs) shedding and death [68]. Caspase activation is important for limiting the intracellular replication of *Salmonella*.

Inflammasome activation is one important pathway during the infection of *Salmonella* in the intestinal epithelial cells [38]. The infection of *Salmonella typhimurium* can also lead to the activation of Caspase 4, and Caspase 4 can limit the replication of *S. typhimurium* in the cells [117]. Activation of caspase 4 can lead to the noncanonical activation of the inflammasome pathway [117].

Salmonella can activate apoptosis of *Salmonella*-infected macrophages using effectors encoded in pathogenicity island-1 through both intrinsic and extrinsic pathways [118]. Cell death induced by the infected cells gives the bacteria an opportunity to be released and infect further cells. *Salmonella* can induce cell death in macrophages through several mechanisms. Immediate cell death can be induced by the type III secretion system (T3SS) of *Salmonella*. Or, the macrophages harboring *Salmonella* can be further phagocytosed by neighboring macrophages. Bacteria are released from dead cells and phagocytized by local macrophages, enabling another cycle of intracellular replication and cell-to-cell spread [114]. Ultimately, this cycle helps ensure the intracellular survival and persistent infection of phagocyte populations with *Salmonella*.

6. Chronic and Systemic Infection of *Salmonella Typhimurium*

Supershedders are the hosts responsible for the host-to-host transmission and reoccurrence of *S. typhimurium* since supershedders shed the bacteria in their feces. Foxp3⁺ Regulatory T cells play a role in the persistent infection of *Salmonella* [119]. Foxp3⁺ Treg ablation early after infection will accelerate bacterial eradication [119]. This indicated that immune regulatory T cells function in the early stages of infection to establish a persistent *Salmonella* infection [119].

Monack et al. demonstrated that *Salmonella* can persist in the MLNs of mice for up to one year. Macrophages in the MLNs can be the reservoirs of the bacteria. Voedisch et al. suggested that the MLN represents a restrictive site for the growth and dissemination of *Salmonella* [33]. In mice whose mesenteric lymph nodes have been surgically excised, the colonization of *Salmonella* in the liver and spleen is increased [33]. In such mice, *Salmonella* forms nonreplicating "persisters" in macrophages [120]. Persisters are in a state of dormant infection that is tolerant to drug treatment [121].

Indeed, they have resistance to antibiotics and can eventually reactivate and begin to replicate once more [122]. Persister cells are one important reason for relapsed infections. Persisters facilitate the chronic infection with *S. typhimurium*. Persisters can undermine the host immune response [123]. These persisters can reprogram the macrophages they dominate [123]. After exposure to ciprofloxacin, a fluoroquinolone antibiotic, *Salmonella enterica* persisters form unstable small colony variants. These phenotypes help the bacteria survive in the face of environmental stress or antibiotic treatments.

Salmonella persister cells are important components of biofilms [124]. Biofilm formation is an important strategy for persistent bacterial infections [125]. Forming biofilm can confer the bacteria survival advantages. Biofilm formation on gallstones is important for the chronic carriage of *Salmonella*. Antibiotic therapy efficiency is compromised in patients with a biofilm in the gall bladder. *Salmonella* infection in the gall bladder can induce the destruction of the epithelial cell integrity.

Biofilms are just one strategy for the bacteria to survive harsh environments. Even without animal reservoirs, biofilms can help *Salmonella* spp. to survive in the environment until uptake into a new host. However, the *Salmonella Typhimurium* ST313 strain which can cause blood stream infections and is typically seen in Sub Saharan Africa [126], has poor biofilm-forming ability and cannot survive long outside a host [127].

Except in antigen-presenting cells, *Salmonella* achieves a persistent infection in epithelial cells [128] by remaining in a dormant state. Luk et al. found that *Salmonella* can live in a dormant state in the vesicular compartment, different from the *Salmonella*-containing vacuoles (SCV). Contrary to macrophages, *Salmonella* in epithelial cells can express *Salmonella* Pathogenicity Island 2 (SPI-2) virulence factors. This report is the first to describe another persistent infection state and mechanism for *S. typhimurium* [128].

The *Salmonella* SPI2 effector SseI (also called SrfH) binds with host factor IQ motifs containing GTPase activating protein 1 (IQGAP1). SseI has been reported to mediate long-term systemic infections [60]. Pseudogenization of SseI leads to rapid systemic dissemination of *Salmonella typhimurium* through migratory dendritic cells [129]. In the sub-Saharan African *Salmonella typhimurium* strain ST313 lineage II, sseI is lost by pseudogenization. ST313 can disseminate from the gut to mesenteric lymph nodes (MLNs) via CD11b⁺ migratory dendritic cells (DCs) [129]. However, recovery of the gene function by expressing functional SseI in ST313 isolates reduces the dissemination of the bacteria [129].

The interplay between the host immune system and pathogens is a complex process during chronic infections. Dendritic cells and macrophages are important reservoirs for the bacteria that enable long-term survival. *Helicobacter pylori*, *Mycobacterium tuberculosis*, and *Salmonella enterica* all survive inside antigen-presenting cells (APCs). The gall bladder, bone marrow [130], and mesenteric lymph nodes are sites that can support persistent infection with *Salmonella*. Persistent infection with *Salmonella* can cause disease

in multiple organs, from gallbladder cancer to pancreatitis. Pancreatitis can be caused by persistent infection of mice with *Salmonella* [14]. Inflammatory, fibrotic, and epithelial responses can be detected in the pancreases of mice persistently infected with *S. typhimurium* [14]. Pancreatic acinar cells can be invaded by *S. typhimurium*.

Salmonella infections are associated with the development of IBD (inflammatory bowel diseases) and colon cancer [131, 132]. One study by Katrin et al. reported that mice with chronic infections with *S. typhimurium* develop severe and persistent intestinal fibrosis and have upregulation of several matrix metalloproteinases (MMPs) [133]. Transforming growth factor- β 1, insulin-like growth factor-I, and type I collagen deposition levels are increased during persistent infection of *S. typhimurium* [134, 135].

As shown in mouse models, chronic infection with *S. typhimurium* increases the susceptibility to intestinal inflammation [136]. The dDextran sulfate sodium (DSS)-induced colitis and interleukin (IL)-10^{-/-} spontaneous inflammation mice models were used in this particular study [137]. Because of persistent infection of *S. typhimurium* in the liver and spleen, these mice are more susceptible to intestinal inflammation. This indicated *S. typhimurium* persistent infection might be related to the accelerated onset of IBD (inflammatory bowel diseases) of the host [137].

Various studies support the mesenteric lymph nodes as a site that harbors *Salmonella* to sustain a chronic infection [138]. *Salmonella* can persist in the hemophagocytic macrophages of MLN. Removal of MLN increases the bacterial burdens in mice, however, indicating that another reservoir of *Salmonella* exists other than MLN [138]. Bacteria can be cultured from the liver tissue of chronically infected mice [139]. Liver macrophages are shifted to the M2 phenotype during persistent infection. An immune response balance exists during chronic infection with *Salmonella*, for example, the proinflammatory IFN γ and antiinflammatory signals IL-10. This balance allows the bacteria to survive in the persistent infection sites [139].

The cytokine Interleukin-22 (IL-22) can help the colonization of *Salmonella* by suppressing other commensal bacteria [140]. IL-22 can function in tissue repair and host defense; it is induced during pathogen infection. Behnsen et al. reported that IL-22 can suppress the intestinal microbiota [140]. IL-22 suppresses commensal *Enterobacteriaceae* and boosts the colonization of *Salmonella*. Binding of bacteria with APCs will induce the release of cytokine IL-23; IL-23 induces IL-17 and IL-22 release [141, 142]. In IL-22^{-/-} mice has higher *E. coli* burden and reduced *Salmonella* colonization in the intestine than wide type mice. IL-22 can induce the antimicrobial proteins lipocalin-2 and calprotectin release to inhibit the growth of commensal microbiota. This mechanism is exploited by *Salmonella* to outcompete intestinal microbiota [140].

7. Concluding Remarks

Achieving a better understanding of the pathogenesis of *Salmonella* will provide further insights into key host-pathogen interactions that affect persistent bacterial

infections. Understanding the detailed mechanisms and the specific host cell types involved in *Salmonella* infections may help guide the future development of therapeutic interventions. Understanding the mechanisms of *Salmonella* persistent infection will enable researchers to improve upon current treatment strategies, especially for asymptotically infected patients. Treating chronically infected patients will help reduce the reservoirs for the bacteria and limit the transmission of the disease.

Conflicts of Interest

The author declares that they have no conflicts of interest.

Acknowledgments

This study was funded by a local grant from the Tongji Hospital of the Huazhong University of Science and Technology.

References

- [1] P. Li, Q. Liu, H. Luo et al., "O-serotype conversion in salmonella typhimurium induces protective immune responses against invasive non-typhoidal Salmonella infections," *Frontiers in Immunology*, vol. 8, p. 1647, 2017.
- [2] S. Issenhuth-Jeanjean, P. Roggentin, M. Mikoleit et al., "Supplement 2008–2010 (no. 48) to the white-Kauffmann-le minor scheme," *Research in Microbiology*, vol. 165, no. 7, pp. 526–530, 2014.
- [3] M. E. M. S. Ohl and S. I. Miller, "Salmonella: a model for bacterial pathogenesis," *Annual Review of Medicine*, vol. 52, no. 1, pp. 259–274, 2001.
- [4] S. E. Majowicz, J. Musto, E. Scallan et al., "The global burden of nontyphoidal Salmonella gastroenteritis," *Clinical Infectious Diseases*, vol. 50, no. 6, pp. 882–889, 2010.
- [5] M. Kosek, C. Bern, and R. L. Guerrant, "The global burden of diarrhoeal disease, as estimated from studies published between 1992 and 2000," *Bulletin of the World Health Organization*, vol. 81, no. 3, pp. 197–204, 2003.
- [6] E. J. Threlfall, A. Bone, D. Murdoch, N. Banatvala, B. I. Shoismatulloev, and L. R. Ward, "Epidemic ciprofloxacin-resistant *Salmonella typhi* in Tajikistan," *The Lancet*, vol. 351, no. 9099, p. 339, 1998.
- [7] M. M. Levine, R. E. Black, and C. Lanata, "Precise estimation of the numbers of chronic carriers of *Salmonella typhi* in Santiago, Chile, an endemic area," *The Journal of Infectious Diseases*, vol. 146, pp. 724–726, 1982.
- [8] T. M. Vogelsang and J. Bøe, "Temporary and chronic carriers of *Salmonella typhi* and salmonella paratyphi B," *Journal of Hygiene*, vol. 46, no. 3, pp. 252–261, 1948.
- [9] G. Nath, Y. K. Singh, P. Maurya, A. K. Gulati, R. C. Srivastava, and S. K. Tripathi, "Does Salmonella Typhi primarily reside in the liver of chronic typhoid carriers?" *Journal of Infection in Developing Countries*, vol. 4, no. 4, pp. 259–261, 2010.
- [10] R. W. Crawford, R. Rosales-Reyes, L. Ramirez-Aguilar Mde, O. Chapa-Azuela, C. Alpuche-Aranda, and J. S. Gunn, "Gallstones play a significant role in Salmonella spp. gallbladder colonization and carriage," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 9, pp. 4353–4358, 2010.

- [11] H. Schiøler, E. D. Christiansen, G. Høybye, S. N. Rasmussen, and J. Greibe, "Biliary calculi in chronic salmonella carriers and healthy controls: a controlled study," *Scandinavian Journal of Infectious Diseases*, vol. 15, no. 1, pp. 17–19, 1983.
- [12] J. S. Gunn, J. M. Marshall, S. Baker, S. Dongol, R. C. Charles, and E. T. Ryan, "Salmonella chronic carriage: epidemiology, diagnosis, and gallbladder persistence," *Trends in Microbiology*, vol. 22, no. 11, pp. 648–655, 2014.
- [13] U. Dutta, P. K. Garg, R. Kumar, and R. K. Tandon, "Typhoid carriers among patients with gallstones are at increased risk for carcinoma of the gallbladder," *American Journal of Gastroenterology*, vol. 95, no. 3, pp. 784–787, 2000.
- [14] K. E. DelGiorno, J. W. Tam, J. C. Hall, G. Thotakura, H. C. Crawford, and A. W. M. van der Velden, "Persistent salmonellosis causes pancreatitis in a murine model of infection," *PLoS One*, vol. 9, no. 4, Article ID e92807, 2014.
- [15] M. R. Wilmes-Riesenberg, B. Bearson, R. Curtis, and R. CuRtis, "Role of the acid tolerance response in virulence of *Salmonella typhimurium*," *Infection and Immunity*, vol. 64, no. 4, pp. 1085–1092, 1996.
- [16] S. I. K. A. Miller, A. M. Kukral, and J. J. Mekalanos, "A two-component regulatory system (phoP phoQ) controls *Salmonella typhimurium* virulence," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 86, no. 13, pp. 5054–5058, 1989.
- [17] F. Bb, "Molecular and cellular mechanisms of *Salmonella* pathogenesis," *Current Topics in Microbiology and Immunology*, vol. 192, pp. 163–185, 1994.
- [18] C. M. A. Khan, "The dynamic interactions between *Salmonella* and the microbiota, within the challenging niche of the gastrointestinal tract," *International Scholarly Research Notices*, vol. 2014, Article ID 846049, 23 pages, 2014.
- [19] H. D. Que Ju and D. J. Hentges, "Effect of streptomycin administration on colonization resistance to *Salmonella typhimurium* in mice," *Infection and Immunity*, vol. 48, no. 1, pp. 169–174, 1985.
- [20] M. Barthel, S. Hapfelmeier, L. Quintanilla-Martinez et al., "Pretreatment of mice with streptomycin provides a *Salmonella enterica* serovar typhimurium colitis model that allows analysis of both pathogen and host," *Infection and Immunity*, vol. 71, no. 5, pp. 2839–2858, 2003.
- [21] M. K. Sandra Y Wotzka, M. Kreuzer, L. Maier et al., "*Escherichia coli* limits salmonella typhimurium infections after diet shifts and fat-mediated microbiota perturbation in mice," *Nat Microbiol*, vol. 4, no. 12, pp. 2164–2174, 2019.
- [22] N. J. A. Jabado, A. Jankowski, V. Picard, S. Grinstein, P. Gros, and P. Gros, "Natural resistance to intracellular infections: natural resistance-associated macrophage protein 1 (Nramp1) functions as a pH-dependent manganese transporter at the phagosomal membrane," *Journal of Experimental Medicine*, vol. 192, no. 9, pp. 1237–1248, 2000.
- [23] B. Stecher, G. Paesold, M. Barthel et al., "Chronic *Salmonella enterica* serovar typhimurium-induced colitis and cholangitis in streptomycin-pretreated nramp1+/+ mice," *Infection and Immunity*, vol. 74, no. 9, pp. 5047–5057, 2006.
- [24] B. D. Jones, N. Ghorri, and S. Falkow, "*Salmonella typhimurium* initiates murine infection by penetrating and destroying the specialized epithelial M cells of the peyer's patches," *Journal of Experimental Medicine*, vol. 180, no. 1, pp. 15–23, 1994.
- [25] J. P. Kraehenbuhl and M. R. Neutra, "Epithelial M cells: differentiation and function," *Annual Review of Cell and Developmental Biology*, vol. 16, no. 1, pp. 301–332, 2000.
- [26] S. C. Corr, C. C. Gahan, and C. Hill, "M-cells: origin, morphology and role in mucosal immunity and microbial pathogenesis," *FEMS Immunology and Medical Microbiology*, vol. 52, no. 1, pp. 2–12, 2008.
- [27] A. Tahoun, S. Mahajan, E. Paxton et al., "Salmonella transforms follicle-associated epithelial cells into M cells to promote intestinal invasion," *Cell Host & Microbe*, vol. 12, no. 5, pp. 645–656, 2012.
- [28] K. P. K. Schneider, K. G. Potter, and C. F. Ware, "Lymphotoxin and LIGHT signaling pathways and target genes," *Immunological Reviews*, vol. 202, no. 1, pp. 49–66, 2004.
- [29] S. Ding, Y. Song, K. F. Brulois et al., "Retinoic acid and lymphotoxin signaling promote differentiation of human intestinal M cells," *Gastroenterology*, vol. 159, no. 1, pp. 214–226, 2020.
- [30] D. A. Giles, S. Zahner, P. Krause et al., "The tumor necrosis factor superfamily members TNFSF14 (LIGHT), lymphotoxin beta and lymphotoxin beta receptor interact to regulate intestinal inflammation," *Frontiers in Immunology*, vol. 9, p. 2585, 2018.
- [31] J. Sudhamsu, J. Yin, E. Y. Chiang, M. A. Starovasnik, J. L. Grogan, and S. G. Hymowitz, "Dimerization of LT β R by LT α_1 β_2 is necessary and sufficient for signal transduction," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 110, no. 49, pp. 19896–19901, 2013.
- [32] A. M. K. Fütterer, K. Mink, M. H. Kosco-Vilbois, K. Pfeffer, and K. Pfeffer, "The lymphotoxin beta receptor controls organogenesis and affinity maturation in peripheral lymphoid tissues," *Immunity*, vol. 9, no. 1, pp. 59–70, 1998.
- [33] S. Voedisch, C. Koenecke, S. David et al., "Mesenteric lymph nodes confine dendritic cell-mediated dissemination of *Salmonella enterica* serovar typhimurium and limit systemic disease in mice," *Infection and Immunity*, vol. 77, no. 8, pp. 3170–3180, 2009.
- [34] C. Cheminay, M. Schoen, M. Hensel et al., "Migration of *Salmonella typhimurium* -harboring bone marrow-derived dendritic cells towards the chemokines CCL19 and CCL21," *Microbial Pathogenesis*, vol. 32, no. 5, pp. 207–218, 2002.
- [35] M. J. Worley, G. S. Nieman, K. Geddes, and F. Heffron, "*Salmonella typhimurium* disseminates within its host by manipulating the motility of infected cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 47, pp. 17915–17920, 2006.
- [36] J. A. Wroblewska, Y. Zhang, H. Tang, X. Guo, C. Nagler, and Y. X. Fu, "Cutting edge: lymphotoxin signaling is essential for clearance of *Salmonella* from the gut lumen and generation of anti-*Salmonella* protective immunity," *The Journal of Immunology*, vol. 198, no. 1, pp. 55–60, 2017.
- [37] G. Gonzalez-Escobedo and J. S. Gunn, "Gallbladder epithelium as a niche for chronic *Salmonella* carriage," *Infection and Immunity*, vol. 81, no. 8, pp. 2920–2930, 2013.
- [38] L. A. C. S. Knodler, S. Crowley, H. Yang et al., "Noncanonical inflammasome activation of caspase-4/caspase-11 mediates epithelial defenses against enteric bacterial pathogens," *Cell Host & Microbe*, vol. 16, no. 2, pp. 249–256, 2014.
- [39] L. A. Knodler, B. A. Vallance, J. Celli et al., "Dissemination of invasive salmonella via bacterial-induced extrusion of mucosal epithelia," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 41, pp. 17733–17738, 2010.
- [40] L. Lou, P. Zhang, R. Piao, and Y. Wang, "Salmonella pathogenicity island 1 (SPI-1) and its complex regulatory network," *Frontiers in Cellular and Infection Microbiology*, vol. 9, p. 270, 2019.

- [41] A. J. Baumler, K. Zhang, A. Riba et al., "Minimal SPI1-T3SS effector requirement for salmonella enterocyte invasion and intracellular proliferation in vivo," *PLoS Pathogens*, vol. 14, no. 3, Article ID e1006925, 2018.
- [42] M. A. Bakowski, V. Braun, and J. H. Brummell, "Salmonella-containing vacuoles: directing traffic and nesting to grow," *Traffic*, vol. 9, no. 12, pp. 2022–2031, 2008.
- [43] J. C. D. Jantsch, D. Chikkaballi, and M. Hensel, "Cellular aspects of immunity to intracellular *Salmonella enterica*," *Immunological Reviews*, vol. 240, no. 1, pp. 185–195, 2011.
- [44] P. Malik-Kale, C. E. Jolly, S. Lathrop, S. Winfree, C. Luterbach, and O. Steele-Mortimer, "Salmonella—at home in the host cell," *Frontiers in Microbiology*, vol. 2, p. 125, 2011.
- [45] A. L. Swart and M. Hensel, "Interactions of *Salmonella enterica* with dendritic cells," *Virulence*, vol. 3, pp. 660–667, 2012.
- [46] J. Guignot, E. Caron, C. Beuzon et al., "Microtubule motors control membrane dynamics of salmonella-containing vacuoles," *Journal of Cell Science*, vol. 117, no. 7, pp. 1033–1045, 2004.
- [47] A. Takeuchi, "Electron microscope studies of experimental salmonella infection. I. penetration into the intestinal epithelium by *Salmonella typhimurium*," *American Journal of Pathology*, vol. 50, no. 1, pp. 109–136, 1967.
- [48] A. J. T. R. Bäuml, R. M. Tsohis, and F. Heffron, "Contribution of fimbrial operons to attachment to and invasion of epithelial cell lines by *Salmonella typhimurium*," *Infection and Immunity*, vol. 64, pp. 1862–1865, 1996.
- [49] P. Cossart and P. J. Sansonetti, "Bacterial invasion: the paradigms of enteroinvasive pathogens," *Science*, vol. 304, no. 5668, pp. 242–248, 2004.
- [50] J. Hanisch, T. E. Stradal, and K. Rottner, "A novel contractility pathway operating in salmonella invasion," *Virulence*, vol. 3, no. 1, pp. 81–86, 2012.
- [51] J. C. Patel and J. E. Galan, "Manipulation of the host actin cytoskeleton by *Salmonella*-all in the name of entry," *Current Opinion in Microbiology*, vol. 8, no. 1, pp. 10–15, 2005.
- [52] M. Rosselin, I. Virlogeux-Payant, C. Roy et al., "Rck of *Salmonella enterica*, subspecies enterica serovar Enteritidis, mediates Zipper-like internalization," *Cell Research*, vol. 20, no. 6, pp. 647–664, 2010.
- [53] M. A. Lambert and S. G. J. Smith, "The PagN protein mediates invasion via interaction with proteoglycan," *FEMS Microbiology Letters*, vol. 297, no. 2, pp. 209–216, 2009.
- [54] E. J. W. L. Heffernan, L. Wu, S. Okamoto, J. Fierer, D. G. Guiney, and D. G. Guiney, "Specificity of the complement resistance and cell association phenotypes encoded by the outer membrane protein genes rck from *Salmonella typhimurium* and ail from *Yersinia enterocolitica*," *Infection and Immunity*, vol. 62, no. 11, pp. 5183–5186, 1994.
- [55] M. A. Lambert and S. G. Smith, "The PagN protein of *Salmonella enterica* serovar typhimurium is an adhesin and invasin," *BMC Microbiology*, vol. 8, no. 1, p. 142, 2008.
- [56] D. Kaur and A. Mukhopadhyaya, "Outer membrane protein OmpV mediates salmonella entericaserovar typhimurium adhesion to intestinal epithelial cells via fibronectin and $\alpha 1\beta 1$ integrin," *Cellular Microbiology*, vol. 22, no. 5, Article ID e13172, 2020.
- [57] K. Takeda, T. Kaisho, and S. Akira, "Toll-like receptors," *Annual Review of Immunology*, vol. 21, no. 1, pp. 335–376, 2003.
- [58] S. Hapfelmeier, B. Stecher, M. Barthel et al., "The *Salmonella* pathogenicity island (SPI)-2 and SPI-1 type III secretion systems allow *Salmonella* serovar typhimurium to trigger colitis via MyD88-dependent and MyD88-independent mechanisms," *The Journal of Immunology*, vol. 174, no. 3, pp. 1675–1685, 2005.
- [59] E. Jennings, T. L. M. Thurston, and D. W. Holden, "Salmonella SPI-2 type III secretion system effectors: molecular mechanisms and physiological consequences," *Cell Host & Microbe*, vol. 22, no. 2, pp. 217–231, 2017.
- [60] M. Hensel, L. M. McLaughlin, G. R. Govoni et al., "The *Salmonella* SPI2 effector SseI mediates long-term systemic infection by modulating host cell migration," *PLoS Pathogens*, vol. 5, no. 11, Article ID e1000671, 2009.
- [61] M. A. Jepson, B. Kenny, and A. D. Leard, "Role of sipA in the early stages of *Salmonella typhimurium* entry into epithelial cells," *Cellular Microbiology*, vol. 3, no. 6, pp. 417–426, 2001.
- [62] D. Zhou, M. S. Mooseker, and J. E. Galán, "Role of the *S. typhimurium* actin-binding protein SipA in bacterial internalization," *Science*, vol. 283, no. 5410, pp. 2092–2095, Article ID 101126, 1999.
- [63] M. Raffatellu, R. P. Wilson, D. Chessa et al., "SipA, SopA, SopB, SopD, and SopE2 contribute to *Salmonella enterica* serotype typhimurium invasion of epithelial cells," *Infection and Immunity*, vol. 73, no. 1, pp. 146–154, 2005.
- [64] S. A. Fattinger, M. E. Sellin, and W. D. Hardt, "Epithelial inflammasomes in the defense against *Salmonella* gut infection," *Current Opinion in Microbiology*, vol. 59, pp. 86–94, 2021.
- [65] J. E. Galan, "Salmonella typhimurium and inflammation: a pathogen-centric affair," *Nature Reviews Microbiology*, vol. 19, no. 11, pp. 716–725, 2021.
- [66] V. Feuillet, S. Medjane, I. Mondor et al., "Involvement of Toll-like receptor 5 in the recognition of flagellated bacteria," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, pp. 12487–12492, 2006.
- [67] A. M. Gram, J. A. Wright, R. J. Pickering et al., "Salmonella flagellin activates NAIP/NLRC4 and canonical NLRP3 inflammasomes in human macrophages," *The Journal of Immunology*, vol. 206, no. 3, pp. 631–640, 2021.
- [68] S. M. Crowley, X. Han, J. M. Allaire et al., "Intestinal restriction of *Salmonella Typhimurium* requires caspase-1 and caspase-11 epithelial intrinsic inflammasomes," *PLoS Pathogens*, vol. 16, no. 4, Article ID e1008498, 2020.
- [69] A. K. B. Iwasaki and B. L. Kelsall, "Unique functions of CD11b⁺, CD8 α ⁺, and double-negative Peyer's patch dendritic cells," *The Journal of Immunology*, vol. 166, no. 8, pp. 4884–4890, 2001.
- [70] I. H. P. Maric, P. G. Holt, J. Bienenstock, and J. Bienenstock, "Class II MHC antigen (Ia)-bearing dendritic cells in the epithelium of the rat intestine," *The Journal of Immunology*, vol. 156, no. 4, pp. 1408–1414, 1996.
- [71] C. H. S. Ruedl and S. Hubele, "Maturation of Peyer's patch dendritic cells in vitro upon stimulation via cytokines or CD40 triggering," *European Journal of Immunology*, vol. 27, no. 6, pp. 1325–1330, 1997.
- [72] B. L. S. W. Kelsall and W. Strober, "Distinct populations of dendritic cells are present in the subepithelial dome and T cell regions of the murine Peyer's patch," *Journal of Experimental Medicine*, vol. 183, no. 1, pp. 237–247, 1996.
- [73] A. K. B. Iwasaki and B. L. Kelsall, "Localization of distinct Peyer's patch dendritic cell subsets and their recruitment by chemokines macrophage inflammatory protein (Mip)-3 α , mip-3 β , and secondary lymphoid organ chemokine," *Journal of Experimental Medicine*, vol. 191, no. 8, pp. 1381–1394, 2000.

- [74] C. W. Cutler, R. Jotwani, and B. Pulendran, "Dendritic cells: immune saviors or Achilles' heel?" *Infection and Immunity*, vol. 69, pp. 4703–4708, 2001.
- [75] S. Tuladhar and T. D. Kanneganti, "NLRP12 in innate immunity and inflammation," *Molecular Aspects of Medicine*, vol. 76, Article ID 100887, 2020.
- [76] C.-L. Ye, L. Qiao, X. Li et al., "*Salmonella typhimurium* interacts CD209 receptors to promote host dissemination and infection," *Infection and Immunity*, vol. 87, 2019.
- [77] T. B. Geijtenbeek, D. S. Kwon, R. Torensma et al., "DC-SIGN, a dendritic cell-specific HIV-1-Binding protein that enhances trans-infection of T cells," *Cell*, vol. 100, no. 5, pp. 587–597, 2000.
- [78] T. B. H. Geijtenbeek and Y. van Kooyk, "Pathogens target DC-SIGN to influence their fate DC-SIGN functions as a pathogen receptor with broad specificity," *Acta Pathologica, Microbiologica et Immunologica Scandinavica*, vol. 111, no. 7-8, pp. 698–714, 2003.
- [79] D. McDonald, L. Wu, S. M. Bohks, V. N. KewalRamani, D. Unutmaz, and T. J. Hope, "Recruitment of HIV and its receptors to dendritic cell-T cell junctions," *Science*, vol. 300, no. 5623, pp. 1295–1297, 2003.
- [80] L. Tailleux, N. Pham-Thi, A. Bergeron-Lafaurie et al., "DC-SIGN induction in alveolar macrophages defines privileged target host cells for mycobacteria in patients with tuberculosis," *PLoS Medicine*, vol. 2, no. 12, 2005.
- [81] Y. X. He, C. L. Ye, P. Zhang et al., "Yersinia pseudotuberculosis exploits CD209 receptors for promoting host dissemination and infection," *Infection and Immunity*, vol. 87, no. 1, Article ID e00654-18, 2019.
- [82] J. Klena, P. Zhang, O. Schwartz, S. Hull, and T. Chen, "The core lipopolysaccharide of *Escherichia coli* is a ligand for the dendritic-cell-specific intercellular adhesion molecule non-integrin CD209 receptor," *Journal of Bacteriology*, vol. 187, pp. 1710–1715, 2005.
- [83] K. Yang, Y. He, C. G. Park et al., "Yersinia pestis interacts with SIGNR1 (CD209b) for promoting host dissemination and infection," *Frontiers in Immunology*, vol. 10, p. 96, 2019.
- [84] K. Yang, C. G. Park, C. Cheong et al., "Host langerin (CD207) is a receptor for *Yersinia pestis* phagocytosis and promotes dissemination," *Immunology & Cell Biology*, vol. 93, no. 9, pp. 815–824, 2015.
- [85] P. Zhang, O. Schwartz, M. Pantelic et al., "Dc-SIGN (DC-SIGN (CD209) recognition of Neisseria gonorrhoeae is circumvented by lipooligosaccharide variation," *Journal of Leukocyte Biology*, vol. 79, no. 4, pp. 731–738, 2006.
- [86] P. Zhang, M. Skurnik, S. S. Zhang et al., "Human dendritic cell-specific intercellular adhesion molecule-grabbing non-integrin (CD209) is a receptor for *Yersinia pestis* that promotes phagocytosis by dendritic cells," *Infection and Immunity*, vol. 76, no. 5, pp. 2070–2079, 2008.
- [87] P. Zhang, S. Snyder, P. Feng et al., "Role of N-acetylglucosamine within core lipopolysaccharide of several species of gram-negative bacteria in targeting the DC-SIGN (CD209)," *The Journal of Immunology*, vol. 177, no. 6, pp. 4002–4011, 2006.
- [88] S. S. Zhang, C. G. Park, P. Zhang et al., "Plasminogen activator Pla of *Yersinia pestis* utilizes murine DEC-205 (CD205) as a receptor to promote dissemination," *Journal of Biological Chemistry*, vol. 283, no. 46, pp. 31511–31521, 2008.
- [89] C. Cheminay, A. Möhlenbrink, and M. Hensel, "Intracellular *Salmonella* Inhibit antigen presentation by dendritic cells," *The Journal of Immunology*, vol. 174, no. 5, pp. 2892–2899, 2005.
- [90] N. Lapaque, J. L. Hutchinson, D. C. Jones et al., "Salmonella regulates polyubiquitination and surface expression of MHC class II antigens," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 33, pp. 14052–14057, 2009.
- [91] J. Farache, I. Koren, I. Milo et al., "Luminal bacteria recruit CD103+ dendritic cells into the intestinal epithelium to sample bacterial antigens for presentation," *Immunity*, vol. 38, no. 3, pp. 581–595, 2013.
- [92] S. Uematsu, M. H. Jang, N. Chevrier et al., "Detection of pathogenic intestinal bacteria by Toll-like receptor 5 on intestinal CD11c+ lamina propria cells," *Nature Immunology*, vol. 7, no. 8, pp. 868–874, 2006.
- [93] S. Uematsu and S. Akira, "Immune responses of TLR5 (+) lamina propria dendritic cells in enterobacterial infection," *Journal of Gastroenterology*, vol. 44, no. 8, pp. 803–811, 2009.
- [94] M. Rescigno, "CCR6 (+) dendritic cells: the gut tactical-response unit," *Immunity*, vol. 24, no. 5, pp. 508–510, 2006.
- [95] R. M. Salazar-Gonzalez, J. H. Niess, D. J. Zammit et al., "CCR6-mediated dendritic cell activation of pathogen-specific T cells in Peyer's patches," *Immunity*, vol. 24, no. 5, pp. 623–632, 2006.
- [96] J. H. B. S. Niess, S. Brand, L. Landsman et al., "CX3CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance," *Science*, vol. 307, no. 5707, pp. 254–258, 2005.
- [97] S. A. N. F. Hopkins, I. E. Corthesy-Theulaz, and J. P. Kraehenbuhl, "A recombinant *Salmonella typhimurium* vaccine strain is taken up and survives within murine Peyer's patch dendritic cells," *Cellular Microbiology*, vol. 2, no. 1, pp. 59–68, 2000.
- [98] J. S. Gunn, S. I. Miller, and S. Miller, "PhoP-PhoQ activates transcription of pmrAB, encoding a two-component regulatory system involved in *Salmonella typhimurium* antimicrobial peptide resistance," *Journal of Bacteriology*, vol. 178, no. 23, pp. 6857–6864, 1996.
- [99] M. Rescigno, M. Urbano, B. Valzasina et al., "Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to sample bacteria," *Nature Immunology*, vol. 2, no. 4, pp. 361–367, 2001.
- [100] A. J.-C. J. Vazquez-Torres, J. Jones-Carson, A. J. Bäumlner et al., "Extraintestinal dissemination of Salmonella by CD18-expressing phagocytes," *Nature*, vol. 401, no. 6755, pp. 804–808, 1999.
- [101] K. I. Uchiya and T. Nikai, "Salmonella pathogenicity island 2-dependent expression of suppressor of cytokine signaling 3 in macrophages," *Infection and Immunity*, vol. 73, no. 9, pp. 5587–5594, 2005.
- [102] O. L. C. Wijburg, C. P. Simmons, R. A. Strugnell, and R. A. Strugnell, "Dual role for macrophages in vivo in pathogenesis and control of murine *Salmonella enterica* var. typhimurium infections," *European Journal of Immunology*, vol. 30, no. 3, pp. 944–953, 2000.
- [103] C. M. R. E. Alpuche-Aranda, E. L. Racoosin, S. I. Miller, and S. I. Miller, "Salmonella stimulate macrophage macrophagocytosis and persist within spacious phagosomes," *Journal of Experimental Medicine*, vol. 179, no. 2, pp. 601–608, 1994.
- [104] A. Vazquez-Torres, Y. Xu, J. Jones-Carson et al., "Salmonella pathogenicity island 2-dependent evasion of the phagocyte NADPH oxidase," *Science*, vol. 287, no. 5458, pp. 1655–1658, 2000.
- [105] M. Gogoi, M. M. Shreenivas, and D. Chakravorty, "Hoodwinking the big-eater to prosper: the salmonella-

- macrophage paradigm," *Journal of Innate Immunity*, vol. 11, no. 3, pp. 289–299, 2019.
- [106] C. M. S. J. Alpuche Aranda, J. A. Swanson, S. I. Miller, and S. I. Miller, "Salmonella typhimurium activates virulence gene transcription within acidified macrophage phagosomes," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 89, no. 21, pp. 10079–10083, 1992.
- [107] S. L. S. Eriksson, S. Lucchini, M. Rhen, J. C. Hinton, and J. C. D. Hinton, "Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*," *Molecular Microbiology*, vol. 47, no. 1, pp. 103–118, 2003.
- [108] K. Lähteenmäki, P. Kyllönen, L. Partanen, and T. K. Korhonen, "Antiprotease inactivation by *Salmonella enterica* released from infected macrophages," *Cellular Microbiology*, vol. 7, no. 4, pp. 529–538, 2004.
- [109] D. M. Monack, A. Mueller, and S. Falkow, "Persistent bacterial infections: the interface of the pathogen and the host immune system," *Nature Reviews Microbiology*, vol. 2, no. 9, pp. 747–765, 2004.
- [110] N. A. Eisele, T. Ruby, A. Jacobson et al., "Salmonella require the fatty acid regulator PPAR δ for the establishment of a metabolic environment essential for long-term persistence," *Cell Host & Microbe*, vol. 14, no. 2, pp. 171–182, 2013.
- [111] A. E. Saliba, L. Li, A. J. Westermann et al., "Single-cell RNA-seq ties macrophage polarization to growth rate of intracellular *Salmonella*," *Nature Microbiology*, vol. 2, Article ID 16206, 2016.
- [112] R. M. Roop and C. C. Caswell, "Bacterial persistence: finding the "sweet spot"," *Cell Host & Microbe*, vol. 14, no. 2, pp. 119–120, 2013.
- [113] T. H. M. Pham, S. M. Brewer, T. Thurston et al., "Salmonella-driven polarization of granuloma macrophages antagonizes TNF-mediated pathogen restriction during persistent infection," *Cell Host & Microbe*, vol. 27, no. 1, pp. 54–67, 2020.
- [114] D. G. Guiney, "The role of host cell death in *Salmonella* infections," *Current Topics in Microbiology and Immunology*, vol. 289, 2005.
- [115] D. M. H. D. Monack, D. Hersh, D. Bouley, A. Zychlinsky, S. Falkow, and S. Falkow, "Salmonella exploits caspase-1 to colonize Peyer's patches in a murine typhoid model," *Journal of Experimental Medicine*, vol. 192, no. 2, pp. 249–258, 2000.
- [116] E. A. Miao and J. V. Rajan, "Salmonella and caspase-1: a complex interplay of detection and evasion," *Frontiers in Microbiology*, vol. 2, p. 85, 2011.
- [117] M. K. Holly, X. Han, E. J. Zhao et al., "*Salmonella enterica* infection of murine and human enteroid-derived monolayers elicits differential activation of epithelium-intrinsic inflammasomes," *Infection and Immunity*, vol. 88, no. 7, 2020.
- [118] H. H. Lin, H. L. Chen, C. C. Weng, R. P. Janapatla, C. L. Chen, and C. H. Chiu, "Activation of apoptosis by salmonella pathogenicity island-1 effectors through both intrinsic and extrinsic pathways in *Salmonella*-infected macrophages," *Journal of Microbiology, Immunology, and Infection*, vol. 54, no. 4, pp. 616–626, 2021.
- [119] T. M. Johanns, J. M. Ertelt, J. H. Rowe, and S. S. Way, "Regulatory T cell suppressive potency dictates the balance between bacterial proliferation and clearance during persistent *Salmonella* infection," *PLoS Pathogens*, vol. 6, no. 8, Article ID e1001043, 2010.
- [120] S. Helaine, A. M. Cheverton, K. G. Watson, L. M. Faure, S. A. Matthews, and D. W. Holden, "Internalization of *Salmonella* by macrophages induces formation of non-replicating persisters," *Science*, vol. 343, no. 6167, pp. 204–208, 2014.
- [121] K. Lewis, "Persister cells," *Annual Review of Microbiology*, vol. 64, no. 1, pp. 357–372, 2010.
- [122] R. A. Fisher, B. Gollan, and S. Helaine, "Persistent bacterial infections and persister cells," *Nature Reviews Microbiology*, vol. 15, no. 8, pp. 453–464, 2017.
- [123] D. A. C. Stapels, P. W. S. Hill, A. J. Westermann et al., "Salmonella persisters undermine host immune defenses during antibiotic treatment," *Science*, vol. 362, no. 6419, pp. 1156–1160, 2018.
- [124] M. Abdallah, C. Benoliel, D. Drider, P. Dhulster, and N. E. Chihib, "Biofilm formation and persistence on abiotic surfaces in the context of food and medical environments," *Archives of Microbiology*, vol. 196, no. 7, pp. 453–472, 2014.
- [125] U. Romling and C. Balsalobre, "Biofilm infections, their resilience to therapy and innovative treatment strategies," *Journal of Internal Medicine*, vol. 272, no. 6, pp. 541–561, 2012.
- [126] N. A. Feasey, G. Dougan, R. A. Kingsley, R. S. Heyderman, and M. A. Gordon, "Invasive non-typhoidal salmonella disease: an emerging and neglected tropical disease in Africa," *The Lancet*, vol. 379, no. 9835, pp. 2489–2499, 2012.
- [127] G. Ramachandran, K. Aheto, M. E. Shirliff, S. M. Tennant, and J. Kaper, "Poor biofilm-forming ability and long-term survival of invasive *Salmonella* Typhimurium ST313," *Pathogens and Disease*, vol. 74, no. 5, Article ID ftw049, 2016.
- [128] C. H. Luk, C. Valenzuela, M. Gil et al., "Salmonella enters a dormant state within human epithelial cells for persistent infection," *PLoS Pathogens*, vol. 17, no. 4, Article ID e1009550, 2021.
- [129] S. E. Carden, G. T. Walker, J. Honeycutt et al., "Pseudogenization of the secreted effector gene sseI confers rapid systemic dissemination of *S. Typhimurium* ST313 within migratory dendritic cells," *Cell Host & Microbe*, vol. 21, no. 2, pp. 182–194, 2017.
- [130] F. S. J. Niedergang, J. C. Sirard, J. P. Kraehenbuhl, and J. P. Kraehenbuhl, "Entry and survival of *Salmonella typhimurium* in dendritic cells and presentation of recombinant antigens do not require macrophage-specific virulence factors," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 26, pp. 14650–14655, 2000.
- [131] L. Mughini-Gras, M. Schaapveld, J. Kramers et al., "Increased colon cancer risk after severe salmonella infection," *PLoS One*, vol. 13, no. 1, Article ID e0189721, 2018.
- [132] R. B. M. Lu, M. Bosland, Y. G. Zhang, I. Kato, J. Sun, and J. Sun, "Presence of *Salmonella* AvrA in colorectal tumor and its precursor lesions in mouse intestine and human specimens," *Oncotarget*, vol. 8, no. 33, pp. 55104–55115, 2017.
- [133] K. Ehrhardt, N. Steck, R. Kappelhoff et al., "Persistent *Salmonella enterica* serovar typhimurium infection induces protease expression during intestinal fibrosis," *Inflammatory Bowel Diseases*, vol. 25, no. 10, pp. 1629–1643, 2019.
- [134] G. A. Grassl, Y. Valdez, K. S. Bergstrom, B. A. Vallance, and B. B. Finlay, "Chronic enteric salmonella infection in mice leads to severe and persistent intestinal fibrosis," *Gastroenterology*, vol. 134, no. 3, pp. 768–780, 2008.
- [135] L. E. Månsson, M. Montero, M. Zarepour et al., "MyD88 signaling promotes both mucosal homeostatic and fibrotic responses during *Salmonella*-induced colitis," *American Journal of Physiology—Gastrointestinal and Liver Physiology*, vol. 303, no. 3, pp. G311–G323, 2012.

- [136] B. M. Schultz, C. A. Paduro, G. A. Salazar et al., "A potential role of salmonella infection in the onset of inflammatory bowel diseases," *Frontiers in Immunology*, vol. 8, p. 191, 2017.
- [137] B. M. Schultz, G. A. Salazar, C. A. Paduro et al., "Persistent *Salmonella enterica* serovar typhimurium infection increases the susceptibility of mice to develop intestinal inflammation," *Frontiers in Immunology*, vol. 9, p. 1166, 2018.
- [138] A. J. Griffin, L. X. Li, S. Voedisch, O. Pabst, and S. J. McSorley, "Dissemination of persistent intestinal bacteria via the mesenteric lymph nodes causes typhoid relapse," *Infection and Immunity*, vol. 79, no. 4, pp. 1479–1488, 2011.
- [139] J. R. Kurtz, W. Nieves, D. Bauer et al., "Salmonella persistence and host immunity is dictated by the anatomical microenvironment," *Infect Immun*, vol. 88, 2020.
- [140] J. Behnsen, S. Jellbauer, C. P. Wong et al., "The cytokine IL-22 promotes pathogen colonization by suppressing related commensal bacteria," *Immunity*, vol. 40, no. 2, pp. 262–273, 2014.
- [141] C. Blaschitz and M. Raffatellu, "Th17 cytokines and the gut mucosal barrier," *Journal of Clinical Immunology*, vol. 30, no. 2, pp. 196–203, 2010.
- [142] S. A. Khader and R. Gopal, "IL-17 in protective immunity to intracellular pathogens," *Virulence*, vol. 1, no. 5, pp. 423–427, 2010.