



Research Article

Clinical and Prognostic Value of PET/CT Imaging with Combination of ^{68}Ga -DOTATATE and ^{18}F -FDG in Gastroenteropancreatic Neuroendocrine Neoplasms

Panpan Zhang,¹ Jiangyuan Yu,² Jie Li,¹ Lin Shen ,¹ Nan Li,² Hua Zhu,² Shizhen Zhai,² Yan Zhang,² Zhi Yang ,² and Ming Lu ,¹

¹Department of Gastrointestinal Oncology, Key Laboratory of Carcinogenesis and Translational Research, Ministry of Education, Peking University School of Oncology, Beijing Cancer Hospital and Institute, Beijing, China

²Department of Nuclear Medicine, Key Laboratory of Carcinogenesis and Translational Research, Ministry of Education, Peking University School of Oncology, Beijing Cancer Hospital and Institute, Beijing, China

Correspondence should be addressed to Zhi Yang; pekyz@163.com and Ming Lu; qiminglu.mail@126.com

Received 25 July 2017; Revised 22 October 2017; Accepted 16 January 2018; Published 26 February 2018

Academic Editor: Chongzhao Ran

Copyright © 2018 Panpan Zhang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. To evaluate the clinical and prognostic value of PET/CT with combination of ^{68}Ga -DOTATATE and ^{18}F -FDG in gastroenteropancreatic neuroendocrine neoplasms (GEP-NENs). **Method.** 83 patients of GEP-NENs who underwent ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT were enrolled between June 2013 and December 2016. Well-differentiated (WD) NETs are divided into group A ($\text{Ki-67} < 10\%$) and group B ($\text{Ki-67} \geq 10\%$), and poorly differentiated (PD) NECs are defined as group C. The relationship between PET/CT results and clinicopathological characteristics was retrospectively investigated. **Result.** For groups A/B/C, the sensitivities of ^{68}Ga -DOTATATE and ^{18}F -FDG were 78.8%/83.3%/37.5% and 52.0%/72.2%/100.0%. A negative correlation between Ki-67 and SUV_{\max} of ^{68}Ga -DOTATATE ($R = -0.415$; $P \leq 0.001$) was observed, while a positive correlation was noted between Ki-67 and SUV_{\max} of ^{18}F -FDG ($R = 0.683$; $P \leq 0.001$). 62.5% (5/8) of patients showed significantly more lesions in the bone if ^{68}Ga -DOTATATE was used, and 22.7% (5/22) of patients showed more lymph node metastases if ^{18}F -FDG was used. **Conclusions.** The sensitivity of dual tracers was correlated with cell differentiation, and a correlation between Ki-67 and both SUV_{\max} of PET-CTs could be observed. ^{68}Ga -DOTATATE is suggested for WD-NET and ^{18}F -FDG is probably suitable for patients with $\text{Ki-67} \geq 10\%$.

1. Introduction

Gastroenteropancreatic neuroendocrine neoplasms (GEP-NENs) are a heterogeneous group of neoplasms that arise from cells of the endocrine system [1]. GEP-NENs are rare and present many clinical challenges. Because of their unpredictable biologic behaviors, the diagnosis usually takes place only after the condition has become advanced. Treatment regimens rely mainly on histological grading via biopsy; however, tumor heterogeneity cannot be fully assessed by tumor biopsy [2, 3]. Although Ki-67 staining has been shown to have prognostic significance in GEP-NENs, pitfalls may lead clinicians to misjudge the tumor grades. Specifically, first, the current gold-standard method, in which 2000 cells

are counted, is heavily dependent on the skill and expertise of the reporting pathologist [4]. Second, the limited tissue in some cases may impede accurate assessment of Ki-67 given the potential for heterogeneity of Ki-67 expression within tumors. Finally, the Ki-67 index may vary over time in the same patient, with changes possible both in response to treatment [5] and over the progression of the disease [6]. We believe that combined dual-tracer PET/CT imaging offers distinct advantages to overcome the above pitfalls.

DOTA-peptides specifically bind to somatostatin receptors 2, 3, and 5 and are usually overexpressed on the surfaces neuroendocrine cells [7, 8]. ^{68}Ga -DOTATATE has been shown to be useful for staging, restaging, surveillance, determining SSTR-based therapy, and monitoring responses

to treatments in NENs [9, 10]. ^{18}F -FDG is a glucose analogue, and PET/CT imaging with this tracer has been shown to be correlated with NENs aggressiveness. The presence of increased glucose in NENs highlights an increased propensity for invasion and metastasis, and PET imaging with ^{18}F -FDG accordingly has higher sensitivity, especially in aggressive and high-grade tumors [11, 12]. Therefore, we believe that PET/CT imaging with combination of ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT is a highly efficient whole-body imaging method, and it could be complementary to conventional imaging methods.

A large number of previous studies have evaluated the diagnostic accuracy of both tracers in the presence of a relative shortage of information, regarding the correlation to pathological findings and prognostic value. Only a few studies have compared the clinical impact of both ^{68}Ga -DOTATATE and ^{18}F -FDG PET tracers on NENs [13, 14]. The present study aimed to determine the clinical value of the complementary PET/CT imaging method in a large histologically proven NENs population.

2. Materials and Methods

We analyzed the data from 83 (50 males and 33 females) consecutive patients with pathologically proven NENs who underwent contemporaneous PET/CT imaging with ^{68}Ga -DOTATATE and ^{18}F -FDG between June 2013 and December 2016. ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT scans were performed within an interval of no more than 2 weeks. No patients were treated during this interval. All NENs were classified according to the histopathological reports, which are based on recent consensus statements of the European Neuroendocrine Tumor Society. According to the grade of differentiation, proliferation index (Ki-67), and mitotic count, the well-differentiated (WD) neoplasms are herein defined as NET and graded G1 (Ki-67 $\leq 2\%$) or G2 (Ki-67 3–20%) and G3a (Ki-67 $> 20\%$); the poorly differentiated (PD) neoplasms are defined as NEC and G3b. G2 patients were further divided into 2 groups as G2a (3–9%) and G2b (10–20%).

2.1. PET/CT Acquisition. Patients fasted for at least 6 h before PET/CT scan. Images were acquired 1 h after injection of 3.7 MBq/kg ^{18}F -FDG or 1 h after the injection of 100–200 MBq ^{68}Ga -DOTATATE. Whole-body scan (brain to mid-thigh) was performed with the patient in the supine position. CT exposure factors for all scans were 120 kV and 100 mA. PET/CT images were reported in consensus by two experienced nuclear medicine physicians who were blinded to the findings of the structural imaging. Any nonphysiological focus of ^{68}Ga -DOTATATE or ^{18}F -FDG uptake greater than the normal liver background was considered positive. At the same time, CT imaging was used to differentiate between lesions and physiological uptake. The maximum standardized uptake value (SUV_{\max}) of primary and metastatic lesions was calculated. SUV_{\max} generated from each patient was used in the final analysis. The SUV ratio of the tumor relative to the maximal liver uptake was calculated by dividing SUV_{\max} of the tumor by SUV_{\max} of the liver ($\text{SUV}_{T/L}$). The ratio between

TABLE 1: Patient characteristics.

Characteristics	NET (%)	NEC (%)	All (%)
Gender			
Female (<i>n</i>)	26 (51.0)	7 (21.9)	33 (39.7)
Male (<i>n</i>)	25 (49.0)	25 (78.1)	50 (60.3)
Primary sites (<i>n</i>)			
Pancreas	19 (37.3)	8 (25.0)	27 (32.5)
Gastrointestinal	24 (47.0)	19 (59.4)	43 (51.8)
Primary unknown	8 (15.7)	5 (15.6)	13 (15.7)
Metastatic sites (<i>n</i>)			
Liver	36 (70.6)	13 (40.6)	49 (59.0)
Lymph nodes	21 (41.2)	27 (84.4)	48 (57.8)
Bone	9 (17.6)	10 (31.3)	19 (22.9)
Lung	4 (7.8)	1 (3.1)	5 (6.0)
Other	6 (11.7)	7 (21.9)	13 (15.7)

NET, neuroendocrine tumor; NEC, neuroendocrine carcinoma.

SUV_{\max} of ^{68}Ga -DOTATATE and that of ^{18}F -FDG (SUV_{\max} ratio) was also calculated.

2.2. Statistical Analysis. Descriptive analyses are presented using mean and SD for normally distributed variables, but median, minimum, and maximum values were used for those that were nonnormally distributed. Analyses were performed using SPSS (version 21.0; IBM). The paired Student's *t*-test was used for the related nonnormally distributed variables. Correlation of SUV_{\max} values of ^{68}Ga -DOTATATE and ^{18}F -FDG with Ki-67 index was assessed using Spearman's correlation coefficient. Overall survival (OS) was calculated from the date of diagnosis to the last day of follow-up or death. Kaplan-Meier survival analysis was performed to assess the prognostic value regarding OS, and differences between groups were analyzed using the log-rank test. $P < 0.05$ was considered as statistical significance.

3. Results

3.1. Patient Characteristics. A total of 83 (50 males and 33 females) patients were included in the study, with a median age of 56 years (range: 27–77 years). The primary tumors were located in the pancreas in 27 patients (32.5%), gastrointestinal tract in 43 patients (51.8%), and unknown locations in 13 patients (15.7%). Among the subjects, 48 patients (57.8%) had lymph node involvement and 60 patients (72.3%) had distant metastases. Pathological evaluation showed that 51 patients (61.4%) had WD-NET and 32 patients (38.6%) had PD-NEC. Among WD-NETs, there were 14 patients (16.9%) in G1 ($\leq 2\%$), 19 patients (22.9%) in G2a (3–9%), 9 patients (10.8%) in G2b (10–20%), and 28 patients (33.7%) in G3a ($> 20\%$) (Table 1).

3.2. Complementary PET/CT Qualitative Evaluation

3.2.1. Comparison in Sensitivity. For all patients, ^{68}Ga -DOTATATE was positive in 53 cases and negative in 30 cases, while ^{18}F -FDG PET/CT was positive in 62 cases and

TABLE 2: Sensitivity and uptake of ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT for different primary sites and grades.

	Sensitivity (%)		SUV_{\max} (mean \pm SD)	
	^{68}Ga -DOTATATE	^{18}F -FDG PET/CT	^{68}Ga -DOTATATE	^{18}F -FDG PET/CT
Primary lesion				
Gastrointestinal tract	55.8%	74.4%	16.75 ± 2.62	7.56 ± 0.87
Pancreas	85.2%	66.7%	29.87 ± 4.77	6.51 ± 0.78
WD NET	80.4%	58.8%	28.87 ± 3.52	4.51 ± 0.45
Gastrointestinal NET	75.0%	54.2%	22.68 ± 2.77	3.71 ± 0.45
Pancreatic NET	89.5%	52.6%	31.19 ± 4.25	5.13 ± 0.93
PD NEC	37.5%	100.0%	10.86 ± 1.78	11.46 ± 0.75
Gastrointestinal NEC	31.6%	100.0%	9.26 ± 1.25	12.44 ± 1.11
Pancreatic NEC	75.0%	100.0%	18.23 ± 5.93	10.23 ± 0.67

WD, well-differentiated; PD, poorly differentiated; NET, neuroendocrine tumor; NEC, neuroendocrine carcinoma.

negative in 21 cases. Overall, ^{18}F -FDG assessment was found to have a better sensitivity (74.7%) compared with ^{68}Ga -DOTATATE (63.8%), although it is not statistically significant ($P = 0.593$). Remarkably, we found that the sensitivity of dual tracers (94.0%) was significantly higher than that with ^{68}Ga -DOTATATE or ^{18}F -FDG alone ($P < 0.01$). For NET and NEC, the sensitivity was 80.39% (41/51) and 37.5% (12/32) with ^{68}Ga -DOTATATE and was 58.82% (30/51) and 100% (32/32) with ^{18}F -FDG. Separating from the primary sites, the sensitivity of ^{68}Ga -DOTATATE in pancreatic NET was higher than that in gastrointestinal NET (NET: 89.5% versus 75.0%, $P = 0.034$; NEC: 75.0% versus 31.6%, $P = 0.027$). However, ^{18}F -FDG showed no difference between pancreatic and gastrointestinal NENs in terms of sensitivity.

3.2.2. Complementary PET/CT Semiquantitative Evaluation. Table 2 illustrates the sensitivities and SUV_{\max} in correlation with primary sites and grades. The uptake values of ^{68}Ga -DOTATATE in PanNENs were significantly higher than those in GI-NEN (29.87 \pm 4.78 versus 16.76 \pm 2.62, $P = 0.011$). The values of ^{18}F -FDG PET/CT in PanNEN had a trend toward a lower SUV than that in GI-NEN (6.51 \pm 0.77 versus 7.57 \pm 0.86, $P = 0.067$) (Figure 1).

3.2.3. Comparison in Different Pathological Groups. The sensitivity of ^{68}Ga -DOTATATE and ^{18}F -FDG in G1/G2a/G2b/G3a/G3b was 78.6%/73.3%/88.9%/77.8%/37.5% and 50.0%/52.6%/66.7%/77.8%/100.0%. ^{68}Ga -DOTATATE imaging provided a sensitivity of >73% in WD-NET (G1-G3a) and only 37.5% in PD-NEC (G3b). For G1 and G2a (Ki-67 < 10%), a statistically significant positive correlation between Ki-67 and the sensitivity with ^{18}F -FDG could be found, and the sensitivity in G1 and G2a (Ki-67 < 10%) was about 50%, increasing dramatically when the Ki-67 index was over 10%. The sensitivity in G3a reached 77.8% and that in G3b was 100%. The patients were divided into 3 groups: group A (G1 + G2a), group B (G2b + G3a), and group C (G3b). With this grouping, the sensitivity of ^{68}Ga -DOTATATE and ^{18}F -FDG in group A/B/C was 78.8%/83.3%/37.5% and 52.0%/72.2%/100.0%. Importantly, we found that the sensitivities of imaging with dual tracers in groups A/B/C

were 84.8%/100%/100%, which were significantly higher than that with the single tracer. There was a significant negative correlation between Ki-67 and ^{68}Ga -DOTATATE SUV_{\max} ($R = -0.415$; $P \leq 0.001$), while a positive correlation was noted between Ki-67 and ^{18}F -FDG SUV_{\max} value ($R = 0.683$; $P \leq 0.001$). Moreover, ^{68}Ga -DOTATATE $\text{SUV}_{T/L}$ showed a negative correlation with Ki-67 index ($R = -0.357$; $P = 0.001$). However, ^{18}F -FDG $\text{SUV}_{T/L}$ was positively correlated with Ki-67 index ($R = 0.617$; $P \leq 0.001$) (Figure 1).

3.2.4. Concordant and Discordant Findings. When combining the results of the dual-tracer PET/CT, 37 patients were positive in both tracers, and 16 patients were ^{68}Ga -DOTATATE-positive and ^{18}F -FDG-negative, while 25 patients were ^{18}F -FDG-positive and ^{68}Ga -DOTATATE-negative. 5 patients were negative in both tracers (Table 3).

^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT findings were concordant in 37 patients with 25 WD-NETs and in 12 PD-NECs. Only 1 patient was diagnosed as localized duodenal NEN. The remaining 36 patients had regional lymph node metastasis, distant metastasis, or both mainly occurring in liver or bone. Of 27 patients with liver metastases, 8 patients (29.6%) examined via ^{68}Ga -DOTATATE showed heterogeneity in SSTR expression. Of 8 patients with bone metastases, imaging findings of 5 patients (62.5%) demonstrated that ^{68}Ga -DOTATATE highlighted more bone lesions than ^{18}F -FDG PET/CT. Of 22 patients with lymph node involvement, the dual tracers with 5 patients (22.7%) showed that ^{18}F -FDG findings could be more prominent than those with ^{68}Ga -DOTATATE (Figure 2).

5 patients showed negative results in both two tracers, with a histological diagnosis of group A. 1 patient was diagnosed as duodenal NET G2 (Ki-67: 3%) with multiple liver metastases (largest lesion: 7.2 cm \times 4.7 cm). The remaining 4 patients had rectal or gastric NET (G1/2) with lesions smaller than 5 mm.

3.3. Treatment and Follow-Up. Of the 83 patients, 26 performed radical surgery; 57 (31 NET and 26 NEC) unresectable patients were treated with palliative surgery, SSA, chemotherapy, and TACE. The median follow-up was 21




FIGURE 1: Comparison of SUV_{max} of PET/CT according to primary sites ((a) and (b)) and tumor grade ((c) and (d)). The sensitivity of group A (G1 + G2a = Ki-67 < 10%), group B (G2b + G3a = well-differentiated neoplasms with Ki-67 ≥ 10%), and group C (G3b = poorly differentiated neoplasms with Ki-67 > 20%) in PET/CT imaging.

months (in the range of 2–62 months). During the follow-up period, 9 (6 NEC and 3 NET) patients died of the progressive disease. Unresectable patients with positive results solely with ^{18}F -FDG showed the worst prognosis, while those positive solely with ^{68}Ga -DOTATATE showed the best prognosis.

For unresectable patients with NET ($n = 31$), being ^{68}Ga -DOTATATE-negative was associated with worse prognosis (HR: 10.4; 95% CI: 1.5–78.2; $P \leq 0.001$), and being ^{18}F -FDG-positive tended to be correlated with a worse prognosis (HR: 3.6; 95% CI: 0.7–9.8; $P = 0.158$). For unresectable patients

TABLE 3: Concordant and discordant findings.

⁶⁸ Ga-DOTATATE	Positive	Positive	Negative	Negative
¹⁸ F-FDG PET/CT	Positive	Negative	Positive	Negative
Primary lesion (<i>n</i>)				
Pancreas	14	9	4	0
Gastrointestinal tract	18	6	14	5
CUP	5	1	7	0
Metastatic sites				
Liver	27	11	9	1
Lymph node	22	7	18	0
Bone	8	2	8	0
Ki-67 range				
Group A (<i>n</i> = 33)	15 (45.5%)	11 (33.3%)	2 (6.1%)	5 (15.1%)
Group B (<i>n</i> = 18)	10 (55.5%)	5 (27.8%)	3 (16.7%)	0
Group C (<i>n</i> = 32)	12 (37.5%)	0	20 (62.5%)	0

CUP, Cancer of unknown primary.




FIGURE 2: A 37-year-old women with pancreatic NEC G3 (Ki-67 = 80%) and lymph node, liver, and bone metastases, from whom the primary lesion has been resected.¹⁸F-FDG PET/CT showed more liver lesions, while ⁶⁸Ga-DOTATATE detected more bone lesions. ((A) and (a)) Liver lesions showed heterogeneity in SSTR expression. ((B) and (b) and (C) and (c)) ¹⁸F-FDG PET/CT failed to show bone metastases in rib and lumbar vertebra.

with NEC (*n* = 26), being ⁶⁸Ga-DOTATATE-negative also tended to be associated with a worse prognosis (HR: 2.4; 95% CI: 0.3–5.4; *P* = 0.382) (Figure 3).

4. Discussion

⁶⁸Ga-DOTATATE and ¹⁸F-FDG PET/CT play a crucial role in the diagnosis and clinical management of NENs with morphologic and functional information. ⁶⁸Ga-DOTATATE was found to be superior to ¹⁸F-FDG in WD-NET, whereas

¹⁸F-FDG was more sensitive in PD-NEC [15]. Considering the costs of molecular imaging, choosing the selected patients for the specific PET/CT imaging is of vital importance. Analyzing the results produced using both tracers for different grades and primary sites could be a balanced approach. The aim of the present study was to compare ⁶⁸Ga-DOTATATE and ¹⁸F-FDG PET/CT in GEP-NENs and to investigate the relationship between the complementary PET/CT results and histopathologic findings in clinical and prognostic values in a large, histologically confirmed NEN population.




FIGURE 3: The overall survival of unresectable patients detected with dual tracers: (a) Kaplan-Meier survival curve for unresectable patients, (A) positive for both tracers, (B) ^{68}Ga -DOTA-TATE only, and (C) ^{18}F -FDG only; (b) unresectable NET patients with ^{68}Ga -DOTA-TATE results (positive or negative); (c) unresectable NET patients with ^{18}F -FDG results (positive or negative); (d) unresectable NEC patients with ^{68}Ga -DOTA-TATE results (positive or negative).

^{68}Ga -DOTATATE imaging and ^{18}F -FDG PET/CT imaging have been compared in several studies which have been shown to have variable sensitivities in detecting NENs with a relatively small number of patients. Naswa et al. [16] reported that the sensitivity of ^{68}Ga -DOTANOC and ^{18}F -FDG was 91.4% and 42.5%, respectively. Koukouraki et al. [17] demonstrated that the sensitivity of ^{68}Ga -DOTATOC and ^{18}F -FDG was 90% and 68%, respectively, and ^{68}Ga -DOTANOC was more sensitive in the detection of primary sites or metastasis than ^{18}F -FDG [18]. Notably the patients included in this study were mainly WD-NETs with lower glucose metabolism. In the present study, the sensitivity of ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT was 63.85% and 74.70%, respectively. Subgroup analysis showed that the sensitivity of ^{68}Ga -DOTATATE was mainly correlated with the degree of differentiation, instead of correlation with

Ki-67 index. The sensitivity of ^{18}F -FDG showed a positive correlation with the Ki-67 index and differentiation. From ^{18}F -FDG imaging, the sensitivity was under 53% for patients with Ki-67 < 10% and 100% for PD-NEC. SUV_{max} of patients with NEC was low even for those who had positive results under ^{68}Ga -DOTATATE. We also observed that patients with Ki-67 < 10% showed low uptake in ^{18}F -FDG PET/CT. On consideration of the weak significance of ^{68}Ga -DOTATATE for PD-NEC and ^{18}F -FDG PET/CT for lower-grade NET, our study demonstrated that ^{18}F -FDG is more suitable for patients with Ki-67 $\geq 10\%$, and ^{68}Ga -DOTATATE is less advantageous in PD-NEC and should be tailored to the individual patients.

Our study demonstrated that PET/CT uptake was statistically significantly different between subgroups of GEP-NENs according to grading. The cohort was separated into




FIGURE 4: PET/CT imaging and treatment regime.

3 groups: A (G1 + G2a), B (G2b + G3a), and C (G3b). Group B showed higher sensitivity for ¹⁸F-FDG than group A, and the median SUV_{max} increased significantly, indicating relatively quick proliferation rate associated with Ki-67 index. Strosberg [19] proposed chemotherapy as a treatment option for tumors with Ki-67 ≥ 10%, especially those with higher ¹⁸F-FDG activity. Such tumors showed high proliferative capacity and aggressive behavior and are recommended for chemotherapy after PRRT or somatostatin therapy. In this way, ¹⁸F-FDG PET/CT may be suitable for identifying patients with aggressive conditions in group B who could benefit from chemotherapy. Thus, PET/CT imaging may establish the missing link between histopathologic findings and the treatment regimen.

Based on our results, dual tracers' assessment is recommended for WD-NET with Ki-67 ≥ 10% (G2b and G3a). ⁶⁸Ga-DOTATATE and ¹⁸F-FDG were complementary in detecting lesions and dual-trace PET/CT showed an advantage in the assessment of SSA, PRRT, and chemotherapy. We proposed that WD-NET patients with Ki-67 ≥ 10% should be examined using dual tracers upon diagnosis. We also suggest that ⁶⁸Ga-DOTATATE should be performed solely in WD-NET patients with Ki-67 < 10% and ¹⁸F-FDG is sufficient for PD NEC. Moreover, repeated PET/CT is warranted when disease progresses rapidly, considering the heterogeneous expression and complementary findings to histopathology (Figure 4).

We investigated the correlations between dual tracers and Ki-67 index. SUV_{max} or SUV_{T/L} was positively correlated with Ki-67 index with respect to ¹⁸F-FDG PET/CT and negatively correlated with Ki-67 index with respect to ⁶⁸Ga-DOTATATE. SUV_{max} of PET/CT may be a suitable biomarker for evaluation of the biological behavior of NENs. The relationship between SUV_{T/L} and Ki-67 index was weak, but it showed a consistency with SUV_{max}. However, SUV_{max} may be influenced by technical elements, while SUV_{T/L} could reduce the differences attributable to equipment and operation to some extent.

GEP-NENs are a heterogeneous group of neoplasms that display great variability in biological behaviors and clinical outcomes [20]. This requires accurate diagnostic techniques for precise staging and choice of therapy. The standard

grading is mainly based on the immunohistochemistry of the proliferation marker Ki-67. However, there are many lesions with variable tracer uptake at different parts of the tumor. Especially within the same organ, these lesions may prevent the biopsy from being a comprehensive reflection of tumor heterogeneity *in vivo*, leading to inaccurate Ki-67 index results. Therefore, dual-trace PET/CT, which is a whole-body noninvasive alternative, is warranted to overcome the shortcomings of histopathologic grading. The findings of this study included the ability of ⁶⁸Ga-DOTATATE to detect heterogeneity in tumors and variable expression in primary sites. SUV_{max} of PanNEN was higher than that of GI-NENs, which was consistent with prior studies [21, 22], where the investigators found higher levels of messenger RNA expression of SSTR2 and SSTR5 in pancreatic than in gastrointestinal NENs. In this way, ⁶⁸Ga-DOTATATE is more sensitive in PanNEN, but it may miss some lesions in GI-NEN. However, SUV_{max} of PanNEN and GI-NEN showed no difference for ¹⁸F-FDG, which is complementary to ⁶⁸Ga DOTATATE. Another finding of the present study was that ⁶⁸Ga DOTATATE has a superb ability to detect heterogeneity in metastatic lesions, which is beneficial for us to select the optimal protocol. We believe that complementary PET/CT can evaluate the tumors' heterogeneity and influence treatment options.

The morphological findings and Ki-67 index are considered important prognostic markers in NENs. One major limitation of these histopathological parameters as prognostic markers is the requirement for tissue sampling, which is not always feasible. A few studies have demonstrated the prognostic value of ¹⁸F-FDG and ⁶⁸Ga-DOTATATE PET/CT in patients with NENs. ¹⁸F-FDG was an independent predictor of PFS [23, 24]. ¹⁸F-FDG SUV_{max} > 3 was demonstrated to be independent predictor of disease progression, superior to conventional prognostic factors such as Ki-67 index and serum CgA [25]. SSTR expression was found to be a positive prognostic factor, and it was therefore expected that SSTR-base PET/CT would also have prognostic value in NENs [26]. Sharma et al. [27] demonstrated the prognostic value of SUV_{max} measured with ⁶⁸Ga DOTATATE in 37 patients with NET. SUV_{max} ≥ 14.5 was found to distinguish patients with progressive and those with nonprogressive disease. A major limitation of the prior studies was that most of the patients included were WD-NET, which may involve selection bias. Our study enrolled 57 unresectable patients (31 NET and 26 NEC). Unresectable patients positive with ¹⁸F-FDG alone showed the worst prognosis, while those positive with ⁶⁸Ga-DOTATATE alone had the best prognosis. NET patients with predominant ¹⁸F-FDG uptake and a negative ⁶⁸Ga-DOTATATE scans had worse prognosis. There is a strong association between high ¹⁸F-FDG uptake and worse outcome even in patients with WD-NETs. However, PD-NEC with negative ⁶⁸Ga-DOTATATE uptake may lead to worse prognosis. The present study has several limitations. First, our study is retrospective, and the second limitation is that the follow-up time was not long enough to assess treatment response, considering the relatively inert biological behavior of NENs.

Noninvasive dual-tracer imaging with ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT seems promising as an alternative to tissue sampling, due to its capacity to reflect two different aspects of tumor biology, SSTR expression and glucose metabolism, respectively. Accordingly, imaging with dual tracer is recommended for WD-NET patients with Ki-67 $\geq 10\%$, providing information for selection of SSA, PRRT, and chemotherapy. Taking the advantages of dual-tracer imaging in detecting lesions is useful for accurate clinical management. Dual-tracer imaging also shows a clearly linear correlation between SUV_{max} and Ki-67 index. On consideration of heterogeneous expression and complementary findings to histopathology, our results suggested that repeated dual-tracer imaging is warranted to evaluate dynamic biological behavior and prognosis.

Abbreviations

GEP-NENs:	Gastroenteropancreatic neoplasms
PD:	Poorly differentiated
WD:	Well-differentiated
OS:	Overall survival
NET:	Neuroendocrine tumor
NEC:	Neuroendocrine carcinoma
PRRT:	Peptide receptor radionuclide therapy

Conflicts of Interest

The authors declare that there are no conflicts of interest that could be perceived as prejudicing the impartiality of the research reported.

Authors' Contributions

Panpan Zhang and Jiangyuan Yu contributed equally to this work. Panpan Zhang and Jiangyuan Yu wrote the manuscript and analyzed the data. Jie Li, Lin Shen, Nan Li, Hua Zhu, Shizhen Zhai, and Yan Zhang collected patient data and technical information. Zhi Yang and Ming Lu reviewed and helped revise the manuscript.

References

- [1] R. Garcia-Carbonero, J. Capdevila, G. Crespo-Herrero et al., "Incidence, patterns of care and prognostic factors for outcome of gastroenteropancreatic neuroendocrine tumors (GEP-NETs): results from the national cancer registry of Spain (RGETNE)," *Annals of Oncology*, vol. 21, no. 9, pp. 1794–1803, 2010.
- [2] D. S. Klimstra, I. R. Modlin, N. V. Adsay et al., "Pathology reporting of neuroendocrine tumors: Application of the delphic consensus process to the development of a minimum pathology data set," *The American Journal of Surgical Pathology*, vol. 34, no. 3, pp. 300–313, 2010.
- [3] G. Carliniante, P. Baccarini, D. Beretti et al., "Ki-67 cytological index can distinguish well-differentiated from poorly differentiated pancreatic neuroendocrine tumors: A comparative cytohistological study of 53 cases," *Virchows Archiv*, vol. 465, no. 1, pp. 49–55, 2014.
- [4] G. Rindi, G. Klöppel, A. Couvelard et al., "TNM staging of midgut and hindgut (neuro) endocrine tumors: A consensus proposal including a grading system," *Virchows Archiv*, vol. 451, no. 4, pp. 757–762, 2007.
- [5] J. C. Yao, A. T. Phan, D. Z. Chang et al., "Efficacy of RAD001 (everolimus) and octreotide LAR in advanced low- to intermediate-grade neuroendocrine tumors: results of a phase II study," *Journal of Clinical Oncology*, vol. 26, no. 26, pp. 4311–4318, 2008.
- [6] S. Singh, J. Hallet, C. Rowsell, and C. H. L. Law, "Variability of Ki67 labeling index in multiple neuroendocrine tumors specimens over the course of the disease," *European Journal of Surgical Oncology*, vol. 40, no. 11, pp. 1517–1522, 2014.
- [7] E. Grande, J. J. Díez, V. Pachón, and A. Carrato, "Advances in the therapy of gastroenteropancreatic-neuroendocrine tumours (GEP-NETs)," *Clinical and Translational Oncology*, vol. 12, no. 7, pp. 481–492, 2010.
- [8] M. Miederer, S. Seidl, A. Buck et al., "Correlation of immunohistopathological expression of somatostatin receptor 2 with standardised uptake values in ^{68}Ga -DOTATOC PET/CT," *European Journal of Nuclear Medicine and Molecular Imaging*, vol. 36, no. 1, pp. 48–52, 2009.
- [9] W. A. P. Breeman, E. de Blois, H. Sze Chan, M. Konijnenberg, D. J. Kwekkeboom, and E. P. Krenning, " ^{68}Ga -labeled DOTA-peptides and ^{68}Ga -labeled radiopharmaceuticals for positron emission tomography: current status of research, clinical applications, and future perspectives," *Seminars in Nuclear Medicine*, vol. 41, no. 4, pp. 314–321, 2011.
- [10] N. Naswa, P. Sharma, A. Kumar et al., "Gallium-68-DOTANOC PET/CT of patients with gastroenteropancreatic neuroendocrine tumors: A prospective single-center study," *American Journal of Roentgenology*, vol. 197, no. 5, pp. 1221–1228, 2011.
- [11] I. Kayani, J. B. Bomanji, A. Groves et al., "Functional imaging of neuroendocrine tumors with combined PET/CT using ^{68}Ga -DOTATATE (Dota-DPhE₁, Tyr₃-octreotate) and ^{18}F -FDG," *Cancer*, vol. 112, no. 11, pp. 2447–2455, 2008.
- [12] R. Abgral, S. Leboulleux, D. Déandreibs et al., "Performance of 18fluorodeoxyglucose-positron emission tomography and somatostatin receptor scintigraphy for high Ki67 ($\geq 10\%$) well-differentiated endocrine carcinoma staging," *The Journal of Clinical Endocrinology & Metabolism*, vol. 96, no. 3, pp. 665–671, 2011.
- [13] D. H. Simsek, S. Kuyumcu, C. Turkmen et al., "Can complementary ^{68}Ga -dotataate and ^{18}F -FDG PET/CT establish the missing link between histopathology and therapeutic approach in gastroenteropancreatic neuroendocrine tumors?" *Journal of Nuclear Medicine*, vol. 55, no. 11, pp. 1811–1817, 2014.
- [14] E. Panagiotidis, A. Alshammari, S. Michopoulos et al., "Comparison of the impact of ^{68}Ga -DOTATATE and ^{18}F -FDG PET/CT on clinical management in patients with Neuroendocrine tumors," *Journal of Nuclear Medicine*, vol. 58, no. 1, pp. 91–96, 2017.
- [15] M. H. Squires, N. Volkan Adsay, D. M. Schuster et al., "Octreoscan Versus FDG-PET for Neuroendocrine Tumor Staging: A Biological Approach," *Annals of Surgical Oncology*, vol. 22, no. 7, pp. 2295–2301, 2015.
- [16] N. Naswa, P. Sharma, S. K. Gupta et al., "Dual tracer functional imaging of gastroenteropancreatic neuroendocrine tumors using ^{68}Ga -DOTA-NOC PET-CT and ^{18}F -FDG PET-CT: Competitive or complimentary?" *Clinical Nuclear Medicine*, vol. 39, no. 1, pp. e27–e34, 2014.
- [17] S. Koukouraki, L. G. Strauss, V. Georgoulias, M. Eisenhut, U. Haberkorn, and A. Dimitrakopoulou-Strauss, "Comparison of the pharmacokinetics of ^{68}Ga -DOTATOC and $[^{18}\text{F}]$ FDG in


- patients with metastatic neuroendocrine tumors scheduled for ^{90}Y -DOTATOC therapy," *European Journal of Nuclear Medicine and Molecular Imaging*, vol. 33, no. 10, pp. 1115–1122, 2006.
- [18] V. Ambrosini, D. Campana, L. Bodei et al., "68Ga-DOTANOC PET/CT clinical impact in patients with neuroendocrine tumors," *Journal of nuclear medicine : official publication, Society of Nuclear Medicine*, vol. 51, no. 5, pp. 669–673, 2010.
- [19] J. R. Strosberg, "Systemic treatment of gastroenteropancreatic neuroendocrine tumors (GEP-NETs): Current approaches and future options," *Endocrine Practice*, vol. 20, no. 2, pp. 167–175, 2014.
- [20] Z. Yang, L. H. Tang, and D. S. Klimstra, "Effect of tumor heterogeneity on the assessment of Ki67 labeling index in well-differentiated neuroendocrine tumors metastatic to the liver: Implications for prognostic stratification," *The American Journal of Surgical Pathology*, vol. 35, no. 6, pp. 853–860, 2011.
- [21] D. Campana, V. Ambrosini, R. Pezzilli et al., "Standardized uptake values of 68Ga-DOTANOC PET: a promising prognostic tool in neuroendocrine tumors," *Journal of Nuclear Medicine*, vol. 51, no. 3, pp. 353–359, 2010.
- [22] D. O'Toole, A. Saveanu, A. Couvelard et al., "The analysis of quantitative expression of somatostatin and dopamine receptors in gastro-entero-pancreatic tumours opens new therapeutic strategies," *European Journal of Endocrinology*, vol. 155, no. 6, pp. 849–857, 2006.
- [23] A. Sundin and A. Rockall, "Therapeutic monitoring of gastroenteropancreatic neuroendocrine tumors: The challenges ahead," *Neuroendocrinology*, vol. 96, no. 4, pp. 261–271, 2012.
- [24] E. Garin, F. Le Jeune, A. Devillers et al., "Predictive value of 18F-FDG PET and somatostatin receptor scintigraphy in patients with metastatic endocrine tumors," *Journal of Nuclear Medicine*, vol. 50, no. 6, pp. 858–864, 2009.
- [25] T. Binderup, U. Knigge, A. Loft, B. Federspiel, and A. Kjaer, " ^{18}F -fluorodeoxyglucose positron emission tomography predicts survival of patients with neuroendocrine tumors," *Clinical Cancer Research*, vol. 16, no. 3, pp. 978–985, 2010.
- [26] H. S. Kim, H. S. Lee, and W. H. Kim, "Clinical significance of protein expression of cyclooxygenase-2 and somatostatin receptors in gastroenteropancreatic neuroendocrine tumors," *Cancer Research and Treatment*, vol. 43, no. 3, pp. 181–188, 2011.
- [27] P. Sharma, N. Naswa, S. S. KC et al., "Comparison of the prognostic values of ^{68}Ga -DOTANOC PET/CT and ^{18}F -FDG PET/CT in patients with well-differentiated neuroendocrine tumor," *European Journal of Nuclear Medicine and Molecular Imaging*, vol. 41, no. 12, pp. 2194–2202, 2014.




**The Scientific
World Journal**




Gastroenterology
Research and Practice



MEDIATORS
of
INFLAMMATION




Journal of
Diabetes Research



Disease Markers



Journal of
Immunology Research



PPAR Research



Hindawi

Submit your manuscripts at
www.hindawi.com




International Journal of
Endocrinology




BioMed
Research International




Journal of
Ophthalmology



Stem Cells
International




eCAM
Evidence-Based
Complementary and
Alternative Medicine




Journal of
Obesity



Journal of
Oncology




Computational and
Mathematical Methods
in Medicine




Behavioural
Neurology



Parkinson's
Disease



AIDS
Research and Treatment



Oxidative Medicine and
Cellular Longevity