

A Mathematical Model for the Dynamics of Hepatitis C

R. AVENDAÑO^a, L. ESTEVA^{b,*}, J.A. FLORES^b, J.L. FUENTES ALLEN^c, G. GÓMEZ^b and JE. LÓPEZ-ESTRADA^b

^aEducación Médica, Centro Médico Nacional Siglo XXI, IMSS, 06720, Mexico, D.F.; ^bDepto. De Matemáticas, Fac. de Ciencias, UNAM, 04510, Mexico, D.F.; ^cHospital de Infectología, Centro Médico la Raza, IMSS, Mexico, D.F.

(Received 27 September 2000; In final form 21 May 2001)

We formulate a model to describe the dynamics of hepatitis C virus (HCV) considering four populations: uninfected liver cells, infected liver cells, HCV and T cells. Analysis of the model reveals the existence of two equilibrium states, the uninfected state in which no virus is present and an endemically infected state, in which virus and infected cells are present. There exists a threshold condition that determines the existence and stability of the endemic equilibrium. We discuss the efficacy of the therapy methods for hepatitis C in terms of the threshold parameter. Success of the therapy could possibly be predicted from the early viral dynamics in the patients.

Keywords: Hepatitis C; Therapy; Threshold; Uninfected steady state; Endemically infected steady state

INTRODUCTION

Infection with hepatitis C virus (HCV) represents a public health problem with an alarming prevalence (2–15%) throughout the world (Neumann *et al.*, 1998). The existence of hepatitis C was not appreciated until 1975, when the application of recently developed diagnostic test for hepatitis A and B revealed that many cases were neither hepatitis A nor hepatitis B. The causative agent was identified in 1989 (Purcell, 1994).

The HCV is commonly transmitted via blood and blood products. Its transmission by other routes as unprotected sex, perinatal transmission from infected mother to offspring, etc. have been proposed but remain controversial and probably of minor importance.

The incubation period of hepatitis C averages 50 days. Acute hepatitis C is generally a mild disease with a mortality rate of 1%. However, more than 50% of acute cases progress to chronicity, and some of them will eventually evolve to cirrhosis or hepatocellular carcinoma, or both (Purcell, 1994).

In acute infection, the most common symptom is fatigue. However, the majority of cases (up to 90%) are asymptomatic. This makes the diagnosis of hepatitis C very difficult.

The current treatment for hepatitis C consists in the application of interferon (IFN) α -2b with dose from 3 to 15 million international units (mIU). However, the

treatment with IFN is successful in only 11–30% cases. In Neumann *et al.* (1998) an analysis of the efficacy of IFN- α therapy is presented. In addition, it has been reported that IFN used in combination with ribavirin (another antiviral agent) is more effective than the treatment with only IFN (Purcell, 1994). No vaccine is available for hepatitis C, since a major obstacle to vaccine development is the probability of extensive antigenic variation between different strains (Lemon and Brown, 1995).

Appropriate mathematical models can be helpful to answer biologically important questions concerned with pathogenesis, the dynamics of the immune response and effectiveness of drug treatment. Models to understand the immune response to persistent virus and effectiveness of drug therapy have been formulated by several authors. Thus, Nowak and Bangham (1996) used a simple mathematical approach to explore the effects of individual variation in immune responsiveness on virus load and diversity. They found that a better indicator of CTL responsiveness is the equilibrium virus load, rather than the abundance of virus specific CTLs. Nowak *et al.* (1996) formulated a model that provided a quantitative understanding of HBV replication dynamics. Their analysis had implication for the optimal timing of drug treatment and immunotherapy in chronic HBV infection. Payne *et al.* (1996) formulated a model of hepatitis B virus infection to address important features of the infection, namely the

*Corresponding author. E-mail: lesteva@servidor.unam.mx

wide manifestations of the infection and the age dependence thereof, and the typically long delay before the development of virus-induced liver cancer. Bonhoeffer *et al.* (1997) analyzed the dynamics of virus populations, the role of the immune system and resistance of drug therapy in limiting virus abundance in infections with HIV or hepatitis B. Neumann *et al.* (1998) used a mathematical model to analyze the efficacy of treatment with IFN- α therapy.

In this paper, we formulate and analyze a model for the HCV dynamics. Our model is closely related to the models proposed in Nowak and Bangham (1996), Nowak *et al.* (1996), Payne *et al.* (1996), Bonhoeffer *et al.* (1997) and Neumann *et al.* (1998), but here we consider the immune response by adding the virus-specific T cell population, and we make a global analysis of the model equations. As in Nowak and Bangham (1996), Nowak *et al.* (1996) and Bonhoeffer *et al.* (1997), we find a threshold parameter R_0 (the basic reproductive number of the virus) which determines the dynamical behavior of the infection. This parameter is further used to account for the efficacy of hepatitis C therapy.

THE MODEL

Before the formulation of the model we remark some facts about the immune response to hepatitis C. Antibodies, cytokines, natural killer cells and T cells are essential components of a normal immune response to virus. For HCV, infected individuals generally develop antibodies reactive with the core (C) protein as well as several nonstructural protein antigens of HCV. However, there is no evidence that HCV antibodies, even when present in high serum titers protect against new cell infections or progression of the disease (Lemon and Brown, 1995).

On the other hand, CD8 + cytotoxic T lymphocytes have been identified in the liver of chronically infected humans and chimpanzees (Lemon and Brown, 1995). These cells are activated by a signal given by the virus to the immune system, either on the surface of the infected cells or on antigen-presenting cells. However, the relative contribution of T cells response to immunity and to disease pathogenesis remains uncertain. It is apparent that they are not capable of eliminating the infection (Lemon and Brown, 1995).

Here, we will consider only T cells response. One of the questions that we want to address by mathematical models is how important is this response on the dynamics of the infection.

Our model contains four variables: healthy liver cells H_s or target cells, infected liver cells H_i , virus load V , and CD8 + cytotoxic T cells. The assumptions are the following.

Healthy liver cells H_s are produced at a constant rate β_s and die at a constant rate μ_s ; H_s cells become infected at a rate proportional to the product of H_s and V , with constant of proportionality k , and once infected die with

a constant rate μ_i ; T cells kill infected cells H_i at a rate proportional to the product of H_i and T , with constant of proportionality δ .

Even when the acute HCV infection appears to be lytic, for chronic HCV infection it is not completely clear whether the virus is intrinsically cytopathic in infected hepatocytes. However, it appears more likely that the liver damage is immunologically mediated, as in chronic hepatitis B. Hepatocellular damage is probably initiated by the activation of virus-specific cytotoxic T cells (Lemon and Brown, 1995). Then it is reasonable to assume that the average life time of infected cells ($1/\mu_i$) is shorter than the average life-time of healthy cells ($1/\mu_s$). Thus, in the following we will assume $\mu_i \geq \mu_s$.

Hepatitis C virions are produced inside the infected cells at a rate of p virions per infected cell per day. On the other hand, viruses die at a *per capita* constant rate μ_v .

In the presence of HCV, supply of new T cells is given by

$$\beta_T V \left(1 - \frac{T}{T_{\max}} \right),$$

where β_T is the rate of growth of T cells, T_{\max} is the maximum T cell population level. On the other hand T cells die at a *per capita* constant rate μ_T .

These assumptions lead to the following differential equations:

$$\begin{aligned} \dot{H}_s &= \beta_s - kH_s V - \mu_s H_s & \dot{H}_i &= kH_s V - \delta H_i T - \mu_i H_i \\ \dot{V} &= pH_i - \mu_v V & \dot{T} &= \beta_T V \left(1 - \frac{T}{T_{\max}} \right) - \mu_T T \end{aligned} \quad (1)$$

All parameters in the model are positive. It is a simple matter to verify that Eq. (1) satisfy the existence and uniqueness conditions. Moreover, the region

$$\begin{aligned} \Omega &= \{(H_s, H_i, V, T) \in \mathbb{R}_+^4 \mid H_s + H_i \leq H_M, V \leq V_M, T \\ &\leq T_M\} \end{aligned}$$

where $H_M = \beta_s / \mu_s$, $V_M = (p / \mu_v) H_M$, and $T_M = (\beta_T / \mu_T) V_M$ with $\mu_T^* = \mu_T + (\beta_T / T_{\max}) V_M$, is positively invariant for system (1), because the vector field on the boundary does not point to the exterior. Therefore, solutions starting in Ω will remain there for $t \geq 0$. In the following we will assume that initial conditions are always given in Ω .

Remark. We observe that H_M is the maximum number of cells in a healthy liver, therefore V_M is maximum virus load supported by an organism. On the other hand $T_M < T_{\max}$ represents the maximum number of T cells generated in an individual with hepatitis C.

EQUILIBRIUM SOLUTIONS

We now show that in Ω there are two possible steady states, one with no virus present, an *uninfected steady*

state, and another with a constant level of virus, an *endemically infected steady state*.

The equilibrium solutions of (1) must satisfy the following algebraic equations.

$$\begin{aligned} 0 &= \beta_s - kH_s V - \mu_s H_s & 0 &= kH_s V - \delta H_i T - \mu_i H_i \\ 0 &= p H_i - \mu_V V & 0 &= \beta_T V \left(1 - \frac{T}{T_{\max}}\right) - \mu_T T. \end{aligned} \quad (2)$$

From the first, third and fourth equations of (2), it can be seen that the equilibrium points satisfy the following relations

$$\begin{aligned} H_s^* &= \frac{\beta_s}{kV^* + \mu_s}, \\ H_i^* &= \frac{\mu_V V^*}{p}, \\ T^* &= \frac{\beta_T T_{\max} V^*}{\beta_T V^* + \mu_T T_{\max}}. \end{aligned}$$

If $V^* = 0$, we obtain the uninfected steady state solution

$$I_0 = \left(\frac{\beta_s}{\mu_s}, 0, 0, 0 \right), \quad (3)$$

in which there is no infection. Consequently, all hepatic cells are healthy and $H_s^* = \beta_s/\mu_s$ is the number of liver cells in a healthy individual.

If $V^* \neq 0$, then substituting H_s^* , H_i^* , and T^* in the second equation of system (2), we obtain after some calculations that V^* must satisfy the following quadratic equation

$$r(V^*) = AV^{*2} + BV^* + C, \quad (4)$$

with coefficients given by

$$\begin{aligned} A &= k\beta_T \mu_V (\delta T_{\max} + \mu_i), \\ B &= -k\beta_s \beta_T p + \delta \beta_T \mu_s \mu_V T_{\max} + k\mu_i \mu_V \mu_T T_{\max} \\ &\quad + \beta_T \mu_i \mu_s \mu_V, \\ C &= \mu_i \mu_s \mu_V \mu_T T_{\max} - k\beta_s p \mu_T T_{\max}. \end{aligned}$$

Now, we see conditions such that Eq. (4) has a solution $0 < V^* < V_M$.

First, note that

$$\begin{aligned} r(V_M) &= \frac{k\beta_s^2 \beta_T p^2 \delta T_{\max}}{\mu_s^2 \mu_V} + \frac{k\beta_s^2 \beta_T p^2}{\mu_s \mu_V} \left[\frac{\mu_i}{\mu_s} - 1 \right] \\ &\quad + \delta \beta_s \beta_T p T_{\max} + \beta_s \beta_T p \mu_i + \mu_i \mu_s \mu_V \mu_T T_{\max} \\ &\quad + k\beta_s p \mu_T T_{\max} \left[\frac{\mu_i}{\mu_s} - 1 \right] \end{aligned}$$

and

$$\begin{aligned} i(V_M) &= \frac{2k\beta_s \beta_T p \delta T_{\max}}{\mu_s} + k\beta_s \beta_T p \left[\frac{2\mu_i}{\mu_s} - 1 \right] \\ &\quad + \delta \beta_T \mu_s \mu_V T_{\max} + k\mu_i \mu_V \mu_T T_{\max} + \beta_T \mu_i \mu_s \mu_V \end{aligned}$$

are bigger than zero since $\mu_i \geq \mu_s$, and all the coefficients are non-negative. Then, the existence of positive solutions of Eq. (4) will depend on the sign of $r(0)$ and $i(0)$. We have the following cases: (a) $r(0) > 0$. In this case it is easy to see that $r(0) = C > 0$ implies $i(0) = B > 0$, therefore Eq. (4) has no solutions $0 < V^* < V_M$. (b) $r(0) = 0$. In this other case we have that $i(0) > 0$, and therefore the only positive root is $V^* = 0$. (c) $r(0) < 0$. Clearly in this case there exists a unique root $0 < V^* < V_M$.

From (2) it is easy to verify that if $0 < V^* < V_M$, then the non-trivial equilibrium $I_1 = (H_s^*, H_i^*, V^*, T^*) \in \Omega$. Note also that $C < 0$ is equivalent to $\frac{k\beta_s p}{\mu_i \mu_s \mu_V} > 1$. Then, summarizing we have the following theorem.

THEOREM 1. *Let R_0 be given by*

$$R_0 \equiv \frac{k\beta_s p}{\mu_i \mu_s \mu_V} \quad (5)$$

If $R_0 \leq 1$, then $I_0 = ((\beta_s/\mu_s), 0, 0, 0)$ is the only equilibrium point in Ω ; if $R_0 > 1$ then the endemically infected equilibrium

$$\begin{aligned} I_1 &= (H_s^*, H_i^*, V^*, T^*) \\ &= \left(\frac{\beta_s}{kV^* + \mu_s}, \frac{\mu_V V^*}{p}, V^*, \frac{\beta_T T_{\max} V^*}{\beta_T V^* + \mu_T T_{\max}} \right) \end{aligned} \quad (6)$$

will also lie in Ω .

The threshold parameter R_0 defined by (5) is called the *basic reproductive number of the virus*. The value of this parameter plays a central role in the dynamics of system (1) with important implications in the treatment of hepatitis C.

The parameter R_0 has an interesting biological meaning: assume that an initial virus load V_0 is introduced in a healthy organism with β_s/μ_s healthy liver cells. These viruses produce in average, $(k\beta_s/\mu_s \mu_V)V_0$ infected cells during their lifespan. Since each infected cell produces p/μ_i virions during its lifespan, $(k\beta_s p/\mu_i \mu_V \mu_s)V_0 = R_0 V_0$ is the average number of new virions produced by the initial virus load V_0 in a healthy organism.

It is worthy to observe that R_0 depends only on six of the ten parameters in the model.

STABILITY OF THE UNINFECTED STEADY STATE

In this section, we study the stability properties of the trivial or *uninfected* equilibrium state I_0 . The Jacobian

matrix $J(H_s, H_i, V, T)$ of system (1) is given by

$$\begin{pmatrix} -kV - \mu_s & 0 & -kH_s & 0 \\ kV & -\delta T - \mu_i & kH_s & -\delta H_i \\ 0 & p & -\mu_V & 0 \\ 0 & 0 & \beta_T \left(1 - \frac{T}{T_{\max}}\right) & -\frac{\beta_T V}{T_{\max}} - \mu_T \end{pmatrix}. \quad (7)$$

Then, the local stability of I_0 is governed by the eigenvalues of the matrix

$$J(I_0) = \begin{pmatrix} -\mu_s & 0 & -k\frac{\beta_s}{\mu_s} & 0 \\ 0 & -\mu_i & k\frac{\beta_s}{\mu_s} & 0 \\ 0 & p & -\mu_V & 0 \\ 0 & 0 & \beta_T & -\mu_T \end{pmatrix}, \quad (8)$$

which clearly are $-\mu_s$, $-\mu_T$ and the roots of the quadratic equation

$$\lambda^2 + (\mu_i + \mu_V)\lambda + \mu_i\mu_V \left(1 - \frac{k\beta_s p}{\mu_s\mu_V\mu_i}\right) = 0. \quad (9)$$

The other two eigenvalues of $J(I_0)$ have negative real part if and only if the coefficients of (9) are positive, and this occurs if and only if $R_0 < 1$. Therefore I_0 is locally asymptotically stable for $R_0 < 1$. We can actually show that it is globally asymptotically stable in Ω for $R_0 \leq 1$. To prove this, we use the Lyapunov function

$$U(H_s, H_i, V, T) = pH_i + \mu_i V. \quad (10)$$

The orbital derivative of U is given by

$$\dot{U} = -\mu_i\mu_V \left[1 - \frac{kpH_s}{\mu_i\mu_V}\right] V - p\delta H_i T. \quad (11)$$

Since $H_s \leq \beta_s/\mu_s$, the expression inside the bracket in (11) is non-negative for $R_0 \leq 1$ and therefore $\dot{U} \leq 0$ in Ω . The subset where $\dot{U} = 0$ is defined by the equations

$$V = 0 \quad H_i T = 0 \quad \text{if } R_0 < 1$$

$$V = 0 \text{ or } H_s = \frac{\beta_s}{\mu_s}, \quad H_i T = 0 \quad \text{if } R_0 = 1.$$

From inspection of system (1), it can be seen that the maximum invariant set contained in $\dot{U} = 0$ is the plane $V = 0$, $H_i = 0$. In this set system (1) becomes

$$\dot{H}_s = \beta_s - \mu_s H_s \quad \dot{H}_i = 0 \quad \dot{V} = 0 \quad \dot{T} = -\mu_T T$$

which implies that solutions started there tend to the equilibrium I_0 as t goes to infinity. Therefore, applying LaSalle–Lyapunov Theorem (Hale, 1969) it follows that I_0 is locally stable and all trajectories starting in Ω approach I_0 . Summarizing, we have proven the following

THEOREM 2. *The uninfected steady state I_0 of system (1) is globally asymptotically stable in the region Ω .*

STABILITY OF THE ENDEMICALLY INFECTED STATE

For $R_0 > 1$, the equilibrium I_0 becomes an unstable hyperbolic point, and the endemically infected equilibrium, I_1 emerges in the region Ω . The local stability of I_1 is given by the Jacobian of (1) evaluated in this point:

$$\begin{pmatrix} -kV^* - \mu_s & 0 & -kH_s^* & 0 \\ kV^* & -\delta T^* - \mu_i & kH_s^* & -\delta H_i^* \\ 0 & p & -\mu_V & 0 \\ 0 & 0 & \beta_T \left(1 - \frac{T^*}{T_{\max}}\right) & -\frac{\beta_T V^*}{T_{\max}} - \mu_T \end{pmatrix},$$

that can be rewritten as

$$\begin{pmatrix} -\frac{\beta_s}{H_s^*} & 0 & -kH_s^* & 0 \\ kV^* & -\frac{kH_s^* p}{\mu_V} & kH_s^* & -\delta H_i^* \\ 0 & p & -\mu_V & 0 \\ 0 & 0 & \frac{\mu_T T^*}{V^*} & -\frac{\beta_T V^*}{T^*} \end{pmatrix}$$

when we take into account the identities:

$$kV + \mu_s = \frac{\beta_s}{H_s^*}, \quad \delta T^* + \mu_i = \frac{kH_s^* V^*}{H_i^*},$$

$$\frac{V^*}{H_i^*} = \frac{p}{\mu_V}, \quad \delta T^* + \mu_i = \frac{kH_s^* p}{\mu_V},$$

$$\beta_T \left(1 - \frac{T^*}{T_{\max}}\right) = \frac{\mu_T T^*}{V^*}, \quad \frac{\beta_T V^*}{T_{\max}} + \mu_T = \frac{\beta_T V^*}{T^*},$$

which are obtained from system (3). Hence, the characteristic polynomial of the linearized system is given by

$$P(\lambda) = \det \begin{pmatrix} -\frac{\beta_s}{H_s^*} - \lambda & 0 & -kH_s^* & 0 \\ kV^* & -\frac{kH_s^* p}{\mu_V} - \lambda & kH_s^* & -\delta H_i^* \\ 0 & p & -\mu_V - \lambda & 0 \\ 0 & 0 & \frac{\mu_T T^*}{V^*} & -\frac{\beta_T V^*}{T^*} - \lambda \end{pmatrix}$$

After some calculations we obtain

$$P(\lambda) = \lambda^4 + a_1\lambda^3 + a_2\lambda^2 + a_3\lambda + a_4 \quad (12)$$

with

$$\begin{aligned} a_1 &= \mu_V + \frac{\beta_s}{H_s^*} + \frac{\beta_T V^*}{T^*} + \frac{kH_s^* p}{\mu_V}, \\ a_2 &= \frac{\beta_s}{H_s^*} \left[\frac{\beta_T V^*}{T^*} + \left(\mu_V + \frac{kH_s^* p}{\mu_V} \right) \right] \\ &\quad + \frac{\beta_T V^*}{T^*} \left(\mu_V + \frac{kH_s^* p}{\mu_V} \right), \\ a_3 &= \delta\mu_V \mu_T T^* + k^2 p H_s^* V^* \\ &\quad + \frac{\beta_s \beta_T V^*}{H_s^* T^*} \left(\mu_V + \frac{kH_s^* p}{\mu_V} \right), \\ a_4 &= \delta\mu_V \mu_T T^* \frac{\beta_s}{H_s^*} + \frac{k^2 p \beta_T H_s^* (V^*)^2}{T^*}. \end{aligned} \quad (13)$$

Using the Routh–Hurwitz criteria (Gantmacher, 1960), the local stability of the endemic equilibrium I_1 will be established if we show that

$$\Delta_3 = \begin{vmatrix} a_1 & 1 & 0 \\ a_3 & a_2 & a_1 \\ 0 & a_4 & a_3 \end{vmatrix} = (a_1 a_2 - a_3) a_3 - a_1^2 a_4 > 0, \quad (14)$$

since the coefficients a_1, a_2, a_3 and a_4 of the characteristic polynomial $P(\lambda)$ are all positive.

To see that condition (14) is satisfied, it is convenient to adopt the following notation:

$$\begin{aligned} A &= \frac{\beta_s}{H_s^*}, \quad B = \frac{\beta_T V^*}{T^*}, \quad C = \frac{kH_s^* p}{\mu_V}, \quad D = \delta\mu_V \mu_T T^* \\ E &= k\mu_V V^*, \quad F = C + \mu_V, \quad G = \frac{\beta_T V^*}{T_{\max}}. \end{aligned}$$

In terms of the variables above we rewrite the coefficients a_1, a_2, a_3 and a_4 as

$$\begin{aligned} a_1 &= A + B + F, \\ a_2 &= AB + aF + BF, \\ a_3 &= ABF + CE + D, \\ a_4 &= AD + BCE. \end{aligned}$$

After tedious calculations we obtain the following expression

$$\begin{aligned} &(a_1 a_2 - a_3) a_3 - a_1^2 a_4 \\ &= A^3 B^2 F + A^2 B D + A^2 C E F + A^2 B^3 F \\ &\quad + B^2 D F + A^3 (B F^2 - D) + B^3 (A F^2 - C E) \\ &\quad + 2A^2 B (B F^2 - D) - A B^2 C E + A B F (B F^2 - D) \\ &\quad + A B F (A F^2 - C E) - A^2 D F - B^2 C E F \\ &\quad + C E (A F^2 - C E) + D (B F^2 - D) - 2C E D. \end{aligned} \quad (15)$$

On the other hand, from the equations in equilibrium (3) we obtain the relations:

$$A = \frac{E}{\mu_V} + \mu_s, \quad D = C\mu_V \mu_T - \mu_i \mu_T \mu_V,$$

and from them it is easy to see that the following inequalities are satisfied

$$A F^2 > 2E C, \quad B F^2 > 2D, \quad B F^2 > B C^2. \quad (16)$$

Now, from (16) we have the following inequalities

$$\begin{aligned} &2A^2 B (B F^2 - D) - A B^2 C E \\ &= A^2 B (B F^2 - 2D) + A B^2 (A F^2 - C E) > 0 \end{aligned} \quad (17)$$

$$\begin{aligned} &A B F (B F^2 - D) + A B F (A F^2 - C E) - A^2 D F \\ &\quad - B^2 C E F \\ &= A B F^3 (A + B) - A B F (C E + D) - F (A^2 D \\ &\quad + B^2 C E) \\ &> A B F^3 (A + B) - A B F (C E + D) \\ &\quad - F \left(\frac{A^2 B F^2}{2} + \frac{B^2 A F^2}{2} \right) \\ &= \frac{A B F^3 (A + B)}{2} - A B F (C E + D) \\ &= A B F \left(\frac{A F^2}{2} - C E \right) + A B F \left(\frac{B F^2}{2} - D \right) > 0, \end{aligned} \quad (18)$$

and

$$\begin{aligned} &D (B F^2 - D) + C E (A F^2 - C E) - 2C D E \\ &> D^2 + (C E)^2 - 2C D E = (D - C E)^2 \geq 0 \end{aligned} \quad (19)$$

From (17)–(19) it follows inequality (14). Therefore we have proved the following theorem.




FIGURE 1 Numerical solution of model (1). The graphs show healthy liver cells, infected liver cells virus load and T cells versus time. The parameters in the simulation are: $\beta_s/\mu_s = 5,000$, $\mu_s = \mu_T = 0.02/\text{day}$, $\mu_i = 0.5/\text{day}$, $\mu_v = 5/\text{day}$, $k = 0.00003$, $p = 100$ virions per millimeter per cell per day, $\delta = 0.00001$, $\beta_T = 0.0003$. In this case $R_0 = 0.6$.

THEOREM 3. For $R_0 >$ the endemically infected state I_1 is in Ω and it is locally asymptotically stable.

DISCUSSION

Starting with a description of healthy and infected liver cells H_s , H_i , virus load V and virus specific T cells, we have developed a model for hepatitis C dynamics. While our model is overly simple in that it does not account for the immune response to HCV infection or mechanisms of cell death other than killing by cytotoxic T cells, it has some interesting predictions.

The basic reproductive number of the virus, R_0 has been used largely in understanding the persistence of viral infections within individuals (see for example Nowak,

1996 and Bonhoeffer *et al.*, 1997), and in the population. For this model, $R_0 = \beta_s k p / \mu_i \mu_s \mu_v$ and it has to be above one for successful chronic HCV infection. If $R_0 \leq 1$, then the level of virus load and infected cells will monotonically decrease and ultimately be eliminated. This decrease may be due to the fact, that virus does not infect enough cells, or infected cells die without producing a sufficient number of viral progeny. In this aspect, the model is similar to epidemiological models in which infected individuals must infect at least a critical number of individuals for an epidemic to occur. As in epidemiological models, we have two steady states, an uninfected steady state where the virus, infected cells and reactive T cells are not present; and an endemically infected steady state where all four populations of the model are maintained.




FIGURE 2 Numerical solution of model (1). The graphs show healthy liver cells, infected liver cells, virus load and T cells versus time. The parameters in the simulation are: $\beta_s/\mu_s = 5,000$, $\mu_s = \mu_T = 0.02/\text{day}$, $\mu_i = 0.5/\text{day}$, $\mu_v = 5/\text{day}$, $k = 0.00003$, $p = 200$ virions per millimeter per cell per day, $\delta = 0.00001$, $\beta_T = 0.0003$. In this case $R_0 = 1.2$.

Figures. 1 and 2 illustrate the typical behavior of the healthy and infected liver cells, virus load and T cells initially positive. In Fig. 1 $R_0 = 0.6$, whereas in Fig. 2, $R_0 = 1.2$. Notice that temporal courses of the liver cells and the virus load present damped oscillations.

One interesting feature of our model is that R_0 does not depend on the T cell immune response of the organism. It appears that activation of virus-specific T cells are more related with the liver damage in chronic hepatitis C (Lemon and Brown, 1995), and it is probably that the effect of the immune response is reflected in the mortality rate of infected cells μ_i .

Clinical studies has been done to document the efficacy of recombinant IFN- α in the treatment of chronic hepatitis C (Lemon and Brown, 1995; Poynard *et al.*, 1998). Although IFN has been approved for treatment of chronic hepatitis C, it has been showed that it is successful in only

11–30% of the cases (Neumann *et al.*, 1998), and the mechanism of action is not well understood. In a recent paper, Neumann *et al.* (1998) analyzed this problem. They hypothesized that IFN acts by blocking the production or release of virions rather than by blocking the novo infections. Using mathematical analysis coupled with clinical studies, they corroborated their hypothesis, and also they estimated the efficacy of IFN therapy, that is, the percentage of HCV production blocked by different doses of IFN. Furthermore, they estimated the infected cell rate.

In this paper, we use a more theoretical point of view to analyze the effects of the current treatment with IFN. It is clear that in order to control the disease (i.e. the virus load declines to zero), we have to reduce R_0 below the value one, and this can be achieved reducing k or p below critical values or increasing μ_v or μ_i above critical values. It is worth to note that R_0 is directly proportional to k and p , and inversely




FIGURE 3 V^* vs. k , p , respectively. The initial values of the parameters for $V^* = 0$ are: $\beta_s/\mu_s = 5,000$, $\mu_s = \mu_T = 0.02/\text{day}$, $\mu_i = 0.5/\text{day}$, $\mu_V = 5/\text{day}$, $k = 0.00003$, $p = 100$ virions per millimeter per cell per day, $\delta = 0.00001$, $\beta_T = 0.0003$.

proportional to μ_i , μ_V , which implies that increasing the mortality of the infected cells and the free virus could be a mechanism to achieve a fast decrease in the virus load. Of course, this will depend on the range of the parameters.

Combination therapies that reduce the rate of “de novo” infections k , production rate of virions p and at the same time rise the mortality of infected cells or virus, could be much more effective than the current therapy with IFN, which is assumed that essentially blocks production of new virions. Recent reports have suggested that ribavirin in combination with IFN is more effective than the treatment with only IFN (see Poynard *et al.*, 1998). Studies on combination therapy are under way, but results are still inconclusive.

The model can be used to study the relation between the parameters involved in the infection and the endemic virus load. Endemic virus load is directly correlated with k and p

(Fig. 3); and inversely correlated to μ_i and μ_V (Fig. 4). These correlations suggest that immune control through faster killing of infected cells or free virus may have an important role in lowering HCV load.

Also, Figs. 3 and 4 show that the magnitude of the variation virus load is sensible to the values of the parameters. Virus load increases for small values of k and p , but saturates for large values of these parameters. On the other hand, for small values of μ_i and μ_V , virus load decreases rapidly, but for larger values it remains almost constant. This suggests that strong or weak response to treatments depends on the state of viral, infection and immune parameters, and the success of the therapy could possibly be predicted from the early knowledge of these parameters in the patients.

Substituting the endemic values $V^*/H_i^* = \frac{p}{\mu_V}$ in the expression $R_0 = 1$ we obtain the equivalent threshold




FIGURE 4 V^* vs. μ_i and μ_V respectively. The initial values of the parameters for $V^* = 0$ are: $\beta_s/\mu_s = 5,000$, $\mu_s = \mu_T = 0.02/\text{day}$, $\mu_i = 0.5/\text{day}$, $\mu_V = 5/\text{day}$, $k = 0.00003$, $p = 100$ virions per millimeter per cell per day, $\delta = 0.00001$, $\beta_T = 0.0003$.

condition for HCV infection

$$\frac{V^*}{\bar{H}_i^*} = \frac{\mu_s}{k} \quad (20)$$

where $\bar{H}_i^* = \frac{H_i^*}{\beta_s/\mu_s}$ is the proportion of infected cells. Theoretically, this means that in order to clear the infection, the quotient of the virus load and the proportion of infected cells has to be less or equal than the right hand side of Eq. (20). Now, V^* and \bar{H}_i^* could be obtained from clinical tests and k and μ_s could be estimated from data (see Neumann *et al.*, 1998) for an estimation of μ_s) in order to test the feasibility of this model.

Acknowledgements

We are grateful to the anonymous referees for their careful reading that helped us to improve the paper. LEP acknowledges support from CONACYT grant.

References

- Bonhoeffer, S., May, R.M., Shaw, G.M. and Nowak, M.A. (1997) "Virus dynamics and drug therapy", *Proc. Natl Acad. Sci. USA* **94**, 6971–6976.
- Gantmacher, F.R. (1960) *The Theory of Matrices* **Vol. 2**.
- Hale, J.K. (1969) *Ordinary Differential Equations* (Wiley, New York).
- Lemon, S.M. and Brown, E.A. (1995) "Hepatitis C Virus", In: Mandell, G.L., Bennet, J.E. and Dolin, R., eds, *Principles and Practice of Infectious Diseases*, 4th Ed. (Churchill Livingstone Inc., New York), pp 1474–1483.
- Neumann, A.U., Lam, N.P., Dahari, H., Gretch, D.R., Wiley, T.E., Layden, T.J. and Perelson, A.S. (1998) "Hepatitis C viral dynamics in vivo and the antiviral efficacy of interferon- α therapy", *Science* **282**, 103–107.
- Nowak, M.A. and Bangham, C.R.M. (1996) "Population dynamics of immune responses to persistent viruses", *Science* **272**, 74–79.
- Nowak, M.A., Bonhoeffer, S., Hill, A.M., Boehme, R., Thomas, H.C. and McDade, H. (1996) "Viral dynamics in hepatitis B virus infection", *Proc. Natl Acad. Sci. USA* **93**, 4398–4402.
- Payne, R.J.H., Nowak, M.A. and Blumberg, B.S. (1996) "The dynamics of hepatitis B virus infection", *Proc. Natl Acad. Sci. USA* **93**, 6542–6546.

Poynard, T., Marcellin, P., Lee, S.S., Niederau, Ch., Minuk, G.S., Ideo, G., Bain, V., Heathcote, J., Zeuzmen, S., Trepo, Ch. and Albrecht, J. (1998) "Randomized trail of interferon α -2b plus ribavirin for 48 weeks or 24 weeks versus interferon α 2b plus placebo for

48 weeks for treatment of chronic infection with hepatitis C virus", *Lancet* **352**, 1426–1432.
Purcell, R.H. (1994) "Hepatitis viruses: Changing patterns of human disease", *Proc. Natl Acad. Sci. USA* **91**, 2401–2406.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

