

Research Article

Dynamics Analysis and Simulation of a Modified HIV Infection Model with a Saturated Infection Rate

Qilin Sun¹ and Lequan Min^{1,2}

¹ School of Automation and Electrical Engineering, University of Science and Technology Beijing, Beijing 100083, China

² School of Mathematics and Physics, University of Science and Technology Beijing, Beijing 100083, China

Correspondence should be addressed to Lequan Min; minlequan@gmail.com

Received 17 December 2013; Accepted 28 January 2014; Published 23 March 2014

Academic Editor: Yang Wei

Copyright © 2014 Q. Sun and L. Min. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper studies a modified human immunodeficiency virus (HIV) infection differential equation model with a saturated infection rate. It is proved that if the basic virus reproductive number R_0 of the model is less than one, then the infection-free equilibrium point of the model is globally asymptotically stable; if R_0 of the model is more than one, then the endemic infection equilibrium point of the model is globally asymptotically stable. Based on the clinical data from HIV drug resistance database of Stanford University, using the proposed model simulates the dynamics of the two groups of patients' anti-HIV infection treatment. The numerical simulation results are in agreement with the evolutions of the patients' HIV RNA levels. It can be assumed that if an HIV infected individual's basic virus reproductive number $R_0 < 1$ then this person will recover automatically; if an antiretroviral therapy makes an HIV infected individual's $R_0 < 1$, this person will be cured eventually; if an antiretroviral therapy fails to suppress an HIV infected individual's HIV RNA load to be of unpredictable level, the time that the patient's HIV RNA level has achieved the minimum value may be the starting time that drug resistance has appeared.

1. Introduction

The human immunodeficiency virus (HIV) mainly targets a host's $CD4^+$ T cells. Chronic HIV infection causes gradual depletion of the $CD4^+$ T cell pool and thus progressively compromises the hosts immune response to opportunistic infections, leading to Acquired Immunodeficiency Syndrome (AIDS) [1].

In recent years, there is much work done on HIV infection from different points of view, such as pathology [2], microbiology [3], and mathematics [4–7]. Mathematical models have become essential tools to make assumptions, suggest new experiments, or help easily explain complex processes [8]. The basic mathematical model widely used for studying the dynamics of HIV infection has the following form [4, 9]:

$$\begin{aligned}\dot{x} &= \lambda - d_1x - k_1xv, \\ \dot{y} &= k_1xv - d_2y, \\ \dot{v} &= ay - d_3v,\end{aligned}\tag{1}$$

where $x(t)$, $y(t)$, and $v(t)$ are the number of uninfected cells, infected cells, and free virus, respectively. Uninfected cells are produced at a constant rate λ , die at rate d_1x , and become infected at rate k_1xv . Infected cells are produced at rate k_1xv and die at rate d_2y . Free virus is produced from infected cells at rate ay and dies at rate d_3v .

Equation (1) has a basic virus reproductive number $R_0 = ak_1\lambda/(d_1d_2d_3)$. According to Nowak and Bangham [4], R_0 is defined as the number of newly infected cells arising from any one infected cell; if R_0 is smaller than 1, then in the beginning of the infection, each virus infected cell produces on average less than one newly infected cell. Thus, the infection cannot spread, and the system returns to the uninfected state; if R_0 is larger than 1, then initially each virus infected cell produces on average more than one newly infected cell. The infected cell population will increase, whereas the uninfected cell population will decline and therefore provide less opportunity for the virus to infect new cells.

There is a discussion about the process of the HIV RNA transcribing into DNA: when an HIV enters a resting $CD4^+$ T cell, the HIV RNA may not be completely reverse transcribed

into DNA [10]. A proportion of resting infected cells can revert to the uninfected state before the viral genome is integrated into the genome of the lymphocyte [11].

Recently, some mathematical models of HIV infection have been proposed based on the assumption that a fraction of infected $CD4^+$ T cells return to the uninfected class [12–14]. Srivastava and Chandra [13] have considered a model with three populations: uninfected $CD4^+$ T cells (x), infected $CD4^+$ T cells (y), and HIV population (v). The model has the following form:

$$\begin{aligned}\dot{x} &= \lambda - d_1x - k_1xv + py, \\ \dot{y} &= k_1xv - (d_2 + p)y, \\ \dot{v} &= ay - d_3v,\end{aligned}\quad (2)$$

where the meanings of the variables $x(t)$, $y(t)$, and $v(t)$ and the parameters λ , d_1 , k_1 , d_2 , a , and d_3 are the same as those given in (1). The term py is the rate of infected cells in the latent stage reverting to the uninfected class. Equation (2) also has a basic virus reproductive number $R_0 = ak_1\lambda/(d_3d_1(p + d_2))$. They have proved that if $R_0 \leq 1$, the infection-free steady state of (2) is globally asymptotically stable; if $R_0 > 1$, the endemic steady state of (2) is globally asymptotically stable [13].

In (2), the mass action term k_1xv used to model infection of $CD4^+$ T cells by free virions is biologically problematic for several reasons. Firstly, since λ/d_1 represents the total number of $CD4^+$ T cells in the basic virus reproductive number $R_0 = ak_1\lambda/(d_3d_1(p + d_2))$, this causes R_0 to depend upon the total number of $CD4^+$ T cells in vivo. This implies the dubious prediction that individuals with more $CD4^+$ T cells will be more easily infected than individuals with less $CD4^+$ T cells. Secondly, the rate of HIV infection is assumed to be bilinear by the mass action term k_1xv . However, the actual incidence rate is probably not linear over the entire range of virus $v(t)$ and uninfected $CD4^+$ T cells $x(t)$ [15–17].

On biological grounds, during primary HIV infection, the rate of virus infection should be approximately proportionate to the virus load k_1v because of a small amount of viral load with respect to a large number of $CD4^+$ T cells. However, since the total number of healthy $CD4^+$ T cells in vivo is limited, the HIV infection will approach saturation with more and more virus produced. In this case, it is more reasonable to assume that the rate of virus infection should be approximately proportionate to the number of healthy $CD4^+$ T cells k_1x .

Based on the argument above, this paper describes an amended model. In this model, we use a saturated infection rate $k_1xv/(x + v)$ to replace the mass action term k_1xv in (2). Under the formulation of this saturated infection rate, the basic virus reproductive number R_0 is independent of the total number of $CD4^+$ T cells. Meanwhile, the actual incidence rate is not linear over the entire range of virus $v(t)$ and uninfected $CD4^+$ T cells $x(t)$ any more. The global stabilities of the infection-free state and the endemic infection state of the modified HIV infection model have been discussed. Based on the clinical data from HIV drug resistance database of Stanford University, using the proposed model simulates

the dynamics of two groups of patients' anti-HIV infection treatment, and then make long-term predictions for the two groups' anti-HIV infection treatment, respectively.

The rest of this paper is organized as follows. Section 2 introduces a modified model and discusses the boundedness of the solutions of the model. Sections 3 and 4 discuss the global stability of the infection-free state and the endemic infection state of the modified HIV infection model, respectively. Section 5 simulates the dynamics of two groups of patients' anti-HIV infection treatment. Section 6 summarizes this paper.

2. Modified HIV Infection Model

2.1. The Modified HIV Infection Model. Based on (2), our modified HIV infection model has the following form:

$$\begin{aligned}\dot{x} &= \lambda - d_1x - \frac{k_1xv}{x + v} + py, \\ \dot{y} &= \frac{k_1xv}{x + v} - (d_2 + p)y, \\ \dot{v} &= ay - d_3v,\end{aligned}\quad (3)$$

where the meanings of the variables $x(t)$, $y(t)$, and $v(t)$ and the parameters λ , d_1 , k_1 , d_2 , p , a , and d_3 are the same as those given in (2). Equation (3) has two steady states:

(1) The infection-free steady state

$$Q_1 = (x_0, 0, 0) \quad (4)$$

represents the virus infection free. Q_1 is called infection-free equilibrium point. Here,

$$x_0 = \frac{\lambda}{d_1}. \quad (5)$$

(2) The endemic infected steady state

$$Q_2 = (\bar{x}, \bar{y}, \bar{v}) \quad (6)$$

represents persistent virus infection. Q_2 is called endemic infection equilibrium point. Here,

$$\begin{aligned}\bar{x} &= \frac{\lambda R_0}{k_1(R_0 - 1) + d_1R_0 - p(R_0 - 1)R_0(d_3/a)}, \\ \bar{y} &= \frac{d_3}{a}(R_0 - 1)\bar{x}, \quad \bar{v} = \frac{a\bar{y}}{d_3} = (R_0 - 1)\bar{x}.\end{aligned}\quad (7)$$

Here,

$$R_0 = \frac{ak_1}{d_3(d_2 + p)}. \quad (8)$$

Since the total rate of disappearance of infected cells is $d_2 + p$, infected cells live on average for time $1/(d_2 + p)$. Each infected cell produces virus at rate a . Thus, each infected cell produces

on average a total of $a/(d_2 + p)$ viruses. Since virus dies at rate d_3 per virion, each virus survives on average for time $1/d_3$. During the time $1/d_3$, each virus infects on average $k_1 x_0/(x_0 + v_0)d_3$ cells, where x_0 and v_0 are the preinfection target cells' density and viruses' density, respectively. Thus, the total number of cells infected by the $a/(d_2 + p)$ viruses is $ak_1 x_0/(x_0 + v_0)d_3(d_2 + p)$. According to (4) and (5), $x_0 = \lambda/d_1$ and $v_0 = 0$ at the preinfection steady state. Then one can obtain that the total number of cells infected by each infected cell is $ak_1/d_3(d_2 + p)$. Hence, R_0 is the basic virus reproductive number of (3) which is independent of the total number of the uninfected $CD4^+$ T cells.

According to (4), (6), and (7), if $R_0 \leq 1$, then Q_1 is the unique infection-free equilibrium point; if $R_0 > 1$, then, in addition to the infection-free equilibrium point, (3) has another equilibrium point Q_2 .

2.2. Boundedness of Solutions. It is easy to show that the solutions of (3) with initial conditions $x(0) > 0$, $y(0) > 0$, and $v(0) > 0$ have all positive components for $t > 0$. Hence, one begins the analysis of (3) by observing the nonnegative octant

$$D = \{(x, y, v) \in R_+^3 : x \geq 0, y \geq 0, v \geq 0\}. \quad (9)$$

According to the first two equations of (3), one can get

$$\dot{x} + \dot{y} = \lambda - d_1 x - d_2 y \leq \lambda - d(x + y), \quad (10)$$

$$d = \min(d_1, d_2),$$

and then

$$x + y \leq \frac{\lambda}{d}. \quad (11)$$

So $x(t)$ and $y(t)$ are bounded. From the last equation of (3), it follows that

$$\dot{v} = ay - d_3 v \leq \frac{a\lambda}{d} - d_3 v, \quad (12)$$

and then

$$v \leq \frac{a\lambda}{dd_3}. \quad (13)$$

So $v(t)$ are bounded. Hence there is a bounded subset of D :

$$\Omega = \left\{ (x, y, v) \in R_+^3 : 0 \leq x + y \leq \frac{\lambda}{d}, 0 \leq v \leq \frac{a\lambda}{dd_3} \right\} \quad (14)$$

such that any solution trajectory $(x(t), y(t), v(t))$ of (3) with initial value $(x(0), y(0), v(0))$ in Ω will keep in the subset Ω .

According to (7), $\bar{x} > 0$, $\bar{y} > 0$, and $\bar{v} > 0$. One can get that the endemic infection equilibrium point Q_2 exists in the interior of Ω :

$$\Omega^0 = \left\{ (x, y, v) \in \Omega : 0 < x, 0 < y, 0 < x + y < \frac{\lambda}{d}, \right. \quad (15)$$

$$\left. 0 < v < \frac{a\lambda}{dd_3} \right\}.$$

Therefore, the stability of the endemic infection equilibrium point Q_2 only needs to be discussed in Ω^0 . \square

3. Stability of the Infection-Free Equilibrium Point Q_1

In this section, we discuss locally asymptotical stability and globally asymptotical stability of the infection-free equilibrium point Q_1 of (3).

3.1. Locally Asymptotical Stability of the Infection-Free Equilibrium Point Q_1

Theorem 1. *If $R_0 = ak_1/(d_3(d_2 + p)) < 1$, then the infection-free equilibrium point Q_1 of (3) is locally asymptotically stable. If $R_0 > 1$, then the infection-free equilibrium point Q_1 is unstable.*

Proof. The Jacobi matrix of (3) at an arbitrary point is given by

$$J(x, y, v) = \begin{bmatrix} -d_1 - a_1 & p & -a_2 \\ a_1 & -p - d_2 & a_2 \\ 0 & a & -d_3 \end{bmatrix}, \quad (16)$$

where $a_1 = k_1 v^2/(x + v)^2$ and $a_2 = k_1 x^2/(x + v)^2$.

Substituting the equilibrium point Q_1 into matrix (16) gives

$$J_{Q_1} = \begin{bmatrix} -d_1 & p & -k_1 \\ 0 & -p - d_2 & k_1 \\ 0 & a & -d_3 \end{bmatrix}. \quad (17)$$

The corresponding eigenequation is

$$|\lambda E - J_{Q_1}| = \begin{vmatrix} \lambda + d_1 & -p & k_1 \\ 0 & \lambda + p + d_2 & -k_1 \\ 0 & -a & \lambda + d_3 \end{vmatrix} = 0. \quad (18)$$

Solving

$$|\lambda E - J_{Q_1}| = (\lambda + d_1)[(\lambda + p + d_2)(\lambda + d_3) - ak_1] = 0 \quad (19)$$

gives

$$\lambda_1 = -d_1 < 0, \quad (20)$$

$$(\lambda + d_2 + p)(\lambda + d_3) - ak_1 k_3 = 0. \quad (21)$$

Equation (21) can be written as

$$\lambda^2 + (d_2 + p + d_3)\lambda + d_3(p + d_2) - ak_1 = 0. \quad (22)$$

\square

Solving equation (22) gives

$$\begin{aligned}\lambda_2 &= \frac{-(d_2 + p + d_3) - \sqrt{(d_2 + p + d_3)^2 - 4[d_3(p + d_2) - ak_1]}}{2} \\ &= \frac{-(d_2 + p + d_3) - \sqrt{(d_2 + p + d_3)^2 - 4d_3(p + d_2)(1 - R_0)}}{2}, \\ \lambda_3 &= \frac{-(d_2 + p + d_3) + \sqrt{(d_2 + p + d_3)^2 - 4[d_3(p + d_2) - ak_1]}}{2} \\ &= \frac{-(d_2 + p + d_3) + \sqrt{(d_2 + p + d_3)^2 - 4d_3(p + d_2)(1 - R_0)}}{2}.\end{aligned}\quad (23)$$

If $R_0 < 1$, then $\lambda_2 < 0$ and $\lambda_3 < 0$. Hence the infection-free equilibrium point Q_1 is locally asymptotically stable. If $R_0 > 1$, then $\lambda_3 > 0$ such that the infection-free equilibrium point Q_1 is unstable.

3.2. Globally Asymptotical Stability of the Infection-Free Equilibrium Point Q_1

Theorem 2. *If $R_0 < 1$, then the infection-free equilibrium point Q_1 of (3) is globally asymptotically stable in Ω .*

Proof. Define a global Lyapunov function by

$$V_1(x, y, v) = y + \frac{(d_2 + p)v}{a}. \quad (24)$$

The derivative of $V_1(x, y, v)$ along the positive solutions of (3) is

$$\begin{aligned}\dot{V}_1 &= \dot{y} + \frac{(d_2 + p)\dot{v}}{a} \\ &= \frac{k_1 x v}{x + v} - d_2 y - p y + (d_2 + p)y \\ &\quad - \frac{d_3(d_2 + p)v}{a} \\ &\leq k_1 v - \frac{d_3(d_2 + p)v}{a} \\ &= \left[\frac{ak_1}{d_3(d_2 + p)} - 1 \right] \frac{d_3(d_2 + p)}{a} v \\ &= (R_0 - 1) \frac{d_3(d_2 + p)}{a} v.\end{aligned}\quad (25)$$

If $R_0 < 1$, then $\dot{V}_1 \leq 0$ holds in Ω . Moreover, $\dot{V}_1 = 0$ if and only if $v = 0$. Hence, the largest compact invariant set in Ω is

$$E_1 = \{(x, y, v) \in \Omega \mid \dot{V}_1 = 0\} = \{(x, y, v) \in \Omega \mid v = 0\}. \quad (26)$$

According to the LaSalle's invariance principle, $\lim_{t \rightarrow +\infty} v(t) = 0$. Then one can get limit equations:

$$\begin{aligned}\dot{x} &= \lambda - d_1 x + p y, \\ \dot{y} &= -d_2 y - p y.\end{aligned}\quad (27)$$

Define a global Lyapunov function by

$$V_2(x, y) = x - x_0 - x_0 \ln \frac{x}{x_0} + y, \quad (28)$$

where

$$\lambda = d_1 x_0, \quad Q_1 = (x_0, 0, 0). \quad (29)$$

The derivative of $V_2(x, y)$ along the positive solutions of (27) is

$$\begin{aligned}\dot{V}_2 &= \dot{x} + \dot{y} - \frac{x_0}{x} \dot{x} \\ &= \lambda - d_1 x + p y - d_2 y - p y - \frac{x_0}{x} (\lambda - d_1 x + p y).\end{aligned}\quad (30)$$

Since $\lambda = d_1 x_0$,

$$\begin{aligned}\dot{V}_2 &= d_1 x_0 - d_1 x - \frac{x_0}{x} d_1 x_0 + d_1 x_0 - \frac{x_0}{x} p y - d_2 y \\ &= d_1 x_0 \left[2 - \frac{x}{x_0} - \frac{x_0}{x} \right] - \left(\frac{x_0}{x} p + d_2 \right) y.\end{aligned}\quad (31)$$

Since the arithmetic mean is greater than or equal to the geometric mean, we obtain $2 - (x/x_0) - (x_0/x) \leq 0$.

Therefore, $\dot{V}_2 \leq 0$ holds in E_1 , and $\dot{V}_2 = 0$ if and only if $x = x_0$ and $y = 0$. There is the largest compact invariant set in E_1 :

$$\begin{aligned}E_2 &= \{(x, y, v) \in E_1 \mid \dot{V}_2 = 0\} \\ &= \{(x, y, v) \in E_1 \mid x = x_0, y = 0\} \\ &= \{Q_1\}.\end{aligned}\quad (32)$$

Hence if $R_0 < 1$, all solution paths in Ω approach the infection-free equilibrium point Q_1 . \square

4. Stability of the Endemic Infection Equilibrium Point Q_2

In this section, we analyze local asymptotical stability and global asymptotical stability of the endemic infection equilibrium point Q_2 of (3).

4.1. Locally Asymptotical Stability of the Endemic Infection Equilibrium Point Q_2

Theorem 3. *If $R_0 > 1$, then the endemic infection equilibrium point Q_2 of (3) is locally asymptotically stable.*

Proof. Put the equilibrium point Q_2 into matrix (16); then one obtains

$$J_{Q_2} = \begin{bmatrix} -d_1 - \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & p & -\frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & -p - d_2 & \frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ 0 & a & -d_3 \end{bmatrix},$$

$$|\lambda E - J_{Q_2}| = \begin{vmatrix} \lambda + d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & -p & \frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ -\frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & \lambda + p + d_2 & -\frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ 0 & -a & \lambda + d_3 \end{vmatrix} = 0. \quad (33)$$

Solving the eigenequation of the matrix above, here is

$$\begin{aligned} & \lambda^3 + \underbrace{\left[d_1 + d_2 + d_3 + p + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right]}_{a_1} \lambda^2 \\ & + \underbrace{\left[d_3 (d_2 + p) + \left(d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) (d_2 + p + d_3) - \frac{pk_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} - \frac{ak_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \right]}_{a_2} \lambda \\ & + \underbrace{d_3 (d_2 + p) \left(d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) + \frac{ak_1^2 \bar{x}^2 \bar{v}^2}{(\bar{x} + \bar{v})^2} - \frac{pd_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} - \left(d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) \frac{ak_1 \bar{x}^2}{(\bar{x} + \bar{v})^2}}_{a_3} = 0. \end{aligned} \quad (34)$$

If $R_0 > 1$, one obtains that

$$\begin{aligned} a_1 &= d_1 + d_2 + d_3 + p + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} > 0, \\ a_2 &= d_3 (d_2 + p) + \left(d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) (d_2 + p + d_3) \\ &\quad - \frac{pk_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} - \frac{ak_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ &= \frac{ak_1}{R_0} + d_1 (d_2 + d_3 + p) \\ &\quad + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} (d_2 + d_3) - \frac{ak_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ &= \frac{ak_1}{R_0} - \frac{ak_1 \bar{x}^2}{R_0^2 \bar{x}^2} + d_1 (d_2 + d_3 + p) \\ &\quad + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} (d_2 + d_3) \\ &= \frac{ak_1 (R_0 - 1)}{R_0^2} + d_1 (d_2 + d_3 + p) \\ &\quad + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} (d_2 + d_3) > 0, \\ a_3 &= d_3 (d_2 + p) \left(d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) \\ &\quad + \frac{ak_1^2 \bar{x}^2 \bar{v}^2}{(\bar{x} + \bar{v})^2} - \frac{pd_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \\ &\quad - \left(d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) \frac{ak_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \end{aligned}$$

$$= d_1 d_3 (d_2 + p) - \frac{ad_1 k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} + \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2}$$

$$= d_1 \frac{ak_1 (R_0 - 1)}{R_0^2} + \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} > 0,$$

$$\begin{aligned} a_1 a_2 - a_3 &= \left(d_2 + d_3 + p + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) \\ &\quad \times \frac{ak_1 (R_0 - 1)}{R_0^2} + d_1 (d_2 + d_3 + p) \\ &\quad \times \left(d_2 + d_3 + p + d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) \\ &\quad + \frac{d_2 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \left(d_2 + d_3 + p + d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) \\ &\quad + \frac{d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \left(d_3 + p + d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right) > 0. \end{aligned} \quad (35)$$

Hence all inequalities of the Routh-Hurwitz criterion are satisfied. Therefore, the endemic infection equilibrium point Q_2 is locally asymptotically stable. \square

4.2. Globally Asymptotical Stability of the Endemic Infection Equilibrium Point Q_2 . In this subsection, we firstly introduce a lemma outlined by Li and Wang [18], and then using this lemma discusses the globally asymptotical stability of the endemic infection equilibrium point Q_2 of (3).

The lemma is briefly summarized as follows.

Let $x \mapsto f(x) \in \mathbb{R}^n$ be a C^1 function for x in an open set $\Gamma \subset \mathbb{R}^n$. Consider the differential system

$$\dot{x} = f(x). \quad (36)$$

Denote by $x(t, x^0)$ the solution to (36) such that $x(0, x^0) = x^0$. Let \bar{x} be an equilibrium point of (36). Li and Wang [18] made the following two basic assumptions:

- (H₁) there exists a compact absorbing set $K \subset \Gamma$;
- (H₂) equation (36) has a unique equilibrium \bar{x} in Γ .

Li and Wang (see Theorem 2.5 in [18]) have given the following lemma.

Lemma 4 (see [18]). *Assume that*

- (1) *assumptions (H₁) and (H₂) hold;*
- (2) *equation (36) satisfies the Poincaré-Bendixson Property;*
- (3) *for each periodic solution $x = p(t)$ to (36) with $p(0) \in \Gamma$, the linear system (the second additive compound system)*

$$\dot{w}(t) = \frac{\partial f^{[2]}}{\partial x}(P(t)) w(t) \quad (37)$$

is asymptotically stable, where $\partial f^{[2]}/\partial x$ is the second additive compound matrix of the Jacobian matrix $\partial f/\partial x$;

- (4) $(-1)^n \det((\partial f/\partial x)(\bar{x})) > 0$.

Then the unique equilibrium \bar{x} is globally asymptotically stable in Γ .

Now one uses Lemma 4 to show the following.

Theorem 5. *If $R_0 > 1$, then the endemic infection equilibrium point Q_2 of (3) is globally asymptotically stable in Ω^0 , where Ω^0 is defined by (15).*

Proof. Based on Lemma 4, the proof of Theorem 5 has been implemented via the following four steps.

(1) For epidemic models and many other biological models where the feasible region is a bounded cone, (H₁) is equivalent to the uniform persistence of the system [19]. By (15), Ω^0 is bounded, so it only needs to show the uniform persistence of (3). According to Proposition 3.3 in

[20], the necessary and sufficient condition for the uniform persistence of (3) is equivalent to the equilibrium point Q_1 being unstable. Theorem 1 has shown that Q_1 is unstable if $R_0 > 1$. Therefore, (3) is uniformly persistent if $R_0 > 1$ so that (H₁) holds if $R_0 > 1$.

Meanwhile, $Q_1 = (x_0, 0, 0)$ by (4), so Q_1 does not exist in Ω^0 . Hence, Q_2 is the unique equilibrium point of (3) in Ω^0 so that (H₂) holds.

The results above verify the condition (1) of Lemma 4.

(2) The Jacobian matrix of (3) is

$$J(x, y, v) = \begin{bmatrix} -d_1 - a_1 & p & -a_2 \\ a_1 & -p - d_2 & a_2 \\ 0 & a & -d_3 \end{bmatrix}, \quad (38)$$

where $a_1 = (k_1 v^2)/(x + v)^2$ and $a_2 = (k_1 x^2)/(x + v)^2$.

If $H = \text{diag}(1, -1, 1)$, then

$$HJH = \begin{bmatrix} -d_1 - \frac{k_1 v^2}{x + v} & -p & -\frac{k_1 x^2}{(x + v)^2} \\ -\frac{k_1 v^2}{(x + v)^2} & -p - d_2 & -\frac{k_1 x^2}{(x + v)^2} \\ 0 & -a & -d_3 \end{bmatrix}, \quad (39)$$

and one can obtain that HJH has nonpositive off-diagonal elements in Ω^0 . Therefore (3) is competitive in Ω^0 . It is known that 3-dimensional competitive systems have the Poincaré-Bendixson Property [21]. Hence, (3) satisfies the Poincaré-Bendixson Property. This verifies condition (2) of Lemma 4.

(3) Let $P(t) = (x(t), y(t), v(t))$ be a periodic solution in Ω^0 .

According to [22], if $B = (b_{ij})$ is a 3×3 matrix, then the second additive compound $B^{[2]}$ of B is

$$B^{[2]} = \begin{bmatrix} b_{11} + b_{22} & b_{23} & -b_{13} \\ b_{32} & b_{11} + b_{33} & b_{12} \\ -b_{31} & b_{21} & b_{22} + b_{33} \end{bmatrix}. \quad (40)$$

The Jacobian matrix of (3) is

$$J(x, y, v) = \begin{bmatrix} -d_1 - a_1 & p & -a_2 \\ a_1 & -p - d_2 & a_2 \\ 0 & a & -d_3 \end{bmatrix}, \quad (41)$$

where $a_1 = k_1 v^2/(x + v)^2$ and $a_2 = k_1 x^2/(x + v)^2$.

And then the second additive compound matrix of the Jacobian matrix of (3) is given by

$$J^{[2]} = \begin{bmatrix} -d_1 - \frac{k_1 v^2}{(x + v)^2} - p - d_2 & \frac{k_1 x^2}{(x + v)^2} & \frac{k_1 x^2}{(x + v)^2} \\ a & -d_1 - \frac{k_1 v^2}{(x + v)^2} - d_3 & p \\ 0 & \frac{k_1 v^2}{(x + v)^2} & -p - d_2 - d_3 \end{bmatrix}, \quad (42)$$

and the second additive compound system of (3) along the periodic solution $P(t) = (x(t), y(t), v(t))$ is

$$\begin{aligned}\dot{w}_1 &= \left(-d_1 - \frac{k_1 v^2}{(x+v)^2} - p - d_2\right) w_1 \\ &\quad + \frac{k_1 x^2}{(x+v)^2} w_2 + \frac{k_1 x^2}{(x+v)^2} w_3, \\ \dot{w}_2 &= a w_1 + \left(-d_1 - \frac{k_1 v^2}{(x+v)^2} - d_3\right) w_2 + p w_3, \\ \dot{w}_3 &= \frac{k_1 v^2}{(x+v)^2} w_2 + (-p - d_2 - d_3) w_3.\end{aligned}\quad (43)$$

Define a global Lyapunov function by

$$V(w_1, w_2, w_3, P) = \left\| \left(w_1, \frac{y(t)}{v(t)} w_2, \frac{y(t)}{v(t)} w_3 \right) \right\|, \quad (44)$$

where $\|\cdot\|$ is the norm in set D defined by

$$\|(w_1, w_2, w_3)\| = \sup \{|w_1|, |w_2| + |w_3|\}. \quad (45)$$

Suppose that the solution $P(t)$ is periodic of least period $\omega > 0$ and that $P(0) \in \Omega^0$. According to [23], (3) is uniformly persistent, if there exists a positive constant μ such that

$$\begin{aligned}\liminf_{t \rightarrow \infty} x(t) &\geq \mu, & \liminf_{t \rightarrow \infty} y(t) &\geq \mu, \\ \liminf_{t \rightarrow \infty} v(t) &\geq \mu.\end{aligned}\quad (46)$$

Step (1) has shown that (3) is uniformly persistent if $R_0 > 1$. Hence, there always exists a positive constant μ which satisfies (46). The orbit of $P(t)$ remains at a positive distance from the boundary of Ω by the uniform persistence, and one can obtain that

$$y(t) \geq \mu, \quad v(t) \geq \mu \quad \text{for large enough } t. \quad (47)$$

Since $v < a\lambda/(dd_3)$ by (15),

$$V(w_1, w_2, w_3, P) \geq \frac{\mu dd_3}{a\lambda} \|(w_1, w_2, w_3)\|, \quad (48)$$

for all $(w_1, w_2, w_3) \in R^3$.

Along a solution (w_1, w_2, w_3) of (43), $V(w_1, w_2, w_3, P)$ becomes

$$V(w_1, w_2, w_3, P) = \sup \left\{ |w_1|, \frac{y(t)}{v(t)} (|w_2| + |w_3|) \right\}. \quad (49)$$

The right-hand derivative of $V(t)$ along the positive solution of (43) is

$$\begin{aligned}D_+ |w_1| &\leq \left(-d_1 - \frac{k_1 v^2}{(x+v)^2} - p - d_2\right) |w_1| \\ &\quad + \frac{k_1 x^2}{(x+v)^2} (|w_2| + |w_3|), \\ D_+ |w_2| &\leq a |w_1| + \left(-d_1 - \frac{k_1 v^2}{(x+v)^2} - d_3\right) |w_2| + p |w_3| \\ D_+ |w_3| &\leq \frac{k_1 v^2}{(x+v)^2} |w_2| + (-p - d_2 - d_3) |w_3|.\end{aligned}\quad (50)$$

Therefore

$$\begin{aligned}D_+ \frac{y}{v} (|w_2| + |w_3|) &= \frac{\dot{y}v - y\dot{v}}{v^2} (|w_2| + |w_3|) + \frac{y}{v} D_+ (|w_2| + |w_3|) \\ &\leq \frac{y}{v} \left(\frac{\dot{y}}{y} - \frac{\dot{v}}{v} \right) (|w_2| + |w_3|) \\ &\quad + \frac{y}{v} (a |w_1| + (-d_1 - d_3) |w_2| + (-d_2 - d_3) |w_3|) \\ &\leq \frac{ay}{v} |w_1| + \frac{y}{v} (|w_2| + |w_3|) \\ &\quad \times \left(\frac{\dot{y}}{y} - \frac{\dot{v}}{v} - d_3 - \min(d_1, d_2) \right), \\ D_+ V(t) &\leq \sup \{g_1(t), g_2(t)\} V(t),\end{aligned}\quad (51)$$

where

$$\begin{aligned}g_1(t) &= -d_1 - p - d_2 - \frac{k_1 v^2}{(x+v)^2} + \frac{k_1 vx^2}{y(x+v)^2} \leq \frac{\dot{y}}{y} - d_1, \\ g_2(t) &= \frac{ay}{v} + \frac{\dot{y}}{y} - \frac{\dot{v}}{v} - d_3 - \min(d_1, d_2) \\ &= \frac{\dot{y}}{y} - \min(d_1, d_2).\end{aligned}\quad (52)$$

Denote $d = \min(d_1, d_2)$, and then

$$\sup \{g_1(t), g_2(t)\} \leq \frac{\dot{y}}{y} - d. \quad (53)$$

By (52) and Gronwall's inequality, one obtains

$$V(t) \leq V(0) y(t) e^{-dt} \leq \frac{V(0) e^{-dt} \lambda}{d}. \quad (54)$$

$V(t) \rightarrow 0$ when $t \rightarrow \infty$, and then $(w_1, w_2, w_3) \rightarrow 0$ when $t \rightarrow \infty$ by (48). The second additive compound system is asymptotically stable. This verifies the condition (3) of Lemma 4.

(4) The Jacobi matrix of (3) at the endemic infection equilibrium Q_2 is

$$J_{Q_2} = \begin{bmatrix} -d_1 - \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & p & -\frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & -p - d_2 & \frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ 0 & a & -d_3 \end{bmatrix}, \quad (56)$$

and then

$$\begin{aligned} \det(J_{Q_2}) &= \begin{vmatrix} -d_1 - \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & p & -\frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} & -p - d_2 & \frac{k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \\ 0 & a & -d_3 \end{vmatrix} \\ &= - \left[d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right] (d_2 + p) d_3 - \frac{a k_1^2 \bar{x}^2 \bar{v}^2}{(\bar{x} + \bar{v})^4} \\ &\quad + d_3 p \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} + \frac{a k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} \left[d_1 + \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \right] \\ &= -d_1 (d_2 + p) d_3 - \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} (d_2 + p) d_3 \\ &\quad - \frac{a k_1^2 \bar{x}^2 \bar{v}^2}{(\bar{x} + \bar{v})^4} + d_3 p \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} + \frac{a d_1 k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} + \frac{a k_1^2 \bar{x}^2 \bar{v}^2}{(\bar{x} + \bar{v})^4} \\ &= d_1 \left[\frac{a k_1 \bar{x}^2}{(\bar{x} + \bar{v})^2} - (d_2 + p) d_3 \right] - \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \\ &\quad - \frac{p d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} + d_3 p \frac{k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \\ &= a d_1 k_1 \left[\left(\frac{\bar{x}}{\bar{x} + \bar{v}} \right)^2 - \frac{1}{R_0} \right] - \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2}. \end{aligned} \quad (57)$$

According to (7), $\bar{v} = (R_0 - 1)\bar{x}$, and then one can obtain

$$\begin{aligned} \det(J_{Q_2}) &= a d_1 k_1 \left[\frac{1}{R_0^2} - \frac{1}{R_0} \right] - \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2} \\ &= \frac{a d_1 k_1 (1 - R_0)}{R_0^2} - \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2}. \end{aligned} \quad (58)$$

Since J_{Q_2} is a 3×3 matrix, one gets $n = 3$. Then

$$(-1)^3 \det(J_{Q_2}) = -\frac{a d_1 k_1 (1 - R_0)}{R_0^2} + \frac{d_2 d_3 k_1 \bar{v}^2}{(\bar{x} + \bar{v})^2}. \quad (59)$$

If $R_0 > 1$, then $(-1)^3 \det(J_{Q_2}) > 0$ holds in Ω^0 . This verifies condition (4) of Lemma 4.

Hence, if $R_0 > 1$, then the endemic infection equilibrium point Q_2 is globally asymptotically stable in Ω^0 by Lemma 4. \square

5. Numerical Simulation

In the first subsection, we determine some parameter values of an anti-HIV infection treatment model based on (3). In the second subsection, using the anti-HIV infection treatment model simulates the dynamics of the Group I's anti-HIV infection treatment. In the third subsection, using the anti-HIV infection treatment model simulates the dynamics of the Group II's anti-HIV infection treatment. In the fourth subsection, we make long-term predictions for the two groups' anti-HIV infection treatment, respectively.

5.1. Modeling. Baxter et al. [24] have reported a randomized study of antiretroviral management based on plasma genotypic antiretroviral resistance testing in HIV patients failing therapy, which was enrolled from 14 units of the Terry Beirn Community Programs for Clinical Research on AIDS and the Walter Reed Army Medical Center (see the HIV drug resistance database of Stanford University [25]). These patients were failing virologically on a combination antiretroviral regimen containing protease inhibitors (PI) and nucleoside reverse transcriptase inhibitors (NRTI) [24]. The patients were seen at 4, 8, and 12 weeks. At each follow-up visit, changes in antiretroviral treatment were recorded and the tested items included patients' plasma CD4⁺ T cells counts and plasma HIV-1 RNA levels by the Chiron 2.0 bDNA assay [24].

In the following subsections, we select, from [24, 25], two group patients' mean uninfected CD4⁺ T cells counts and mean HIV RNA levels to simulate and make long-term predictions for the patients' treatment outcomes. Group I consists of 15 patients. Group II consists of 13 patients. The two groups of patients received the same PI: ritonavir (RTV) and saquinavir (SQV). Additionally, Group I received NRTI: struvudine (D4T). Group II received NRTI: struvudine (D4T) and dideoxyinosine (DDI) [24, 25].

Based on (3), the anti-HIV infection treatment model has the form

$$\begin{aligned} \dot{x} &= \lambda - d_1 x - \frac{(1-m)k_1 vx}{x+v} + py, \\ \dot{y} &= \frac{(1-m)k_1 vx}{x+v} - d_2 y - py, \\ \dot{v} &= (1-n)ay - d_3 v, \end{aligned} \quad (60)$$

where m, n ($0 \leq m, n \leq 1$) are the efficacy variables of the treatment.

The infection-free equilibrium point Q_1 of (60) is the same as that defined by (4):

$$Q_1 = \left(\frac{\lambda}{d_1}, 0, 0 \right). \quad (61)$$




FIGURE 1: Outcomes of the treatment efficacy of Group I. Circles: the clinical data; solid line: the numerical simulation of (60). (a) Mean uninfected CD4⁺ T cells counts. (b) Mean HIV RNA levels.

The endemic infection equilibrium point Q_2 of (60) is given by

$$Q_2 = (\bar{x}, \bar{y}, \bar{v}), \quad (62)$$

where,

$$\begin{aligned} \bar{x} &= \frac{\lambda R_0}{(1-m)k_1(R_0-1) + d_1 R_0 - p(R_0-1)R_0(d_3/(1-n)a)}, \\ \bar{y} &= \frac{d_3}{(1-n)a}(R_0-1)\bar{x}, \quad \bar{v} = \frac{(1-n)a\bar{y}}{d_3} = (R_0-1)\bar{x}, \end{aligned} \quad (63)$$

and R_0 is the basic virus reproductive number of (60):

$$R_0 = \frac{(1-n)(1-m)ak_1}{d_3(d_2+p)}. \quad (64)$$

Determine the parameter value ranges of (60).

- (1) Naive CD4⁺ T cells decayed with an average half-life of 50 days [26]. Therefore, one obtains

$$d_1 = \frac{-\ln(0.5)}{50}. \quad (65)$$

Hence $d_1 \approx 0.0139$. Because the apoptosis of CD4⁺ T cells is raised by HIV infection [27, 28], d_1 should be more than 0.0139 during the simulation.

- (2) Since healthy individuals have an average of 830/ μ L CD4⁺ T cells [29],

$$\lambda = d_1 \times 830 \approx 0.0139 \times 830 = 11.5370. \quad (66)$$

- (3) Since the cells that produce the virus are also short-lived, with a half-life of approximately 1.2 days [30], one obtains

$$d_2 = \frac{-\ln(0.5)}{1.2} \approx 0.5776. \quad (67)$$

- (4) Since the half-life of HIV-1 in the plasma appears to be only 1 to 2 days [30], one selects

$$d_3 = \frac{-\ln(0.5)}{1.5} \approx 0.4621. \quad (68)$$

TABLE 1: Parameter values and R_0 at different weeks.

Weeks	d_1	m	n	R_0
0~4	0.033	0.53	0.84	0.6148
4~12	0.038	0.45	0.76	1.0792

- (5) Because only a small fraction of cells in the eclipse phase will revert to the uninfected state, it assumes that $p = 0.01$ [12]. Hence one obtains

$$p = 0.01. \quad (69)$$

- (6) According to reference [31], one can determine the other parameter value ranges as follows:

$$\begin{aligned} k_1 &\in [2.5 \times 10^{-5}, 0.5], \quad a \in [2, 1250], \\ m &\in [0, 1], \quad n \in [0, 1]. \end{aligned} \quad (70)$$

In each group, there was one patient whose clinical data was not complete. Therefore we do not conclude the two patients' clinical data in the following simulations.

5.2. *The Mean Dynamics Simulation of Group I's Anti-HIV Infection Treatment.* Using the other 14 patients' clinical data determines the equation parameter values as follows:

$$k_1 = 6 \times 10^{-2}, \quad a = 37. \quad (71)$$

The value changes of the parameters d_1 , m , and n and the basic virus reproductive number R_0 are listed in Table 1.

The simulation results of the mean dynamics of anti-HIV infection treatment of Group I are shown in Figure 1. During the first 4 weeks, the treatment reduced the basic virus reproductive number R_0 from 8.1759 to 0.6148. Hence the patients' mean HIV RNA levels decreased rapidly to approach infection-free steady state Q_1 defined by (4) as Theorem 2 predicts.

However, after the 4th week, the resistance to antiretroviral drugs appeared. It made the therapy efficacy parameters m and n decrease from 0.53 to 0.45 and 0.84 to 0.76, respectively. Meanwhile the apoptosis of CD4⁺ T cells was raised by HIV more strongly. Hence d_1 rose from 0.033 to 0.038. The

TABLE 2: Parameter values and R_0 at different weeks.

Weeks	d_1	m	n	R_0
0~4	0.041	0.65	0.68	0.6435
4~12	0.0428	0.55	0.59	1.0600

suboptimal treatment increased R_0 value of Group I from 0.6148 to 1.0792. As a result, the patients' mean HIV RNA levels increased slowly to converge to a new infected steady state Q_2 defined by (62) and (63) as Theorem 5 predicts. On the other hand, observe that the mean uninfected $CD4^+$ T cell counts of Group I increased rapidly in the first 4 weeks and decreased slowly in the following 8 weeks.

5.3. The Mean Dynamics Simulation of Group II's Anti-HIV Infection Treatment. Using 12 patients' clinical data determines the equation parameter values as follows:

$$k_1 = 6 \times 10^{-2}, \quad a = 26. \quad (72)$$

The value changes of the parameters d_1 , m , and n and the basic virus reproductive number R_0 are listed in Table 2.

The simulation results of the mean dynamics of anti-HIV infection treatment of Group II are shown in Figure 2. During the first 4 weeks, the treatment reduced the basic virus reproductive number R_0 from 5.7452 to 0.6435. Hence the patients' mean HIV RNA levels decreased rapidly to approach to infection-free steady state Q_1 defined by (4) as Theorem 2 predicts.

However, after the 4th week, the resistance to antiretroviral drugs appeared. It made the therapy efficacy parameters m and n decrease from 0.65 to 0.55 and 0.68 to 0.59, respectively. Meanwhile the apoptosis of $CD4^+$ T cells was raised by HIV more strongly. Hence d_1 rose from 0.041 to 0.0428. The suboptimal treatment made R_0 value of Group II increase from 0.6435 to 1.0600. As a result, the patients' mean HIV RNA levels increased slowly to converge to a new infected steady state Q_2 defined by (62) and (63) as Theorem 5 predicts. On the other hand, observe that the mean uninfected $CD4^+$ T cell counts of Group II increased rapidly in the first 4 weeks but rose slowly in the following 8 weeks.

5.4. The Long-Term Predictions for the Two Groups' Anti-HIV Infection Treatment. According to 2013 HIV therapy guidelines published by World Health Organization (WHO) [32], viral load is recommended as the preferred monitoring approach to diagnose and confirm antiretroviral treatment failure; treatment failure is defined by a persistently detectable viral load exceeding 1000 copies/mL after at least six months of using antiretroviral drugs. However, HIV RNA levels of the two groups of patients were only tested at 4, 8, and 12 weeks in the study [24]. Therefore, it is necessary to make long-term predictions to detect whether the treatments for the two groups are failure.

Assume that after ending the 12 weeks' antiretroviral treatment testing, the two groups keep receiving the anti-HIV infection treatment for 2 years. During the 2 years, the drug

resistance does not become worse and all parameter values do not change. Using the numerical simulation of Equation (60) makes the long-term predictions for the two groups' anti-HIV infection treatment. The long-term prediction for Group I' anti-HIV infection treatment is shown in Figure 3. Two years' outcomes of the therapy for Group II are shown in Figure 4. Observe that after finishing the 12 weeks' antiretroviral treatment testing, the mean uninfected $CD4^+$ T cells counts of the two groups both decline a little and finally keep at a level larger than the mean baseline values; the mean HIV RNA levels of the two groups both rise a lot to a level less than the mean baseline values but keep exceeding 1000 copies/mL all the time. The long-term predictions suggest that the treatments for the two groups are failure and better anti-HIV infection therapies should be considered.

6. Conclusion

This paper introduces a modified HIV infection differential equation model with a saturated infection rate $k_1 x v / (x + v)$ and the proportion of infected cells reverting to the uninfected state.

The basic virus reproductive number R_0 of the model is independent of a patient's plasma total $CD4^+$ T cell counts λ/d_1 , and the actual incidence rate is not linear over the entire range of virus $v(t)$ and uninfected $CD4^+$ T cells $x(t)$ any more. This suggests that our model is more reasonable than the model proposed by [13].

The modified model has two equilibrium points: infection-free equilibrium point Q_1 and endemic infection equilibrium point Q_2 . This paper discusses the locally asymptotical stabilities and globally asymptotical stabilities of the two equilibrium points, simulates the dynamics of two group patients' anti-HIV infection treatment, and makes long-term predictions for the two groups' anti-HIV infection treatment.

The theoretical results suggest the following.

- (1) If the basic virus reproductive number $R_0 < 1$, then the infection-free equilibrium point Q_1 of (3) is globally asymptotically stable in Ω . This means that if a person with $R_0 < 1$, the person can recover automatically even if infected with a large amount of HIV; if a treatment makes a patient's $R_0 < 1$, the patient will be cured eventually even if infected with a large amount of HIV.

The recent reports on three HIV infected patients have shown that some HIV infected patients may be cured via bone marrow transplants. The Berlin Patient was the first person cured of HIV [33]. After the Berlin Man, two cases reported cured of HIV in Kenya [34]. These reports can make one postulate that most individuals who connect HIV virus will not be infected by it and are not infecting other people. They will recover automatically without any treatment. The fact has not been well recognized since AIDS has been discovered in 1983. Mathematically, such phenomena can be described also via (3) where $k_1 = a = 0$, and thus $R_0 = 0$.




FIGURE 2: Outcomes of the treatment efficacy of Group II. Circles: the clinical data; solid line: the numerical simulation of (60). (a) Mean uninfected CD4⁺ T cells counts. (b) Mean HIV RNA levels.




FIGURE 3: The long-term prediction for the treatment efficacy of Group I. Circles: the clinical data; solid line: the numerical simulation of (60). (a) Mean uninfected CD4⁺ T cells counts. (b) Mean HIV RNA levels.

(2) If the basic virus reproductive number $R_0 > 1$, then the endemic infection equilibrium point Q_2 of (3) is globally asymptotically stable in the interior of Ω . This means that a person with $R_0 > 1$ will have endemic infection even if infected with only one HIV; if a treatment cannot make a patient's $R_0 < 1$, the patient's HIV RNA in vivo cannot be cleared up eventually.

Based on the simulation results, one can propose the following hypotheses.

(1) For a poor HIV treatment response patient, the drug resistance appears when the patient's HIV RNA level reduces to the first lowest level.

This hypothesis may interpret why the two group patients' mean CD4⁺ T cells counts rose and mean HIV RNA levels declined rapidly in the first 4 weeks but contrary in the following weeks (see Tables 1 and 2 and Figures 1 and 2).

(2) Once a patient's drug resistance appears, the patient's HIV in vivo promotes the apoptosis of CD4⁺ T cells more strongly.

This hypothesis may interpret why between 4th week and 8th week the mean HIV RNA levels of the two groups kept a lower level than the first 4 weeks, but Group I's mean CD4⁺ T cells counts started to decrease slowly and Group II's mean CD4⁺ T cells counts rose more slowly than before (see Tables 1 and 2 and Figures 1 and 2).

(3) According to the 2013 HIV therapy guidelines published by WHO [32], treatment failure is defined by a persistently detectable viral load exceeding 1000 copies/mL after at least six months of using antiretroviral drugs. Our long-term numerical simulation predictions suggest that after ending the 12 weeks' antiretroviral treatment [24], the two group patients' mean HIV RNA levels keep exceeding 1000 copies/mL during the additional 2 years' anti-HIV infection treatment. The treatments for the two groups are failure and better anti-HIV infection therapies should be considered. This means that the additional 2 years' treatments are not able to make patients obtain better outcomes. This may interpret why WHO defines that half a year's treatment cannot suppress a patient's HIV level below 1000 copies/mL to be treatment failure.

Highlights

(1) This paper introduces a modified HIV infection (anti-HIV infection therapy) differential equation model with a saturated infection rate. The basic virus reproductive number R_0 of the model is independent of a patient's plasma total CD4⁺ T cell counts λ/d_1 . This suggests that our model is more reasonable than the model proposed by Srivastava and Chandra in 2010 [13] whose basic virus reproductive number R_0 is dependent on λ/d_1 which follows that the more CD4⁺




FIGURE 4: The long-term prediction for the treatment efficacy of Group II. Circles: the clinical data; solid line: the numerical simulation of (60). (a) Mean uninfected $CD4^+$ T cells counts. (b) Mean HIV RNA levels.

T cell counts an individual has, the more easily the individual is infected by HIV.

- (2) This paper proposes and proves two theorems (Theorems 2 and 5) on the globally asymptotical stabilities of the infection-free equilibrium point Q_1 and the endemic infection equilibrium point Q_2 of the modified model.
- (3) This paper points out the implications of the two theorems which are ignored by other similar researches on modelling HIV infection (anti-HIV infection therapy):
 - (a) A person with the basic virus reproductive number $R_0 < 1$ will recover automatically even if the person is infected with a large amount of HIV. If a treatment makes a patient's $R_0 < 1$, this patient will be cured eventually even if infected with a large amount of HIV.
 - (b) A person with $R_0 > 1$ will have endemic infection even if the person is infected with only one HIV. If a treatment cannot make a patient's $R_0 < 1$, the patient's HIV RNA in vivo cannot be cleared up eventually.

The recent reports on three HIV infected patients show that HIV infected patients may be cured via bone marrow transplants (e.g., see: Berlin Patient, first person cured of HIV, may soon have company, Los Angeles Times, July 27, 2012; after Berlin Man, two reported cured of HIV in Kenya, Africa Review, May 6, 2013).

These reports can make one postulate that most individuals who connect HIV virus will not be infected by it and are not infecting other people, who will recover automatically without any treatment. In this case, $R_0 = 0$ where $k_1 = a = 0$. The fact has not been recognized since AIDS has been discovered in 1983.

In the report [35], a small proportion of human immunodeficiency virus type 1 (HIV-1) infected individuals, called elite and viremic controllers, spontaneously control plasma HIV RNA levels to undetectable (elite controller) or <2000 copies/mL (viremic controller) in the absence of antiretroviral therapy.

These phenomena can be interpreted by our Theorem 2: HIV infected people's basic virus reproductive number $R_0 < 1$.

- (4) Based on the simulation results, we can propose the following hypotheses:
 - (a) for a poor HIV treatment response patient, the drug resistance appears when the patient's HIV RNA level reduces to the first lowest level;
 - (b) Once a patient's drug resistance appears, the patient's HIV promotes the apoptosis of $CD4^+$ T cells more strongly;
 - (c) According to the 2013 HIV therapy guidelines published by WHO [32], treatment failure is defined by a persistently detectable viral load exceeding 1000 copies/mL after at least six months of using antiretroviral drugs. Our long-term numerical simulation predictions suggest that after ending the 12 weeks' antiretroviral treatment [24], the two group patients' mean HIV RNA levels keep exceeding 1000 copies/mL during the additional 2 years' anti-HIV infection treatment. The treatments for the two groups are failure and better anti-HIV infection therapies should be considered. This means that the additional 2 years' treatments are not able to make patients obtain better outcomes. This may interpret why WHO defines that half a year's treatment cannot suppress a patient's HIV level below 1000 copies/mL to be treatment failure.

Conflict of Interests

All authors have no financial or other conflict of interests pertaining to this paper.

Acknowledgments

The authors gratefully acknowledge anonymous reviewers for valuable suggestions and Dr. Xiao Chen for her help in this study. The authors would like to thank the HIV drug resistance database of Stanford University which provides the

clinical data on anti-HIV infection treatment. This work is jointly supported by the National Natural Science of China (no. 61074192) and Doctoral Research Funds of University of Science and Technology Beijing (no. 06108126).

References

- [1] L. Wang and M. Y. Li, "Mathematical analysis of the global dynamics of a model for HIV infection of CD4⁺ T cells," *Mathematical Biosciences*, vol. 200, no. 1, pp. 44–57, 2006.
- [2] C. J. Miller and R. J. Shattock, "Target cells in vaginal HIV transmission," *Microbes and Infection*, vol. 5, no. 1, pp. 59–67, 2003.
- [3] T. Bronshtein, N. Toledano, D. Danino, S. Pollack, and M. MacHluf, "Cell derived liposomes expressing CCR5 as a new targeted drug-delivery system for HIV infected cells," *Journal of Controlled Release*, vol. 151, no. 2, pp. 139–148, 2011.
- [4] M. A. Nowak and C. R. M. Bangham, "Population dynamics of immune responses to persistent viruses," *Science*, vol. 272, no. 5258, pp. 74–79, 1996.
- [5] M. A. Nowak, S. Bonhoeffer, G. M. Shaw, and R. M. May, "Anti-viral drug treatment: dynamics of resistance in free virus and infected cell populations," *Journal of Theoretical Biology*, vol. 184, no. 2, pp. 203–217, 1997.
- [6] J. E. Mittler, B. Sulzer, A. U. Neumann, and A. S. Perelson, "Influence of delayed viral production on viral dynamics in HIV-1 infected patients," *Mathematical Biosciences*, vol. 152, no. 2, pp. 143–163, 1998.
- [7] L. Rong and A. S. Perelson, "Modeling HIV persistence, the latent reservoir, and viral blips," *Journal of Theoretical Biology*, vol. 260, no. 2, pp. 308–331, 2009.
- [8] J. M. Orellana, "Optimal drug scheduling for HIV therapy efficiency improvement," *Biomedical Signal Processing and Control*, vol. 6, no. 4, pp. 379–386, 2011.
- [9] P. W. Nelson and A. S. Perelson, "Mathematical analysis of delay differential equation models of HIV-1 infection," *Mathematical Biosciences*, vol. 179, no. 1, pp. 73–94, 2002.
- [10] J. A. Zack, S. J. Arrigo, S. R. Weitsman, A. S. Go, A. Haislip, and I. S. Y. Chen, "HIV-1 entry into quiescent primary lymphocytes: molecular analysis reveals a labile, latent viral structure," *Cell*, vol. 61, no. 2, pp. 213–222, 1990.
- [11] P. Essunger and A. S. Perelson, "Modeling HIV infection of CD4⁺ t-cell subpopulations," *Journal of Theoretical Biology*, vol. 170, no. 4, pp. 367–391, 1994.
- [12] L. Rong, M. A. Gilchrist, Z. Feng, and A. S. Perelson, "Modeling within-host HIV-1 dynamics and the evolution of drug resistance: trade-offs between viral enzyme function and drug susceptibility," *Journal of Theoretical Biology*, vol. 247, no. 4, pp. 804–818, 2007.
- [13] P. K. Srivastava and P. Chandra, "Modeling the dynamics of HIV and CD4⁺ T cells during primary infection," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 2, pp. 612–618, 2010.
- [14] B. Buonomo and C. Vargas-De-León, "Global stability for an HIV-1 infection model including an eclipse stage of infected cells," *Journal of Mathematical Analysis and Applications*, vol. 385, no. 2, pp. 709–720, 2012.
- [15] X. Wang, Y. Tao, and X. Song, "Global stability of a virus dynamics model with Beddington-DeAngelis incidence rate and CTL immune response," *Nonlinear Dynamics*, vol. 66, no. 4, pp. 825–830, 2011.
- [16] X. Song and A. U. Neumann, "Global stability and periodic solution of the viral dynamics," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 281–297, 2007.
- [17] G. Huang, W. Ma, and Y. Takeuchi, "Global properties for virus dynamics model with Beddington-DeAngelis functional response," *Applied Mathematics Letters*, vol. 22, no. 11, pp. 1690–1693, 2009.
- [18] M. Y. Li and L. Wang, "Global stability in some SEIR epidemic models," in *Mathematical Approaches for Emerging and Reemerging Infectious Diseases: Models, Methods, and Theory*, pp. 295–311, Springer Science+Business Media, New York, NY, USA, 2002.
- [19] P. Waltman, "A brief survey of persistence," in *Proceedings of a Conference in Honor of Kenneth Cooke*, pp. 31–40, 1991.
- [20] M. Y. Li, J. R. Graef, L. Wang, and J. Karsai, "Global dynamics of a SEIR model with varying total population size," *Mathematical Biosciences*, vol. 160, no. 2, pp. 191–213, 1999.
- [21] M. W. Hirsch, "Systems of differential equations which are competitive or cooperative, IV: structural stabilities in three dimensional systems," *The SIAM Journal on Mathematical Analysis*, vol. 21, pp. 1225–1234, 1990.
- [22] X. Wang and X. Song, "Global stability and periodic solution of a model for HIV infection of CD4⁺ T cells," *Applied Mathematics and Computation*, vol. 189, no. 2, pp. 1331–1340, 2007.
- [23] P. Waltman, "A brief survey of persistence in dynamical systems," in *Delay Differential Equations and Dynamical Systems*, vol. 1475 of *Lecture Notes in Mathematics*, pp. 31–40, 1991.
- [24] J. D. Baxter, D. L. Mayers, D. N. Wentworth et al., "A randomized study of antiretroviral management based on plasma genotypic antiretroviral resistance testing in patients failing therapy," *AIDS*, vol. 14, no. 9, pp. F83–F93, 2000.
- [25] "Clinical Trial Datasets GART Study," <http://hivdb.stanford.edu/pages/clinicalStudyData/GART.html>.
- [26] C. Bourgeois, Z. Hao, K. Rajewsky, A. J. Potocnik, and B. Stockinger, "Ablation of thymic export causes accelerated decay of naive CD4⁺ T cells in the periphery because of activation by environmental antigen," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 25, pp. 8691–8696, 2008.
- [27] P. U. Cameron, M. Pope, S. Gezelter, and R. M. Steinman, "Infection and apoptotic cell death of CD4⁺ T cells during an immune response to HIV-1-pulsed dendritic cells," *AIDS Research and Human Retroviruses*, vol. 10, no. 1, pp. 61–71, 1994.
- [28] G. Herbein, C. van Lint, J. L. Lovett, and E. Verdin, "Distinct mechanisms trigger apoptosis in human immunodeficiency virus type 1-infected and in uninfected bystander T lymphocytes," *Journal of Virology*, vol. 72, no. 1, pp. 660–670, 1998.
- [29] M. Bofill, G. Janossy, C. A. Lee et al., "Laboratory control values for CD4 and CD8 T lymphocytes. Implications for HIV-1 diagnosis," *Clinical and Experimental Immunology*, vol. 88, no. 2, pp. 243–252, 1992.
- [30] D. R. Kuritzkes, "HIV Pathogenesis and Viral Markers," April 2000, <http://www.medscape.org/viewarticle/421048>.
- [31] Y. Wang, Y. Zhou, J. Wu, and J. Heffernan, "Oscillatory viral dynamics in a delayed HIV pathogenesis model," *Mathematical Biosciences*, vol. 219, no. 2, pp. 104–112, 2009.
- [32] World Health Organization, *Consolidated Guidelines on General HIV Care and the Use of Antiretroviral Drugs for Treating and Preventing HIV Infection: Recommendations for a Public Health Approach*, WHO Press, Geneva, Switzerland, 2013.

- [33] “Berlin Patient, first person cured of HIV, may soon have company,” July 2012, <http://articles.latimes.com/2012/jul/27/news/la-heb-hiv-cure-aids-meeting-20120727>.
- [34] “After Berlin Man, two reported cured of HIV in Kenya,” May 2013, <http://www.africareview.com/Special-Reports/After-Berlin-Man-two-reported-cured-of-HIV-in-Kenya/-/979182/1843388/-/3saq6o/-/index.html>.
- [35] B. Julg, F. Pereyra, M. J. Bouzón et al., “Infrequent recovery of HIV from but robust exogenous infection of activated CD_4^+T cells in HIV elite controllers,” *Clinical Infectious Diseases*, vol. 51, no. 2, pp. 233–238, 2010.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

