

Case Report

Conservative Treatment of Bilateral Impacted Mandibular Canines Traction

Gustavo Armando Ruíz-Mora ^{1,2} **Luis Ernesto Arriola-Guillén** ³
Aron Aliaga-Del Castillo ⁴ **Yalil Augusto Rodríguez-Cárdenas** ^{2,5} **Vinicius Dutra** ⁶
and Mabel Mejía-Milian ²

¹Division of Orthodontics, Faculty of Dentistry, Universidad Nacional de Colombia, Bogota, Colombia

²Division of Oral and Maxillofacial Radiology, School of Dentistry, Universidad Científica del Sur, Lima, Peru

³Division of Orthodontics and Division of Oral and Maxillofacial Radiology, School of Dentistry, Universidad Científica del Sur, Lima, Peru

⁴Department of Orthodontics and Pediatric Dentistry, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

⁵Division of Oral and Maxillofacial Radiology, School of Dentistry, Universidad Nacional de Colombia, Bogota, Colombia

⁶Department of Oral Pathology, Medicine, and Radiology, School of Dentistry, Indiana University, Indianapolis, Indiana, USA

Correspondence should be addressed to Luis Ernesto Arriola-Guillén; luchoarriola@gmail.com

Received 25 March 2023; Revised 8 May 2023; Accepted 12 July 2023; Published 1 August 2023

Academic Editor: Andrea Scribante

Copyright © 2023 Gustavo Armando Ruíz-Mora et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The objectives of the treatment of impacted canines differ according to the characteristics of dental malocclusion. Traction of the tooth is a conservative and viable alternative, which allows for maintaining stability and occlusal function. The following case report describes the treatment of an 11.6-year-old female patient, who presented bilateral impaction of mandibular canines in contact with the midline, mixed dentition in the inter-transitional period, class I angle malocclusion, with light crowding teeth. The treatment comprised three phases. The first phase, verticalization of the mandibular canines in mixed dentition, was performed to pull the impacted bilateral mandibular canines towards the dental arch to achieve their verticalization, maintaining the molar relationship, and the position of the upper and lower incisors. In the second phase, osteogenic rest was planned to relieve post-traction stress while awaiting the replacement of the mixed dentition. Finally, the third phase in permanent dentition was to align and level canines within the arch after extraction of the deciduous canines. For the viability of the permanent mandibular canines in the dental arch, orthosurgical traction was implemented, with a traction system with closed nickel-titanium coil springs with a transitory rigid dental-mucous-supported anchorage device, which allowed control and protection of the adjacent teeth and movements with helical forces of a controlled three-dimensional range. The results of the treatment were adequate, achieving consolidated molar and canine relationships, overjet, overbite, and optimal facial balance.

1. Introduction

Dental impaction in the oral cavity is one of the most defying challenges in clinical practice, with the severity increasing when teeth that are indispensable for function, esthetics, and stability in the occlusion are involved, as in the case of the canines. The eruptive development of the mandibular canines should be monitored from early mixed dentition, evaluating their relationship with the deciduous teeth. The trajectory of the mandibular canine

germ indicates the possibility of impaction [1]. Evaluation with diagnostic imaging is an adequate tool for detecting early anomalies in the eruptive development, aiding in treatment planning, and avoiding displacement of the impacted mandibular canine (IMbC) [2, 3]. Nevertheless, this alteration is commonly an incidental finding identified in routine radiographic images or clinical manifestations suggesting this possible clinical situation [3, 4].

The incidence of impaction of the mandibular canine is low, with the bilateral presentation being even less frequent

and it is more common in females [3–5]. It is related to etiological mechanical or genetic factors, such as abnormal resorption of the deciduous canine or interruption of the development of its eruptive process [2]. Eruptive disturbances may also be influenced by pathologic anomalies, severe space discrepancies, narrow arcs, and delayed eruption, increasing the risk of dental impaction [3, 4]. Many IMbCs are displaced to the midline, producing the transmigration phenomenon [5].

The most common treatment options for IMbC are surgical extraction with the closing of the edentulous space with the premolar as a substitute, maintaining the deciduous canines or substitution with an osteointegrated implant, or by auto-transplantation of the impacted tooth, or surgical exposition and traction to the dental arc [6–8].

Treatment planning for the traction of an impacted tooth must consider several factors, such as the severity of the malposition, treatment time, orthodontic biomechanics, periodontal stability, relationship with adjacent teeth, surgical techniques, adhesive, and traction mechanism, and clinical experience [6, 9, 10]. Several reports on skeletal and dental anchorage mechanisms have described traction attachments, including metallic chains, metallic ligatures, elastic chains, and nickel–titanium closed coil springs, which affect the control, friction, soft tissue health, time, and magnitude of traction force [11–14].

This case report describes a biomechanical alternative performed in three phases from mixed dentition to permanent dentition, in a female patient with mandibular canines that were bilaterally impacted in the midline and displaced towards the buccal side.

2. Case Presentation

2.1. Publication Consent. The patient's mother signed the consent for the publication of this case report, and the patient gave her informed assent.

2.2. Diagnosis and Etiology. A female patient of 11 years and 6 months of age was referred for orthodontic consultation due to bilateral impaction of the mandibular canines. Clinical examination showed facial symmetry, a slightly convex profile, decreased lower facial height, labial competence, and a brachyfacial face type, without functional problems, and with no medical history of relevance. In the intraoral examination, mixed dentition was observed in the inter-transitional period, not coinciding with the chronological age due to a delay in the spontaneous eruption of the premolars and canines, although all the first permanent molar, upper and lower incisors, eight deciduous molars, and both upper and lower deciduous canines were present. The patient also presented a class I angle malocclusion, with no evidence of crowding and coinciding midlines (Figure 1).

In the orthopantomography dental germs of permanent premolars and molars in intraosseous evolution were visualized, and the bilateral ectopic position of the mandibular canines was detected, located meso-angulated, buccal, and in the midline, over the inferior third of the lower incisor roots. The permanency of the deciduous mandibular canines with

no root resorption was confirmed (Figure 1). The severity of the impaction was diagnosed with anatomical mandibular landmarks, adapted from the study by Ericson and Kuroi [14] on maxillary canines. In this case scenario, this report describes the condition of both buccally impacted canines located in the middle third of the incisor roots as measured in the panoramic radiograph. The longitudinal axis of the right canine with relation to the midline presented a 54.1° angle, and with relation to the longitudinal axis of the adjacent lateral incisor of 55° , with a distance from the occlusal plane of 12.4 mm; the left canine presented 55.4° with relation to the midline, 53.8° with the longitudinal axis of the adjacent lateral incisor, and a distance from the occlusal plane of 14 mm, located more buccally than the right canine (Figure 1).

The cephalometric measures indicated a skeletal class I ($ANB = 1.1^\circ$), horizontal mandibular plane, diminished inferior facial height, upper incisors in adequate inclination, and pro-inclination of the lower incisors (Table 1).

Additionally, following the principles and recommendations of ALARA, cone-beam computed tomography (CBCT) images were taken to confirm the exact location of the impacted canines and their relationship with the lower incisors. The tomographic slices confirmed direct contact of the follicles and the incisor roots, without any evident damage, as well as the more buccal position of the left canine. The CBCT images were obtained with an i-CAT scanner (Imaging Sciences International, Hatfield, PA, USA) with the following configuration: 120 kV and 47.7 mA, with an exposure time of 20 seconds, a field of view of $8\text{ cm} \times 8\text{ cm}$, and a voxel size of 0.4 mm. The DICOM file was analyzed with the Dolphin 3D Imaging version 11.9 software (Chatsworth, CA, USA), allowing multiplanar and volumetric reconstruction.

2.3. Treatment Objectives. The objective of the treatment planning, in this case, was divided into three phases. The first phase involved the mixed dentition and was aimed at achieving the traction of both IMbCs into the arch to produce verticalization, whereas maintaining the molar relationship and incisor position. The second phase was planned as an osteogenic rest to alleviate the post-traction stress while awaiting the eruption of the permanent teeth. Finally, the objective of the third phase in the permanent dentition was to align and level both canines in the arch before the extraction of the deciduous canines, consolidating the molar and canine relationships, overjet, overbite, and an optimal facial balance.

2.4. Treatment Alternatives. Several treatment alternatives were considered for the resolution of this case report: (1) extraction of the IMbCs, preserving the deciduous canines; (2) extraction of the lower canines, the lower deciduous canines, and upper first bicuspids; (3) extraction of the lower permanent and deciduous canines, followed by the use of temporary anchorage devices to mesialize the lower posterior teeth; (4) extraction of lower permanent and deciduous canines, then retraction and uprighting of the lower incisor, and use of an orthopedic or functional appliance for stimulating forward growth potential; (5) auto-transplant of the impacted permanent canines; and (6) traction of permanent




FIGURE 1: Initial clinical radiographic records. (a) Facial and intraoral photographs. (b) Lateral X-ray. (c) Cephalometric tracing. (d) Panoramic X-ray: right canine: impactation sector 5 (close to the midline, adapted from Ericson and Kurol [14]); impact level: middle root third; angle between the longitudinal axis of the canine about the midline: 54.1°; angle between the longitudinal axis of the canine and the longitudinal axis of the adjacent lateral: 55°; distance from the cusp of the canine to the occlusal plane: 12.4 mm. Left canine: impactation sector 5 (close to the midline); impact level: middle root third; angle between the longitudinal axis of the canine in relation to the midline: 55.4°; angle between the longitudinal axis of the canine and the longitudinal axis of the adjacent lateral 53.8°; distance from the cusp of the canine to the occlusal plane: 14 mm.

TABLE 1: Cephalometric values.

Measurements	NORM	SD	Pre-treatment	Started the second phase	Final
SNA (°)	82	3.5	88	84.2	84
SNB (°)	80	3	86.9	82.2	82.7
ANB (°)	2°	2.4	1.1	2	2.3
OP-SN (°)	14.4	2.5	11.4	11.4	9.2
SN-GoGn (°)	32.9	5.2	20.1	20.6	18.6
FMA (°)	23.9	4.5	19.8	14.5	11.4
ENA-ME (mm)	66.7	4.1	54.01	52.10	53.20
U1-L1 (°)	130	6.0	118.4	112.7	109.8
U1-PP (°)	110	5	109.2	113.4	116.8
U1.NA (°), U1-NA (mm)	22.8, 4.3	5.7, 2.7	19.4, 4	26.5, 4.9	28.2, 5.9
L1.NB (°), L1-NB (mm)	25.3, 4.0	6.0, 1.8	42.5, 2.6	38.8, 4.4	39.8, 6.4
IMPA (L1-MP) (°)	95	7.0	112.4	113.8	115.8
Overjet (mm)	2.5	2.5	3.7	3.1	2.3
Overbite (mm)	2.5	2.0	4.8	3.1	1.5
Upper lip to E-plane (mm)	0	2.0	1.73	0.44	-0.90
Lower lip to E-plane (mm)	0	2.0	2.32	-0.91	-0.69
Nasolabial angle (°)	95	5.0	88.90	75.08	67.00

TABLE 2: Three phases of the therapy for the case report.

Phase I: active traction	Phase II: osteogenic rest	Phase III: full fixed orthodontic appliances
Goal: Upright and traction of both impacted mandibular canines.	Goals: Relieve stress from the neighboring tissues.	Goal: Comprehensive orthodontic treatment.
Alignment and leveling: Metallic brackets (0.022" × 0.028" slot) in the mandibular arch including the primary teeth present. CuNiTi 0.016" × 0.022" arch-wire (1 month before surgical exposure).	Maintain permanent canines upright and parallel to primary canines.	Bonding full fixed orthodontic appliance: Include all permanent teeth. Metallic brackets (0.022" × 0.028" slot).
Temporary rigid anchorage device: Lower lingual holding arch (SS, 1.1 mm) associated with interproximal occluso-buccal extensions (SS .028") with hooks in the distal direction.	Removal of the temporary anchorage device.	Extraction of the lower primary canines and immediate retraction, leveling, and alignment of the permanent canines.
Surgical exposure: Incision, flap, osteotomy, removal of dental follicles, clearance of canine crown, bonding of attachments, and NiTi closed coil springs (0.010 mm × 0.036 mm diameter and 13 mm long).	Segmented arch to maintain coil springs tied.	Arch-wire sequence: Lower: CuNiTi 0.016" × 0.022", NiTi 0.017" × 0.025", SS 0.019" × 0 : 025", SS 0.021" × 0.025". Australian .016" for anterior extrusion and finishing.
Activation: Each 6 weeks. Traction force: 100 g.	Monitor root development of permanent teeth.	Upper: CuNiTi 0.016" × 0.022", NiTi 0.017" × 0.025", SS 0.016" × 0.22" (with boot loop for incisor derotation), SS 0.019" × 0.025", SS 0.021" × 0.025".
Phase I time: 5 months.	Phase II time: 13 months.	Phase III time: 18 months.


canines to the dental arch, considering other anchor resources and personal attachments in the traction of the canine.

The different treatment alternatives were presented to the patient's parents, and the final decision was to perform traction of the permanent mandibular canine into the dental arch. Then, a traction system was made of nickel-titanium closed spring coils with a rigid temporal mucodental supported device. This anchorage and traction system allowed control and protection of the adjacent teeth and movements



with helical forces of a controlled three-dimensional (3D) range in the orthogonal axes x , y , and z [15].

2.5. Treatment Progress. The treatment progress was made up of three phases (Table 2). The first phase was in the mixed dentition, the main objective of which was achieving verticalization of both IMbCs.

Metallic brackets were placed on the mandibular teeth, slot 0.022" × 0.028" (Synergy; Rocky Mountain Orthodontics, Denver, CO, USA.), reaching a 0.016 × 0.022 CuNiTi




(a)



(b)

FIGURE 2: Continued.



(c)

FIGURE 2: (a) Fixed device for temporary anchorage. (A) Lingual arch main support in 1.1 mm SS wire. (B) Prefabricated metal bands with lower first molar tubes. (C) Vertical hooks in 0.028" SS wire, with bends towards the distal direction. (b) Trans-surgical radiographic and clinical records. (c) Three months of traction. Radiographic and clinical records.

arch-wire (Synergy; Rocky Mountain Orthodontics), including the deciduous molars. One month before surgical exposure of the canines performed with the closed technique, in which only the NiTi coil spring for traction was exposed, and before suturing, a temporary rigid anchorage device consisting of a lingual arch made with a 1.1 mm stainless steel (SS) arch-wire (Synergy; Rocky Mountain Orthodontics) was cemented in the first permanent molars. This was adapted to the lingual surface of the mandibular teeth, with proximal occluso-buccal extensions (SS 0.028", Dentaurum, Ispringen, Germany) from canine to molar with the hooks in the distal direction on both sides of the dental arch (Figure 2).


The surgical intervention of the closed traction technique involved: (1) incision and flap lift, (2) osteotomy and exposure of dental follicles, (3) elimination of the pericoronary follicles and complete clearance of the permanent canine clinical crown, without exceeding the cemento-enamel junction, (4) isolation and blood control with hemostatic agents and drying, to ensure adhesion of the attachments, (5) placing the two nickel-titanium closed coil spring (Dentos, Daegu, South Korea) of 0.010×0.036 in diameter and 13 mm in length in each impacted canine, respectively, due to the distance exceeded in the activation, and the high resistance of the impacted tooth, and the relation of the coil springs with the anatomical structures and the surrounding dental landmarks. Each coil spring was activated immediately after suturing, with distal movement in the x -axis, vestibular movement in the z -axis, and extrusion in the y -axis. The activation was made every 6 weeks, extending the coil springs 13 mm, until 60% of their length, and by the gradual movement of each canine. Approximately 100 g of force was used to achieve traction of the tooth (Figure 2) [15].

Five months after the initial bracket placement, when the permanent canines were in a good vertical position parallel to the deciduous canines (Figure 3), the temporary anchorage device was removed, and the second treatment phase or osteogenic rest was started to relieve stress from the surrounding bone tissue, whereas maintaining the deciduous canine in place (Figure 3).

The third and final phase of the treatment started 18 months after the first visit, with a collection of clinical and radiography data (Figure 3). In this stage, the patient presented all permanent dentition, and the bracket placement was completed in both dental arches. At this time, the extraction of the deciduous canine was performed to start the immediate alignment of the permanent canines over the alveolus bone post-extraction. The alignment and leveling started using $0.016" \times 0.022"$ CuNiTi arch-wire in both dental arches (Synergy; Rocky Mountain Orthodontics), followed by a $0.017" \times 0.025"$ NiTi arch-wire, and finishing with $0.019" \times 0.025"$ of SS (Synergy; Rocky Mountain Orthodontics). Additionally, an Australian $0.016"$ arch-wire for anterior extrusion in the upper arch was used (Figure 4).


Due to the evident buccal malposition of the canines, there was a cortical reduction. To measure this, tomographic imaging was carried out with the consent of the patient's parents. Based on the transaxial slices and to move the canines in a lingual direction, set bends were made to incorporate the canine roots into the alveolar process (Figure 4).

2.6. Treatment Outcome Assessment. The patient presented good dental alignment, molar class I with adequate lateral and anterior functional guides. The clinical records showed adequate dental arch form, good inter-arch engagement,



(a)

FIGURE 3: Continued.



(b)

FIGURE 3: (a) Images of the first phase. Radiographic and clinical records. (b) Records of the start of the third phase of treatment.

ideal overjet and overbite, and coinciding midlines, as well as an acceptable facial profile with few variations (Figure 5).


The post-treatment panoramic radiograph showed dental alignment, conservation of the integrity of the anterior teeth, and the mandibular canine maintained a good root length. In the cephalometric examination, the mandibular incisor inclination had a similar position, whereas the inclination of the upper incisors had moderately increased (Table 1; Figure 6).

No periodontal disease was observed, with the periodontal probe showing good adherence to the gum (Hufriedy). Furthermore, a beneficial increased crown length


was observed, with moderate cervical exposure of the crown of both canines. A lower lingual fixed retainer was adapted from canine to canine (Figure 5).

3. Discussion

The risk of teeth impaction is greater in patients with late eruption, being more common in females [5, 16]. In this clinical case, a female patient with late mixed dentition was treated. The diagnostic severity of the mandibular canine impaction was determined using anatomical parameters, and references adapted from the study of Ericson and Kurool



(a)



(b)

FIGURE 4: (a) Six months after the start of the third phase. (b) One year after initiating the third phase.

on maxillary canines [14, 17]. The final diagnosis was an ectopic mesioangulated buccal eruption of both permanent mandibular canines, over the midline and in follicular contact with the roots of the lower central incisor. The complexity and severity of this clinical case led to the design of a biomechanical traction and anchorage system, which has not previously been described for IMbCs.

Before and after the traction treatment, 3D images were used for the diagnosis, treatment, and follow-up using different axial, coronal, and sagittal slices, allowing evaluation of the severity of the malposition, location of the ectopic tooth related to the alveolar cortical, the repercussion on

the neighboring tooth, and possible bone losses [18]. The follow-up CBCTs were used to reevaluate the biomechanics of the lingual movement of the roots of the canines inside the bone by in-set bends in the first order and required torque movements.

Different treatment alternatives for IMbCs have been reported [19]. The most common alternative is surgical extraction of the impacted canine [17, 20–24]. Other alternatives are surgical exposure treatments and orthodontic traction [6, 7, 25]. In the present case, some treatment alternatives were considered, mainly extraction of the IMbCs and preservation of the deciduous canines, or the extraction of




FIGURE 5: Extra-oral and intra-oral photographs and final radiographic-cephalometric records.




FIGURE 6: (a) Pre-treatment tomographic images. (b) Intermediate tomographic images showing the absence of buccal cortical at the level of the mandibular canines. (c) Post-treatment transaxial tomographic slices of lower incisors. (d) Superimposition of initial and final phase tracings.

the lower canines, the lower deciduous canines, and replacement of these teeth with bicuspid and advancing the lower posterior sector. However, the possibility of producing a canine guide and the consequent esthetics prevailed over the patient’s choice of traction for the affected teeth. Therefore, the treatment of this case was planned in three phases. In the first phase involving mixed dentition in the first transitional period, traction and verticalization of the IMbCs were performed with a reinforced anchoring device until a parallel position was achieved with the deciduous canines in the arch. In the second phase in the mixed dentition second transitional period, osteogenic rest and extraction of deciduous canines were performed, and finally, third phase involved alignment and leveling of the permanent canines within the alveolar bone with complete orthodontic arches. The extraction of the deciduous canines was postponed until the end of the second phase and the beginning of third phase to take advantage of the osteoid-osteopenic period of the fresh deciduous alveolus in the healing processes.

The effectiveness and need for the use of anchorage resources for the traction of impacted teeth are important. Compared with a temporary anchorage device-type skeletal anchorage, a customized rigid anchor cemented with bands was preferred [11, 26]. Tooth-supported anchorage devices are a viable alternative for controlled traction of bilaterally IMbCs, as they are low-cost, transitory devices that provide protection to the teeth and do not involve tissues compared

with the adverse effects of action and reaction forces, thus avoiding important alterations in the patient’s occlusion and the immersion of the springs in the mucoperiosteum [6, 27, 28]. Additionally, as the treatment has been completed without the use of mini-screws, the risk of root injury was avoided [29–31]. In this treatment alternative, a customized dentomucosal anchorage device with buccal extensions was adopted during the first phase. The anchoring system is efficient when combined with the nickel–titanium closed coil springs, which allow adequate force management and continuous preservation of the periodontal tissues throughout the traction process. Furthermore, they are not susceptible to deformation or fatigue, facilitating the implementation of traction with helical forces controlled in magnitude, direction, and direction, with distribution towards the distal direction in the *x*-axis, vestibular in the *y*-axis, and extrusive in the *z*-axis, providing superelasticity, memory, and a wide range of activation with light continuous forces, in a total traction period of approximately 6 months [12, 28, 32–35].

Although lower incisor resorption due to canine impaction is infrequent [36–38], sequelae due to root resorption have been described in the upper maxilla, but are infrequent in the mandible [15], favoring the selection of orthosurgical traction treatments, maintaining the permanent canines in the dental arch. The 3D imaging showed no root involvement of the incisors despite the severity of impaction and the bilateral frictional movement of the canine follicles on

the root cement of the incisors (Figure 6). Another challenge was periodontal health, which can be affected by various factors, such as the type of surgical procedure selected to expose the crown of the impacted tooth, the eruption of the teeth through the attached gingiva or mucosa, the hygiene methods used, and the anatomical characteristics of the mandibular bone, including reduced bicortical width and alveolar rigidity [39]. It has been shown that surgical procedures closed by the buccal approach, such as that used in this case report, offer better results [40–44]. Nevertheless, coronal elongation of the canines was observed in the patient, with cervical exposure and possible projection of the roots through the attached gingiva margin due to the narrowness of the alveolar bone corridor, for which it was recommended to keep the canine and anteroinferior area in optimal hygiene conditions and monitor the stability of the occlusion.

The cephalometric findings showed proinclination of the upper and lower incisors due to treatment without extraction and the location of the canine impacted in the dental arch, producing an evident increase in the length, perimeter, and width in both arches. Another reason for the proinclination could be the need for canine and anterior functional guides [15, 35].

The length of treatment was prolonged due to predetermination of the chronological and dental age, the conservative biomechanics of verticalization of the canines contacting the incisors, and the three-phase treatment plan with long resting periods for achieving morphologic and functional occlusal stability.

The case report was completed with good posterior occlusion settlement following the three-phase treatment, demonstrating that with light forces acceptable occlusion can be achieved. However, a limitation of this intervention is the great biological variability of the patients, which may not allow the same good results and may lead to the development of unexpected side effects. Therefore, more studies with larger sample sizes are needed to establish the most adequate procedure to perform in patients with impaction of mandibular canines.

4. Conclusion

The presence of bilaterally IMbCs is a diagnostic and therapeutic challenge for orthodontists. Nevertheless, with a good tridimensional plan, the use of biological forces and the implementation of different anchorage and control traction biomechanical methods can achieve optimum results, constituting a viable treatment alternative for orthodontists.

Data Availability

The intra- and extra-oral figures and cephalometric data used to support the findings of this study are included within the article. For any information required, contact the corresponding author or Dr. Gustavo Ruiz responsible for managing the clinical case. garruiz@gmail.com, Private Dental Clinic luchoarriola@gmail.com.

Consent

Written informed consent was obtained from the patient who agreed to take part in the study.

Conflicts of Interest

The author(s) declare(s) that they have no conflicts of interest.

Authors' Contributions

All authors have read and approved the final manuscript content, write-up of the manuscript, and have contributed significantly to the study: GARM, LEAG, and MMM carried out the data collection and write-up of the manuscript; YARC, VD, and AADC participated in the preparation of the manuscript.

References

- [1] Y. Brorsson and J. Naoumova, "Delayed diagnosis of displaced and impacted canines - a prospective longitudinal study," *Acta Odontologica Scandinavica*, vol. 78, no. 3, pp. 165–172, 2020.
- [2] M. Mejía-Milian, L. E. Arriola-Guillén, A. Aliaga-Del Castillo, Y. A. Rodríguez-Cárdenas, and G. A. Ruíz-Mora, "Three-dimensional evaluation of mandibular canine impaction characteristics and their relationship with lower incisor root morphology," *Journal of Clinical and Experimental Dentistry*, vol. 14, no. 10, pp. e791–e802, 2022.
- [3] S. Jain, M. Agrawal, S. Jain, and S. Jain, "Evaluation of the mandibular arch in patients with impacted permanent lower canines," *Australian Orthodontic Journal*, vol. 31, no. 1, pp. 37–41, 2015.
- [4] S. Jain, K. S. Shetty, A. T. Prakash, M. Agrawal, and S. Jain, "Permanent mandibular canine(s) impaction: expansion of our understanding," *Australian Orthodontic Journal*, vol. 30, no. 1, pp. 39–44, 2014.
- [5] G. Sharma and A. Nagpal, "A study of transmigrated canine in an Indian population," *International Scholarly Research Notices*, vol. 2014, Article ID 756516, 2014.
- [6] S. Agarwal, S. Yadav, N. V. Shah, A. Valiathan, F. Uribe, and R. Nanda, "Correction of bilateral impacted mandibular canines with a lip bumper for anchorage reinforcement," *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 143, no. 3, pp. 393–403, 2013.
- [7] S. Cavuoti, G. Matarese, G. Isola, J. Abdolreza, F. Femiano, and L. Perillo, "Combined orthodontic-surgical management of a transmigrated mandibular canine," *The Angle Orthodontist*, vol. 86, no. 4, pp. 681–691, 2016.
- [8] M. H. Aras, K. Halicioğlu, M. S. Yavuz, and M. Çağlaroğlu, "Evaluation of surgical-orthodontic treatments on impacted mandibular canines," *Medicina Oral, Patología Oral y Cirugía Bucal*, vol. 16, no. 7, pp. e925–e928, 2011.
- [9] M. Gunashekhar and M. Rohini, "Transmigration of mandibular canines: a rare case report and review of literature," *Journal of Dentistry for Children (Chicago, Ill.)*, vol. 78, no. 1, pp. 19–23, 2011.
- [10] O. O. Sanu, T. A. Adeyemi, and M. C. Isiekwe, "Incidence of impacted mandibular canine and associated pathologies in an orthodontic patient population in Lagos, Nigeria,"

- Nigerian Quarterly Journal of Hospital Medicine*, vol. 22, no. 4, pp. 291–295, 2012.
- [11] R. C. Almeida, F. A. Carvalho, M. A. Almeida, and J. Capelli Jr., “Orthodontic management of a patient with impacted and transposed mandibular canines,” *World Journal of Orthodontics*, vol. 10, no. 4, pp. 345–349, 2009.
 - [12] C. Manhartberger and W. Seidenbusch, “Force delivery of Ni-Ti coil springs,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 109, no. 1, pp. 8–21, 1996.
 - [13] R. H. Peerlings, “Treatment of a horizontally impacted mandibular canine in a girl with a class II division 1 malocclusion,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 137, no. 4, pp. S154–S162, 2010.
 - [14] S. Ericson and J. Kurol, “Early treatment of palatally erupting maxillary canines by extraction of the primary canines,” *European Journal of Orthodontics*, vol. 10, no. 1, pp. 283–295, 1988.
 - [15] G. A. Ruiz-Mora, L. E. Arriola-Guillen, Y. A. Rodriguez-Cardenas, A. Aliaga-Del Castillo, and H. L. Dias-Da Silveira, “A conservative approach to traction of impacted maxillary canines in adults with severe incisor root resorption,” *Journal of Clinical Orthodontics*, vol. 54, no. 12, pp. 746–759, 2020.
 - [16] Z. Bayramoglu, O. Miloglu, and I. F. Yozgat, “The findings of impacted and transmigrated maxillary and mandibular canines: a retrospective cone beam computed tomography study,” *Minerva Dental and Oral Science*, vol. 72, no. 2, pp. 90–98, 2023.
 - [17] M. R. Joshi, “Transmigrant mandibular canines: a record of 28 cases and a retrospective review of the literature,” *The Angle Orthodontist*, vol. 71, no. 1, pp. 12–22, 2001.
 - [18] S. Baumgaertel, J. M. Palomo, L. Palomo, and M. G. Hans, “Reliability and accuracy of cone-beam computed tomography dental measurements,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 136, no. 1, pp. 19–25, 2009, discussion 25–18.
 - [19] J. Vuchkova and C. S. Farah, “Canine transmigration: comprehensive literature review and report of 4 new Australian cases,” *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics*, vol. 109, no. 4, pp. e46–e53, 2010.
 - [20] J. Rebellato and B. Schabel, “Treatment of a patient with an impacted transmigrant mandibular canine and a palatally impacted maxillary canine,” *The Angle Orthodontist*, vol. 73, no. 3, pp. 328–336, 2003.
 - [21] N. Umashree, A. Kumar, and T. Nagaraj, “Transmigration of mandibular canines,” *Case Reports in Dentistry*, vol. 2013, Article ID 697671, 2013.
 - [22] H. M. Novak, T. Baccetti, L. M. Sigler, and J. A. McNamara Jr., “A controlled study on diagnostic and prognostic measurements of palatally displaced canines on lateral cephalograms,” *Progress in Orthodontics*, vol. 13, no. 1, pp. 42–48, 2012.
 - [23] S. Camilleri and E. Scerri, “Transmigration of mandibular canines—a review of the literature and a report of five cases,” *The Angle Orthodontist*, vol. 73, no. 6, pp. 753–762, 2003.
 - [24] D. M. Abbott, J. A. Svirsky, and B. H. Yarborough, “Transposition of the permanent mandibular canine,” *Oral Surgery, Oral Medicine, and Oral Pathology*, vol. 49, no. 1, p. 97, 1980.
 - [25] K. Gruszka, T. K. Rozylo, I. Rozylo-Kalinowska, K. Denkwicz, and K. Maslowska, “Transmigration of mandibular canine—case report,” *Polish Journal of Radiology*, vol. 79, pp. 20–23, 2014.
 - [26] F. Heravi, H. Shafae, A. Forouzanfar, S. H. Zarch, and M. Merati, “The effect of canine disimpaction performed with temporary anchorage devices (TADs) before comprehensive orthodontic treatment to avoid root resorption of adjacent teeth,” *Dental Press Journal of Orthodontics*, vol. 21, no. 2, pp. 65–72, 2016.
 - [27] G. A. Ruiz-Mora, L. E. Arriola-Guillen, Y. A. Rodriguez-Cardenas, A. Aliaga-Del Castillo, M. Boessio-Vizzotto, and H. L. Dias-Da Silveira, “Changes in alveolar bone morphology after traction of buccally vs palatally unilateral maxillary impacted canines: a cone-beam computed tomography study,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 159, no. 3, pp. 258–270, 2021.
 - [28] Y. A. Rodriguez-Cardenas, G. A. Ruiz-Mora, A. Aliaga-Del Castillo, H. L. Dias-Da Silveira, and L. E. Arriola-Guillen, “Root and alveolar bone changes in first premolars adjacent to the traction of buccal versus palatal maxillary impacted canines,” *PLoS One*, vol. 14, no. 12, article e0226267, 2019.
 - [29] M. A. Montasser and A. Scribante, “Root injury during inter-radicular insertion is the most common complication associated with orthodontic miniscrews,” *The Journal of Evidence-Based Dental Practice*, vol. 22, no. 1, article 101688, 2022.
 - [30] Y. Xin, Y. Wu, C. Chen, C. Wang, and L. Zhao, “Miniscrews for orthodontic anchorage: analysis of risk factors correlated with the progressive susceptibility to failure,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 162, no. 4, pp. e192–e202, 2022.
 - [31] M. F. Sfondrini, P. Gandini, R. Alcozer, P. K. Vallittu, and A. Scribante, “Failure load and stress analysis of orthodontic miniscrews with different transmucosal collar diameter,” *Journal of the Mechanical Behavior of Biomedical Materials*, vol. 87, pp. 132–137, 2018.
 - [32] A. Bezrouk, L. Balsky, M. Smutny et al., “Thermomechanical properties of nickel-titanium closed-coil springs and their implications for clinical practice,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 146, no. 3, pp. 319–327, 2014.
 - [33] S. Yadav, J. Chen, M. Upadhyay, F. Jiang, and W. E. Roberts, “Comparison of the force systems of 3 appliances on palatally impacted canines,” *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 139, no. 2, pp. 206–213, 2011.
 - [34] K. Hayashi, J. Uechi, S. P. Lee, and I. Mizoguchi, “Three-dimensional analysis of orthodontic tooth movement based on XYZ and finite helical axis systems,” *European Journal of Orthodontics*, vol. 29, no. 6, pp. 589–595, 2007.
 - [35] M. H. Bertl, C. Frey, K. Bertl, K. Giannis, A. Gahleitner, and G. D. Strbac, “Impacted and transmigrated mandibular canines: an analysis of 3D radiographic imaging data,” *Clinical Oral Investigations*, vol. 22, no. 6, pp. 2389–2399, 2018.
 - [36] A. Koc, S. Kaya, and W. A. Abdulsalam, “Three-dimensional analysis of impacted maxillary and mandibular canines and evaluation of factors associated with transmigration on cone-beam computed tomography images,” *Journal of Oral and Maxillofacial Surgery*, vol. 79, no. 3, pp. 538.e1–538.e11, 2021.
 - [37] H. Cakir Karabas, I. Ozcan, A. F. Erturk, B. Guray, G. Unsal, and S. N. Senel, “Cone-beam computed tomography evaluation of impacted and transmigrated mandibular canines: a retrospective study,” *Oral Radiology*, vol. 37, no. 3, pp. 403–411, 2021.
 - [38] A. S. Jati, L. Z. Furquim, and A. Consolaro, “Gingival recession: its causes and types, and the importance of orthodontic treatment,” *Dental Press Journal Orthodontics*, vol. 21, no. 3, pp. 18–29, 2016.

- [39] A. Becker, I. Brin, Y. Ben-Bassat, Y. Zilberman, and S. Chaushu, "Closed-eruption surgical technique for impacted maxillary incisors: a postorthodontic periodontal evaluation," *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 122, no. 1, pp. 9–14, 2002.
- [40] S. Chaushu, A. Becker, R. Zeltser, S. Branski, N. Vasker, and G. Chaushu, "Patients' perception of recovery after exposure of impacted teeth: a comparison of closed- versus open-eruption techniques," *Journal of Oral and Maxillofacial Surgery*, vol. 63, no. 3, pp. 323–329, 2005.
- [41] S. Chaushu, I. Brin, Y. Ben-Bassat, Y. Zilberman, and A. Becker, "Periodontal status following surgical-orthodontic alignment of impacted central incisors with an open-eruption technique," *European Journal of Orthodontics*, vol. 25, no. 6, pp. 579–584, 2003.
- [42] C. Chavez-Alvarez, L. E. Arriola-Guillen, Y. A. Rodriguez-Cardenas et al., "Changes in maxillary incisor inclination and position after traction of unilateral vs bilateral maxillary impacted canines in nonextraction treatment: a cone-beam computed tomography study," *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 156, no. 6, pp. 767–778, 2019.
- [43] A. Aquino-Valverde, D. Barrientos-Sanchez, and K. Atoche-Socola, "Orthodontic treatment in impacted maxillary canines. A review of the literature," *Revista Científica Odontológica*, vol. 9, no. 4, article e085, 2021.
- [44] C. Díaz-Reissner, E. Pistilli, R. Cajé, C. Maldonado, E. Jolay, and M. Ferreira-Gaona, "Pronóstico de caninos impactados según su posición en el maxilar superior mediante diferentes análisis radiográficos," *Revista Científica Odontológica*, vol. 10, no. 1, article e096, 2022.