

Case Report

A Case of Recurrent Erysipelas Caused by *Streptococcus mitis* Group

David Nygren, Bo Nilson, and Magnus Rasmussen

Supplement Table S1 MALDI-TOF MS identification and antibiotic susceptibility profile of isolated *S. mitis* and *S. oralis* isolates

	MALDI Biotyper ^a							Specific peaks for <i>S. pneumoniae</i> , <i>S. mitis</i> , and <i>S. oralis</i> ^c					Antibiotic susceptibility profile				
	log (score)		list (score) ^b					m/z value					penicillin	vancomycin	gentamicin	clindamycin	optochin
Strain (Isolate no)	Best match	score value	p	m	o	sp	Best match	4964.3 ^d	5297.6 ^e	5822.5 ^e	6839.1 ^f	6888.9 ^d	MIC (mg/L)	MIC (mg/L)	MIC (mg/L)	SIR ^g	SIR ^g
15-1 (1)	<i>S. mitis</i>	2.28	2	92	26	0	<i>S. mitis</i>	-	-	-	+	-	0.064	0.25	2	S	R
17-1 (2)	<i>S. mitis</i>	2.51	0	90	43	0	<i>S. mitis</i>	-	-	-	+	-	0.125	0.5	2	S	R
17-2 (3)	<i>S. oralis</i>	2.48	0	11	113	0	<i>S. oralis</i>	-	-	+	+	-	0.125	1	16	S	R

^a Bruker MALDI Biotyper database version 7311

^b The list (score) is defined as the sum of weighted log (scores) for each single species according to Harju *et al* (20). p, *S. pneumoniae*; m, *S. mitis*; o, *S. oralis*; sp., any other species in the MALDI Biotyper database.

^c Specific peaks according to Marin *et al* (21)

^d Peak associated with *S. pneumoniae*

^e Peak associated with *S. oralis*

^f Peak associated with *S. mitis* and *S. oralis*

^g S, susceptible; R, resistant