Case Report

A Rare Cause of Stroke in Young Adults: Occlusion of the Middle Cerebral Artery by a Meningioma Postpartum

Stéphane Mathis,¹ Benoît Bataille,² Samy Boucebci,³ Marion Jeantet,⁴ Jonathan Ciron,¹ Laurene Vandamme,¹ and Jean-Philippe Neau¹

¹ Department of Neurology, CHU Poitiers, University of Poitiers, 2 rue de la Milétrie, 86021 Poitiers, France

² Department of Neurosurgery, CHU Poitiers, University of Poitiers, 2 rue de la Milétrie, 86021 Poitiers, France

³ Department of Radiology, CHU Poitiers, University of Poitiers, 2 rue de la Milétrie, 86021 Poitiers, France

⁴ Department of Pathology, CHU Poitiers, University of Poitiers, 2 rue de la Milétrie, 86021 Poitiers, France

Correspondence should be addressed to Stéphane Mathis; stephane.mathis@chu-poitiers.fr

Received 8 July 2013; Accepted 4 September 2013

Academic Editors: J. L. González-Gutiérrez, Y. Iwasaki, and Z. Siddiqi

Copyright © 2013 Stéphane Mathis et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Meningioma is the most common nonglial intracranial primary tumor. It is a slowly growing tumor and presents clinically by causing seizures along with neurological or neuropsychological deficit. However, acute presentation of meningioma is possible. We are reporting a case of cerebral infarction due to a sphenoid wing meningotheial meningioma (with progesterone receptor positivity) leading to an occlusion of the middle cerebral artery (MCA) in a 30-year-old right-handed woman (1 month after childbirth). After surgery, no new neurological event occurred, and she recovered most of her neurological functions. Strokes due to meningioma are a highly rare clinical occurrence but should be given serious consideration, particularly in young patients.

1. Introduction

Meningioma is the most common nonglial intracranial primary tumor, accounting for 13 to 26% of all primary intracranial tumors [1]. This slowly growing tumor arises from the arachnoid cells of the leptomeninges and presents clinically by causing seizures along with neurological or neuropsychological deficit [2]. Meningiomas variably express hormone receptors for progesterone, androgen, estrogen, placenta growth factor, and exogenous hormones [3, 4] and are also 2-3 times more common in women than in men [5].

2. Case Report

We report on a 30-year-old right-handed woman with a personal history of migraine admitted to the Department of Neurology, 1 month after childbirth, for acute onset of headache and complete paresis of her left hemibody, without sensory loss; National Institutes of Health Stroke Scale (NIHSS) was 10. Laboratory examination found no

abnormality. T1-weighted magnetic resonance (MR) imaging with gadolinium disclosed a right sphenoid wing meningioma measuring 35 mm in diameter (Figure 1(a)). Diffusion-weighted MR imaging showed a hyperintense area in the right deep middle cerebral artery (MCA) territory (Figure 1(b)), indicating acute cerebral infarction; MR angiography sequences revealed interrupted arterial flow in the right MCA (Figure 1(c)). Selective angiography of the right carotid artery demonstrated an occlusion of the M1 portion of the right MCA and a rich vascular stain of the tumor, fed primarily by intracavernous branches of the right carotid siphon (Figure 1(d)). No cause for MCA infarct other than the meningioma was found. The tumor was totally removed. The MCA trunk was set in the meningioma, and a thrombus of the right MCA was observed during the surgery. The final pathological diagnosis confirmed a meningotheial meningioma with progesterone receptor positivity (Figure 2). No new neurological event occurred after surgery, and over the next several weeks, the patient recovered most of her neurological functions, with only a moderate spasticity of

FIGURE 1: Brain MRI T1-weighted postgadolinium sequence showing homogenous enhancement of the meningioma (a). Brain MRI diffusion-weighted sequence showing acute deep MCA stroke (b). Brain MR angiography (axial sequence) showing occlusion of the right MCA (arrow) (c). Angiogram of the right internal carotid artery showing complete occlusion of the right MCA (arrow) and vascular blush of the meningioma (*) (d).

the upper and lower limbs. A brain MRI (six months later) showed no recurrence of meningioma.

3. Discussion

If encasement and intracranial arterial obstruction are common in cases of meningioma, strokes only rarely occur in association with this slowly growing tumor. In most of these cases, the different manifestations constitute a transient neurological deficit mimicking transient ischemic attacks probably due to hemodynamic stenoses. It has been suggested that meningeomal compression of the carotid artery may produce transient neurological symptoms including loss of consciousness, hemiparesis, paresthesia, and global transient amnesia [6]. However, complete cerebral infarction due to meningioma is exceedingly rare, and to our knowledge, only 6 other cases have been previously reported [7–10]: among these cases (including ours), 4 were MCA infarcts, 1 was an anterior cerebral artery infarct, 1 was a posterior

watershed stroke, and 1 was a pontine infarct. Most of the patients were under 50 years of age. In our observation, neurological symptoms appear during postpartum period, only one month after childbirth. As is the case, progesterone receptors are expressed in 50 to 80% of human meningeomas [11]. It is well known that the response of meningioma to increased serum progesterone level during the second half of pregnancy may account for accelerated growth of this tumor [12], whereas a classical clinical and radiological regression of meningeomas during postpartum could be observed [13]. Pregnancy certainly played a crucial role in increasing the tumor volume of our patient.

A compressive mechanism is usually suggested as an explanation for this phenomenon. In fact, meningeomas involving the cavernous sinus commonly lead to encasement and narrowing of the internal carotid artery (8 to 100% of cases) [8], and invasion of the internal carotid artery by cavernous sinus meningioma is likewise possible but rare [14]. However, the incidence of meningioma-related cerebral

FIGURE 2: Meningothelial meningioma. Hematoxylin eosin staining section (original magnification $\times 100$) showing characteristic cellular whorls (a). Immunohistochemistry (IHC) sections (original magnification $\times 100$): the Ki 67 labelling index (1/100 *Dako*) indicates a low grade (1-2%) of the meningioma (b); IHC for estrogen receptors (*Ventana*) showing no positive nuclei (c), whereas IHC for progesterone receptors (*Ventana*) showing positive nuclei (d).

ischemia by carotid artery compression is estimated at only 0.13% [6]. The reason why strokes are infrequent in this situation is still unknown, but it has been hypothesized that arteries (with high pressure) may be less compromised than cortical veins and dural sinuses (with low pressure) [6]. In conclusion, strokes due to meningioma are a highly rare clinical occurrence but should be given serious consideration, especially in young patients.

Conflict of Interests

The authors declare that they have no conflict of interests.

References

- [1] M. Bondy and B. Lee Ligon, "Epidemiology and etiology of intracranial meningiomas: a review," *Journal of Neuro-Oncology*, vol. 29, no. 3, pp. 197–205, 1996.
- [2] I. R. Whittle, C. Smith, P. Navoo, and D. Collie, "Meningiomas," *The Lancet*, vol. 363, no. 9420, pp. 1535–1543, 2004.
- [3] S. Pravdenkova, O. Al-Mefty, J. Sawyer, and M. Husain, "Progesterone and estrogen receptors: opposing prognostic indicators in meningiomas," *Journal of Neurosurgery*, vol. 105, no. 2, pp. 163–173, 2006.
- [4] E. B. Claus, P. M. Black, M. L. Bondy et al., "Exogenous hormone use and meningioma risk: what do we tell our patients?" *Cancer*, vol. 110, no. 3, pp. 471–476, 2007.
- [5] P. M. Black, "Hormones, radiosurgery and virtual reality: new aspects of meningioma management," *Canadian Journal of Neurological Sciences*, vol. 24, no. 4, pp. 302–306, 1997.
- [6] R. J. Komotar, S. C. Keswani, and R. J. Wityk, "Meningioma presenting as stroke: report of two cases and estimation of incidence," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 74, no. 1, pp. 136–137, 2003.
- [7] S. Bito and S. Sakaki, "Case of middle fossa meningioma with rapid development of symptoms due to obstruction of the middle cerebral artery," *Nō to Shinkei*, vol. 23, no. 8, pp. 953–957, 1971.
- [8] S. Heye, G. Maleux, J. van Loon, and G. Wilms, "Symptomatic stenosis of the cavernous portion of the internal carotid artery due to an irresectable medial sphenoid wing meningioma: treatment by endovascular stent placement," *American Journal of Neuroradiology*, vol. 27, no. 7, pp. 1532–1534, 2006.
- [9] M. S. Manzano-Palomo and J. A. Egidio, "Ictus pontino secundario a meningioma de foramen magno," *Revista de Neurologia*, vol. 40, no. 11, pp. 668–670, 2005.
- [10] J. Masuoka, F. Yoshioka, H. Ohgushi, M. Kawashima, and T. Matsushima, "Meningioma manifesting as cerebral infarction," *Neurologia Medico-Chirurgica*, vol. 50, no. 7, pp. 585–587, 2010.
- [11] A. Brandis, S. Mirzai, M. Tatagiba et al., "Immunohistochemical detection of female sex hormone receptors in meningiomas: correlation with clinical and histological features," *Neurosurgery*, vol. 33, no. 2, pp. 212–218, 1993.
- [12] Y. Saitoh, Y. Oku, S. Izumoto, and J. Go, "Rapid growth of a meningioma during pregnancy: relationship with estrogen

and progesterone receptors. Case report," *Neurologia Medico-Chirurgica*, vol. 29, no. 5, pp. 440–443, 1989.

- [13] J. S. Smith, A. Quiñones-Hinojosa, M. Harmon-Smith, A. W. Bollen, and M. W. McDermott, "Sex steroid and growth factor profile of a meningioma associated with pregnancy," *Canadian Journal of Neurological Sciences*, vol. 32, no. 1, pp. 122–127, 2005.
- [14] M. E. Shaffrey, V. V. Dolenc, G. Lanzino, W. P. Wolcott, and C. I. Shaffrey, "Invasion of the internal carotid artery by cavernous sinus meningiomas," *Surgical Neurology*, vol. 52, no. 2, pp. 167–171, 1999.