

## Case Report

# Paratubal Cystectomy in a Pregnant Woman Using the Single-Incision Laparoscopic Surgery (SILS) Technique

Luke William Nelson  and Elliot MacKenzie

Dunedin Public Hospital, 201 Great King Street, Dunedin 9010, New Zealand

Correspondence should be addressed to Luke William Nelson; [luke.nelson@southernhb.govt.nz](mailto:luke.nelson@southernhb.govt.nz)

Received 4 November 2021; Accepted 13 June 2022; Published 14 July 2022

Academic Editor: Maria Grazia Porpora

Copyright © 2022 Luke William Nelson and Elliot MacKenzie. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Introduction.* The proliferation of prenatal ultrasound has enhanced the detection of adnexal masses during pregnancy. The presentation necessitates a clear approach to investigation and treatment that balances both maternal and fetal risk. Laparoscopy is a safe approach to surgical management in the pregnant patient, and SILS may contribute to minimising perioperative complications. *Case Presentation.* We present the case of a 21-year-old female in her second trimester of pregnancy presenting with a large 20 cm right adnexal cyst. We proceeded with laparoscopic cystectomy via the SILS technique. There were no intraoperative complications, and she recovered well postoperatively. *Conclusion.* Laparoscopic resection of adnexal lesions is safe during pregnancy and should be favoured over the open approach. SILS minimises incision sites and has potential for reduction in perioperative morbidity.

## 1. Introduction

The proliferation of prenatal ultrasound has enhanced the detection of adnexal masses during pregnancy. With an incidence of 2%, the presentation necessitates a clear approach to investigation and treatment that balances both maternal and fetal risk [1]. In determining the appropriate management strategy, namely, intervention vs. conservative, there are considerations idiosyncratic to the pregnant patient that should be accounted for. The absence of malignant features on imaging and stable size may obviate the need for intervention and allow for conservative management [2].

Historical concerns regarding safety of laparoscopy in pregnancy have now largely been assuaged. There still exist considerations unique to the pregnant patient. For example, the gravid uterus may be susceptible to puncture during blind insufflation [3].

Single-incision laparoscopic surgery (SILS) is an emerging approach to laparoscopy. It confers improved cosmesis and, by limiting the surgery to a single-entry point, reduces

the potential for infection and bleeding [4]. Its chief disadvantages include a greater technical challenge and longer operative duration [5].

We present a case report of a 21-year-old pregnant woman who underwent a right paratubal cystectomy at 18 weeks gestation using a SILS approach.

## 2. Case Presentation

Our patient was a 21-year-old female who presented with a wanted pregnancy at 14 + 3 weeks gestation complicated by abdominal pain and vomiting. She had a medical history of 2 previous first trimester miscarriages and a raised body mass index of 33. She was taking appropriate folic acid supplementation. On examination, she had a palpable mass in the right lower abdominal quadrant, with associated tenderness and guarding. A vaginal examination was unremarkable, and the cervical os was closed. She had normal blood pressure and pulse. Her serum hCG was 15900.




FIGURE 1: Transverse ultrasound view of the right adnexa, demonstrating the cyst. Note its homogenous echotexture.

She proceeded to have a pelvic ultrasound which demonstrated a single live intrauterine pregnancy consistent with a gestational age of 14 + 3 weeks and a large right adnexal mass measuring 201 × 148 × 73 mm (1129 cc), likely arising from the right ovary. There were no septations, solid components, or abnormal vascularity noted (Figure 1).

She was admitted overnight for analgesia and antiemesis and discharged the next day following improvement in her symptoms. A planned MRI was conducted as an outpatient when she was 17 + 2 weeks gestation (Figure 2). This demonstrated interval growth of the cyst from the previous ultrasound scan and concurred with the presumptive diagnosis of a large right simple cyst measuring 190 × 170 × 100 mm (1680 cc).

The patient was referred to our gynaecological-oncology multidisciplinary team. It was determined that the cyst was benign in appearance, but surgical resection was necessitated. Cystectomy was favoured over drainage, for concerns regarding reaccumulation.

We proceeded to take her to theatre for laparoscopic right adnexal cystectomy at 18 + 2 weeks gestation. The operation was conducted under general anaesthesia by endotracheal intubation in Trendelenburg position with a urinary catheter in situ. Due to her gravid state, no transvaginal uterine manipulation was performed. Following skin preparation and application of sterile drapes, we made a 15 mm incision at the umbilicus. A SILS port (Applied Medical Gelseal) was placed at the umbilicus, through which we insufflated the peritoneal cavity with carbon dioxide. Via the SILS port, a 30-degree laparoscope and 2 working instruments were inserted. A large right adnexal mass extending to reach the liver was visualised and determined to be paratubal in origin (Figure 3).

Both ovaries and left fallopian tube were normal. Peritoneal washings were taken. A sharp incision was made on the cyst capsule followed by water-jet hydrodissection using a standard laparoscopic suction irrigation system under pressure. Following near complete dissection within its capsule, leaving base attached to maintain orientation, the cyst was punctured sharply with laparoscopic scissors under suction tip and suction immediately placed into cyst cavity. 1300 millilitres of clear serous fluid were drained without spill.

Patient size, gravid uterus, and instrument length then limited reach deep into pelvis from the umbilicus position to allow cyst excision and control of haemostasis at its base via the SILS port alone; hence, an accessory 5 mm port was placed in the left flank. We then proceeded to complete the excision of the cyst capsule using Ligasure bipolar device, delivering it extracorporeally via the SILS port. The right-sided fallopian tube was elongated and partially twisted. We untwisted the right fallopian tube following the cystectomy. Otherwise, both the right and left fallopian tubes and ovaries were normal and left intact bilaterally (Figure 4).

Pneumoperitoneum was released, and the SILS port was removed from the abdomen. The rectus sheath at the umbilicus was sutured with 0 Vicryl, and overlying skin was sutured with 3/0 Monocryl. The left-sided accessory port site was repaired with 3/0 Monocryl.

The estimated blood loss was approximately 75 millilitres, and no intraoperative complications arose. She had an uneventful postoperative course and was discharged the next day.

Histology confirmed the diagnosis of a benign serous cystadenoma. The patient was followed up at 22 + 2 weeks gestation, and her pregnancy was progressing well.

### 3. Discussion

The finding of an ovarian cyst during pregnancy presents a dilemma to the obstetrician. Should it be necessitated, surgery must be used judiciously and timed so that fetal risk is minimised. Amongst such patients, there is a preponderance of functional cysts. This is reflected by a 70% spontaneous resolution rate by the 2<sup>nd</sup> trimester [6]. Nevertheless, 1-6% of adnexal masses in pregnancy are malignant and necessitate timely resection and staging [7].

**3.1. Diagnosis and Selection for Surgery.** Ultrasound is the mainstay of investigation of ovarian masses in pregnancy, allowing the obstetrician to stratify patients according to features suggestive of malignancy [8]. Tumour markers such as the glycoprotein CA-125 are of reduced applicability in the pregnant patient and may rise as a consequence of pregnancy itself [9, 10]. MRI can provide a useful adjunct to




FIGURE 2: Sagittal view (MRI) of the right adnexal cyst. Note its simple appearance.




FIGURE 3: The partially dissected right adnexal cyst.




FIGURE 4: The right adnexa following complete excision of the cyst. Note the normal right ovary and gravid uterus.

diagnosis, through further characterisation of mass morphology [11]. Both methods are safe and do not expose the fetus to the effects of ionising radiation [12].

Our patient had radiographic features highly suggestive of a benign simple cyst. However, imaging alone cannot fully

replace pathological staging [13]. As such, pregnant patients with features of ovarian malignancy are a group who require surgical intervention. Alongside increasing the suspicion of malignancy, ovarian masses that are large (>10 cm) or increasing in size pose a risk of torsion or obstructed labour and are an indication for surgery [14]. Another important consideration is the risk of emergency surgery in the event of ovarian torsion and its apparent predisposition towards preterm birth [15].

Such risks must be balanced with the risks inherent to the surgery itself, both to fetus and mother. Delaying surgery until the second trimester affords time for resolution of functional cysts and avoids spontaneous first trimester miscarriage being falsely linked to the operation. Furthermore, organogenesis has largely concluded as well as the pregnancy's reliance upon the corpus luteum for progesterone [1, 16, 17].

In keeping with current practice, we performed the operation in the patient's second trimester for the aforementioned reasons. Additionally, operating before the third trimester reduces the technical burden that a large gravid uterus may place upon the surgeon. Our management included review by our gynaecologic-oncology multidisciplinary team. This is concordant with Vernooij et al. [18] who determined that mean survival time for ovarian malignancy is improved through consultation with a gynaecological oncology service [18].

**3.2. Laparoscopy and Pregnancy.** Multiple studies have corroborated the safety of laparoscopic management of adnexal masses in pregnant patients [19–21]. Compared to an open approach, laparoscopy may mitigate the risks of thromboembolism and maternal hypoventilation by offering earlier mobilisation and less reliance upon postoperative opioid analgesia, respectively [22]. Concerns regarding the risk of preterm birth when operating in the third trimester may be unfounded given that the preterm birth rate ranges between 7.3 and 11.7% in the general population [23, 24].

**3.3. Pneumoperitoneum.** An animal study performed by Barnard et al. [25] demonstrated reduced placental perfusion in the presence of maternal pneumoperitoneum [25]. Encouragingly, fetal perfusion and blood gas values were not adversely affected. These findings are yet to be confirmed or refuted by human studies. In light of its uncertain significance, multiple sources advocate for limiting the operating pneumoperitoneum to below 15 or even 12 mmHg [26–28]. Intra-abdominal pressures should be titrated to account for the already deleterious effect that the gravid uterus has upon visualisation [29].

**3.4. Electrosurgical Instrumentation.** Several case series have incorporated electrosurgical instruments into their laparoscopic surgeries on pregnant women without operative complications [15, 30, 31]. We used the bipolar tissue sealing device Ligasure to excise the cyst, without intraoperative or postoperative complication. The envelopment of the fetus in amniotic fluid is believed to be protective from energy-related injury. As with any application of electrosurgery, care

should be taken to avoid inadvertent trauma. This is especially salient in the pregnant patient in whom there is a paucity of evidence regarding its usage [16].

**3.5. Uterine Manipulation and Tocolysis.** Cervical manipulation of the pregnant uterus has the propensity to induce premature contractions, and as such, is contraindicated. Intra-abdominal surgery, particularly in the third trimester, may lead to uterine contractions [32]. However, Walsh et al. [33] demonstrated no reduction in preterm birth with prophylactic tocolysis [33]. As such, we did not manipulate the uterus during the course of our surgery, and tocolysis was not employed.

**3.6. SILS.** SILS is a novel approach to minimal access surgery. A single-entry point is placed, typically at the umbilicus, through which the laparoscope and all instruments are inserted [34]. Its efficacy and safety have been demonstrated across the spectrum of surgical disciplines [35].

Prospective studies comparing SILS to standard laparoscopic management of adnexal lesions are lacking. Several studies have demonstrated the feasibility and safety of SILS for adnexal lesions [36, 37]. A recent retrospective study demonstrated longer operative times when SILS was applied to adnexal lesions [38], whilst shorter operating times have been described by other applications of the technique [39]. Nevertheless, the potential for improved cosmesis and perioperative outcome should not be discounted based on scarcity of evidence.

Delay in widespread adoption of SILS may be in part due to its technical challenges. Convergence of all instruments through one port can reduce the working space available [40]. Furthermore, movement of the camera can be restricted by its proximity to the working instruments [4].

The applicability of SILS to the treatment of a variety of adnexal diseases has been established. Marcelli et al. [41] exhibited good success rates for SILS salpingectomy for ectopic pregnancy. Despite prolonging operative time, patients treated with SILS had shorter hospital stays than those who underwent conventional multiport laparoscopy [41]. Similarly, Loh et al. (2017) found outcomes from SILS management of ectopic pregnancy to be at least equivalent with conventional laparoscopy [42]. Dursun et al. [43] successfully treated 14 women with benign adnexal masses using SILS in combination with standard laparoscopic instruments. Optimisation of instrument and surgeon position compensated for difficulties encountered with instrument collision [43]. In the future, this issue may be avoided entirely with the advent of instruments designed specifically for SILS. Common to existing research surrounding SILS, these studies were marred by small sample sizes. Xiao et al. (2020) have since demonstrated the feasibility of SILS in laparoscopic management of adnexal disease, myomectomy, and cervical cerclage, specifically in the pregnant woman [44].

The importance of cosmesis cannot be understated given that the population who undergo laparoscopy for benign gynaecological disease largely consist of young women. By virtue of a single incision site, SILS has the potential to max-

imise patient satisfaction postoperatively [45]. Additionally, utilising the umbilicus as the sole incision site limits the potential for immediate and long-term postoperative pain [46].

In our case, due to locomotive restrictions and instrument length size, we were compelled to place an additional port into the left flank to remove the cyst capsule at the end. This was further compounded by the inability to manipulate the uterus due to its gravid state. Our experience is echoed in other reports on the usage of SILS in adnexal cysts. Rezai et al. [47] noted the difficulty faced through not being able to alter the gravid uterus' impact upon the operating field [47]. Also, 24% of patients in a study by Huang et al. [38] required the addition of an accessory port [38]. We were still able to excise the lesion using the SILS port in tandem with the accessory port, avoiding the placement of an additional trocar into the abdomen. It could also be argued that additional port placement after a large cyst decompression in the presence of a gravid uterus gives a safety margin on trauma to the uterus derived from an increased space created as in our case. Bariatric surgery size instruments could be utilised if the concern is instrument size alone. We did not have these available to us.

Ross et al. [48] proposed several potential drawbacks to the use of SILS. The presence of preexisting pelvic adhesions and excessive extra- and intraperitoneal fat may exacerbate the intrinsic difficulties with instrument triangulation [48]. This may render previous surgery and a raised BMI relative contraindications to SILS. However, in our experience, the elevated BMI of the patient did not substantially alter the use of SILS. Moreover, the availability of longer instruments may have averted the need to place an accessory port. This is a technical consideration which could be ameliorated with the development and dispersion of instruments specific to the SILS technique. On the discovery of extensive adhesions or an abdomen that is not amenable to SILS, there is always the recourse to convert to standard laparoscopy.

There is reasonable concern regarding the propensity of SILS to cause umbilical hernia due to the larger incision size for entry. The overall risk of port site hernia is a difficult entity to quantify, a systematic review estimated its prevalence as 0.5% for all laparoscopic surgery [49]. Gunderson et al. (2013) looked specifically at SILS, and adjusting for confounding variables, derived an umbilical hernia rate of 0.5% [50]. Nevertheless, there is not significant literature to reliably refute the hypothetical increased risk of herniation with SILS. In our case, we made a 15 mm incision at the umbilicus. We feel that given that this is only marginally larger than the Hasson entry technique incision the potential benefits of SILS outweigh the theoretical increased risk of umbilical hernia [51]. Overall, the application of SILS reduces the total number of incision sites and potential locations for herniation. Furthermore, fewer trocar insertions expose the patient to less risk of intra-abdominal trauma, bleeding, and pain.

Potential issues with instrument triangulation and obtaining adequate fulcrum in order to manipulate tissue adequately were countered by the flexibility of the Gelseal SILS port. This flexibility affords the surgeon adequate

instrument articulation; hence, we do not foresee significant challenges in intracorporeal suturing. Furthermore, the presence of a large volume cyst did not restrict our use of SILS as we decompressed the cyst prior to its removal. We expect that with the increasing prevalence of SILS, adaptations to its unique challenges will proliferate.

#### 4. Conclusion

Laparoscopic resection of adnexal lesions is safe during pregnancy and should be favoured over the open approach. SILS minimises incision sites and has potential for reduction in perioperative morbidity. The scope to add additional ports if needed makes SILS a feasible primary approach to laparoscopic cystectomy in the pregnant patient.

#### Data Availability

The data presented in this case report is available from the corresponding author upon request.

#### Consent

Informed consent has been obtained from the patient.

#### Disclosure

This study was performed as part of employment at our institution: Dunedin Public Hospital.

#### Conflicts of Interest

There are no conflicts of interest to declare.

#### References

- [1] J. Haan, M. Verheecke, and F. Amant, "Management of ovarian cysts and cancer in pregnancy," *Facts, Views and Vision in Obstetrics and Gynaecology*, vol. 7, no. 1, pp. 25–31, 2015.
- [2] K. M. Schmeler, W. W. Mayo-Smith, J. F. Peipert, S. Weitzen, M. D. Manuel, and M. E. Gordinier, "Adnexal masses in pregnancy: surgery compared with observation," *Obstetrics and Gynaecology*, vol. 105, 5, Part 1, pp. 1098–1103, 2005.
- [3] M. B. Reedy, H. L. Galan, W. E. Richards, C. K. Preece, P. A. Wetter, and T. J. Kuehl, "Laparoscopy during pregnancy. A survey of laparoendoscopic surgeons," *Journal of Reproductive Medicine*, vol. 42, no. 1, pp. 33–38, 1997.
- [4] N. Greaves and J. Nicholson, "Single incision laparoscopic surgery in general surgery: a review," *Annals of the Royal College of Surgeons of England*, vol. 93, no. 6, pp. 437–440, 2011.
- [5] S. A. Antoniou, O. O. Koch, G. A. Antoniou et al., "Meta-analysis of randomized trials on single-incision laparoscopic versus conventional laparoscopic appendectomy," *American Journal of Surgery*, vol. 207, no. 4, pp. 613–622, 2014.
- [6] R. L. Giuntoli, R. S. Vang, and R. E. Bristow, "Evaluation and management of adnexal masses during pregnancy," *Clinical Obstetrics and Gynaecology*, vol. 49, no. 3, pp. 492–505, 2006.
- [7] K. E. Webb, K. Sakhel, S. P. Chauhan, and A. Z. Abuhamad, "Adnexal mass during pregnancy: a review," *American Journal of Perinatology*, vol. 32, no. 11, pp. 1010–1016, 2015.
- [8] P. Aggarwal and S. Kehoe, "Ovarian tumours in pregnancy: a literature review," *European Journal of Obstetrics, Gynaecology and Reproductive Biology*, vol. 155, no. 2, pp. 119–124, 2011.
- [9] L. Rucker, Z. Walker, and B. Casey, "Case report: management of an early third trimester large adnexal mass in a singleton pregnancy," *Obstetrics and Gynaecology Cases-Reviews*, vol. 8, no. 1, p. 190, 2021.
- [10] G. S. Leiserowitz, "Managing ovarian masses during pregnancy," *Obstetrical and Gynaecological Survey*, vol. 61, no. 7, pp. 463–470, 2006.
- [11] I. A. Yakasai and L. A. Bappa, "Diagnosis and management of adnexal masses in pregnancy," *Journal of Surgical Technique and Case Report*, vol. 4, no. 2, pp. 79–85, 2012.
- [12] J. G. Ray, M. J. Vermeulen, A. Bharatha, W. J. Montanera, and A. L. Park, "Association between MRI exposure during pregnancy and fetal and childhood outcomes," *Journal of the American Medical Association*, vol. 316, no. 9, pp. 952–961, 2016.
- [13] A. Shaaban and M. Rezvani, "Ovarian cancer: detection and radiologic staging," *Clinical Obstetrics and Gynaecology*, vol. 52, no. 1, pp. 73–93, 2009.
- [14] M. K. Montes de Oca, S. K. Dotters-Katz, J. A. Kuller, and R. A. Previs, "Adnexal masses in pregnancy," *Obstetrical and Gynaecological Survey*, vol. 76, no. 7, pp. 437–450, 2021.
- [15] G. S. R. Lee, S. Y. Hur, J. C. Shin, S. P. Kim, and S. J. Kim, "Elective vs. conservative management of ovarian tumors in pregnancy," *International Journal of Gynaecology and Obstetrics*, vol. 85, no. 3, pp. 250–254, 2004.
- [16] E. Ball, N. Waters, N. Cooper et al., "Evidence-based guideline on laparoscopy in pregnancy," *Facts, Views and Visions in Obstetrics and Gynaecology*, vol. 11, no. 1, pp. 5–25, 2019.
- [17] T. W. Sadler, *Langman's Medical Embryology*, Lippincott Williams and Wilkins, Philadelphia, 11th edition, 2010.
- [18] F. Vernooij, P. Heintz, E. Witteveen, and Y. van der Graaf, "The outcomes of ovarian cancer treatment are better when provided by gynecologic oncologists and in specialized hospitals: a systematic review," *Gynaecologic Oncology*, vol. 105, no. 3, pp. 801–812, 2007.
- [19] P. Mathevet, K. Nessah, D. Dargent, and G. Mellier, "Laparoscopic management of adnexal masses in pregnancy: a case series," *European Journal of Obstetrics, Gynaecology and Reproductive Biology*, vol. 108, no. 2, pp. 217–222, 2003.
- [20] A. G. Rizzo, "Laparoscopic surgery in pregnancy: long-term follow-up," *Journal of Laparoendoscopic and Advanced Surgical Techniques*, vol. 13, no. 1, pp. 11–15, 2003.
- [21] Y. Lenglet, H. Roman, B. Rabishong et al., "Laparoscopic management of adnexal masses during pregnancy," *Gynaecology, Obstetrics and Fertility*, vol. 34, no. 2, pp. 101–106, 2006.
- [22] S. Morton, N. Cooper, J. Walravens-Evans, M. S. Nair, A. Karthikeyan, and W. Yoong, "Operative laparoscopy in advanced pregnancy beyond 20 weeks," *The Obstetrician and Gynaecologist*, vol. 22, no. 3, pp. 237–241, 2020.
- [23] National Institute for Health and Care Excellence, "Preterm labour and birth," 2015, October 2021 <https://www.nice.org.uk/guidance/ng25>.
- [24] K. B. Markham and M. Klebanoff, "Prevention of preterm birth in modern obstetrics," *Clinics in Perinatology*, vol. 41, no. 4, pp. 773–785, 2014.
- [25] J. M. Barnard, D. C. Chaffin, S. Droste, A. Tierney, and T. Phernetton, "Fetal response to carbon dioxide

- pneumoperitoneum in the pregnant ewe,” *Obstetrics and Gynaecology*, vol. 85, no. 5, pp. 669–674, 1995.
- [26] Society of American Gastrointestinal and Endoscopic Surgeons, *Guidelines for the Use of Laparoscopy during Pregnancy*, 2017, October 2021 <https://www.sages.org/publications/guidelines/guidelines-for-diagnosis-treatment-and-use-of-laparoscopy-for-surgical-problems-during-pregnancy/>.
- [27] M. Candiani, S. Maddalena, M. Barbieri, S. Izzo, D. Alberico, and S. Ronzoni, “Adnexal masses in pregnancy: fetomaternal blood flow indices during laparoscopic surgery,” *Journal of Minimally Invasive Gynaecology*, vol. 19, no. 4, pp. 443–447, 2019.
- [28] M. Ko, T. Lai, and S. Chen, “Laparoscopic management of complicated adnexal masses in the first trimester of pregnancy,” *Fertility and Sterility*, vol. 92, no. 1, pp. 283–287, 2009.
- [29] D. Nasioudis, D. Tsilimigras, and K. P. Economopoulos, “Laparoscopic cholecystectomy during pregnancy: a systematic review of 590 patients,” *International Journal of Surgery*, vol. 27, pp. 165–175, 2016.
- [30] S. H. Park, M. I. Park, J. S. Choi, J. H. Lee, H. O. Kim, and H. Kim, “Laparoscopic appendectomy performed during pregnancy by gynecological laparoscopists,” *European Journal of Obstetrics, Gynaecology and Reproductive Biology*, vol. 148, no. 1, pp. 44–48, 2010.
- [31] J. C. Chung, G. S. Cho, E. J. Shin, H. C. Kim, and O. P. Song, “Clinical outcomes compared between laparoscopic and open appendectomy in pregnant women,” *Canadian Journal of Surgery*, vol. 56, no. 5, pp. 341–346, 2013.
- [32] F. R. Nezhat, S. Tazuke, C. H. Nezhat, D. S. Seidman, D. R. Phillips, and C. R. Nezhat, “Laparoscopy during pregnancy: a literature review,” *Journal of The Society of Laparoscopic and Robotic Surgeons*, vol. 1, no. 1, pp. 17–27, 1997.
- [33] C. A. Walsh, T. Tang, and S. R. Walsh, “Laparoscopic versus open appendectomy in pregnancy: a systematic review,” *International Journal of Surgery*, vol. 6, no. 4, pp. 339–344, 2008.
- [34] S. Saeed and S. Miraj, “Single-incision laparoscopy surgery: a systematic review,” *Electronic Physician*, vol. 8, no. 10, pp. 3088–3095, 2016.
- [35] E. Chouillard, S. Alsabah, R. Daher et al., “Single-incision laparoscopy could be better than standard laparoscopy in right colectomy for cancer,” *Journal of Laparoendoscopic and Advanced Surgical Techniques*, vol. 26, no. 5, pp. 371–378, 2016.
- [36] P. F. Escobar, M. A. Bedaiwy, A. N. Fader, and T. Falcone, “Laparoendoscopic single-site (LESS) surgery in patients with benign adnexal disease,” *Fertility and Sterility*, vol. 93, no. 6, pp. 2074.e7–2074.e10, 2010.
- [37] M. A. Bedaiwy, D. Sheyn, L. Eghdami et al., “Laparoendoscopic single-site surgery for benign ovarian cystectomies,” *Gynaecologic and Obstetric Investigation*, vol. 79, no. 3, pp. 179–183, 2015.
- [38] K. J. Huang, K. T. Lin, C. J. Wu, Y. X. Li, W. C. Chang, and B. C. Sheu, “Single incision laparoscopic surgery using conventional laparoscopic instruments versus two-port laparoscopic surgery for adnexal lesions,” *Scientific Reports*, vol. 11, no. 1, p. 4118, 2021.
- [39] T. Shiraishi, T. Tominaga, T. Nonaka et al., “A learning curve in using organ retractor for single-incision laparoscopic right colectomy,” *Scientific Reports*, vol. 11, no. 1, p. 6546, 2021.
- [40] P. Garg, S. Misra, J. D. Thakur, and J. Song, “Single incision laparoscopic surgery ovarian cystectomy in large benign ovarian cysts using conventional instruments,” *Journal of Minimal Access Surgery*, vol. 7, no. 4, pp. 232–235, 2011.
- [41] M. Marcelli, C. Lamourdedieu, A. Lazard, L. Cravello, M. Gamerre, and A. Agostini, “Salpingectomy for ectopic pregnancy by transumbilical single-site laparoscopy with the SILS system,” *European Journal of Obstetrics, Gynaecology and Reproductive Biology*, vol. 167, no. 1, pp. 67–70, 2012.
- [42] A. Loh, M. P. Torrizo, and Y. W. Ng, “Single incision laparoscopic surgery for surgical treatment of tubal ectopic pregnancy: a feasible alternative to conventional laparoscopy,” *Journal of Gynaecologic Surgery*, vol. 33, no. 2, pp. 61–67, 2017.
- [43] P. Dursun, T. Tezcaner, H. B. Zeyneloglu, I. Alyazici, A. Haberal, and A. Ayhan, “Adnexal masses treated using a combination of the SILS port and noncurved straight laparoscopic instruments: Turkish experience and review of the literature,” *Minimally Invasive Surgery*, vol. 2013, Article ID 836380, 8 pages, 2013.
- [44] J. Xiao, K. Fu, K. Duan, J. Wang, S. Sunkara, and X. Guan, “Pregnancy-preserving Laparoendoscopic Single-site Surgery for Gynecologic Disease: A Case Series,” *Journal of Minimally Invasive Gynaecology*, vol. 27, no. 7, pp. 1588–1597, 2020.
- [45] F. R. Borle, B. Mehra, and A. Singh, “Comparison of cosmetic outcome between single-incision laparoscopic cholecystectomy and conventional laparoscopic cholecystectomy in rural Indian population: a randomized clinical trial,” *Indian Journal of Surgery*, vol. 77, no. S3, pp. 877–880, 2015.
- [46] B. S. M. Chern, S. Lakhotia, C. K. Khoo, and A. Y. M. Siow, “Single incision laparoscopic surgery in gynecology: Evolution, current trends, and future perspectives,” *Gynaecology and Minimally Invasive Therapy*, vol. 1, no. 1, pp. 9–18, 2012.
- [47] S. Rezai, A. C. Hughes, E. Wang et al., “Laparoendoscopic single-site cystectomy in pregnancy for a benign Mullerian serous cystadenofibroma: a case report with review of literature,” *Obstetrics and Gynaecology International Journal*, vol. 9, no. 4, pp. 234–236, 2018.
- [48] S. B. Ross, W. C. Clark, C. A. Morton, and A. S. Rosemurgy, “Access for Laparoendoscopic Single Site Surgery,” *Diagnostic and Therapeutic Endoscopy*, vol. 2010, Article ID 943091, 7 pages, 2010.
- [49] H. A. Swank, I. M. Mulder, C. F. Chapelle, J. B. Reitsma, J. F. Lange, and W. A. Bemelman, “Systematic review of trocar-site hernia,” *British Journal of Surgery*, vol. 99, no. 3, pp. 315–323, 2012.
- [50] C. C. Gunderson, J. Knight, J. Ybanez-Morano et al., “The Risk of Umbilical Hernia and Other Complications with Laparoendoscopic Single-Site Surgery,” *Journal of Minimally Invasive Gynaecology*, vol. 19, no. 1, pp. 40–45, 2012.
- [51] A. K. Sangrasi, A. I. Memon, M. M. Memon, M. R. Abbasi, A. A. Laghari, and J. N. Qureshi, “A safe quick technique for placement of the first access port for creation of pneumoperitoneum,” *Journal of the Society of Laparoscopic and Robotic Surgeons*, vol. 15, no. 4, pp. 504–508, 2011.