

Case Report

Management of a Complex Supracondylar Periprosthetic Femur Fracture with Intramedullary Strut Allograft and Bilateral Locking Plates

Rita Moukarzel,¹ Dany Aouad ,² Mohammad Daher,³ Wendy Ghanem,² Hady Ezzeddine,² and George El Rassi²

¹Medical School, Lebanese American University Medical Center, Lebanese American University, P.O. Box 13-5053, Chouran Beirut: 1102 2801, Lebanon

²Department of Orthopedic Surgery and Traumatology, Saint Georges University Medical Center, Balamand University, P.O.Box 166378, Achrafieh, Beirut 1100 2807, Lebanon

³Faculty of Medicine, University Saint Joseph, P.O. Box 17-5208, Mar Mikhael, Beirut, Lebanon

Correspondence should be addressed to Dany Aouad; dany_aouad@hotmail.com

Received 15 August 2021; Accepted 8 November 2021; Published 17 November 2021

Academic Editor: Werner Kolb

Copyright © 2021 Rita Moukarzel et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supracondylar periprosthetic femoral fractures occurring above total knee replacements have been considered a rare entity. However, they continue to increase in frequency with the increasing number of arthroplasties and the improvement in morbidity and mortality in the concerned patient population. The management of periprosthetic distal femoral fractures is a challenging orthopedic problem. In this brief communication, a case of 49-year-old woman with rheumatoid arthritis who sustained a low distal comminuted periprosthetic femoral fracture is presented. Her fracture was eventually managed with an intramedullary fibular strut allograft and bilateral locking plate placement reaching satisfactory healing and restoration of alignment. The primary aim of this report is to provide insight into this novel technique as a successful alternative to other standard surgical options.

1. Introduction

Supracondylar periprosthetic femoral fractures have been considered a rare complication of total knee replacements. However, it is no surprise that they continue to increase in frequency alongside the rising number of total knee arthroplasties performed yearly and the improved quality of life and life expectancy of the concerned patient population [1].

These fractures generally occur in the geriatric population aged above 60 years with associated osteopenic or osteoporotic bone quality [2]. Commonly, the fracture is secondary to minimal low-velocity trauma [3]. In the setting of total knee arthroplasties, the supracondylar distal femur is the most frequent location for periprosthetic fractures [4], which are specifically associated with comminution [5]. Soininvaara et al. reported up to 25.5% rapid bone loss in

the distal femoral bone in the first 6-month period posttotal knee arthroplasty [6]. In fact, management with bisphosphonates after total knee arthroplasty helps decrease the periprosthetic osteopenia [7].

Distal periprosthetic femoral fractures can be managed either conservatively or surgically. Nevertheless, they are all managed surgically in patients tolerable of anesthesia who are otherwise ambulatory in order to prevent the complications of conservative management [8].

This review shares the management of a complicated periprosthetic distal femoral fracture with a novel combination surgical technique that uses an intramedullary fibular strut allograft supplemented with bilateral plate-and-screw placement. This method was used on a 49-year-old woman, with rheumatoid arthritis, who sustained a low distal comminuted periprosthetic femoral fracture about a total knee




FIGURE 1: Preoperative A-P (a) and lateral (b) radiographs showing a complex left supracondylar distal femoral fracture adjacent to a total knee arthroplasty.

arthroplasty (TKA). The fracture was initially managed with surgical open reduction and internal fixation (ORIF) with lateral plate placement but failed to achieve appropriate union and alignment. Ultimately, the fracture was successfully managed with an intramedullary fibular strut allograft with bilateral locking plate placement.

The effective use of retrograde intramedullary fibular strut allograft is proposed to augment plate fixation especially when the distal portion is small for instrument insertion or when the bone stock is low with increased comminution. The primary aim of this article is to provide insight into this novel surgical technique as a satisfactory alternative to traditional treatment options for managing periprosthetic distal femoral fractures, particularly in osteopenic patients.

2. Case Report

This is a case of a 49-year-old female patient, known to have juvenile rheumatoid arthritis currently managed with methotrexate and adalimumab. The patient had undergone a bilateral total knee replacement ten years prior to presentation.

She initially presented after sustaining a low-energy fall from standing height. Plain radiographs were done and showed a supracondylar displaced periprosthetic fracture of the left distal femur (Figure 1).

The fracture was comminuted with segmental bone defect in the distal femoral region. The patient underwent surgical open reduction and internal fixation (ORIF) with bone graft and lateral plate-and-screw (Figure 2).

The surgery was performed with neither intraoperative nor direct postoperative complications, and the patient was discharged one week postoperatively. The patient was followed with a series of radiographs to assess for satisfactory healing, alignment, and stable construct fixation. One month postoperatively, a follow-up radiograph showed fracture reduction with start of callus formation (Figure 3).

Two months postoperatively, follow-up radiographs (Figure 4) showed nonunion with severe comminution in the distal femoral region characterized by significantly low bone stock, which was also seen in the radiographs that followed in seven and nine months postoperatively (Figure 5).

A computed tomography scan with 3D reconstruction was done one year after the surgery showing failure of hardware, nonunion, and malalignment (Figure 6).

Consequently, an elective surgical repair was scheduled. The primary lateral plate was removed, and an intramedullary fibular allograft was introduced through the intercondylar region into the medullary cavity supplemented with lateral plate fixation (lateral curved LCP 8 hole condylar plate (Synthes, Beirut, Lebanon)) and medial




FIGURE 2: Postoperative lateral (a) and A-P (b, c) radiographs done in the OR, showing reduction of fracture status post management with ORIF and lateral plate-and-screw fixation.

minimally invasive (MIS) plate (medial curved LCP 18 hole condylar plate (Synthes, Beirut Lebanon)) fixation. Demineralized bone matrix (DBM) and bone morphogenic protein-2 (BMP-2) were also inserted.

Follow-up radiographs (Figure 7) done one month postoperatively showed stable fixation and alignment with beginning of healing and callus formation.

In addition, further follow-up imaging done at three, five, and seven months postoperatively (Figure 8) showed improvement in healing.

The patient progressively improved to a painless, full range of motion mobilization of the knee. She suffered no postoperative complications of graft rejection, infection, or mechanical instability on weightbearing.




FIGURE 3: Postoperative follow-up A-P radiographs done one month postoperatively showing reduced fracture and beginning of callus formation in the distal femoral region.




FIGURE 4: Postoperative follow-up A-P and lateral radiographs done two months postoperatively showing nonunion with severe comminution in the distal femoral region which is also associated with low bone stock.

Ten months after the surgery, the patient was admitted for bilateral plate removal with bone graft placed in screw holes. Cultures obtained two months later revealed negative tissue and serum results, and plain radiographs (Figure 9) showed complete healing of the fracture with a satisfactory alignment.

3. Discussion

Supracondylar periprosthetic femoral fractures occur usually in geriatric populations [2] due to low velocity traumas such as a fall from standing height [3]. Osteoporotic and osteopenic bone is the major risk factors for




FIGURE 5: Postoperative follow-up AP and lateral radiographs done nine months after the surgery still showing nonunion, comminution, and low bone stock (yellow arrows).




FIGURE 6: Postoperative follow-up with 3D reconstruction done one year postoperatively showing fracture nonunion (yellow circle), malalignment, and plate breakage (yellow arrow).

such injuries [2]. Other predisposing factors include female gender, rheumatoid arthritis, chronic steroid use, neurological diseases and recurrent falls, intraoperative anterior femo-

ral notching, and revision arthroplasty [1, 3], among which risk factors leading to low bone density considered more important [4].



FIGURE 7: Postoperative follow-up radiographs done one month after revision surgery showing satisfactory alignment.

Prognostically, periprosthetic femoral shaft fractures are associated with better postoperative outcomes compared to periprosthetic distal femoral fractures which carry high postoperative morbidity and mortality [1]. Hence, periprosthetic distal femoral fractures are technically challenging to manage, particularly in elderly osteopenic patients with associated thin cortices and loss of bone stock [9]. The deficient bone density in the distal femoral region as well as the low fracture localization adjacent to an arthroplasty component [2] obligates the modification of traditional fixation techniques. Specifically, low fractures with a little distal osteopenic fragment would impede strong fixation [10].

Although the management of these fractures has been increasingly widely practiced, postoperative complications continue to be high in these patients regardless of the technique used. These include reduced knee range of motion, residual fracture malunion, nonunion, malalignment, infection, and perioperative death [3]. Therefore, the treatment goals must comprise painless healing, early restoration of range of motion and weight-bearing, uncomplicated fracture union, radiographic alignment restoration, and return to preinjury function [1]. Reestablishing appropriate bone stock and ensuring prosthesis mechanical stability are also essential considerations to safeguard treatment success [1].

Preoperative evaluation must take into account the anatomic fracture site with respect to the arthroplasty component, prosthesis stability and type, local bone stock, bone displacement, and patient's preinjury ambulatory status and past medical history [1, 3] in order to decide on the most suitable management technique.

This injury can both be managed conservatively and surgically. Conservative treatment primarily entails cast



FIGURE 8: Postoperative follow-up AP radiograph of both knees nine months after revision surgery showing improvement in healing and satisfactory alignment.

immobilization and is usually reserved for undisplaced fracture types [2]. Nonconservative surgical treatment options include open reduction and internal fixation (ORIF), external fixation, anterograde or retrograde intramedullary nailing, internal fixation with locking plates, and distal femoral replacement, among others [5]. The latter must be considered after considering all treatment options due to its disastrous complications [11].

Surgical management is considered superior to conservative treatment because the latter is associated with prolonged immobilization and carries increased risks of nonunion and reoperation [2]. Although nonsurgical management avoids perioperative complications, Moran et al. mentions that 12-40% of the cases managed nonoperatively resulted in nonunion, and 15-30% of cases required reoperation [12]. Also, progression from nondisplaced to displaced fractures ensued in many cases, which required close radiologic follow-up [2]. Therefore, in most cases, orthopedists opt for surgical fixation as the primary option for managing these fractures. In most cases, ORIF with conventional plate fixation is performed as it presents a safe and minimally invasive surgical technique. In cases where the fracture is close to the arthroplasty component anteriorly, retrograde intramedullary nail combined with plate-and-screw fixation is used [8]. However, in cases like the one presented above, where the fracture is too distal and the metaphyseal region is severely




FIGURE 9: Postoperative radiographs two months after hardware removal showing complete healing of the fracture with satisfactory alignment.

comminuted and osteopenic for plate placement or nail insertion, using an intramedullary bulk allograft can be of aid. The intramedullary graft provides stable reduction and fixation for subsequent plate insertion. A multicenter experience by Rollo et al. showed that a combination of both strut allografts and plating may be the most efficient method of treatment of femoral periprosthetic fractures [13]. It is even more efficient than the treatment with minimally invasive plate osteosynthesis [14].

The aim of treating a periprosthetic fracture above TKA and severe osteopenia in a weightbearing bone must primarily target mechanical stability of the fixation construct. Intraoperatively, the fibular allograft was initially introduced into the distal femoral medullary cavity through the intercondylar notch at the arthroplasty construct [5]. This ensured adequate fixation of the fracture for placement of the lateral locking plate and the minimally invasive medial MIS plate. This technique, however, is only performed by highly skilled orthopedic surgeons able to manipulate allografts for major reconstruction [5].

Ultimately, although this surgical technique presents prognostic and mechanical advantages to overcome these fractures, it certainly is not free of limitations. Just like any other operation, fracture reduction does carry with it a risk of nonunion, malalignment, and infection. Moreover, the use of an allograft might present with graft complications, like host-graft rejection and disease transmission, for which one must safeguard appropriate consideration [15].

4. Conclusion

In conclusion, patients presenting with a low periprosthetic distal femoral fracture in the setting of severe osteopenia and comminution should be assessed promptly for displacement, anatomic site of fracture, and arthroplasty component in order to ensure the management technique with the best

outcome. Using an intramedullary fibular strut allograft is believed to augment the stability supplemented by bilateral plate insertion, and that it is a feasible alternative in managing these complicated fractures. The proposed technique is worth putting into practice as it provides good postoperative outcomes and improved quality of life in the concerned patient population.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Consent

Full consent from the patient was obtained for publishing this article and images.

Conflicts of Interest

The authors declare no conflict of interest regarding the publication of this article.

References

- [1] W. M. Ricci, "Periprosthetic femur fractures," *Journal of Orthopaedic Trauma*, vol. 29, no. 3, pp. 130–137, 2015.
- [2] D. A. Quinzi, S. Childs, J. S. Lipof, S. P. Sojin, and B. F. Ricciardi, "The treatment of periprosthetic distal femoral fractures after total knee replacement," *JBJS Reviews*, vol. 8, no. 9, pp. 1–12, 2020.
- [3] E. T. Su, H. Dewal, and P. E. CesareDi, "Periprosthetic femoral fractures above total knee replacements," *Journal of the American Academy of Orthopaedic Surgeons*, vol. 12, no. 1, pp. 12–20, 2004.

- [4] C. L. Boulton and E. K. Rodriguez, "treatment of periprosthetic fractures after total knee arthroplasty," *Current Orthopaedic Practice*, vol. 20, no. 1, pp. 58–64, 2009.
- [5] A. Kumar, I. Chambers, G. Maistrelli, and P. Wong, "Management of periprosthetic fracture above total knee arthroplasty using intramedullary fibular allograft and plate fixation," *The Journal of Arthroplasty*, vol. 23, no. 4, pp. 554–558, 2008.
- [6] T. Soininvaara, H. Kröger, J. S. Jurvelin, H. Miettinen, O. Suomalainen, and E. Alhava, "Measurement of bone density around Total knee arthroplasty using fan-beam dual energy X-ray absorptiometry," *Calcified Tissue International*, vol. 67, no. 3, pp. 267–272, 2000.
- [7] M. Shi, L. Chen, H. Wu et al., "Effect of bisphosphonates on periprosthetic bone loss after total knee arthroplasty: a meta-analysis of randomized controlled trials," *BMC Musculoskeletal Disorders*, vol. 19, no. 1, pp. 177–178, 2018.
- [8] A. Nauth, B. Ristevski, T. Bégué, and E. H. Schemitsch, "Periprosthetic distal femur fractures: current concepts," *Journal of Orthopaedic Trauma*, vol. 25, Supplement 2, pp. S82–S85, 2011.
- [9] H. Lindahl, G. Garellick, H. Regné, P. Herberts, and H. Malchau, "Three hundred and twenty-one periprosthetic femoral fractures," *The Journal of Bone & Joint Surgery*, vol. 88, no. 6, pp. 1215–1222, 2006.
- [10] S. Kalyanasundaram, B. Pankapilly, J. Varughese, G. Jacob, and V. K. Menon, "Hybrid stabilization of periprosthetic distal femoral fractures following total knee replacement: technique and results," *Techniques in Orthopaedics*, vol. 32, no. 2, pp. 126–130, 2017.
- [11] G. Rollo, P. Pichierri, P. Grubor et al., "The challenge of non-union and malunion in distal femur surgical revision," *Medicinski Glasnik*, vol. 16, 2019.
- [12] M. C. Moran, "Femoral periprosthetic fractures: nonoperative treatment," in *Surgical Techniques in Total Knee Arthroplasty*, G. R. Scuderi and A. J. Tria, Eds., pp. 545–552, Springer, New York, 2002.
- [13] G. Rollo, E. M. Bonura, G. Huri et al., "Standard plating vs. cortical strut and plating for periprosthetic knee fractures: a multicentre experience," *Medicinski Glasnik*, vol. 17, no. 1, 2020.
- [14] G. Rollo, M. Ronga, E. M. Bonura et al., "Surgical treatment of multifragmentary segmental femur shaft fractures with orif and bone graft versus mipo: a prospective control-group study," *Medicinski Glasnik*, vol. 17, no. 2, pp. 334–344, 2020.
- [15] T. Snoap, M. Jaykel, and J. Roberts, "Antegrade endosteal fibular strut augmentation for periprosthetic femoral fracture above stemmed total knee arthroplasty," *Current Orthopaedic Practice*, vol. 28, no. 1, pp. 104–107, 2017.