

Case Report

Vulva Fibroadenoma Associated with Lactating Adenoma in a 26-Year-Old Nigerian Female

C. C. Anunobi,¹ F. J. N. Obiajulu,¹ A. A. F. Banjo,¹ and A. O. E. Okonkwo²

¹ Department of Anatomic and Molecular Pathology, Lagos University Teaching Hospital, PMB 12003, Lagos 101014, Nigeria

² Angel Hospital, Lagos, Nigeria

Correspondence should be addressed to C. C. Anunobi; dozieanunobi@yahoo.com

Received 3 June 2013; Accepted 28 July 2013

Academic Editors: A. Agaimy and A. Usubutun

Copyright © 2013 C. C. Anunobi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Vulva lactating adenoma is rare and may be misdiagnosed as adenocarcinoma in frozen section and aspiration cytology if breast tissue is not anticipated. **Objective.** To raise the awareness of lactating vulva ectopic breast lesion among clinicians and pathologists. **Case Report.** We present a case of vulva fibroadenoma associated with lactating adenoma in a 26-year-old Nigerian female. **Conclusion.** The rarity of vulva ectopic breast tissue can present a diagnostic challenge for both the clinician and the anatomical pathologist. Once excisional biopsy is done and the lesion confirmed histologically, the anxious patient can be reassured.

1. Introduction

Ectopic breast tissue is found along the primitive embryonic milk lines which extend from the axilla to the medial side of the groin [1]. Normally in humans, only a small part of the milk line persists in the mid-thoracic region. Incomplete involution anywhere outside this midthoracic region and along the primitive milk streak can result in ectopic mammary tissue. The reported incidence of ectopic breast in women is 2–6%, but only a few of such case reports arising from the vulva exist in the literature [2]. Deaver reported that Hartung in 1872 described the first fully formed mammary gland in the vulva [3]. Ectopic breast tissue responds to hormones and may undergo benign or malignant process as those of normally located breast tissue [4]. Although lactational changes can occur in ectopic breast, lactating adenoma in the vulva is extremely rare [4].

We describe a case of ectopic vulva breast tissue presenting as fibroadenoma associated with lactating adenoma in a breast feeding woman.

2. Case Report

A 26-year-old G₄P₃⁺¹ woman presented to a peripheral hospital with a painless vulva swelling of 6 months duration in

the second trimester of her pregnancy. The swelling was first noticed as a small nodule that grew appreciably in size during pregnancy and postpartum. Physical examination revealed a 4 cm × 2 cm mass located at the right lip of the labium majus. She delivered a set of twins after which the mass was excised and sent to the Department of Anatomic and Molecular Pathology of Lagos, University Teaching Hospital, Idi-Araba, Lagos for histological analysis.

On gross examination, the specimen was an encapsulated piece of grayish white tissue measuring 3 cm × 2.5 cm × 2 cm. Cut section showed bulging lobulated grayish white surface (Figure 1).

The specimen was processed and evaluated with haematoxylin and eosin staining.

Histological sections showed an encapsulated breast tissue consisting of combined proliferation of epithelial and mesenchymal elements. The fibrocellular stroma proliferates around the tubular ducts. The ducts are of different sizes and shapes and are lined by inner columnar epithelial cells and outer myoepithelial cells. An area of lactating adenoma is seen with glands showing secretory activity. Large alveolar spaces separated by fine fibrovascular trabeculae are seen (Figures 2 and 3).

Immunohistochemical studies were also done and the cells reacted positively for ER and PR (Figures 4 and 5).



FIGURE 1: Encapsulated oval tissue with lobulated grayish white cut surface.




FIGURE 2: (H&E $\times 100$) Encapsulated growth (fibroadenoma) consisting of compressed glands and ducts within a dense fibrocollagenous stroma.




FIGURE 3: ($\times 100$) Encapsulated breast lesion showing dilated glands and ducts with secretory activity.




FIGURE 4: ($\times 400$) Ductal cells with strong estrogen receptor positivity.




FIGURE 5: ($\times 400$) Ductal cells with weak progesterone receptor positivity.

Avidin biotin complex type method with antibody dilution of 1:100 was used.

A diagnosis of fibroadenoma associated with lactating adenoma occurring in a vulva ectopic breast tissue was made. The postoperative period was uneventful.

3. Discussion

Ectopic breast tissue in the vulva is a rare lesion [5]. This might be related to the fact that the vulva is located at the inferior end of the primitive mammary ridge. The lesions may undergo benign or malignant changes, as they often respond to hormones as those of normally located breast [4]. Dhaoui et al. stated that about 50 cases of ectopic benign and 20 malignant breast lesions have been reported in the literature while lactating adenoma in the vulva is extremely rare [4]. Only four cases of lactating adenoma have been reported [4]. Most ectopic lactating adenomas have been noted within the age of 23 to 39 years, with measurement of 0.8 and 4 cm [6, 7]. The index patient is 26 years old and the mass measured an average of 3.5 cm.

The common origin of the ectopic breast with that of the breast at normal location can be explained by the effect of pregnancy and lactation on the ectopic breast tissue as noted in the index patient whose swelling grew significantly through pregnancy and the period of lactation. This overlapping effect of pregnancy and lactation on ectopic breast has been described by some authors [4, 6].

An alternative theory suggests that this tissue represents the normal constituents of the anogenital area that probably do not represent true breast tissue [8, 9]. These glands are now referred to as anogenital mammary-like glands (AGMLG) by some authors [10]. Research work on the largest series of these lesions done by Kazakov et al. [10] was published in 2010. It included among others fibroadenoma of the vulva with lactational changes in a 30-year-old female [10].

The cytologic features of lactational lesions of ectopic breast pose special problems in diagnosis by fine needle aspiration [6]. Vulva lactating adenoma may be misdiagnosed as adenocarcinoma in frozen section and aspiration cytology if breast tissue is not anticipated [6].

To the best of our knowledge, this is the first case report of vulva fibroadenoma associated with lactating adenoma in a young African woman.

Complete excision of ectopic breast tissue in the vulva should be considered since the lesion can undergo malignant transformation [11, 12].

4. Conclusion

The rarity of vulva ectopic breast tissue can present a diagnostic challenge for both the clinician and the anatomical pathologist. It is our opinion that this case report will raise the awareness of vulva ectopic breast lesion among clinicians. Once excisional biopsy is done and the lesion confirmed histologically, the anxious patient can be reassured.

References

- [1] S. Pathak and J. Preston, "A rare case of multiple accessory breast tissue in the axillae, lower abdomen and vulval areas," *Journal of Obstetrics and Gynaecology*, vol. 27, no. 5, pp. 531–533, 2007.
- [2] D. V. Kazakov, D. V. Spagnolo, D. Kacerovska, and M. Michal, "Lesions of anogenital mammary-like glands: an update," *Advances in Anatomic Pathology*, vol. 18, no. 1, pp. 1–28, 2011.
- [3] J. B. Deaver and J. McFarland, *The Breast: Its Anomalies, Its Disease and Their Treatment*, P. Blackistion's Sons, Philadelphia, Pa, USA, 1917.
- [4] A. Dhaoui, H. Nfoussi, N. Kchir, and S. Haouet, "Vulvar lactating adenoma associated to a fibroadenoma: common neoplasms in an uncommon site," *The Pan African Medical Journal*, vol. 13, article 47, 2012, <http://www.panafrican-med-journal.com>.
- [5] N. N. Mahajan and L. Choudhrie, "Fibroadenoma of ectopic breast tissue in the vulva: a case report," *Journal of Obstetrics and Gynecology of India*, vol. 60, no. 6, pp. 529–530, 2010.
- [6] M. F. O'Hara and D. L. Page, "Adenomas of the breast and ectopic breast under lactational influences," *Human Pathology*, vol. 16, no. 7, pp. 707–712, 1985.
- [7] M. E. Reeves and A. Tabuenca, "Lactating adenoma presenting as a giant breast mass," *Surgery*, vol. 127, no. 5, pp. 586–588, 2000.
- [8] S. C. J. van der Putte, "Anogenital "sweat" glands: histology and pathology of a gland that may mimic mammary glands," *American Journal of Dermatopathology*, vol. 13, no. 6, pp. 557–567, 1991.
- [9] S. C. J. van der Putte, "Mammary-like glands of the vulva and their disorders," *International Journal of Gynecological Pathology*, vol. 13, no. 2, pp. 150–160, 1994.
- [10] D. V. Kazakov, D. V. Spagnolo, C. J. Stewart et al., "Fibroadenoma and phyllodes tumors of anogenital mammary-like glands: a series of 13 neoplasms in 12 cases, including mammary-type juvenile fibroadenoma, fibroadenoma with lactation changes, and neurofibromatosis-associated pseudoangiomatous stromal hyperplasia with multinucleated giant cells," *American Journal of Surgical Pathology*, vol. 34, no. 1, pp. 95–103, 2010.
- [11] G. Lopes, T. DeCesare, G. Ghurani et al., "Primary ectopic breast cancer presenting as a vulvar mass," *Clinical Breast Cancer*, vol. 7, no. 3, pp. 278–279, 2006.
- [12] D. A. Farcy, D. Rabinowitz, and M. Frank, "Ectopic glandular breast tissue in a lactating young woman," *Journal of Emergency Medicine*, vol. 41, no. 6, pp. 627–629, 2011.

