Case Report

Nasal Chondromesenchymal Hamartoma: Rare Case Report in an Elderly Patient and Brief Review of Literature

Kanish Mirchia and Rana Naous

Department of Pathology, SUNY Upstate Medical University, Syracuse, NY 13210, USA

Correspondence should be addressed to Kanish Mirchia; mirchiak@upstate.edu

Received 29 May 2018; Accepted 24 September 2018; Published 14 October 2018

Academic Editor: Stefan Pambuccian

Copyright © 2018 Kanish Mirchia and Rana Naous. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hamartomas are considered a mixture of nonneoplastic tissue, which may be indigenous to a different location in the body. As such, they may be epithelial, mesenchymal, or mixed. In the sinonasal region, the following hamartomatous lesions are considered to lie on a spectrum and include respiratory epithelial adenomatoid hamartoma (REAH), chondro-osseous respiratory epithelial adenomatoid hamartoma (COREAH), and nasal chondromesenchymal hamartoma (NCMH). To our knowledge, less than 50 cases of sinonasal hamartomas have been reported in the English literature so far with NCMH being very rare and primarily a tumor in infancy, with only 2 cases reported in individuals older than 16 years of age. We report a highly unusual case of a NCMH in the right maxillary sinus of a 70-year-old female.

1. Case Report

A 70-year-old female presented with a two-year history of slowly growing, nonpainful maxillary sinus mass. She has a history of chronic maxillary sinusitis corresponding to presentation of the mass, with the first episode reported in 2014. Computed tomography (CT) imaging revealed an erosive right maxillary sinus mass (2.5 x 2.1 cm) with bony destruction.

Surgical excision of the right maxillary sinus mass revealed a fragmented, white, vaguely nodular, and whorled lesion. Histological examination revealed fragments of respiratory-type epithelium with focal cystic invagination and associated squamous metaplasia [Figure 1]. The underlying stroma consisted of a variably cellular, benign spindle cell proliferation with an associated background of hyalinization [Figure 2], calcification and ossification [Figure 3], and focal chondroid change [Figure 4] in a vague lobule-like arrangement. Focal areas of aneurysmal and cystic changes [Figure 5] were seen which would provide an explanation for the clinically noted enlargement since hamartomas by definition would be expected to have a much lower rate of growth. The intrinsic slow-growing nature is also supported by the deficit of mitotic activity even in the highly cellular/spindled regions

of the lesion (less than 1/10 hpf). Areas with haphazard arrangement of nerve bundles within the collagenous stroma [Figure 6] were also noted. Immunohistochemical stains were positive for SMA [Figures 7(a) and 7(b)] in the spindle cells and negative for CK AE1/AE3, EMA, CD34, Stat6, ERG/FLI-1, Mucin 4, S-100, Sox-10, and desmin [Figure 8]; ruling out perineurioma, solitary fibrous tumor, a vascular neoplasm, Evans tumor, a benign peripheral nerve sheath tumor, or a myogenic neoplasm. The overall findings were suggestive of a hamartomatous lesion, most likely a nasal chondromesenchymal hamartoma. The absence of submucosal glandular proliferation, myxoid stroma, or mucinous metaplasia in the lining epithelium lowers the likelihood of other neoplastic hamartomatous lesions such as COREAH.

2. Discussion

Nasal chondromesenchymal hamartomas are most commonly seen in the nasal cavity of children less than 3 months old, with less common involvement of the paranasal sinuses [2]. As per one review [1], mean age for NCMH was 9.6 years. Review of the English PubMed literature reveals 43 cases [Table 1] of NCMH previously published, with our case being

FIGURE 1: Area of respiratory lining epithelium with squamous metaplastic change (arrowheads).

FIGURE 2: Focal areas of stroma displaying hyalinization (arrowheads).

FIGURE 3: Focal areas displaying calcification and ossification surrounded by variably spindled stroma.

FIGURE 4: Chondroid regions which support the hamartomatous nature of the lesion. Inset shows area at low-power with spatial relation of components, including surface ciliated epithelium (*) and bone (within dashed lines).

FIGURE 5: Variably dilated cystic regions within the lesion.

FIGURE 6: Disorganized bundles of nervous tissue interspersed within collagenous stroma.

FIGURE 7: Immunohistochemistry for smooth muscle actin (SMA) showing positive staining in the spindled lesional cells.

FIGURE 8: Lesional cells are negative for pan-cytokeratin (ae1/ae3), desmin, Sox-10, and S100. CD34 and ERG/FLI-1 highlight vascular endothelial cells.

TABLE 1: Brief review of cases of nasal chondromesenchymal hamartomas reported in the English literature. Some cases also reported in older review articles [1].

Age	Sex	Follow-up (Asymptomatic)	Site	Pertinent Information	Study	Year
5 days	M	2 years	Nasal cavity	-	[2] McDermott	1998
12 days	F	< 16 months	Nasal cavity	Intracranial extension	[2] McDermott	1998
14 days	M	-	Nasal cavity Ethmoid Sinus	Intracranial extension Residual tumor	[2] McDermott	1998
2 months	M	18 months	Nasal cavity	Intracranial extension	[2] McDermott	1998
3 months	F	2 years	Nasal cavity Ethmoid Sinus	Intracranial extension Residual tumor	[2] McDermott	1998
3 months	M	4 years	Nasal cavity	-	[2] McDermott	1998
7 years	<i>M</i>	-	<i>Nasal cavity Sphenoid sinus</i>	PPB, multiple recurrences	[2] McDermott	1998
4 months	M	13 years	Nasal cavity	Intracranial extension	[3] Kato	1999
0 days	M	5 years	Nasal cavity Sphenoid sinus Ethmoid sinus	Orbital compression	[4] Hsueh	2001
9 months	M	9 months	Nasal cavity	-	[4] Hsueh	2001
16 years	<i>M</i>	<i>8 months</i>	<i>Nasal cavity</i>	<i>3-month history</i>	[5] Alrawi M	2003
5 months	M	-	Nasal cavity	Orbital compression	[6] Kim B	2004
11 years	<i>M</i>	-	<i>Nasal cavity Ethmoid sinus</i>	<i>8-month history</i>	[7] Norman ES	2004
1 year	M	-	Nasal cavity	Orbital extension Residual tumor	[8] Shet T	2004
11 years	<i>M</i>	-	<i>Nasal cavity Ethmoid sinus</i>	-	[9] Ozolek JA	2005
17 years	<i>F</i>	-	<i>Nasal cavity</i>	-	[9] Ozolek JA	2005
25 years	<i>M</i>	-	<i>Nasal cavity Maxillary sinus</i>	Bilateral NCMH <i>Intracranial aneurysms</i>	[9] Ozolek JA	2005
69 years	<i>F</i>	-	<i>Nasal cavity Ethmoid sinus</i>	-	[9] Ozolek JA	2005
11 years	<i>M</i>	<i>2 months</i>	<i>Nasal cavity</i>	-	[10] Low SE	2006
15 years	<i>F</i>	<i>6 months</i>	<i>Nasal cavity</i>	Bilateral NCMH PPB	[11] Johnson C	2007
7 months	M	18 months	Nasal cavity	Orbital compression	[12] Silkiss RZ	2007
12 months	M	-	Nasal cavity	Orbital compression	[13] Finitis S	2009
19 months	M	10 months	Nasal cavity	Intracranial, orbital extension	[14] Kim JE	2009
2 cases previously reported, both with PPB, multiple recurrences					[15] Priest JR	2010
10 years	<i>F</i>	<i>21 months</i>	<i>Nasal cavity</i>	Bilateral NCMH PPB	[15] Priest JR	2010
11 years	<i>M</i>	<i>4 months</i>	<i>Nasal cavity</i>	PPB	[15] Priest JR	2010
11 years	<i>M</i>	-	-	PPB	[16] Behery RE	2012
8 years	<i>M</i>	<i>6 months</i>	<i>Sphenoid sinus Ethmoid sinus</i>	<i>4-month history</i>	[17] Uzomefuna	2012
14 years	<i>M</i>	<i>4 years</i>	<i>Nasal cavity Maxillary sinus</i>	-	[18] Cho YC	2013
23 years	<i>M</i>	<i>3 months</i>	<i>Nasal cavity Ethmoid Sinus</i>	<i>Orbital extension</i>	[19] Li GY	2013
40 years	<i>F</i>	-	<i>Nasal cavity Ethmoid Sinus</i> <i>Maxillary sinus</i>	Malignant transformation Recurrence	[20] Li Y	2013
9 months	F	-	Nasal cavity Maxillary sinus	Orbital compression	[21] Moon S	2014

TABLE 1: Continued.

Age	Sex	Follow-up (Asymptomatic)	Site	Pertinent Information	Study	Year
14 years	M	-	Nasal cavity	Bilateral NCMH PPB	[22] Obidan AA	2014
6 weeks	F	10 months	Nasal cavity	-	[23] Wang T	2014
5 years	M	3 years	Nasal cavity Ethmoid sinus	4-year history	[23] Wang T	2014
10 months	M	18 months	Nasal cavity	6-month history	[24] Lee CH	2015
49 years	M	2 years 4 years (phone)	Nasal cavity	5-year history	[1] Mason AK	2015
Systematic review					[1] Mason AK	2015
5 years	M	-	Nasal cavity	Previous rhabdomyosarcoma in remission	[25] Avci H	2016
13 years	F	12 months	Nasal cavity	6-month history	[26] Unal A	2016
3 years	M	3 years	Nasal cavity	-	[27] Nakaya M	2017

Index. Cases older than 1 year of age at presentation. **Bilateral/cases associated with pleuropulmonary blastoma.**

the oldest patient reported, and presenting with a tumor in an unusual location.

Our case would lend support to extending the age range for NCMH and considering it in the differential diagnosis of all sinonasal region tumors, irrespective of age, and location in the head and neck region. Despite primarily being a benign lesion, these tumors can present with areas of necrosis and local destruction, including bony invasion. The tumors can be aggressive appearing on imaging, extending into bony structures, including the cranium and/or the orbital cavity, which should not lead away from the diagnosis of this benign lesion. Detailed CT or preferably MRI prior to surgical excision should be performed.

NCMH has been associated with development of pleuropulmonary blastoma (PPB) during infancy. A recent [28] report highlighted the association of NCMH and PPB with DICER1 mutation and various associated entities such as lung cysts, cystic nephroma, renal sarcoma, Wilms tumor, thyroid hyperplasia, and CNS tumors. NCMH in isolation however is a benign lesion with follow-up in patients up to 16 years after excision, except for one reported case with malignant transformation in the literature [20]. Etiologically, it would make sense that cases in adults, such as ours, represent a tissue response to insult, such as chronic sinusitis rather than an inborn germline error (such as a DICER1 mutation).

Whether the presentation of a NCMH at a later age predisposes to malignant transformation due to the long-standing nature of the lesion is up for debate. It could represent a somatic DICER1 mutation rather than a germline mutation, causing the hamartoma to form later in age. Longer follow-up results from the adult cases and routine genetic testing in all NCMH will help provide an answer to these questions.

3. Conclusion

We report an unusual case of NCMH eroding the right maxillary sinus of a 70-year-old female. Although, NCMH is a rare entity with predilection for pediatric age groups, it is important to consider NCMH in the differential diagnosis of nasal/sinonasal masses in adult patients in order to avoid diagnostic errors.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

- [1] K. A. Mason, A. Navaratnam, E. Theodorakopoulou, and P. G. Chokkalingam, "Nasal Chondromesenchymal Hamartoma (NCMH): A systematic review of the literature with a new case report," *Journal of Otolaryngology - Head and Neck Surgery*, vol. 44, no. July, 2015.
- [2] M. B. McDermott, T. B. Ponder, and L. P. Dehner, "Nasal chondromesenchymal hamartoma: An upper respiratory tract analogue of the chest wall mesenchymal hamartoma," *The American Journal of Surgical Pathology*, vol. 22, no. 4, pp. 425–433, 1998.
- [3] K. Kato, R. Ijiri, Y. Tanaka, M. Hara, and K. Sekido, "Nasal chondromesenchymal hamartoma of infancy: The first Japanese case report," *Pathology International*, vol. 49, no. 8, pp. 731–736, 1999.
- [4] C. Hsueh, "Nasal chondromesenchymal hamartoma in children: report of 2 cases with review of the literature," *Archives of Pathology & Laboratory Medicine*, vol. 125, no. 3, p. 400, 2001.
- [5] M. Alrawi, M. McDermott, D. Orr, and J. Russell, "Nasal chondromesenchymal hamartoma presenting in an adolescent," *International Journal of Pediatric Otorhinolaryngology*, vol. 67, no. 6, pp. 669–672, 2003.

- [6] B. Kim, S.-H. Park, H. S. Min, J. S. Rhee, and K. C. Wang, "Nasal chondromesenchymal hamartoma of infancy clinically mimicking meningoencephalocele," *Pediatric Neurosurgery*, vol. 40, no. 3, pp. 136–140, 2004.
- [7] E. S. Norman, S. Bergman, and J. K. Trupiano, "Nasal chondromesenchymal hamartoma: Report of a case and review of the literature," *Pediatric and Developmental Pathology*, vol. 7, no. 5, pp. 517–520, 2004.
- [8] T. Shet, A. Borges, C. Nair, S. Desai, and R. Mistry, "Two unusual lesions in the nasal cavity of infants - A nasal chondromesenchymal hamartoma and an aneurysmal bone cyst like lesion - More closely related than we think?" *International Journal of Pediatric Otorhinolaryngology*, vol. 68, no. 3, pp. 359–364, 2004.
- [9] J. A. Ozolek, "Nasal chondromesenchymal hamartoma in older children and adults: series and immunohistochemical analysis," *Archives of Pathology & Laboratory Medicine*, vol. 129, no. 11, pp. 1444–50, 2005.
- [10] S. E. Low, R. K. Sethi, E. Davies, and J. S. Stafford, "Nasal chondromesenchymal hamartoma in an adolescent," *Histopathology*, vol. 49, no. 3, pp. 321–323, 2006.
- [11] C. Johnson, U. Nagaraj, J. Esguerra, D. Wasdahl, and D. Wurzbach, "Nasal chondromesenchymal hamartoma: Radiographic and histopathologic analysis of a rare pediatric tumor," *Pediatric Radiology*, vol. 37, no. 1, pp. 101–104, 2007.
- [12] R. Z. Silkiss, S. S. Mudvari, and D. Shetlar, "Ophthalmologic presentation of nasal chondromesenchymal hamartoma in an infant," *Ophthalmic Plastic & Reconstructive Surgery*, vol. 23, no. 3, pp. 243–244, 2007.
- [13] S. Finitis, C. Giavroglou, S. Potsi et al., "Nasal chondromesenchymal hamartoma in a child," *CardioVascular and Interventional Radiology*, vol. 32, no. 3, pp. 593–597, 2009.
- [14] J.-E. Kim, H.-J. Kim, H. K. Ji, Y.-H. Ko, and S.-K. Chung, "Nasal chondromesenchymal hamartoma: CT and MR imaging findings," *Korean Journal of Radiology*, vol. 10, no. 4, pp. 416–419, 2009.
- [15] J. R. Priest, G. M. Williams, W. A. Mize, L. P. Dehner, and M. B. McDermott, "Nasal chondromesenchymal hamartoma in children with pleuropulmonary blastoma-A report from the International Pleuropulmonary Blastoma Registry registry," *International Journal of Pediatric Otorhinolaryngology*, vol. 74, no. 11, pp. 1240–1244, 2010.
- [16] R. El Behery, J. Bedrnicek, A. Lazenby et al., "Translocation t(12;17)(q24.1;q21) as the sole anomaly in a nasal chondromesenchymal hamartoma arising in a patient with pleuropulmonary blastoma," *Pediatric and Developmental Pathology*, vol. 15, no. 3, pp. 249–253, 2012.
- [17] V. Uzomefunu, F. Glynn, J. Russell, and M. McDermott, "Nasal chondromesenchymal hamartoma with no nasal symptoms," *BMJ Case Reports*, 2012.
- [18] Y. C. Cho, I. Y. Sung, J. H. Son, and R. Ord, "Nasal chondromesenchymal hamartoma: Report of a case presenting with intraoral signs," *Journal of Oral and Maxillofacial Surgery*, vol. 71, no. 1, pp. 72–76, 2013.
- [19] G.-Y. Li, B. Fan, and Y.-Y. Jiao, "Endonasal endoscopy for removing nasal chondromesenchymal hamartoma extending from the lacrimal sac region," *Canadian Journal of Ophthalmology*, vol. 48, no. 2, pp. e22–e23, 2013.
- [20] M. Sharif and Abdul Jawad, "Interacting generalized dark energy and reconstruction of scalar field models," *Modern Physics Letters A*, vol. 28, no. 38, Article ID 1350180, 15 pages, 2013.
- [21] S. H. Moon and M. M. Kim, "Nasal chondromesenchymal hamartoma with concomitant esotropia in an infant: A case report," *Canadian Journal of Ophthalmology*, vol. 49, no. 1, pp. e30–e32, 2014.
- [22] A. A. Obidan and M. M. Ashoor, "Nasal chondromesenchymal hamartoma in an adolescent with pleuropulmonary blastoma," *Saudi Medical Journal*, vol. 35, no. 8, pp. 876–878, 2014.
- [23] T. Wang, W. Li, X. Wu et al., "Nasal chondromesenchymal hamartoma in young children: CT and MRI findings and review of the literature," *World Journal of Surgical Oncology*, vol. 12, no. 1, p. 257, 2014.
- [24] C. H. Lee, Y. H. Park, J. Y. Kim, and J. H. Bae, "Nasal chondromesenchymal hamartoma causing sleep-disordered breathing in an infant," *International Journal of Clinical and Experimental Pathology*, vol. 8, no. 8, pp. 9643–9646, 2015.
- [25] H. Avcı, Ş. Çomoğlu, E. Öztürk, B. Bilgiç, and Ö. E. Kiyak, "Nasal chondromesenchymal hamartoma: a rare nasal benign tumor," *Kulak burun bogaz ihtisas dergisi : KBB = Journal of ear, nose, and throat*, vol. 26, no. 5, pp. 300–303, 2016.
- [26] A. Ünal, R. O. Kum, Y. Avcı, and D. T. Ünal, "Nasal chondromesenchymal hamartoma, a rare pediatric tumor: Case report," *The Turkish Journal of Pediatrics*, vol. 58, no. 2, pp. 208–211, 2016.
- [27] M. Nakaya, S. Yoshihara, A. Yoshitomi, and S. Baba, "Endoscopic endonasal excision of nasal chondromesenchymal hamartoma with intracranial extension," *European Annals of Otorhinolaryngology, Head and Neck Diseases*, vol. 134, no. 6, pp. 423–425, 2017.
- [28] D. A. Hill, J. Ivanovich, J. R. Priest et al., "DICER1 mutations in familial pleuropulmonary blastoma," *Science*, vol. 325, no. 5943, p. 965, 2009.