

Case Report

Acute Cheilitis Associated with Oseltamivir after Influenzae A Infection

Koji Yokoyama  and Mitsukazu Mamada

Department of Pediatrics, Japanese Red Cross Wakayama Medical Center, Wakayama, Japan

Correspondence should be addressed to Koji Yokoyama; kojiy@kuhp.kyoto-u.ac.jp

Received 13 July 2023; Revised 1 November 2023; Accepted 2 December 2023; Published 6 December 2023

Academic Editor: Alexander K. C. Leung

Copyright © 2023 Koji Yokoyama and Mitsukazu Mamada. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Oseltamivir is a neuraminidase inhibitor used to treat acute influenza A or B in adult and pediatric patients. Adverse reactions are usually mild. Here, we report novel side effects associated with oseltamivir. The patient was an 11-year-old girl who developed lower lip cheilitis and stomatitis, after taking oseltamivir. Her symptoms and signs resolved within 36 h of oseltamivir discontinuation. She has clinically fully recovered and has remained well.

1. Case Presentation

An 11-year-old girl was admitted to our hospital with a 1 day history of lower lip cheilitis and stomatitis. No other part of her body or face had any swelling. She had no history of food, drug allergies, and inhalant abuse. It was the first time this had happened. She had no abdominal pain, no history of allergies or eczema, and no family history of angioedema. She had started taking oseltamivir capsules (75 mg twice daily) 2 days before the swelling began, having visited her primary care doctor with a bout of seasonal influenza with fever, cough, and arthralgia. These symptoms eased gradually while taking oseltamivir. She also took an expectorant (dextromethorphan) and an anti-inflammatory drug (tiaramide hydrochloride) for 2 days after visiting her doctor. She has taken an expectorant and an anti-inflammatory drug many times before without any problems. She was not taking any other prescription medications, including antiepileptic drugs or antibiotics. She was febrile (38.0°C), had a white swollen lip, an ulcer on the labia, and bilateral cervical lymphadenopathy (Figure 1). There were no rash or meningeal signs. She had no vulvar ulcers, erythema nodosum, folliculitis-like rash, or ophthalmological abnormalities. The remainder of the physical examination was unremarkable. Laboratory investigations revealed leukopenia (WBC, $2.03 \times 10^9/L$); neutrophil granulocytes, 64.6%; elevated C-

reactive protein (CRP, 1.48 mg/mL), and no vitamin deficiencies. No significant results were found in comprehensive respiratory virus testing or antibody titer studies. Oseltamivir was discontinued, and she was treated with an expectorant and anti-inflammatory drug. Her symptoms and signs resolved within 36 h of oseltamivir discontinuation. She has been followed up for over a year, no recurrence of symptoms has been observed. The possible reaction of oseltamivir was reported to Chugai Pharmaceutical Co., Ltd. The guardian of the patient must provide written consent for the data to be published.

2. Discussion

The course of this patient raises two important clinical points. First, patients taking oseltamivir can suffer from lower lip cheilitis and stomatitis. Second, these two conditions can occur simultaneously in pediatric patients. Oseltamivir is a neuraminidase inhibitor used to treat acute influenza A or B in adult and pediatric patients. Adverse reactions are usually mild. Oseltamivir-induced oral reactions have been reported, particularly tongue swelling, which has been observed and reported in postmarketing surveillance [1, 2]. Glossal swelling associated with oseltamivir is likely to be an idiosyncratic drug reaction, likely caused by an immune-mediated reaction to metabolites or



FIGURE 1: Image of her white swollen lip and an ulcer on the labia.

by direct cytotoxicity of the drug or its metabolites [1]. An IgE-mediated allergic reaction is also possible. However, the patient's IgE levels were normal, and she had no history of allergic disease. Drug-related cheilitis is one type of reversible cheilitis that can be caused by retinoids and antiviral drugs [3]. The prevalence of influenza and the dispensation of antifu agents are higher in children than in adults [4]. Idiosyncratic reactions can occur even in children (indeed, age may be a factor), and these reactions can be unpredictable and are not fully understood [5]. The pathogenesis of oseltamivir-induced cheilitis and stomatitis is not well understood, and future studies are needed to ascertain how oseltamivir causes oral reactions, whether predisposing factors exist and the actual incidence of this condition. More cases need to be recorded if we are to better understand the pathology of drug-associated oral reactions.

Physicians should pay attention to the various side effects of drugs.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

KY conceived and drafted the manuscript. MM participated in report's design and coordinated and helped draft the manuscript. All the authors have read and approved the final manuscript.

Acknowledgments

We thank bioedit (<https://www.bioedit.kr>) for editing a draft of this manuscript.

References

- [1] T. Kalsi, J. Stevenson, P. Wade, and M. Kinirons, "Tongue swelling in association with oseltamivir (Tamiflu)," *British Medical Journal Case Reports*, vol. 2011, no. 1, Article ID bcr0620114396, 2011.
- [2] A. Scott, R. More, and R. C. Freebairn, "Tongue swelling complicating management of a ventilated patient with acute respiratory distress syndrome secondary to novel influenza A (H1N1)," *Anaesthesia and Intensive Care*, vol. 38, no. 2, pp. 370–372, 2010.

- [3] L. Lugović-Mihić, K. Pilipovic, I. Crnaric, M. Situm, and T. Duvancic, "Differential diagnosis of cheilitis-how to classify cheilitis?" *Acta Clinica Croatica*, vol. 57, no. 2, pp. 342–351, 2018.
- [4] S. Inaida, S. Matsuno, and J. Okumura, "Longitudinal surveillance of influenza in Japan, 2006-2016," *Scientific Reports*, vol. 12, no. 1, Article ID 12026, 2022.
- [5] G. Zaccara, D. Franciotta, and E. Perucca, "Idiosyncratic adverse reactions to antiepileptic drugs," *Epilepsia*, vol. 48, no. 7, pp. 1223–1244, 2007.