Hindawi Publishing Corporation
Discrete Dynamics in Nature and Society
Volume 2009, Article ID 608976, 8 pages
doi:10.1155/2009/608976

Research Article

On the Recursive Sequence $x_n = A + x_{n-k}^p / x_{n-1}^r$

Fangkuan Sun, Xiaofan Yang, and Chunming Zhang

College of Computer Science, Chongqing University, Chongqing 400044, China

Correspondence should be addressed to Xiaofan Yang, xf_yang1964@yahoo.com

Received 15 December 2008; Accepted 7 May 2009

Recommended by Guang Zhang

This paper studies the dynamic behavior of the positive solutions to the difference equation $x_n = A + x_{n-k}^p / x_{n-1}^r$, $n = 1, 2, \ldots$, where A, p, and r are positive real numbers, and the initial conditions are arbitrary positive numbers. We establish some results regarding the stability and oscillation character of this equation for $p \in (0, 1)$.

1. Introduction

In recent years, there has been intense interest in the dynamic behavior of the positive solutions to a class of difference equations of the form

$$x_n = A + \frac{x_{n-k}^p}{x_{n-1}^p}, \quad n \in \mathbb{N}, \tag{1.1}$$

where A and p are positive real numbers. Now, let us make a brief review on the advances in this class of difference equations.

In 1999, Amleh et al. [1] studied the second-order rational difference equation

$$x_n = A + \frac{x_{n-2}}{x_{n-1}}, \quad n \in \mathbb{N}. \tag{1.2}$$

Later, Berenhaut and Stević [2], Stević [3], and El-Owaidy et al. [4] extended this work to the following more general second-order difference equation:

$$x_n = A + \frac{x_{n-2}^p}{x_{n-1}^p}, \quad n \in \mathbb{N}. \tag{1.3}$$

On the other hand, DeVault et al. [5] investigated the following higher-order version of (1.2):

$$x_n = A + \frac{x_{n-k}}{x_{n-1}}, \quad n \in \mathbb{N}. \tag{1.4}$$

By combining (1.3) and (1.4), Berenhaut and Stević [6] examined a larger class of difference equations, which are of the form

$$x_n = A + \frac{x_{n-k}^p}{x_{n-1}^p}, \quad n \in \mathbb{N}. \tag{1.5}$$

Very recently, Berenhaut et al. [7] studied the following generalization of (1.5):

$$x_n = A + \frac{x_{n-k}^p}{x_{n-m}^p}, \quad n \in \mathbb{N}. \tag{1.6}$$

For some related work, the interested reader is referred to [1, 3, 8–19].

Inspired by the previous work and by the work owing to Stević [15], this paper studies the behavior of the recursive equation

$$x_n = A + \frac{x_{n-k}^p}{x_{n-1}^r}, \quad n \in \mathbb{N}. \tag{1.7}$$

We establish some interesting results regarding the stability and oscillation character of this equation for $p \in (0, 1)$.

2. Stability Character

In this section we investigate the stability character of the positive solutions to (1.7).

A point $\overline{x} \in \mathbb{R}$ is an equilibrium point of (1.7) if and only if it is a root for the function

$$g(x) = x - x^{p-r} - A, \tag{2.1}$$

that is,

$$\overline{x} = \overline{x}^{p-r} + A. \tag{2.2}$$

Lemma 2.1. *Let* $0 < p < r + 1$, *then* (1.7) *has a unique equilibrium point* $\overline{x} > 1$.

Proof

Case 1. $p = r$. Then $\overline{x} = A + 1 > 1$.

Case 2. $r < p < r+1$. Then g defined by (2.1) is decreasing on $[0, (p-r)^{1/(r-p+1)}]$ and increasing on $[(p-r)^{1/(r-p+1)}, \infty)$. Since $g(1) = -A$ and $\lim_{x\to\infty} g(x) = \infty$, then g has a unique zero $\overline{x} > 1$.
Case 3. $0 < p < r$. Since g is increasing on $[0, \infty)$, $g(1) = -A$ and $\lim_{x\to\infty} g(x) = \infty$, then g has a unique zero $\overline{x} > 1$.

Lemma 2.2. *Let* $0 < p < r+1$. *Assume that* $\overline{x}$ *is the equilibrium point of* (1.7). *If* $(p+r)^{(p-r)/(r+1-p)}(p+r-1) < A$, *then* $\overline{x}$ *is locally asymptotically stable.*

Proof. By the Linearized Stability Theorem [11], $\overline{x}$ is locally asymptotically stable if and only if $\overline{x}^{r+1-p} > p+r$. A simple calculations shows that

$$g\left((p+r)^{1/(r+1-p)}\right) = (p+r)^{(p-r)/(r+1-p)}(p+r-1) - A < 0, \tag{2.3}$$

where g is defined by (2.1). Then since $\lim_{x\to\infty} g(x) = \infty$, we have $\overline{x} > (p+r)^{1/(r+1-p)}$ and $\overline{x}^{r+1-p} > p+r$. The proof is complete. □

Lemma 2.3. *If* $p \in (0,1)$, *then every positive solution to* (1.7) *is bounded.*

Proof. Note that each $n \in \mathbb{N}$ can be written in the form $lk+i$ for some $l \in \mathbb{N}_0$ and $i \in \{0, 1, \ldots, k-1\}$. From (1.7) and since $x_n > A$ for every $n \geq 0$, we have that

$$x_{lk+i} = A + \frac{x^p_{(l-1)k+i}}{x^r_{lk+i-1}} < A + \frac{x^p_{(l-1)k+i}}{A^r}, \tag{2.4}$$

for every $l \in \mathbb{N}_0$ and $i \in \{0, 1, \ldots, k-1\}$. Let $(u^{(i)}_l)_{l\in\mathbb{N}_0}$ be the solution to the difference equation

$$u^{(i)}_l = A + \frac{\left(u^{(i)}_{l-1}\right)^p}{A^r}, \qquad u^{(i)}_0 = x_{-k+i}. \tag{2.5}$$

From (2.4) and by induction we see that $x_{(l-1)k+i} \leq u^{(i)}_l$, $l \in \mathbb{N}_0$. Hence it is enough to prove that the sequences $(u^{(i)}_l)_{l\geq 0}$, $i \in \{0, 1, \ldots, k-1\}$ are bounded.

Since the function $f(x) = A + x^p/A^r$, $x \in (0, \infty)$ is increasing and concave for $p \in (0,1)$, it follows that there is a unique fixed point $\overline{x}$ of the equation $f(x) = x$ and that the function f satisfies

$$(f(x) - x)(x - \overline{x}) < 0, \quad x \in (0, \infty). \tag{2.6}$$

Using this fact it is easy to see that if $u^{(i)}_l \in (0, \overline{x}]$, the sequence is nondecreasing and bounded from above by $\overline{x}$, and if $u^{(i)}_l \geq \overline{x}$, it is nonincreasing and bounded from below by $\overline{x}$. Hence for every $u^{(i)}_0 \in (0, \infty)$, each of the sequences $u^{(i)}_l$, $i \in \{0, 1, \ldots, k-1\}$ is bounded. The claimed result follows. □

Lemma 2.4 (see [18]). *Let s, t be distinct nonnegative integers. Consider the difference equation*

$$\begin{gathered} x_n = f(x_{n-s}, x_{n-t}), \quad n = 1, 2, 3, \ldots, \\ x_{1-\max(s,t)}, x_{2-\max(s,t)}, \ldots, x_0 \in [a, b]. \end{gathered} \tag{2.7}$$

Suppose f satisfies the following conditions.

(H_1) *$f : [a, b]^2 \to [a, b]$ is a continuous function that is nondecreasing in the first argument and is nonincreasing in the second argument.*

(H_2) *The system*

$$\begin{gathered} x = f(x, y), \\ y = f(y, x) \end{gathered} \tag{2.8}$$

has a unique solution $(\overline{x}, \overline{x}) \in [a, b] \times [a, b]$.

Then $\overline{x}$ is the global attractor of all solutions to (2.7).

Theorem 2.5. *Let $p + r \leq 1$, then the unique equilibrium $\overline{x}$ to* (1.7) *is globally asymptotically stable.*

Proof. By Lemma 2.3, there must exist positive constants P and Q such that $P \leq x_n \leq Q$. Let $f(u, v) = A + u^p / v^r$, $u, v \in [P, Q]$, it is easy to verify that (H_1) holds. In addition, if

$$\begin{gathered} x = A + \frac{x^p}{y^r}, \\ y = A + \frac{y^p}{x^r}, \end{gathered} \tag{2.9}$$

then

$$\frac{x - A}{y - A} = \frac{x^{p+r}}{y^{p+r}}. \tag{2.10}$$

Assume that $x \neq y$, then $x > y$ or $x < y$.

In case $x > y$, we have $(x - A)/(y - A) > x/y \geq x^{p+r}/y^{p+r}$, which contradicts with (2.10).

In case $x < y$, we have $(x - A)/(y - A) < x/y \leq x^{p+r}/y^{p+r}$, again a contradiction.

Thus $x = y = \overline{x}$. By Lemma 2.4, the required result follows. □

Theorem 2.6. *Let $0 < p \leq r < 1$ and $A^{r-p+1} \geq p/r$. Then every positive solution to* (1.7) *converges to the unique equilibrium $\overline{x}$.*

Proof. By Lemma 2.3, every positive solution $\{x_n\}$ to (1.7) is bounded, which implies that there are finite $\liminf x_n = I$ and $\limsup x_n = S$. Assume that $I \neq S$ $(I < S)$. Taking the lim inf and lim sup in (1.7), it follows that

$$A + \frac{I^p}{S^r} \leq I < S \leq A + \frac{S^p}{I^r}. \tag{2.11}$$

From this and $r \in (0,1)$, it follows that

$$AS^r + I^p \leq IS^r < SI^r \leq AI^r + S^p, \tag{2.12}$$

yielding

$$AS^r - S^p < AI^r - I^p. \tag{2.13}$$

Define function $f(x) = Ax^r - x^p$, $x \in (A, \infty)$. Since

$$f'(x) = Arx^{r-1} - px^{p-1} = x^{p-1}\left(Arx^{r-p} - p\right) > x^{p-1}\left(rA^{r-p+1} - p\right) \geq 0, \tag{2.14}$$

we deduce that f is increasing, and thus (2.13) cannot hold. Therefore we have $I = S$, which implies the result. □

Theorem 2.7. *Let* $0 < p < 1$, $r \geq 1$, *and* $A^{r-p+1} \geq r + p - 1$. *Then every positive solution to* (1.7) *converges to the unique equilibrium* $\overline{x}$.

Proof. From (2.11) we have

$$AI^{r-1}S^r + I^{p+r-1} \leq I^r S^r \leq AI^r S^{r-1} + S^{p+r-1}. \tag{2.15}$$

Consequently, we obtain $(AI^{r-1}S^{r-1})(S - I) \leq (S^{r+p-1} - I^{r+p-1})$. Suppose that $I \neq S$, we get

$$AI^{r-1}S^{r-1} \leq \frac{S^{r+p-1} - I^{r+p-1}}{S - I} = (r + p - 1)\gamma^{p+r-2}, \tag{2.16}$$

where $\gamma \in (I, S)$, leading to

$$A^r S^{r-1} \leq AI^{r-1}S^{r-1} \leq (r + p - 1)\gamma^{p+r-2} < (r + p - 1)A^{p-1}S^{r-1}. \tag{2.17}$$

This implies that $A^{r-p+1} < r + p - 1$, which is a contradiction. Hence, $I = S = \overline{x}$. □

3. Oscillation Character

In this section we investigate the oscillation character of the positive solutions to (1.7).

Theorem 3.1. *Let $\{x_n\}_{n=-k}^{\infty}$ be a positive solution to* (1.7)*. Then either $\{x_n\}_{n=-k}^{\infty}$ consists of a single semicycle or $\{x_n\}_{n=-k}^{\infty}$ oscillates about the equilibrium $\overline{x}$ with semicycles having at most $k-1$ terms.*

Proof. Suppose that $\{x_n\}_{n=-k}^{\infty}$ has at least two semicycles. Then there exists $N \geq -k$ such that either $x_N < \overline{x} \leq x_{N+1}$ or $x_{N+1} < \overline{x} \leq x_N$. Assume that $x_N < \overline{x} \leq x_{N+1}$. (The argument for the case $x_{N+1} < \overline{x} \leq x_N$ is similar and is omitted). Now suppose that the positive semicycle beginning with the term x_{N+1} has $k-1$ terms. Then $x_N < \overline{x} \leq x_{N+k-1}$ and so

$$x_{N+k} = A + \frac{x_N^p}{x_{N+k-1}^r} < A + \frac{\overline{x}^p}{\overline{x}^r} = A + \overline{x}^{p-r} = \overline{x}. \tag{3.1}$$

This completes the proof. □

Theorem 3.2. *Suppose that k is even and let $\{x_n\}_{n=-k}^{\infty}$ be a solution to* (1.7)*, which has $k-1$ consecutive semicycles of length one, then every semicycle after this point is of length one.*

Proof. There exists $N \geq -k$ such that either

$$x_N, x_{N+2}, \ldots, x_{N+k-2} < \overline{x} \leq x_{N+1}, x_{N+3}, \ldots, x_{N+k-1} \tag{3.2}$$

or

$$x_{N+1}, x_{N+3}, \ldots, x_{N+k-1} < \overline{x} \leq x_N, x_{N+2}, \ldots, x_{N+k-2}. \tag{3.3}$$

We prove the former case. The proof for the latter is similar and is omitted. Now, we have

$$\begin{aligned} x_{N+k} &= A + \frac{x_N^p}{x_{N+k-1}^r} < A + \frac{\overline{x}^p}{\overline{x}^r} = A + \overline{x}^{p-r} = \overline{x}, \\ x_{N+k+1} &= A + \frac{x_{N+1}^p}{x_{N+k}^r} > A + \frac{\overline{x}^p}{\overline{x}^r} = A + \overline{x}^{p-r} = \overline{x}. \end{aligned} \tag{3.4}$$

The result then follows by induction. □

Lemma 3.3. *Let $0 < p < r+1$. Then* (1.7) *has no nontrivial periodic solutions of (not necessarily prime) period $k-1$.*

Proof. Suppose that $\{x_n\}_{n=-k}^{\infty}$ is a positive solution to (1.7) satisfying $x_{n-1} = x_{n-k}$ for all $n \geq 1$, then $x_n = A + x_{n-k}^p / x_{n-1}^r = A + x_{n-1}^{p-r}$ implies that $x_{n-1} = x_n = \overline{x}$ for all $n > -k$. The proof is complete. □

Theorem 3.4. *Assume that $p \leq r$. Let $\{x_n\}_{n=-k}^{\infty}$ be a positive solution to* (1.7)*, which consists of a single semicycle, then $\{x_n\}_{n=-k}^{\infty}$ converges to the equilibrium $\overline{x}$.*

Proof. Suppose $x_n \geq \overline{x}$ (the case for $x_n \leq \overline{x}$ is similar and is omitted) for all $n \geq -k$, then

$$x_{n+1} = A + \frac{x_{n-(k-1)}^p}{x_n^r} \geq \overline{x} = A + \overline{x}^{p-r}, \tag{3.5}$$

implying that

$$x_{n-(k-1)} \geq \overline{x}^{(p-r)/p} x_n^{r/p} \geq x_n^{(p-r)/p} x_n^{r/p} = x_n, \tag{3.6}$$

and so

$$x_{n-(k-1)} \geq x_n \geq \overline{x} \quad \text{for } n = 1, 2, \ldots. \tag{3.7}$$

From here it is clear that for $i = 0, \ldots, k-2$ there exists α_i such that

$$\lim_{n \to \infty} x_{n(k-1)+i} = \alpha_i. \tag{3.8}$$

But then $\alpha_0, \alpha_1, \ldots, \alpha_{k-2}$ is a periodic solution of (not necessarily prime) period $k-1$. By Lemma 3.3 the result holds. □

Acknowledgments

The author is grateful to the anonymous referees for their valuable comments and suggestions. This work was supported by the Natural Science Foundation of China (no. 10771227) and the Project for New Century Excellent Talents of Educational Ministry of China (no. NCET-05-0759).

References

[1] A. M. Amleh, E. A. Grove, G. Ladas, and D. A. Georgiou, "On the recursive sequence $x_{n+1} = \alpha + x_{n-1}/x_n$," *Journal of Mathematical Analysis and Applications*, vol. 233, no. 2, pp. 790–798, 1999.

[2] K. S. Berenhaut and S. Stević, "The behaviour of the positive solutions of the difference equation $x_n = A + (x_{n-2}/x_{n-1})^p$," *Journal of Difference Equations and Applications*, vol. 12, no. 9, pp. 909–918, 2006.

[3] S. Stević, "On the recursive sequence $x_{n+1} = \alpha + x_{n-1}^p/x_n^p$," *Journal of Applied Mathematics & Computing*, vol. 18, no. 1-2, pp. 229–234, 2005.

[4] H. M. El-Owaidy, A. M. Ahmed, and M. S. Mousa, "On asymptotic behaviour of the difference equation $x_{n+1} = \alpha + x_{n-1}^p/x_n^p$," *Journal of Applied Mathematics & Computing*, vol. 12, no. 1-2, pp. 31–37, 2003.

[5] R. DeVault, C. Kent, and W. Kosmala, "On the recursive sequence $x_{n+1} = p + x_{n-k}/x_n$," *Journal of Difference Equations and Applications*, vol. 9, no. 8, pp. 721–730, 2003.

[6] K. S. Berenhaut and S. Stević, "A note on positive non-oscillatory solutions of the difference equation $x_{n+1} = \alpha + x_{n-k}^p/x_n^p$," *Journal of Difference Equations and Applications*, vol. 12, no. 5, pp. 495–499, 2006.

[7] K. S. Berenhaut, J. D. Foley, and S. Stević, "The global attractivity of the rational difference equation $y_n = A + (y_{n-m}/y_{n-k})^p$," *Proceedings of the American Mathematical Society*, vol. 136, no. 1, pp. 103–110, 2008.

[8] R. M. Abu-Saris and R. DeVault, "Global stability of $y_{n+1} = A + y_n/y_{n-k}$," *Applied Mathematics Letters*, vol. 16, no. 2, pp. 173–178, 2003.

[9] R. DeVault, G. Ladas, and S. W. Schultz, "On the recursive sequence $x_{n+1} = A/x_n + 1/x_{n-1}$," *Proceedings of the American Mathematical Society*, vol. 126, no. 11, pp. 3257–3261, 1998.

[10] R. DeVault, W. Kosmala, G. Ladas, and S. W. Schultz, "Global behavior of $y_{n+1} = (p + y_{n-k})/(qy_n + y_{n-k})$," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, no. 7, pp. 4743–4751, 2001.

[11] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, vol. 256 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993.

[12] M. Saleh and M. Aloqeili, "On the rational difference equation $y_{n+1} = A + y_n/y_{n-k}$," *Applied Mathematics and Computation*, vol. 177, no. 1, pp. 189–193, 2006.

[13] S. Stević, "On monotone solutions of some classes of difference equations," *Discrete Dynamics in Nature and Society*, vol. 2006, Article ID 53890, 9 pages, 2006.

[14] S. Stević, "On the recursive sequence $x_{n+1} = A + x_n^p/x_{n-1}^p$," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 34517, 9 pages, 2007.

[15] S. Stević, "On the recursive sequence $x_{n+1} = A + x_n^p/x_{n-1}^r$," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 40963, 9 pages, 2007.

[16] F. Sun, "On the asymptotic behavior of a difference equation with maximum," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 243291, 6 pages, 2008.

[17] T. Sun and H. Xi, "Global behavior of the nonlinear difference equation $x_{n+1} = f(x_{n-s}, x_{n-t})$," *Journal of Mathematical Analysis and Applications*, vol. 311, no. 2, pp. 760–765, 2005.

[18] X. Yang, Y. Yang, and J. Luo, "On the difference equation $x_n = (p + x_{n-s})/(qx_{n-t} + x_{n-s})$," *Applied Mathematics and Computation*, vol. 189, no. 1, pp. 918–926, 2007.

[19] Y. Yang and X. Yang, "On the difference equation $x_n = (px_{n-s} + x_{n-t})/(qx_{n-s} + x_{n-t})$," *Applied Mathematics and Computation*, vol. 203, no. 2, pp. 903–907, 2008.