

Research Article

Optimal Parametric Iteration Method for Solving Multispecies Lotka-Volterra Equations

Vasile Marinca^{1,2} and Nicolae Herișanu^{1,2}

¹ Faculty of Mechanical Engineering, Polytechnic University of Timișoara, Mihai Viteazu Avenue 1, 300222 Timișoara, Romania

² Center for Advanced and Fundamental Technical Research, Romanian Academy, Timișoara Branch, Mihai Viteazu Avenue 24, 300222 Timișoara, Romania

Correspondence should be addressed to Nicolae Herișanu, herisanu@mec.upt.ro

Received 15 August 2011; Accepted 19 November 2011

Academic Editor: Zengji Du

Copyright © 2012 V. Marinca and N. Herișanu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We apply an analytical method called the Optimal Parametric Iteration Method (OPIM) to multispecies Lotka-Volterra equations. By using initial values, accurate explicit analytic solutions have been derived. The method does not depend upon small parameters and provides us with a convenient way to optimally control the convergence of the approximate solutions. An excellent agreement has been demonstrated between the obtained solutions and the numerical ones. This new approach, which can be easily applied to other strongly nonlinear problems, is very effective and yields very accurate results.

1. Introduction

A substantial amount of research work has been invested in the study of nonlinear systems of differential equations. Systems of nonlinear differential equations arise in many scientific models such as biological systems and are used in various fields as engineering, chemistry, and ecology. In 1925, Lotka [1] developed the motion of an evolutionary system based on two fundamental changes, those involving matter between components of a system and those involving exchanges of energy [2]. Unlike being grounded in chemistry, Lotka believed that these ideas could be applied to any biological system. In 1926, Volterra [3] developed the well-known mathematical models of multispecies interaction. These models, the predator, prey and competition models are known today as Lotka-Volterra models. The environment of the species can be influenced by the effect of food availability, weather conditions, temperature,

mating habits, contact with predators, and other resource or physical environmental quantities. Lotka-Volterra equations describe variations of population densities of few species that compete for the same resources. However, the ecological system is often affected by environmental changes and other human activities. In many practical situations, it is often the case that one of the species maybe suffers a significant loss or increase in density for some reason at some transitory time slots. These models, for instance, they can describe the competing fish species which are exploited by human activities, can also describe the dynamics of normal and tumour cells in a changing environment under the effects of the chemotherapy. These models are also applicable in case we are interested in the existence and stability of tumor-free solution and how treatment affects the interaction of tumour and normal cells. Although simplistic, these few models are still used as the foundation for mathematical models in biology [4]. These models can also describe the time history of a biological system and are used in various fields as engineering, chemistry, biology, or mathematics [5]. In fact the Lotka-Volterra model is one of the most popular ones to demonstrate a simple nonlinear control system. The accurate solutions of the Lotka-Volterra equations may become a difficult task either if the equations are stiff or when the number of species is large.

Nonlinear analytical techniques for solving nonlinear problems have been dominated by the perturbation methods, which have found wide applications in engineering [6]. But perturbation methods have their limitations: perturbation techniques are based on the existence of a small parameter. A majority of nonlinear problems have no small parameters at all. So, it is necessary to develop a kind of new nonlinear analytical methods which does not require small parameters at all.

There exist same alternative methods, such as the variational iteration method [7], the Adomian decomposition method [8], a modified Lindstedt-Poincare method [9], the homotopy analysis method [10], the homotopy perturbation method [11], and the optimal homotopy asymptotic method [12].

In recent years, a growing interest towards the application of iterative techniques in nonlinear problems has appeared in science and engineering. In 1987 Mickens [13] proposed an iterative scheme for nonlinear problems. In 2002 Lim and Wu [14], in 2006 Hu [15], and in 2008 Chen and Liu [16] proposed modified iteration procedures. Also He [17] proposed some iterative and asymptotic methods for nonlinear problems and later, Marinca and Herişanu [18] proposed in 2006 a new iteration method by combining Mickens' and He's iteration methods. Ramos [19] investigated other iterative techniques for nonlinear differential equations. These methods are valid not only for a small parameter, but also for very large parameters and have been used to solve nonlinear both conservative and nonconservative oscillators.

In this paper we propose a new approach to find analytic approximate solution of multispecies Lotka-Volterra models using a new iterative procedure, namely, the Optimal Parametric Iteration Method (OPIM). The efficiency of the proposed procedure is proved since an accurate solution is explicitly analytically achieved in an iterative manner after only one iteration. This new approach involves the presence of a finite number of initially unknown parameters, which are optimally determined, providing a rigorous way to control the convergence of the solutions [20]. Different methodologies have been applied to study Lotka-Volterra models [21–25]. Using this new approach, the obtained approximate solutions rapidly converge to the exact solution. Some alternative ways for mathematical biology problems are presented in [26–28].

2. Formulation and Solution Approach

In order to introduce the OPIM framework for the most general form of systems of nonlinear ODEs, we consider the following system:

$$\begin{aligned}
 \dot{x}_1 &= f_1(x_1, x_2, x_3, \dots, x_s) \\
 \dot{x}_2 &= f_2(x_1, x_2, x_3, \dots, x_s) \\
 &\vdots \\
 \dot{x}_s &= f_s(x_1, x_2, x_3, \dots, x_s) \\
 x_1(0) &= a_1, \quad x_2(0) = a_2, \dots, x_s(0) = a_s,
 \end{aligned} \tag{2.1}$$

where dot denotes derivative with respect to time t , f_i , ($i = 1, 2, \dots, s$) are nonlinear functions.

In the frame of OPIM we define the following iteration formulas:

$$\begin{aligned}
 \dot{x}_{1,n+1} &= f_1(x_{1,n}, x_{2,n}, \dots, x_{s,n}) + m_1(t, C_i) f_{1,x_1}(x_{1,n}, x_{2,n}, x_{s,n}) + n_1(t, C_i) f_{1,x_2}(x_{1,n}, x_{2,n}, \dots, x_{s,n}) \\
 &\quad + \dots + p_1(t, C_i) f_{1,x_s}(x_{1,n}, x_{2,n}, \dots, x_{s,n}) \\
 \dot{x}_{2,n+1} &= f_2(x_{1,n}, x_{2,n}, \dots, x_{s,n}) + m_2(t, C_j) f_{2,x_1}(x_{1,n}, x_{2,n}, x_{s,n}) + n_2(t, C_j) f_{2,x_2}(x_{1,n}, x_{2,n}, \dots, x_{s,n}) \\
 &\quad + \dots + p_2(t, C_j) f_{2,x_s}(x_{1,n}, x_{2,n}, \dots, x_{s,n}) \\
 &\vdots \\
 \dot{x}_{s,n+1} &= f_s(x_{1,n}, x_{2,n}, \dots, x_{s,n}) + m_s(t, C_k) f_{s,x_1}(x_{1,n}, x_{2,n}, x_{s,n}) + n_s(t, C_i) f_{s,x_2}(x_{1,n}, x_{2,n}, \dots, x_{s,n}) \\
 &\quad + \dots + p_s(t, C_k) f_{s,x_s}(x_{1,n}, x_{2,n}, \dots, x_{s,n}),
 \end{aligned} \tag{2.2}$$

where $f_{1,x_i} = \partial f_1 / \partial x_i$, $n = 0, 1, 2, \dots$ and C_i, C_j, \dots, C_k are unknown constants at this moment.

A special case of this general system is the Lotka-Volterra model with three species, which is described by the following system of nonlinear differential equation [29, 30]:

$$\begin{aligned}
 \dot{x} &= x(1 - x - \alpha y - \beta z), \\
 \dot{y} &= y(1 - \beta x - y - \alpha z), \\
 \dot{z} &= z(1 - \alpha x - \beta y - z),
 \end{aligned} \tag{2.3}$$

where α and β are known parameters not necessary small. Initial conditions are

$$x(0) = a, \quad y(0) = b, \quad z(0) = c. \tag{2.4}$$

In this special case, it is clear that $s = 3$ and

$$\begin{aligned}
 x_1 &= x, \quad x_2 = y, \quad x_3 = z, \quad x_4 = x_5 = \dots = 0 \\
 f_1 &= x(1 - x - \alpha y - \beta z), \quad f_2 = y(1 - \beta x - y - \alpha z), \\
 f_3 &= z(1 - \alpha x - \beta y - z), \quad f_4 = f_5 = \dots = 0.
 \end{aligned} \tag{2.5}$$

With the notations (2.5), the iteration formulas (2.2) can be written in the form

$$\begin{aligned}\dot{x}_{n+1} &= f(x_n, y_n, z_n) + m_1(t, C_i) f_x(x_n, y_n, z_n) + n_1(t, C_i) f_y(x_n, y_n, z_n) + p_1(t, C_i) f_z(x_n, y_n, z_n) \\ \dot{y}_{n+1} &= g(x_n, y_n, z_n) + m_2(t, C_j) g_x(x_n, y_n, z_n) + n_2(t, C_j) g_y(x_n, y_n, z_n) + p_2(t, C_j) g_z(x_n, y_n, z_n) \\ \dot{z}_{n+1} &= h(x_n, y_n, z_n) + m_3(t, C_k) h_x(x_n, y_n, z_n) + n_3(t, C_k) h_y(x_n, y_n, z_n) + p_3(t, C_k) h_z(x_n, y_n, z_n),\end{aligned}\tag{2.6}$$

where

$$\begin{aligned}f(x, y, z) &= x(1 - x - \alpha y - \beta z), \\ g(x, y, z) &= y(1 - \beta x - y - \alpha z), \\ h(x, y, z) &= z(1 - \alpha x - \beta y - z),\end{aligned}\tag{2.7}$$

and $f_x = \partial f / \partial x$, $n = 0, 1, 2, \dots$, and $C_i, C_j, C_k, i, j, k = 1, 2, \dots$ are unknown constants at this moment. There are many possibilities to choose the auxiliary functions $m_e(t, C_i)$, $n_e(t, C_j)$ and $p_e(t, C_k)$, $e = 1, 2, 3$. Basically, the shape of m_e , n_e , and p_e must follow the terms appearing in (2.6). Therefore, we try to choose the auxiliary functions so that in (2.6) the products $m_e F_x$ are of the same shape with the function F_x , where $F \in \{f, g, h\}$. The constants C_i, C_j, C_k which appear in the auxiliary functions m_e , n_e , and p_e can be optimally determined via various methods, for example, putting the conditions that the residual functionals J_1, J_2 , and J_3 given, respectively, by

$$\begin{aligned}J_1 &= \int_a^b [\dot{x} - f(x, y, z)]^2 dt, \\ J_2 &= \int_a^b [\dot{y} - g(x, y, z)]^2 dt, \\ J_3 &= \int_a^b [\dot{z} - h(x, y, z)]^2 dt\end{aligned}\tag{2.8}$$

be minimum, that is:

$$\frac{\partial J_i}{\partial C_j} = 0, \quad i = 1, 2, 3, \quad j = 1, 2, \dots, s\tag{2.9}$$

taking into account also the initial conditions. Other methods suitable to determine the constants C_j are the Galerkin method, the collocation method, and so on.

On the other hand, the initial approximations x_0, y_0 , and z_0 and also the auxiliary functions m_e, n_e , and p_e are not unique.

In short, the basic idea of the proposed procedure (OPIM) are the construction a new iteration scheme (2.6), the involvement of the auxiliary functions, and the convergence-control constants C_1, C_2, \dots which lead to an excellent agreement of the approximate solutions with the exact ones.

3. Results and Discussions

In this section we present an example to show the efficiency of the method described in the previous section for solving (2.3).

Taking into account the initial conditions, we choose the initial approximations as functions of the form

$$x_0 = ae^{-t}, \quad y_0 = be^{-t}, \quad z_0 = ce^{-t}. \quad (3.1)$$

For $n = 0$ into (2.6) we can construct the following iteration formulas:

$$\begin{aligned} \dot{x}_1 &= x_0 - x_0^2 - \alpha x_0 y_0 - \beta x_0 z_0 + m_1(t, C_i) [1 - (2x_0 + \alpha y_0 + \beta z_0)] + n_1(t, C_i) (-\alpha x_0) \\ &\quad + p_1(t, C_i) (-\beta x_0), \\ \dot{y}_1 &= y_0 - \beta x_0 y_0 - y_0^2 - \alpha y_0 z_0 + m_2(t, C_j) (-\beta y_0) + n_2(t, C_j) [1 - (\beta x_0 + 2y_0 + \alpha z_0)] \\ &\quad + p_2(t, C_i) (-\alpha y_0), \\ \dot{z}_1 &= z_0 - \alpha x_0 z_0 - \beta y_0 z_0 - z_0^2 + m_3(t, C_k) (-\alpha z_0) + n_3(t, C_k) (-\beta z_0) + p_3(t, C_k) \\ &\quad \times [1 - (\alpha x_0 + \beta y_0 + 2z_0)]. \end{aligned} \quad (3.2)$$

For the auxiliary functions $m_i, n_i, p_i, i = 1, 2, 3$ we choose the expressions

$$\begin{aligned} m_1(t, C_i) &= (C_1 t + C_2) e^{-t}, & n_1(t, C_i) &= C_3 t + C_4, & p_1(t, C_i) &= (C_5 t + C_6) e^{-t} \\ m_2(t, C_j) &= (C_7 t + C_8), & n_2(t, C_j) &= (C_9 t + C_{10}) e^{-t}, & p_2(t, C_j) &= (C_{11} t + C_{12}) e^{-2t} \\ m_3(t, C_k) &= (C_{13} t + C_{14}) e^{-t}, & n_3(t, C_k) &= (C_{15} t + C_{16}) e^{-2t}, & p_3(t, C_k) &= (C_{17} t + C_{18}) e^{-t}. \end{aligned} \quad (3.3)$$

Alternatively, we can consider either other expressions for the initial approximations and the auxiliary function, follows as:

$$\begin{aligned} x_0 &= ae^{-2t}, & y_0 &= b, & z_0 &= ce^{-t}, \\ m_1(t, C'_i) &= C'_1 e^{-2t}, & n_1(t, C'_i) &= (C'_2 t + C'_3) e^{-t}, & p_1(t, C'_i) &= (C'_4 t + C_5) e^{-2t} \\ m_2(t, C'_j) &= (C'_6 t + C'_7) e^{-t}, & n_2(t, C'_j) &= C'_8 e^{-2t}, & p_2(t, C'_j) &= (C'_9 t + C'_{10}) e^{-t} \\ m_3(t, C'_k) &= C'_{11} e^{-t}, & n_3(t, C'_k) &= C'_{12} t + C'_{13}, & p_3(t, C'_k) &= (C'_{14} t + C'_{15}) e^{-2t}, \end{aligned} \quad (3.4)$$

and so on.

Now, substituting (3.1) and (3.3) into (3.2), we obtain the equations in \dot{x}_1 , \dot{y}_1 , and \dot{z}_1 , respectively,

$$\begin{aligned}
\dot{x}_1 &= (-\alpha a C_3 t + a - \alpha a C_4) e^{-t} + \left[(C_1 - \beta a C_5) t + C_2 - \beta a C_6 - (a^2 + \alpha a b + \beta a c) \right] e^{-2t} \\
&\quad - (2a + \alpha b + \beta c) (C_1 t + C_2) e^{-3t} \\
\dot{y}_1 &= [(C_9 t - \beta b C_7) t + b - \beta b C_8 + C_{10}] e^{-t} \\
&\quad - \left[(\beta a + 2b + \alpha c) C_9 t + \beta a b + b^2 + a b c + (\beta a + 2b + \alpha c) C_{10} \right] e^{-2t} - (a b C_{11} t + a b C_{12}) e^{-3t} \\
\dot{z}_1 &= (C_{17} t + C_{18} + c) e^{-t} \\
&\quad - \left\{ [(\alpha a + \beta c + 2c) C_{17} + \alpha c C_{13}] t + \alpha a c + \beta b c + c^2 + \alpha c C_{14} + (\alpha a + \beta c + 2c) C_{18} \right\} e^{-2t} \\
&\quad - (\beta c C_{14} t + \beta c C_{16}) e^{-3t}.
\end{aligned} \tag{3.5}$$

The solutions of (3.5) with the initial conditions

$$x_1(0) = a, \quad y_1(0) = b, \quad z_1(0) = c \tag{3.6}$$

can be written in the form

$$\begin{aligned}
x_1(t) &= a + \alpha a C_3 t e^{-t} + (a - \alpha a C_3 - \alpha a C_4) (1 - e^{-t}) - \frac{1}{2} (C_1 - \beta a C_5) t e^{-2t} \\
&\quad + \frac{1}{4} \left[C_1 - \beta a C_5 + 2(C_2 - \beta a C_6) - 2(a^2 + \alpha a b + \beta a c) \right] (1 - e^{-2t}) + \frac{1}{3} (2a + \alpha b + \beta c) \\
&\quad \times C_1 t e^{-3t} - \frac{1}{9} (2a + \alpha b + \beta c) (C_1 + 3C_2) (1 - e^{-3t}) \\
y_1(t) &= b + (b - \beta b C_8 + C_{10}) (1 - e^{-t}) + (\beta b C_7 - C_9) t e^{-t} \\
&\quad - \frac{1}{4} \left[(\beta a + 2b + \alpha c) C_9 + 2(\beta a b + b^2 + a b c) + 2(\beta a + 2b + \alpha c) C_{10} \right] (1 - e^{-2t}) \\
&\quad + \frac{1}{2} (\beta a + 2b + \alpha c) C_9 t e^{-2t} - \frac{1}{9} a b (C_{11} + 3C_{12}) (1 - e^{-3t}) + \frac{1}{3} a b C_{11} t e^{-3t} \\
z_1(t) &= c + (C_{17} + C_{18} + c) (1 - e^{-t}) - C_{17} t e^{-t} \\
&\quad - \frac{1}{4} \left[2(\alpha a c + \beta b c + c^2) + 2\alpha c C_{14} + 2(\alpha a + \beta c + 2c) C_{18} + \alpha c C_1 + (\alpha a + \beta c + 2c) C_{17} \right] \\
&\quad \times (1 - e^{-2t}) + \frac{1}{2} [\alpha c C_{13} + (\alpha a + \beta c + 2c) C_{17}] t e^{-2t} + \frac{1}{3} \beta c C_{15} t e^{-3t} - \frac{1}{4} \beta c (C_{15} + 3C_{16}) \\
&\quad \times (1 - e^{-3t}).
\end{aligned} \tag{3.7}$$

By substituting (3.7) into (2.3) it results in the residuals:

$$\begin{aligned}
R_1(t, C_i) &= \dot{x}_1 - x_1(1 - \alpha x_1 - \beta y_1 - \beta z_1), \\
R_2(t, C_i) &= \dot{y}_1 - y_1(1 - \beta x_1 - \beta y_1 - \alpha z_1), \\
R_3(t, C_i) &= \dot{z}_1 - z_1(1 - \alpha x_1 - \beta y_1 - \beta z_1).
\end{aligned} \tag{3.8}$$

Making collocation in the arbitrary points $t_i, i = 1, 2, \dots, 6$

$$\begin{aligned} R_1(t_1, C_i) &= R_1(t_2, C_i) = \dots = R_1(t_6, C_i) = 0 \\ R_2(t_1, C_i) &= R_2(t_2, C_i) = \dots = R_2(t_6, C_i) = 0 \\ R_3(t_1, C_i) &= R_3(t_2, C_i) = \dots = R_3(t_6, C_i) = 0, \end{aligned} \quad (3.9)$$

we obtain the optimal values of the constants C_1, C_2, \dots, C_{18} and therefore the solution (3.7) in the first approximation is well determined.

In the case when $a = 0.2, b = 0.3, c = 0.5, \alpha = 0.1, \beta = 0.1, t_1 = 0.3, t_2 = 0.7, t_3 = 1, t_4 = 2, t_5 = 3, t_6 = 5$, from (3.9) we obtain

$$\begin{aligned} C_1 &= 1.669841466, & C_2 &= -0.452081708, & C_3 &= 0.899181851, & C_4 &= 2.361528973, \\ C_5 &= -1.940901217, & C_6 &= 0.582517076, & C_7 &= 1033.008542, & C_8 &= -540.7564632, \\ C_9 &= 30.90993616, & C_{10} &= -14.59841284, & C_{11} &= 29.8078082, & C_{12} &= 94.01456893, \\ C_{14} &= 53.21346396, & C_{15} &= -22.21139178, & C_{16} &= -35.8538265, & C_{17} &= -0.345181562, \\ & & C_{18} &= 0.993490199. \end{aligned} \quad (3.10)$$

By substituting (3.10) into (3.7) we obtain after only one iteration, the following expression for the first-order approximate solution of Lotka-Volterra equations in three species (2.3):

$$\begin{aligned} x_1(t) &= 0.8281811 - (0.452081708t + 1.669841466)e^{-t} + (2.361528973t - 0.899181851)e^{-2t} \\ &\quad + (0.582517076t + 1.940902217)e^{-3t}, \end{aligned} \quad (3.11)$$

$$\begin{aligned} y_1(t) &= 0.83333001 + (0.08032009t - 1.924281055)e^{-t} \\ &\quad + (1.715501457t + 0.342446025)e^{-2t} + (0.298078082t + 1.04850505)e^{-3t}, \end{aligned} \quad (3.12)$$

$$\begin{aligned} z_1(t) &= 0.836102311 + (0.345181562t - 1.148308637)e^{-t} + (0.001606241t + 1.532769033)e^{-2t} \\ &\quad - (0.370189863t + 0.720960396)e^{-3t}. \end{aligned} \quad (3.13)$$

It is easy to verify the accuracy of the obtained solution if we graphically compare the analytical solution with the numerical one. Figures 1, 2, and 3 show the comparison between the present solutions and the numerical integration results obtained by a fourth-order Runge-Kutta method.

It can be seen from Figures 1–3 that the solution obtained by OPIM is in very good agreement with numerical integration results.

4. Conclusions

In this work the Optimal Parametric Iteration Method (OPIM) is employed to propose new analytic solutions for some nonlinear differential equations. The validity of the method is




Figure 1: Comparison between the approximate solution (3.11) and numerical results obtained for (2.3) in case $a = 0.2$, $b = 0.3$, $c = 0.5$, $\alpha = 0.1$, $\beta = 0.1$: — numerical solution; --- approximate solution.




Figure 2: Comparison between the approximate solution (3.12) and numerical results obtained for (2.3) in case $a = 0.2$, $b = 0.3$, $c = 0.5$, $\alpha = 0.1$, $\beta = 0.1$: — numerical solution; --- approximate solution.




Figure 3: Comparison between the approximate solution (3.13) and numerical results obtained for (2.3) in case $a = 0.2$, $b = 0.3$, $c = 0.5$, $\alpha = 0.1$, $\beta = 0.1$: — numerical solution; --- approximate solution.

illustrated on the Lotka-Volterra model with three species. Our procedure is independent of the presence of small parameters. Our construction of iterations is different from other known iteration techniques. In short, the basic new ideas of our method are related to the construction of a new iteration scheme which involves some auxiliary functions m_e , n_e , and p_e , $e = 1, 2, 3$ whose parameters C_1, C_2, \dots (viz. the convergence-control constants) lead to an excellent agreement of the approximate solutions with the exact ones. Our procedure is very effective, explicit, and accurate for nonlinear approximations, rapidly converging to the exact solution after only one iteration. This procedure provides a simple but rigorous way to control and adjust the convergence of the solutions by optimally determining the parameters C_1, C_2, \dots . Our method gives analytic solutions valid globally in time unlike other known methods, for instance. Adomian decomposition method, which unfortunately does not guarantee analytic solutions valid globally in time as proved by Rëpaci [31].

References

- [1] A. J. Lotka, *Elements of Physical Biology*, William and Wilkins, Baltimore, Md, USA, 1925.
- [2] S. E. Kingsland, *Modeling Nature: Episodes in the History of Population Ecology*, University of Chicago Press, Chicago, Ill, USA, 1995.
- [3] V. Volterra, "Variazioni e fluttuazioni del numero d'individui in specie animali conniventi," *Memorie della Reale Accademia Nazionale delle Scienze*, vol. 2-3, pp. 30–111, 1926.
- [4] A. W. Leung, *Systems of Nonlinear Partial Differential Equations, Applications to Biology and Engineering*, Kluwer Academic, Dordrecht, The Netherlands, 1989.
- [5] J. Hofbauer and K. Sigmund, *The Theory of Evolution and Dynamical Systems*, vol. 7, Cambridge University Press, London, UK, 1988.
- [6] A. H. Nayfeh and D. T. Mook, *Nonlinear Oscillations*, John Wiley and Sons, New York, NY, USA, 1979.
- [7] N. Herişanu V. Marinca, "A modified variational iteration method for strongly nonlinear problems," *Nonlinear Science Letters A*, vol. 1, pp. 183–192, 2010.
- [8] G. Adomian, "A review of the decomposition method and some recent results for nonlinear equations," *Computers and Mathematics with Applications*, vol. 21, no. 5, pp. 101–127, 1991.
- [9] C. W. Lim, S. K. Lai, B. S. Wu, W. P. Sun, Y. Yang, and C. Wang, "Application of a modified lindstedt-poincaré method in coupled TDOF systems with quadratic nonlinearity and a constant external excitation," *Archive of Applied Mechanics*, vol. 79, no. 5, pp. 411–431, 2009.
- [10] S. Liao, *Beyond Perturbation: Introduction to the Homotopy Analysis Method*, vol. 2, Chapman & Hall/CRC, 2003.
- [11] J.-H. He, "A coupling method of a homotopy technique and a perturbation technique for non-linear problems," *International Journal of Non-Linear Mechanics*, vol. 35, no. 1, pp. 37–43, 2000.
- [12] V. Marinca and N. Herişanu, "Determination of periodic solutions for the motion of a particle on a rotating parabola by means of the optimal homotopy asymptotic method," *Journal of Sound and Vibration*, vol. 329, no. 9, pp. 1450–1459, 2010.
- [13] R. E. Mickens, "Iteration procedure for determining approximate solutions to nonlinear oscillator equations," *Journal of Sound and Vibration*, vol. 116, no. 1, pp. 185–187, 1987.
- [14] C. W. Lim and B. S. Wu, "A modified Mickens procedure for certain non-linear oscillators," *Journal of Sound and Vibration*, vol. 257, no. 1, pp. 202–206, 2002.
- [15] H. Hu, "Solutions of nonlinear oscillators with fractional powers by an iteration procedure," *Journal of Sound and Vibration*, vol. 294, no. 3, pp. 608–614, 2006.
- [16] Y. M. Chen and J. K. Liu, "A modified Mickens iteration procedure for nonlinear oscillators," *Journal of Sound and Vibration*, vol. 314, no. 3–5, pp. 465–473, 2008.
- [17] J.-H. He, "Iteration perturbation method for strongly nonlinear oscillations," *Journal of Vibration and Control*, vol. 7, no. 5, pp. 631–642, 2001.
- [18] V. Marinca and N. Herişanu, "A modified iteration perturbation method for some nonlinear oscillation problems," *Acta Mechanica*, vol. 184, no. 1–4, pp. 231–242, 2006.
- [19] J. I. Ramos, "On the variational iteration method and other iterative techniques for nonlinear differential equations," *Applied Mathematics and Computation*, vol. 199, no. 1, pp. 39–69, 2008.
- [20] V. Marinca and N. Herişanu, "An optimal iteration method with application to the Thomas-Fermi equation," *Central European Journal of Physics*, vol. 9, no. 3, pp. 891–895, 2011.

- [21] Y.-M. Wang, "Global asymptotic stability of Lotka-Volterra competition reaction-diffusion systems with time delays," *Mathematical and Computer Modelling*, vol. 53, no. 1-2, pp. 337–346, 2011.
- [22] L. Nie, J. Peng, and Z. Teng, "Permanence and stability in multi-species non-autonomous Lotka-Volterra competitive systems with delays and feedback controls," *Mathematical and Computer Modelling*, vol. 49, no. 1-2, pp. 295–306, 2009.
- [23] M. Gyllenberg and P. Yan, "Four limit cycles for a three-dimensional competitive Lotka-Volterra system with a heteroclinic cycle," *Computers and Mathematics with Applications*, vol. 58, no. 4, pp. 649–669, 2009.
- [24] J. Giné, Z. Kadyrsizova, Y. Liu, and V. G. Romanovski, "Linearizability conditions for Lotka-Volterra planar complex quartic systems having homogeneous nonlinearities," *Computers and Mathematics with Applications*, vol. 61, no. 4, pp. 1190–1201, 2011.
- [25] J. Angelova, A. Dishliev, and S. Nenov, "I-optimal curve for impulsive Lotka-Volterra predator-prey model," *Computers and Mathematics with Applications*, vol. 43, no. 10-11, pp. 1203–1218, 2002.
- [26] A. Yıldırım and Z. Pınar, "Application of the exp-function method for solving nonlinear reaction-diffusion equations arising in mathematical biology," *Computers and Mathematics with Applications*, vol. 60, no. 7, pp. 1873–1880, 2010.
- [27] A. Yıldırım and Y. Gülkanat, "Reliable analysis for the nonlinear fractional calculus model of the semilunar heart valve vibrations," *International Journal for Numerical Methods in Engineering*, vol. 26, pp. 1515–1521, 2010.
- [28] A. Gökdoğan, M. Merdan, and A. Yıldırım, "A multistage differential transformation method for approximate solution of Hantavirus infection model," *Communications in Nonlinear Science and Numerical Simulation*, vol. 17, no. 1, pp. 1–8, 2012.
- [29] M. S. H. Chowdhury, I. Hashim, and O. Abdulaziz, "Application of homotopy-perturbation method to nonlinear population dynamics models," *Physics Letters A*, vol. 368, no. 3-4, pp. 251–258, 2007.
- [30] B. Batiha, M. S. M. Noorani, and I. Hashim, "Variational iteration method for solving multispecies Lotka-Volterra equations," *Computers and Mathematics with Applications*, vol. 54, no. 7-8, pp. 903–909, 2007.
- [31] A. Rëpaci, "Nonlinear dynamical systems: on the accuracy of Adomian's decomposition method," *Applied Mathematics Letters*, vol. 3, no. 4, pp. 35–39, 1990.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

