

Research Article

Bifurcation of Limit Cycles of a Class of Piecewise Linear Differential Systems in \mathbb{R}^4 with Three Zones

Yanyan Cheng

Department of Electronics and Information Engineering, Huazhong University of Science and Technology, Wuhan 430074, China

Correspondence should be addressed to Yanyan Cheng; cdymore@126.com

Received 1 March 2013; Accepted 16 April 2013

Academic Editor: Qingdu Li

Copyright © 2013 Yanyan Cheng. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the bifurcation of limit cycles from periodic orbits of a four-dimensional system when the perturbation is piecewise linear with two switching boundaries. Our main result shows that when the parameter is sufficiently small at most, six limit cycles can bifurcate from periodic orbits in a class of asymmetric piecewise linear perturbed systems, and, at most, three limit cycles can bifurcate from periodic orbits in another class of asymmetric piecewise linear perturbed systems. Moreover, there are perturbed systems having six limit cycles. The main technique is the averaging method.

1. Introduction and Statement of the Main Result

Piecewise linear systems are used extensively to model many physical phenomena, such as switching circuits in power electronics [1, 2] and impact and dry frictions in mechanical systems [3]. These systems exhibit not only standard bifurcations but also complicated dynamical phenomena not existing in smooth systems. The study and classification of various kinds of bifurcation phenomena for piecewise linear systems have attracted great attentions since the last century, see, for example, [4, 5] and the references therein.

In recent years, many papers studied the bifurcation of limit cycles and the number and distribution of these limit cycles. Most of them studied the planar piecewise linear system, see for example, [6–9] and the references quoted there. There are also some papers which studied bifurcation of limit cycles of 3D piecewise linear systems [10, 11]. For high-dimensional cases, there are a few papers [12–16]. Especially in [12] the authors studied the bifurcation of limit cycles of a class of piecewise linear systems in \mathbb{R}^4 . They showed that three is an upper bound for the number of limit cycles that bifurcate from periodic orbits.

In this paper, we study the limit cycles bifurcated from periodic orbits of a linear differential system in \mathbb{R}^4 when the perturbation is piecewise linear with two switching

boundaries. We consider two classes of asymmetric perturbation. With the first class of asymmetric perturbation, six is the upper bound for the number of limit cycles bifurcated from periodic orbits, and there are perturbed systems having six limit cycles. With the second class of asymmetric perturbation, three is the upper bound for the number of limit cycles bifurcated from periodic orbits, which generalizes the result of the paper [12].

More precisely, we study the maximum number of limit cycles of the 4-dimensional continuous piecewise linear vector fields with three zones of the form

$$\dot{x} = A_0 x + \varepsilon F(x), \quad (1)$$

for $\varepsilon \neq 0$ sufficiently small real parameter, where

$$A_0 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad (2)$$

and $F: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ is given by

$$F(x) = Ax + \varphi(k^T x)b, \quad (3)$$

with $A \in M_4(\mathbb{R})$, $k, b \in \mathbb{R}^4 \setminus \{0\}$, and $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ the piecewise linear function

if $m_1 < 0 < m_2$,

$$\varphi(x) = \begin{cases} hm_1, & \text{for } x \in (-\infty, m_1), \\ hx, & \text{for } x \in (m_1, m_2), \\ hm_2, & \text{for } x \in (m_2, +\infty); \end{cases} \quad (4)$$

if $m_1 < m_2 < 0$,

$$\varphi(x) = \begin{cases} hm_1 - hm_2, & \text{for } x \in (-\infty, m_1), \\ hx - hm_2, & \text{for } x \in (m_1, m_2), \\ 0, & \text{for } x \in (m_2, +\infty); \end{cases} \quad (5)$$

if $0 < m_1 < m_2$,

$$\varphi(x) = \begin{cases} 0, & \text{for } x \in (-\infty, m_1), \\ hx - hm_1, & \text{for } x \in (m_1, m_2), \\ hm_2 - hm_1, & \text{for } x \in (m_2, +\infty), \end{cases} \quad (6)$$

where $h \in \mathfrak{R} \setminus \{0\}$. The independent variable is denoted by t ; vectors of \mathfrak{R}^4 are column vectors, and k^T denotes a transposed vector.

For $\varepsilon = 0$, system (1) becomes

$$\dot{x}_1 = -x_2, \quad \dot{x}_2 = x_1, \quad \dot{x}_3 = -x_4, \quad \dot{x}_4 = x_3. \quad (7)$$

Our main results are the following.

Theorem 1. *If $m_1 m_2 > 0$, six is the upper bound for the number of limit cycles of system (1) which bifurcate from the periodic orbits of system (7) with ε sufficiently small. Moreover, there are systems of form (1) having six limit cycles.*

Theorem 2. *If $m_1 m_2 < 0$, three is the upper bound for the number of limit cycles of system (1) which bifurcate from the periodic orbits of system (7) with ε sufficiently small. Moreover, there are systems of form (1) having three limit cycles.*

It is worth to note that Theorem 2 generalizes the result of paper [12]. The method for computing the number of limit cycles bifurcated from periodic orbits is the averaging method, which is obtained by Buică and Llibre [17]. By means of the result of paper [18], we can study the stability of the limit cycles of Theorem 1; for more details see Remark 10.

Theorems 1 and 2 will be proved in Section 3. In Section 2, we review the results from the averaging theory necessary for proving these two theorems. Further discussions on the number of limit cycles of the perturbed system are present in Section 4. There is a conclusion given in the last section.

2. First-Order Averaging Method

The aim of this section is to review the first-order averaging method which is obtained by Buică and Llibre [17]. The advantage of this method is that the smoothness assumptions for the vector field of the differential system are minimal.

Theorem 3 (see [17]). *Consider the following differential system:*

$$\dot{x}(t) = \varepsilon H(t, \varepsilon) + \varepsilon^2 R(t, x, \varepsilon), \quad (8)$$

where $H : \mathfrak{R} \times D \rightarrow \mathfrak{R}^n$, $R : \mathfrak{R} \times D \times (-\varepsilon_0, \varepsilon_0) \rightarrow \mathfrak{R}^n$ are continuous functions. T -periodic in the first variable, and D is an open subset of \mathfrak{R}^n . We define $h : D \rightarrow \mathfrak{R}^n$ as

$$h(z) = \int_0^T H(s, z) ds, \quad (9)$$

and assume that

- (i) H and R are locally Lipschitz with respect to x ;
- (ii) for $a \in D$ with $h(a) = 0$, there exists a neighborhood V of a such that $h(z) \neq 0$ for all $z \in \bar{V} \setminus \{a\}$ and $d_B(h, V, 0) \neq 0$.

Then, for $|\varepsilon| > 0$ sufficiently small, there exists an isolated T -periodic solution $\phi(\cdot, \varepsilon)$ of system (8) such that $\phi(\cdot, \varepsilon) \rightarrow a$ as $\varepsilon \rightarrow 0$.

We remind here that $d_B(h, V, a)$ denotes the Brouwer degree of the function h with respect to the set V and the point a , as is defined in [19]. The following fact is useful for the proof of Theorems 1 and 2.

Fact 1. Let $h : D \rightarrow \mathfrak{R}^n$ be a C^1 function, with $h(a) = 0$, where D is an open subset of \mathfrak{R}^n and $a \in D$. Whenever a is a simple zero of h (i.e., $J(a) \neq 0$), there exists a neighborhood V of a such that $h(z) \neq 0$ for all $z \in \bar{V} \setminus \{a\}$. Then, $d_B(h, V, 0) \in \{-1, 1\}$.

3. Proof of Main Theorems

The proof of Theorems 1 and 2 is based on the first-order averaging method presented in the previous section. In order to apply this method, we will first reduce the four parameters of the vector k in the definition of the function $F(x)$ to one, and then we will change the variables in order to transform the system into the standard form for the averaging method. After that, we will calculate the number of its isolated zeros.

Lemma 4. *By a linear change of variables, system (1) can be transformed into the system*

$$\dot{x} = A_0 x + \varepsilon \bar{A} x + \varepsilon \varphi(x_1) \bar{b}, \quad (10)$$

where $\bar{A} \in M_4(\mathfrak{R})$ is an arbitrary matrix and $\bar{b} = (\bar{b}_1, \bar{b}_2, 0, 0)^T$ or $\bar{b} = e_3$.

Proof. A linear change of variables $x = Py$, with P invertible, transforms system (1) into

$$\dot{y} = P^{-1} A_0 P y + \varepsilon P^{-1} A P y + \varepsilon \varphi(k^T P y) P^{-1} b, \quad (11)$$

where $b = (b_1, b_2, b_3, b_4)^T$, $k = (k_1, k_2, k_3, k_4)^T$.

We have to find P invertible which satisfies

$$\begin{aligned} P^{-1} A_0 P &= A_0, \\ k^T P &= e_1^T. \end{aligned} \quad (12)$$

It is easy to obtain that P^{-1} has the following form:

$$P^{-1} = \begin{pmatrix} k_1 & k_2 & k_3 & k_4 \\ -k_2 & k_1 & -k_4 & k_3 \\ p_{31} & p_{32} & p_{33} & p_{34} \\ -p_{32} & p_{31} & -p_{34} & p_{33} \end{pmatrix}. \quad (13)$$

Thus, we have

$$P^{-1}b = \bar{b}, \quad (14)$$

where

$$\begin{aligned} \bar{b}_1 &= \sum_{i=1}^4 k_i b_i, & \bar{b}_2 &= -k_2 b_1 + k_1 b_2 - k_4 b_3 + k_3 b_4, \\ \bar{b}_3 &= \sum_{i=1}^4 p_{3i} b_i, & \bar{b}_4 &= -p_{32} b_1 + p_{31} b_2 - p_{34} b_3 + p_{33} b_4. \end{aligned} \quad (15)$$

If $\bar{b}_1^2 + \bar{b}_2^2 \neq 0$, it is easy to find P^{-1} invertible with p_{31} , p_{32} , p_{33} , p_{34} satisfying

$$\bar{b}_3 = \sum_{i=1}^4 p_{3i} b_i = 0, \quad (16)$$

$$\bar{b}_4 = -p_{32} b_1 + p_{31} b_2 - p_{34} b_3 + p_{33} b_4 = 0.$$

If $\bar{b}_1^2 + \bar{b}_2^2 = 0$, it is easy to find P^{-1} invertible with p_{31} , p_{32} , p_{33} , p_{34} satisfying

$$\bar{b}_3 = \sum_{i=1}^4 p_{3i} b_i = 1, \quad (17)$$

$$\bar{b}_4 = -p_{32} b_1 + p_{31} b_2 - p_{34} b_3 + p_{33} b_4 = 0.$$

Changing variables y to x with $x = y$, then we obtain system (10). \square

The standard form of the averaging method is obtained by changing variables (x_1, x_2, x_3, x_4) to (r, θ, ρ, s) with

$$\begin{aligned} x_1 &= r \cos \theta, & x_2 &= r \sin \theta, \\ x_3 &= \rho \cos (\theta + s), & x_4 &= \rho \sin (\theta + s). \end{aligned} \quad (18)$$

Thus, system (10) is transformed into the following system:

$$\begin{aligned} \frac{dr}{d\theta} &= \varepsilon H_1(\theta, r, \rho, s) + \varepsilon^2 O(1), \\ \frac{d\rho}{d\theta} &= \varepsilon H_2(\theta, r, \rho, s) + \varepsilon^2 O(1), \\ \frac{ds}{d\theta} &= \varepsilon H_3(\theta, r, \rho, s) + \varepsilon^2 O(1), \end{aligned} \quad (19)$$

where H_1 , H_2 , and H_3 are given by

$$\begin{aligned} H_1 &= \cos \theta F_1 + \sin \theta F_2, \\ H_2 &= \cos (\theta + s) F_3 + \sin (\theta + s) F_4, \end{aligned}$$

$$\begin{aligned} H_3 &= \frac{1}{r} \sin \theta F_1 - \frac{1}{r} \cos \theta F_2 - \frac{1}{\rho} \sin (\theta + s) F_3 \\ &\quad + \frac{1}{\rho} \cos (\theta + s) F_4 \end{aligned} \quad (20)$$

and for every $i = 1, 2, \dots, 4$,

$$\begin{aligned} F_i &= a_{i1} r \cos \theta + a_{i2} r \sin \theta + a_{i3} \rho \cos (\theta + s) + a_{i4} \rho \sin (\theta + s) \\ &\quad + \varphi(r \cos \theta) \bar{b}_i, \end{aligned} \quad (21)$$

where a_{ij} are elements of the matrix \bar{A} of Lemma 4.

We take ε_0 sufficiently small, m arbitrarily large and

$$D_m = \left\{ (r, \rho, s) \mid (r, \rho, s) \in \left(\frac{1}{m}, m \right)^2 \times S \right\}. \quad (22)$$

Then, the vector of system (19) is well defined and continuous on $S \times D_m \times (-\varepsilon_0, \varepsilon_0)$. Moreover, the system is 2π -periodic with respect to variable θ and locally Lipschitz with respect to variables (r, ρ, s) . Our next step is to find the corresponding function $h : D \rightarrow \mathfrak{R}^3$, $h = (h_1, h_2, h_3)$, where

$$h_i(r, \rho, s) = \int_0^{2\pi} H_i(r, \theta, \rho, s) d\theta, \quad (23)$$

for $i = 1, 2, 3$.

In order to calculate the exact expression of h , we denote

$$I_1(r) = \int_0^{2\pi} \varphi(r \cos \theta) \cos \theta d\theta, \quad (24)$$

$$I_2(r) = \int_0^{2\pi} \varphi(r \cos \theta) \sin \theta d\theta, \quad (25)$$

for each $r > 0$, where φ is the piecewise linear function given by (4)–(6). Without loss of generality, we assume that the slope h of φ is positive.

Lemma 5. *The integrals I_1 and I_2 given by (24)–(25), respectively, have the following expressions:*

$$I_2(r) = 0, \quad \forall r > 0, m_1, m_2, \quad (26)$$

and

(1) if $0 < m_1 < m_2$,

$$I_1(r) = \begin{cases} 0, & \text{if } 0 < r \leq m_1, \\ J(r, m_1), & \text{if } m_1 < r < m_2, \\ K(r, m_1, m_2), & \text{if } r \geq m_2; \end{cases} \quad (27)$$

(2) if $m_1 < m_2 < 0$,

$$I_1(r) = \begin{cases} 0, & \text{if } 0 < r \leq |m_2|, \\ -J(r, m_2), & \text{if } |m_2| < r < |m_1|, \\ K(r, m_1, m_2), & \text{if } r \geq |m_1|; \end{cases} \quad (28)$$

(3) if $m_1 < 0 < m_2$ and $|m_1| < |m_2|$,

$$I_1(r) = \begin{cases} \pi hr, & \text{if } 0 < r \leq |m_1|, \\ \pi hr + J(r, m_1), & \text{if } |m_1| < r < m_2, \\ \pi hr + K(r, m_1, m_2), & \text{if } r \geq m_2; \end{cases} \quad (29)$$

(4) if $m_1 < 0 < m_2$ and $|m_1| > |m_2|$,

$$I_1(r) = \begin{cases} \pi hr, & \text{if } 0 < r \leq m_2, \\ \pi hr - J(r, m_2), & \text{if } m_2 < r < |m_1|, \\ \pi hr + K(r, m_1, m_2), & \text{if } r \geq |m_1|; \end{cases} \quad (30)$$

(5) if $m_1 < 0 < m_2$ and $|m_1| = |m_2| = m$,

$$I_1(r) = \begin{cases} \pi hr, & \text{if } 0 < r \leq m, \\ \pi hr - 2J(r, m), & \text{if } r > m, \end{cases} \quad (31)$$

where

$$J(r, m_i) = hr \left(\arctan \frac{\sqrt{r^2 - m_i^2}}{m_i} - \frac{m_i \sqrt{r^2 - m_i^2}}{r^2} \right), \quad (32)$$

for $i = 1, 2$, and

$$K(r, m_1, m_2) = hr \left(\arctan \frac{\sqrt{r^2 - m_1^2}}{m_1} - \arctan \frac{\sqrt{r^2 - m_2^2}}{m_2} - \frac{m_1 \sqrt{r^2 - m_1^2}}{r^2} + \frac{m_2 \sqrt{r^2 - m_2^2}}{r^2} \right). \quad (33)$$

The proof of this lemma is given in the appendix.

Remark 6. If $m_1 < 0 < m_2$ and $|m_1| = |m_2|$, system (1) can be transformed into the system which is studied in the paper [12].

Lemma 7. If $m_1 m_2 > 0$, one defines $\bar{m} = \max(|m_1|, |m_2|)$, $\underline{m} = \min(|m_1|, |m_2|)$ and consider the equation $I_1(r) = cr$, $r > 0$ with I_1 given by (24), and c is a real parameter. Then,

- (1) if $c < 0$ or $c > h(\arctan(\bar{m}/\underline{m}) - \arctan(\underline{m}/\bar{m}))$, the equation has no solutions;
- (2) if $c = 0$, then the interval $(0, \underline{m}]$ is a continuum of solutions;
- (3) if $c = h(\arctan(\bar{m}/\underline{m}) - \arctan(\underline{m}/\bar{m}))$, there is an unique solution $r^* = \sqrt{\bar{m}^2 + \underline{m}^2}$;
- (4) if $c \in (0, h(\arctan(\bar{m}/\underline{m}) - \arctan(\underline{m}/\bar{m})))$, there are two solutions $r_1^* < \sqrt{\bar{m}^2 + \underline{m}^2}$ and $r_2^* > \sqrt{\bar{m}^2 + \underline{m}^2}$.

Proof. If $0 < m_1 < m_2$, we have $\bar{m} = m_2$ and $\underline{m} = m_1$. It is easy to see that all $r \in (0, \underline{m}]$ are a solution if $c = 0$. If $c \neq 0$ changing the variable $u = \sqrt{r^2 - m_1^2}/m_1$ and defining $g(u) = I_1(r)/hr$, we obtain the equivalent equation

$$g(u) = \begin{cases} \arctan u - \frac{u}{1+u^2}, & \text{if } u \in \left(0, \frac{\sqrt{m_2^2 - m_1^2}}{m_1}\right), \\ \arctan u - \frac{u}{1+u^2} - \arctan \frac{\sqrt{(1+u^2)m_1^2 - m_2^2}}{m_2} + \frac{m_2 \sqrt{(1+u^2)m_1^2 - m_2^2}}{m_1^2(1+u^2)}, & \text{if } u \in \left(\frac{\sqrt{m_2^2 - m_1^2}}{m_1}, +\infty\right) \end{cases} \quad (34)$$

with simple computation; we find that the function g is strictly monotonically increasing of variable u when $u \in (0, m_2/m_1)$ and strictly monotonically decreasing when $u \in (m_2/m_1, +\infty)$. The function g gets to the maximal value $g_{\max} = \arctan(m_2/m_1) - \arctan(m_1/m_2)$ when $u = m_2/m_1$. Also we have $f(u) \rightarrow 0$ as $u \rightarrow 0$ and $u \rightarrow +\infty$. The proof is similar if $m_1 < m_2 < 0$. \square

Lemma 8. If $m_1 m_2 < 0$, one defines $\underline{m} = \min(|m_1|, |m_2|)$ and consider the equation $I_1(r) = cr$, $r > 0$ with I_1 given by (24), and c is a real parameter. Then,

- (1) if $c < 0$ or $c > \pi h$, the equation has no solutions;
- (2) if $c = \pi h$, then the interval $(0, \underline{m}]$ is a continuum of solutions;
- (3) if $c \in (0, \pi h)$, there is an unique solution r^* .

Proof. We only consider the case when $|m_1| < |m_2|$; the proof is similar when $|m_1| > |m_2|$ and $|m_1| = |m_2|$. It is easy to see that all $r \in (0, |m_1|]$ are a solution if $c = \pi h$. If $c \neq 0$ changing the variable $u = \sqrt{r^2 - m_1^2}/m_1$ and defining $g(u) = I_1(r)/hr$, we obtain the equivalent equation

$$g(u) = \begin{cases} \pi + \arctan u - \frac{u}{1+u^2}, & \text{if } u \in \left[\frac{m_2^2 - m_1^2}{m_1}, 0\right), \\ \pi + \arctan u - \frac{u}{1+u^2} - \arctan \frac{\sqrt{(1+u^2)m_1^2 - m_2^2}}{m_2} + \frac{m_2 \sqrt{(1+u^2)m_1^2 - m_2^2}}{m_1^2(1+u^2)}, & \text{if } u \in \left(-\infty, \frac{\sqrt{m_2^2 - m_1^2}}{m_1}\right). \end{cases} \quad (35)$$

With simple computation, we find that the function g is strictly monotonically increasing of variable u . It is easy to know $f(u) \rightarrow \pi$ as $u \rightarrow 0$ and $f(u) \rightarrow 0$ as $u \rightarrow -\infty$. \square

With Lemma 5, we obtain the expressions for the components of function h ,

$$h_1(r, \rho, s) = c_1 r + c_2 \rho \cos s + c_3 \rho \sin s + \bar{b}_1 I_1(r),$$

$$\begin{aligned}
h_2(r, \rho, s) &= c_7 \rho + c_5 r \cos s + c_6 r \sin s + \bar{b}_3 \cos s I_1(r), \\
h_3(r, \rho, s) &= c_4 + (c_3 \cos s - c_2 \sin s) \frac{\rho}{r} + (c_6 \cos s - c_5 \sin s) \frac{r}{\rho} \\
&\quad - \bar{b}_3 \sin s \frac{I_1(r)}{\rho} - \bar{b}_2 \frac{I_1(r)}{r},
\end{aligned} \tag{36}$$

where c_i are constants that depend linearly on a_{ij}

$$\begin{aligned}
c_1 &= (a_{11} + a_{22}) \pi, & c_2 &= (a_{13} + a_{24}) \pi, \\
c_3 &= (a_{14} - a_{23}) \pi, & c_4 &= (a_{43} + a_{12} - a_{34} - a_{21}) \pi, \\
c_5 &= (a_{31} + a_{42}) \pi, & c_6 &= (a_{41} - a_{32}) \pi, \\
c_7 &= (a_{33} + a_{44}) \pi.
\end{aligned} \tag{37}$$

According to Theorem 3 and Fact 1, for each simple zero (r^*, ρ^*, s^*) of (36) there is an isolated 2π -periodic solution $\phi(\cdot, \varepsilon)$ of system (19) with $|\varepsilon| \neq 0$ sufficiently small such that $\phi(\cdot, \varepsilon) \rightarrow (r^*, \rho^*, s^*)$ as $\varepsilon \rightarrow 0$. Any isolated 2π -periodic solution of system (19) with $|\varepsilon| \neq 0$ sufficiently small corresponds to a limit cycle of system (10). Thus, the most important task is to calculate the number of the simple zeros of function h . We solve the two first equations of (36), then, we get

$$I_1(r) = \frac{k_2(s)}{d(s)} \rho, \quad r = \frac{k_1(s)}{d(s)} \rho, \tag{38}$$

where

$$\begin{aligned}
d(s) &= (\bar{b}_1 c_5 - \bar{b}_3 c_1) \cos s + \bar{b}_1 c_6 \sin s, \\
k_1(s) &= \bar{b}_3 \cos s (c_2 \cos s + c_3 \sin s) + \bar{b}_1 c_7, \\
k_2(s) &= c_1 c_7 - c_2 c_5 \cos^2 s - c_3 c_6 \sin^2 s \\
&\quad - (c_2 c_6 + c_3 c_5) \sin s \cos s.
\end{aligned} \tag{39}$$

Substituting (38) into the third equation, we obtain

$$h_3(r, \rho, s) = \frac{f(s)}{d(s) k_1(s)} = 0, \tag{40}$$

where

$$\begin{aligned}
f(s) &= c_4 d(s) k_1(s) + (c_3 \cos s - c_2 \sin s) d^2(s) \\
&\quad + (c_6 \cos s - c_5 \sin s) k_1^2(s) \\
&\quad - \bar{b}_3 \sin s k_1(s) k_2(s) - \bar{b}_2 d(s) k_2(s).
\end{aligned} \tag{41}$$

It is necessary to study the zeros of f instead of the zeros of h .

Lemma 9. *The function $f : [0, 2\pi) \rightarrow \mathbb{R}$ given by formula (41) can have at most six isolated zeros, and they appear in pairs $\{s^*, s^* + \pi(\bmod 2\pi)\}$.*

Proof. Substituting $\cos s = x$ and $\sin s = -\sqrt{1-x^2}$ in $f(s) = 0$ we get

$$D_1 x + D_3 x^3 + (D_0 + D_2 x^2) \sqrt{1-x^2} = 0, \tag{42}$$

where

$$\begin{aligned}
D_0 &= \bar{b}_1^2 (c_2 c_6^2 - c_4 c_6 c_7 + c_5 c_7^2) + \bar{b}_1 \bar{b}_2 (c_3 c_6^2 - c_1 c_6 c_7), \\
D_1 &= \bar{b}_1^2 (2c_2 c_5 c_6 - c_3 c_6^2 - c_4 c_5 c_7 - c_6 c_7^2) + \bar{b}_3^2 (c_1 c_3 c_7 - c_6 c_3^2) \\
&\quad + \bar{b}_1 \bar{b}_2 (2c_2 c_5 c_6 + c_3 c_5^2 - c_1 c_5 c_7), \\
D_2 &= \bar{b}_1^2 (c_2 c_5^2 - c_2 c_6^2 - 2c_3 c_5 c_6) \\
&\quad + \bar{b}_3^2 (c_1 c_2 c_7 + c_2 c_1^2 - c_1 c_3 c_4 - 2c_2 c_3 c_6) \\
&\quad + \bar{b}_1 \bar{b}_2 (2c_2 c_5 c_6 + c_3 c_5^2 - c_3 c_6^2), \\
D_3 &= \bar{b}_1^2 (c_3 c_6^2 - c_3 c_5^2 - 2c_2 c_5 c_6) \\
&\quad + \bar{b}_3^2 (c_3^2 c_6 - c_3 c_1^2 - c_1 c_2 c_4 - c_1 c_3 c_7 - c_6 c_2^2) \\
&\quad + \bar{b}_1 \bar{b}_2 (c_2 c_5^2 - 2c_3 c_5 c_6 - c_2 c_6^2).
\end{aligned} \tag{43}$$

When we consider the case $\cos s = x$ and $\sin s = -\sqrt{1-x^2}$, $f(s) = 0$ becomes

$$D_1 x + D_3 x^3 - (D_0 + D_2 x^2) \sqrt{1-x^2} = 0. \tag{44}$$

It follows that we have to find solutions of (42) or (44) in the interval $[-1, 1]$. This is equivalent to

$$D_1 x + D_3 x^3 - (D_0 + D_2 x^2)^2 (1-x^2) = 0 \tag{45}$$

which is the polynomial equation

$$\begin{aligned}
&(D_3^2 + D_2^2) x^6 + (2D_1 D_3 + 2D_0 D_2 - D_2^2) x^4 \\
&\quad + (D_1^2 + D_0^2 - 2D_0 D_2) x^2 - D_0^2 = 0.
\end{aligned} \tag{46}$$

This equation can have at most six roots in the interval $[-1, 1]$. Then, $f(s) = 0$ has at most six solutions $s \in [0, 2\pi)$. Since $f(s + \pi) = -f(s)$ for all $s \in [0, 2\pi)$, it is clear that if s^* is a zero of f then $s^* + \pi(\bmod 2\pi)$ is also a zero. \square

The functions $f(s)$, $d(s)$, $k_1(s)$, and $k_2(s)$ have the properties $f(s + \pi) = -f(s)$, $d(s + \pi) = -d(s)$, $k_1(s + \pi) = k_1(s)$, and $k_2(s + \pi) = k_2(s)$. So, we have

$$\frac{k_1(s)}{d(s)} > 0 \implies \frac{k_1(s + \pi)}{d(s + \pi)} < 0. \tag{47}$$

Thus, the equation $h_3 = 0$ at most three zeros that satisfy $k_1(s)/d(s) > 0$. With Lemma 7 for a fixed s^* , we at most find two isolated value of r^* from $I_1(r)/r = k_2(s^*)/d(s^*)$. With Lemma 8 for a fixed s^* , we at most find one isolated value of r^* from $I_1(r)/r = k_2(s^*)/d(s^*)$. For fixed s^* and fixed r^* , $\rho/r = k_1(s^*)/d(s^*)$ gives at most one isolated value for ρ^* . Thus, we conclude that if $m_1 m_2 > 0$ the maximum number of limit cycles for system (1) is six, and if $m_1 m_2 < 0$ the maximum number of limit cycles for system (1) is three.

Remark 10. Using the main result of [18], the stability of the limit cycles associated with the solution (r^*, ρ^*, s^*) is given by the eigenvalues of the matrix

$$\frac{\partial (h_1, h_2, h_3)}{\partial (r, \rho, s)} \Big|_{(r, \rho, s) = (r^*, \rho^*, s^*)}. \quad (48)$$

In order to show that there exist examples with exactly six limit cycles, we consider the following values of the coefficients:

$$\begin{aligned} c_1 &= -1, & c_2 &= \frac{-4\sqrt{3}}{3}, & c_3 &= 0, \\ c_4 &= \frac{-4\sqrt{3}}{3}, & c_5 &= 0, & c_6 &= 1, & c_7 &= -1, \\ \bar{b}_1 &= 1, & \bar{b}_2 &= \bar{b}_3 = \bar{b}_4 = 0, & h &= 6, \\ m_1 &= 1, & m_2 &= \sqrt{3}. \end{aligned} \quad (49)$$

More precisely, the system has the following form:

$$\dot{x} = A_0 x + \varepsilon A x + \varepsilon \varphi(x_1) \bar{b}, \quad (50)$$

where

$$\varphi(x_1) = \begin{cases} 0, & \text{for } x_1 \in (-\infty, 1), \\ 6x_1 - 6, & \text{for } x_1 \in (1, \sqrt{3}), \\ 6\sqrt{3} - 6, & \text{for } x_1 \in (\sqrt{3}, +\infty), \end{cases}$$

$$A_0 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad (51)$$

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{14} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & -a_{31} & a_{43} & a_{44} \end{pmatrix}, \quad \bar{b} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix},$$

and $a_{ij} \in \mathfrak{R}$ satisfy

$$\begin{aligned} (a_{11} + a_{22})\pi &= -1, & (a_{13} + a_{24})\pi &= \frac{-4\sqrt{3}}{3}, \\ (a_{43} + a_{12} - a_{34} - a_{21})\pi &= \frac{-4\sqrt{3}}{3}, \\ (a_{41} - a_{32})\pi &= 1, & (a_{33} + a_{44})\pi &= -1. \end{aligned} \quad (52)$$

It is easy to know $6(\arctan \sqrt{3} - \arctan(\sqrt{3}/3)) = \pi$. Computing the six solutions of $f(s) = 0$, we get $\{\pi/6, \pi/3, \pi/2, 7\pi/6, 4\pi/3, 3\pi/2\}$. The values of $d(s^*)$, $k_1(s^*)$ and $k_2(s^*)$ are given in Table 1.

There are three values of s^* that satisfy $k_1(s^*)/d(s^*) > 0$ and $0 < k_2(s^*)/k_1(s^*) < \pi$. These three solutions are $\{\pi/6, \pi/3, \pi/2\}$.

The six values of solution s^* , r^* , ρ^* and the value of the Jacobian at the solution (r^*, ρ^*, s^*) are given in Table 2.

TABLE 1: The values of s^* , $d(s^*)$, $k_1(s^*)$, and $k_2(s^*)$.

s^*	$k_1(s^*)$	$d(s^*)$	$k_2(s^*)$
$\pi/6$	1	1/2	2
$\pi/3$	1	$\sqrt{3}/2$	2
$\pi/2$	1	1	1
$7\pi/6$	1	-1/2	2
$4\pi/3$	1	$-\sqrt{3}/2$	2
$3\pi/2$	1	-1	1

TABLE 2: The values of solution s^* , r^* , ρ^* and the Jacobian $Jh(r^*, \rho^*, s^*)$.

s^*	r^*	ρ^*	$Jh(r^*, \rho^*, s^*)$
$\pi/6$	1.484	0.742	11.957
$\pi/6$	4.139	2.07	-4.688
$\pi/3$	1.484	1.285	-3.27
$\pi/3$	4.139	3.585	-1.144
$\pi/2$	1.254	1.254	5.732
$\pi/2$	8.672	8.672	-0.964

4. Conclusion

In this paper, we have studied the limit cycles bifurcated from periodic orbits of a linear differential system in \mathfrak{R}^4 when the perturbation is piecewise linear with two switching boundaries. We considered two classes of asymmetric perturbation. We have found that the perturbed system could have at most six limit cycles with one class of the asymmetric perturbation and three limit cycles with the other class of asymmetric perturbation, which generalized the result of paper [12].

Appendix

The Proof of Lemma 5

Case 1 ($0 < m_1 < m_2$). We have $|r \sin \theta| \leq m_1$ and $|r \cos \theta| \leq m_1$ for all $\theta \in [0, 2\pi)$ if $0 < r \leq m_1$. Then, $\varphi(r \cos \theta) = 0$ for every θ . Thus,

$$I_1(r) = I_2(r) = 0. \quad (A.1)$$

We now fix $m_1 < r < m_2$ and consider $\theta_c \in (0, \pi/2)$ which satisfies $r \cos \theta_c = m_1$. Then, we have

$$\begin{aligned} I_1(r) &= \int_0^{\theta_c} (hr \cos \theta - hm_1) \cos \theta d\theta \\ &\quad + \int_{2\pi-\theta_c}^{2\pi} (hr \cos \theta - hm_1) \cos \theta d\theta \\ &= hr\theta_c - hm_1 \sin \theta_c, \\ I_2(r) &= \int_0^{\theta_c} (hr \cos \theta - hm_1) \sin \theta d\theta \\ &\quad + \int_{2\pi-\theta_c}^{2\pi} (hr \cos \theta - hm_1) \sin \theta d\theta \\ &= 0. \end{aligned} \quad (A.2)$$

We now fix $r \geq m_2$ and consider $0 < \tilde{\theta}_c < \theta_c < \pi/2$ which satisfies $r \cos \tilde{\theta}_c = m_2$. Then, we have

$$\begin{aligned} I_1(r) &= \int_{-\tilde{\theta}_c}^{\tilde{\theta}_c} (hm_2 - hm_1) \cos \theta d\theta \\ &\quad + \int_{\tilde{\theta}_c}^{\theta_c} (hr \cos \theta - hm_1) \cos \theta d\theta \\ &\quad + \int_{2\pi-\tilde{\theta}_c}^{2\pi-\theta_c} (hr \cos \theta - hm_1) \cos \theta d\theta, \\ I_2(r) &= \int_{-\tilde{\theta}_c}^{\tilde{\theta}_c} (hm_2 - hm_1) \sin \theta d\theta \\ &\quad + \int_{\tilde{\theta}_c}^{\theta_c} (hr \cos \theta - hm_1) \sin \theta d\theta \\ &\quad + \int_{2\pi-\tilde{\theta}_c}^{2\pi-\theta_c} (hr \cos \theta - hm_1) \sin \theta d\theta. \end{aligned} \quad (\text{A.3})$$

With simple computation, we get

$$\begin{aligned} I_1(r) &= hm_2 \sin \tilde{\theta}_c - hm_1 \sin \theta_c + hr\theta_c - hr\tilde{\theta}_c, \\ I_2(r) &= 0, \end{aligned} \quad (\text{A.4})$$

where

$$\sin \theta_c = \frac{\sqrt{r^2 - m_1^2}}{r}, \quad \sin \tilde{\theta}_c = \frac{\sqrt{r^2 - m_2^2}}{r}, \quad (\text{A.5})$$

$$\theta_c = \arctan \frac{\sqrt{r^2 - m_1^2}}{m_1}, \quad \tilde{\theta}_c = \arctan \frac{\sqrt{r^2 - m_2^2}}{m_2}. \quad (\text{A.6})$$

Case 2 ($m_1 < m_2 < 0$). We have $|r \sin \theta| \leq |m_2|$ and $|r \cos \theta| \leq |m_2|$ for all $\theta \in [0, 2\pi]$ if $0 < r \leq |m_2|$. Then, $\varphi(r \cos \theta) = 0$ for every θ . Thus,

$$I_1(r) = I_2(r) = 0. \quad (\text{A.7})$$

We fix now $|m_2| < r < |m_1|$ and consider $\theta_c \in (\pi/2, \pi)$ which satisfies $r \cos \theta_c = m_2$. Then, we have

$$\begin{aligned} I_1(r) &= \int_{\theta_c}^{2\pi-\theta_c} (hr \cos \theta - hm_2) \cos \theta d\theta \\ &= \pi hr - hr\theta_c + hm_2 \sin \theta_c, \\ I_2(r) &= \int_{\theta_c}^{2\pi-\theta_c} (hr \cos \theta - hm_2) \sin \theta d\theta = 0. \end{aligned} \quad (\text{A.8})$$

We now fix $r \geq |m_1|$ and consider $\tilde{\theta}_c \in (\pi/2, \pi)$ which satisfies $r \cos \tilde{\theta}_c = m_1$. Obviously, $\theta_c < \tilde{\theta}_c$. Then, we have

$$\begin{aligned} I_1(r) &= \int_{\theta_c}^{\tilde{\theta}_c} (hr \cos \theta - hm_2) \cos \theta d\theta \\ &\quad + \int_{\tilde{\theta}_c}^{2\pi-\tilde{\theta}_c} (hm_1 - hm_2) \cos \theta d\theta \\ &\quad + \int_{2\pi-\tilde{\theta}_c}^{2\pi-\theta_c} (hr \cos \theta - hm_2) \cos \theta d\theta, \\ I_2(r) &= \int_{\theta_c}^{\tilde{\theta}_c} (hr \cos \theta - hm_2) \sin \theta d\theta \\ &\quad + \int_{\tilde{\theta}_c}^{2\pi-\tilde{\theta}_c} (hm_1 - hm_2) \sin \theta d\theta \\ &\quad + \int_{2\pi-\tilde{\theta}_c}^{2\pi-\theta_c} (hr \cos \theta - hm_2) \sin \theta d\theta. \end{aligned} \quad (\text{A.9})$$

With simple computation, we get

$$\begin{aligned} I_1(r) &= hm_2 \sin \theta_c - hm_1 \sin \tilde{\theta}_c - hr\theta_c + hr\tilde{\theta}_c, \\ I_2(r) &= 0, \end{aligned} \quad (\text{A.10})$$

where

$$\begin{aligned} \sin \theta_c &= \frac{\sqrt{r^2 - m_2^2}}{r}, & \sin \tilde{\theta}_c &= \frac{\sqrt{r^2 - m_1^2}}{r}, \\ \theta_c &= \pi + \arctan \frac{\sqrt{r^2 - m_2^2}}{m_2}, & \tilde{\theta}_c &= \pi + \arctan \frac{\sqrt{r^2 - m_1^2}}{m_1}. \end{aligned} \quad (\text{A.11})$$

Case 3 ($m_1 < 0 < m_2$ and $|m_1| < |m_2|$). We have $|r \sin \theta| \leq |m_1|$ and $|r \cos \theta| \leq |m_1|$ for all $\theta \in [0, 2\pi]$ if $0 < r \leq |m_1|$. Then, $\varphi(r \cos \theta) = hr \cos \theta$ for every θ . Thus,

$$\begin{aligned} I_1(r) &= \int_0^{2\pi} hr \cos^2 \theta d\theta = \pi hr, \\ I_2(r) &= \int_0^{2\pi} hr \cos \theta \sin \theta d\theta = 0. \end{aligned} \quad (\text{A.12})$$

We fix now $|m_1| < r < |m_2|$ and consider $\theta_c \in (\pi/2, \pi)$ which satisfies $r \cos \theta_c = m_1$. Then, we have

$$\begin{aligned} I_1(r) &= \int_{-\theta_c}^{\theta_c} hr \cos^2 \theta d\theta \\ &\quad + \int_{\theta_c}^{2\pi-\theta_c} hm_1 \cos \theta d\theta = hr\theta_c - hm_1 \sin \theta_c, \\ I_2(r) &= \int_{-\theta_c}^{\theta_c} hr \sin \theta \cos \theta d\theta + \int_{\theta_c}^{2\pi-\theta_c} hm_1 \sin \theta d\theta = 0. \end{aligned} \quad (\text{A.13})$$

We now fix $r \geq m_2$ and consider $\tilde{\theta}_c \in (0, \pi/2)$ which satisfies $r \cos \tilde{\theta}_c = m_2$. Then, we can write

$$I_1(r) = \int_{-\tilde{\theta}_c}^{\tilde{\theta}_c} hm_2 \cos \theta d\theta + \int_{\tilde{\theta}_c}^{\theta_c} hr \cos^2 \theta d\theta \quad (\text{A.14})$$

$$+ \int_{\theta_c}^{2\pi-\theta_c} hm_1 \cos \theta d\theta + \int_{2\pi-\theta_c}^{2\pi-\tilde{\theta}_c} hr \cos^2 \theta d\theta,$$

$$I_2(r) = \int_{-\tilde{\theta}_c}^{\tilde{\theta}_c} hm_2 \sin \theta d\theta + \int_{\tilde{\theta}_c}^{\theta_c} hr \cos \theta \sin \theta d\theta \quad (\text{A.15})$$

$$+ \int_{\theta_c}^{2\pi-\theta_c} hm_1 \sin \theta d\theta + \int_{2\pi-\theta_c}^{2\pi-\tilde{\theta}_c} hr \cos \theta \sin \theta d\theta.$$

With simple computation, we get

$$I_1(r) = hm_2 \sin \tilde{\theta}_c - hm_1 \sin \theta_c + hr\theta_c - hr\tilde{\theta}_c, \quad (\text{A.16})$$

$$I_2(r) = 0, \quad (\text{A.17})$$

where

$$\sin \theta_c = \frac{\sqrt{r^2 - m_1^2}}{r}, \quad \sin \tilde{\theta}_c = \frac{\sqrt{r^2 - m_2^2}}{r}, \quad (\text{A.18})$$

$$\theta_c = \pi + \arctan \frac{\sqrt{r^2 - m_1^2}}{m_1}, \quad \tilde{\theta}_c = \arctan \frac{\sqrt{r^2 - m_2^2}}{m_2}. \quad (\text{A.19})$$

Case 4 ($m_1 < 0 < m_2$ and $|m_1| > |m_2|$). We have $|r \sin \theta| \leq m_2$ and $|r \cos \theta| \leq m_2$ for all $\theta \in [0, 2\pi)$ if $0 < r \leq m_2$. Then, $\varphi(r \cos \theta) = hr \cos \theta$ for every θ . Thus,

$$I_1(r) = \int_0^{2\pi} hr \cos^2 \theta d\theta = \pi hr, \quad (\text{A.20})$$

$$I_2(r) = \int_0^{2\pi} hr \cos \theta \sin \theta d\theta = 0.$$

We fix now $m_2 < r < |m_1|$ and consider $\theta_c \in (0, \pi/2)$ which satisfies $r \cos \theta_c = m_2$. Then, we have

$$I_1(r) = \int_{-\theta_c}^{\theta_c} hm_2 \cos \theta d\theta + \int_{\theta_c}^{2\pi-\theta_c} hr \cos^2 \theta d\theta$$

$$= \pi hr - hr\theta_c + hm_2 \sin \theta_c,$$

$$I_1(r) = \int_{-\theta_c}^{\theta_c} hm_2 \sin \theta d\theta + \int_{\theta_c}^{2\pi-\theta_c} hr \cos \theta \sin \theta d\theta = 0. \quad (\text{A.21})$$

We now fix $r \geq |m_1|$ and consider $\tilde{\theta}_c \in (\pi/2, \pi)$ which satisfies $r \cos \tilde{\theta}_c = m_1$. Then, we have

$$I_1(r) = \int_{-\theta_c}^{\theta_c} hm_2 \cos \theta d\theta + \int_{\theta_c}^{\tilde{\theta}_c} hr \cos^2 \theta d\theta$$

$$+ \int_{\tilde{\theta}_c}^{2\pi-\tilde{\theta}_c} hm_1 \cos \theta d\theta + \int_{2\pi-\tilde{\theta}_c}^{2\pi-\theta_c} hr \cos^2 \theta d\theta,$$

$$I_2(r) = \int_{-\theta_c}^{\theta_c} hm_2 \sin \theta d\theta + \int_{\theta_c}^{\tilde{\theta}_c} hr \cos \theta \sin \theta d\theta$$

$$+ \int_{\tilde{\theta}_c}^{2\pi-\tilde{\theta}_c} hm_1 \sin \theta d\theta + \int_{2\pi-\tilde{\theta}_c}^{2\pi-\theta_c} hr \cos \theta \sin \theta d\theta. \quad (\text{A.22})$$

With simple computation, we get

$$I_1(r) = hm_2 \sin \theta_c - hm_1 \sin \tilde{\theta}_c - hr\theta_c + hr\tilde{\theta}_c, \quad (\text{A.23})$$

$$I_2(r) = 0,$$

where

$$\sin \theta_c = \frac{\sqrt{r^2 - m_2^2}}{r}, \quad \sin \tilde{\theta}_c = \frac{\sqrt{r^2 - m_1^2}}{r},$$

$$\theta_c = \arctan \frac{\sqrt{r^2 - m_2^2}}{m_2}, \quad \tilde{\theta}_c = \pi + \arctan \frac{\sqrt{r^2 - m_1^2}}{m_1}. \quad (\text{A.24})$$

Case 5 ($m_1 < 0 < m_2$ and $|m_1| = |m_2| = m$). We have $|r \sin \theta| \leq m$ and $|r \cos \theta| \leq m$ for all $\theta \in [0, 2\pi)$ if $0 < r \leq m$. Then, $\varphi(r \cos \theta) = hr \cos \theta$ for every θ . Thus,

$$I_1(r) = \int_0^{2\pi} hr \cos^2 \theta d\theta = \pi hr, \quad (\text{A.25})$$

$$I_2(r) = \int_0^{2\pi} hr \cos \theta \sin \theta d\theta = 0.$$

We fix now $r > m$ and consider $\theta_c \in (0, \pi/2)$ which satisfies $r \cos \theta_c = m$. Then, we have

$$I_1(r) = \int_{-\theta_c}^{\theta_c} hm \cos \theta d\theta + \int_{\theta_c}^{\pi-\theta_c} hr \cos^2 \theta d\theta$$

$$- \int_{\pi-\theta_c}^{\pi+\theta_c} hm \cos \theta d\theta + \int_{\pi+\theta_c}^{2\pi-\theta_c} hr \cos^2 \theta d\theta,$$

$$I_1(r) = \int_{-\theta_c}^{\theta_c} hm \sin \theta d\theta + \int_{\theta_c}^{\pi-\theta_c} hr \cos \theta \sin \theta d\theta$$

$$- \int_{\pi-\theta_c}^{\pi+\theta_c} hm \sin \theta d\theta + \int_{\pi+\theta_c}^{2\pi-\theta_c} hr \cos \theta \sin \theta d\theta. \quad (\text{A.26})$$

With simple computation, we get

$$I_1(r) = \pi hr + 2hm \sin \theta_c - 2hr\theta_c, \quad (\text{A.27})$$

$$I_2(r) = 0,$$

where

$$\sin \theta_c = \frac{\sqrt{r^2 - m^2}}{r}, \quad \theta_c = \arctan \frac{\sqrt{r^2 - m^2}}{m}. \quad (\text{A.28})$$

This completes the proof of the lemma.

Acknowledgment

This work is supported in part by the National Natural Science Foundation of China (1097 2082).

References

- [1] J. Llibre and M. A. Teixeira, "On the stable limit cycle of a weight-driven pendulum clock," *European Journal of Physics*, vol. 31, no. 1, pp. 249–254, 2010.
- [2] M. Vidyasagar, *Nonlinear Systems Analysis*, Poelltice-Hall, 2nd edition, 1993.
- [3] M. di Bernardo, C. J. Budd, A. R. Champneys, and P. Kowalczyk, *Piecewise-Smooth Dynamical Systems*, vol. 163 of *Applied Mathematical Sciences*, Springer, London, UK, 2008, Theory and applications.
- [4] V. Carmona, E. Freire, E. Ponce, and F. Torres, "Bifurcation of invariant cones in piecewise linear homogeneous systems," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 15, no. 8, pp. 2469–2484, 2005.
- [5] S. Huan and X.-S. Yang, "Generalized Hopf bifurcation emerged from a corner in general planar piecewise smooth systems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 75, no. 17, pp. 6260–6274, 2012.
- [6] C. Christopher and C. Li, *Limit Cycles in Differential Equations*, Birkhauser, Boston, Mass, USA, 2007.
- [7] S.-M. Huan and X.-S. Yang, "On the number of limit cycles in general planar piecewise linear systems," *Discrete and Continuous Dynamical Systems A*, vol. 32, no. 6, pp. 2147–2164, 2012.
- [8] J. Li, "Hilbert's 16th problem and bifurcations of planar polynomial vector fields," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 13, no. 1, pp. 47–106, 2003.
- [9] J. Llibre and A. Makhlof, "Bifurcation of limit cycles from a two-dimensional center inside \mathbb{R}^4 ," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 3-4, pp. 1387–1392, 2010.
- [10] E. Freire, E. Ponce, and J. Ros, "The focus-center-limit cycle bifurcation in symmetric 3D piecewise linear systems," *SIAM Journal on Applied Mathematics*, vol. 65, no. 6, pp. 1933–1951, 2005.
- [11] V. Carmona, S. Fernández-García, F. Fernández-Sánchez, E. García-Medina, and A. E. Teruel, "Reversible periodic orbits in a class of 3D continuous piecewise linear systems of differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 75, no. 15, pp. 5866–5883, 2012.
- [12] A. Buică and J. Llibre, "Bifurcation of limit cycles from a four-dimensional center in control systems," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 15, no. 8, pp. 2653–2662, 2005.
- [13] C. A. Buzzi, J. Llibre, and J. C. Medrado, "On the limit cycles of a class of piecewise linear differential systems in \mathbb{R}^4 with two zones," *Mathematics and Computers in Simulation*, vol. 82, no. 4, pp. 533–539, 2011.
- [14] C. A. Buzzi, J. Llibre, J. C. Medrado, and J. Torregrosa, "Bifurcation of limit cycles from a centre in \mathbb{R}^4 in resonance 1:N," *Dynamical Systems*, vol. 24, no. 1, pp. 123–137, 2009.
- [15] P. T. Cardin, T. de Carvalho, and J. Llibre, "Bifurcation of limit cycles from an n -dimensional linear center inside a class of piecewise linear differential systems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 75, no. 1, pp. 143–152, 2012.
- [16] J. Llibre, E. Ponce, and J. Ros, "Algebraic determination of limit cycles in a family of three-dimensional piecewise linear differential systems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 17, pp. 6712–6727, 2011.
- [17] A. Buică and J. Llibre, "Averaging methods for finding periodic orbits via Brouwer degree," *Bulletin des Sciences Mathématiques*, vol. 128, no. 1, pp. 7–22, 2004.
- [18] A. Buică, J. Llibre, and O. Makarenkov, "Asymptotic stability of periodic solutions for nonsmooth differential equations with application to the nonsmooth van der Pol oscillator," *SIAM Journal on Mathematical Analysis*, vol. 40, no. 6, pp. 2478–2495, 2009.
- [19] F. E. Browder, "Fixed point theory and nonlinear problems," *American Mathematical Society*, vol. 9, no. 1, pp. 1–39, 1983.

