Brief Communication

Isolation of CYP3A4 Inhibitors from the Black Cohosh (*Cimicifuga racemosa*)

Sachiko Tsukamoto, Maki Aburatani and Tomihisa Ohta

Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

Recent investigation on drug interaction has shown that some foods and herbal medicines increase the oral availability of a variety of CYP3A4 substrates, which is caused by the reduction of CYP3A4 in intestinal epithelium. During the course of our investigation on CYP3A4 interaction, we found that the commercially available dietary supplement made from black cohosh (*Cimicifuga racemosa*) showed CYP3A4 inhibition. Black cohosh has been used for the treatment of menopausal and post-menopausal symptoms as a dietary supplement. Bioassay-guided isolation from the supplement afforded six active principles, which were identified as cycloartanoid triterpene glycosides.

Keywords: black cohosh – *Cimicifuga racemosa* – CYP3A4 – cytochrome P450 – inhibition – triterpene glycoside

Introduction

The rhizomes of Cimicifuga species (Ranunculaceae) are used as Traditional Chinese Medicines. Among Cimicifuga species, C. heracleifolia, C. dahurica and C. foetida are listed in the Chinese Pharmacopoeia and used as an anti-inflammatory, antipyretic and analgesic remedy (1). Cimicifuga racemosa, commonly known as black cohosh, is a herb used among Native Americans to treat a variety of ailments, including diarrhea, sore throat and rheumatism. It is best known for its health benefit in treating menopausal disorders (2). It is used therapeutically to reduce the frequency and intensity of hot flushes (3), and in the USA the extract of C. racemosa is available for sale as a dietary supplement used for the treatment of menopausal and postmenopausal symptoms (4). Cytochrome P450 (CYP) enzymes belong to heme-containing monooxygenases and constitute three families, CYP1, CYP2 and CYP3 (5), and these enzymes are recognized to be responsible for drug metabolism, carcinogenesis and degradation of xenobiotics. Among a family of CYP enzymes, CYP3A4 is the most abundant enzyme in human liver microsomes and intestinal epithelium; ~30% of the total CYP was suggested to be CYP3A4 (6), and >50% of clinically used drugs are oxidized by CYP3A4 (7,8). It is widely known that concomitant oral administration of several foods and herbs affects drug metabolism in humans by inhibiting CYP3A4 activity. Among them, the inhibition by grapefruit juice has been well studied and it is reported that concomitant intake of the juice alters the pharmacokinetics of various drugs, including cyclosporin (9,10), midazolam (11), dihydropyridine-type calcium channel blockers (12) and triazolam (13). In the course of our study on CYP inhibitors from foods, we have reported the isolation and structural elucidation of CYP inhibitors from grapefruit (Citrus paradisii) juice (14-16), white pepper, Piper nigrum (17,18) and the strawberry fruit, Fragaria ananassa (19). Recently, we found that the commercially available black cohosh, C. racemosa Nutt, showed CYP3A4 inhibition. Here we report the isolation, structural identification and CYP inhibitory activity of constituents from black cohosh.

Methods

General Methods

Nuclear magnetic resonance (NMR) spectra were recorded on a JEOL GSX-500 NMR spectrometer in pyridine- d_5 . Mass spectra were measured on a JEOL SX-102 mass spectrometer.

© The Author (2005). Published by Oxford University Press. All rights reserved.

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that: the original authorship is properly and fully attributed; the Journal and Oxford University Press are attributed as the original place of publication with the correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oupjournals.org

For reprints and all correspondence: Tomihisa Ohta, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa, 920-1192 Japan. Tel: +81-76-234-4468; Fax: +81-76-264-6241; E-mail: ohta@p.kanazawa-u.ac.jp

Isolation of Triterpene Glycosides 1–6 from *C. racemosa*

Commercially available black cohosh, C. racemosa Nutt, was generously provided by from Tokiwa Phytochemical Co., Ltd. (Japan). The powder of black cohosh (20 g) was suspended in water and extracted with ethyl acetate (EtOAc). The organic layer was then partitioned between hexane and 90% methanol (MeOH)-H₂O. The polar fraction (6.1 g), which exhibited significant CYP3A4 inhibition, 44% at 5 mg/ml, was subjected to silica gel chromatography with EtOAc/MeOH followed by octadecyl silica gel (ODS) chromatography with MeOH/H2O and then acetonitrile (CH₃CN)/H₂O to afford a potent CYP3A4 inhibitory fraction. The purification by ODS high-performance liquid chromatography (HPLC) with CH₃CN/H₂O gave six triterpene glycosides 1–6. On the basis of ¹H NMR spectra in C_5D_5N and fast atom bombardment mass spectrometry (FABMS), the six triterpene glycosides were identified as cimiracemoside H (1, 2.0 mg, 0.01% yield) (19), 26-deoxyactein (2, 1.1 mg, 0.0055%) (20,21), 23-O-acetylshengmanol 3-O-β-D-xylopyranoside (3, 1.9 mg, 0.0095%) (22), actaeaepoxide 3-O-β-D-xylopyranoside (4, 1.1 mg, 0.0055%) (23), 25-O-acetylcimigenol 3-O-α-L-arabinopyranoside (5, 5.9 mg, 0.029%) (24) and 25-O-acetylcimigenol 3-O-β-D-xylopyranoside (6, 4.2 mg, 0.021%) (25,26) (Figure 1).

Assay of CYP3A4 Inhibition

CYP3A4 activity was based on nifedipine oxidation. Various amounts (0–10 μ M, final concentration) of samples in 1 μ l of dimethylsulfoxide (DMSO) were added to 192 μ l of solution containing 100 mM phosphate buffer (pH 7.4) containing 50 μ M nifedipine (Wako Pure Chemical Industries, Ltd, Osaka, Japan), 5 mM glucose-6-phosphate (Oriental Yeast Co., Ltd, Tokyo, Japan), 0.5 mM β -NADP⁺ (Oriental Yeast Co., Ltd), 0.5 mM MgCl₂ and 4.3 µg/ml glucose-6-phosphate dehydrogenase (Oriental Yeast Co., Ltd) and incubated at 37°C for 5 min. CYP3A4 (Gentest Co., Woburn, MA) was also pre-incubated in 7 µl of the buffer at 37°C for 5 min and added to the sample solution. After the incubation at 37°C for 1 h, the reaction was quenched by the addition of 100 µl of MeOH. After adding 3.7 µg of 6-methoxycarbonyl-5-methyl-7-(2-nitrophenyl)-4,7-dihydrofuro[3,4-b]pyridin-1-(3H)-one in 1 μ l of DMSO as an internal standard, the reaction mixture was extracted with 1 ml of ether, and the ether layer was evaporated. The residue was dissolved in 100 μ l of MeOH, and an aliquot (20 μ l) was analyzed by reverse-phase HPLC (column, TSK-gel ODS-120T, 4.6 mm i.d. \times 150 mm; mobile phase, 64% MeOH-H₂O; flow rate, 1.0 ml/min; detection, UV 254 nm); retention times: 2.9 min for the internal standard, 4.0 min for the nifedipine metabolite (nifedipine pyridine) and 5.5 min for nifedipine. The IC₅₀ value, the concentration required for 50% inhibition of CYP3A4 activity, was calculated from the data of duplicate measurements.

Results

Bioassay-guided Isolation of Triterpene Glycosides 1–6 from *C. racemosa*

The CYP inhibition was tested using human CYP3A4 as shown in Methods. Commercially available supplement of *C. racemosa* was extracted with EtOAc, and the organic layer was then partitioned between hexane and 90% MeOH-H₂O. The polar fraction exhibited significant CYP inhibition. Forty-four percent inhibition at 5 mg/ml was as potent as that by ketoconazole, 58% at 5 µg/ml. Bioassay-guided fractionation afforded six triterpene glycosides **1–6**. On the basis of their ¹H NMR spectra in C₅D₅N and FABMS, they were identified as cimiracemoside H (**1**, 0.01% yield) (20), 26-deoxyactein (**2**, 0.0055%) (21,22),



Figure 1. Structures of CYP inhibitors isolated from C. racemosa

Table 1. CYP3A4 inhibition of C. racemosa metabolites

Compound	IC ₅₀ (mM)
1	0.48
2	0.10
3	0.11
4	7.78
5	1.4
6	0.83
Ketoconazole ^a	0.00011

^aA typical CYP3A4 inhibitor.

23-*O*-acetylshengmanol 3-*O*-β-D-xylopyranoside (**3**, 0.095%) (23), actaeaepoxide 3-*O*-β-D-xylopyranoside (**4**, 0.0055%) (24), 25-*O*-acetylcimigenol 3-*O*-α-L-arabinopyranoside (**5**, 0.0029%) (25) and 25-*O*-acetylcimigenol 3-*O*-β-D-xylopyranoside (**6**, 0.0021%) (26,27). The CYP inhibitory activities of **1–6** are shown in Table 1, which indicates moderate IC₅₀ values.

Discussion

Bailey *et al.* demonstrated that the blood concentration of antihypertensive drugs such as nifedipine was kept at a high level with grapefruit juice co-administered to mask the taste of alcohol. Since that time there has been a great deal of interest in food–drug interactions because of the clinical risks associated with changes in the bioavailability or metabolic rate of clinically administered drugs. In spite of the increasing use of herbal remedies and nutraceuticals, information on the drug interaction that occurs with them including foodstuffs, is still insufficient to understand the clinical risks.

We have studied drug interactions with foodstuffs such as grapefruit juice and pepper, isolating several potent CYP3A4 inhibitors in each material. The fact that a foodstuff contains more than one inhibitor suggests that the clinical effect of drug interaction with herbs and foodstuffs could be better understood by studying the mixture of inhibitors and/or an extract of these.

During our study on CYP3A4 inhibitors contained in food, we found that the supplement of C. racemosa exhibited potent inhibition. The polar fraction from the extract showed 44% inhibition at 5 mg/ml, which was as potent as the inhibition produced by ketoconazole, 58% at 5 µg/ml. We clarified the main constituents, cognate triterpene glycosides, as the CYP3A4 inhibitory principles. To date, highly potent CYP3A4 inhibitors in food have been reported, e.g. paradisins A and B (IC₅₀, 0.07 and 0.07 μ M) from grapefruit juice (15), dipiperamide A (IC₅₀, 0.18 μ M) from white pepper (17) and gomisin C (IC₅₀, 0.254 μ M) from Schisandra fruit (29). Although the IC₅₀ value of each isolate was moderate, the high content of a series of cognates in the supplement could result in the exhibition of significant CYP inhibition. Care should be taken to avoid concomitant intake of black cohosh with any kind of medication because of the possibility of increasing the bioavailable concentration of drugs in the blood by the downregulation suppression of CYP3A4. Medical practitioners therefore should pay more attention in order to avoid any trouble with diet containing black cohosh.

The IC₅₀, 0.027 mg/ml, of the extract of black cohosh was very low in spite of the weak activities of the isolated compounds. The IC₅₀ value indicates that 40 mg of the black cohosh extract could be estimated to inhibit the metabolism of roughly half a dose of nifedipine for 1 day.

Evaluation of the IC_{50} of both constituents and the extract may be useful for clinical study of drug interactions of herbs and foodstuffs predicted to be a clinical risk, especially for the study of the mechanism of action and the pharmacokinetics.

Acknowledgments

The authors are grateful to Tokiwa Phytochemical Co., Ltd (Japan) for the generous gift of black cohosh.

References

- New Medicinal College of Jiangsu (ed.). Dictionary of Chinese Materia Medica, Shanghai Scientific and Technological Press, Shanghai, 1977: p. 451.
- Lieberman S, A review of the effectiveness of *Cimicifuga racemosa* (black cohosh) for the symptoms of menopause. J Women's Health 1998;7:525–29.
- McKenna D, Jones K, Humphrey S, Hughes K. Black cohosh: efficacy, safety, and use in clinical and preclinical applications. *Altern Ther* 2001;7:93–100.
- Borrelli F, Izzo AA, Ernst E. Pharmacological effects of *Cimicifuga* racemosa. Life Sci 2003;73:1215–29.
- Nelson DR, Koymans L, Kamataki T, Stegeman JJ, Feyereisen R, Waxman DJ, Waterman MR, Gotoh O, Coon MJ, Estabrook RW, Gunsalus IC, Nebert DW. P450 superfamily: update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics* 1996;6:1–42.
- Shimada T, Yamazaki H, Mimura M, Inui Y, Guengerish FP. Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* 1994;270:414–23.
- Guengerish FP. Role of cytochrome P450 enzymes in drug–drug interactions. Adv Pharmacol 1997;43:7–35.
- Rendic S, DiCarlo FJ. Human cytochrom P450 enzymes: a status report summarizing their reactions, substrates, inducers, and inhibitors. *Drug Metab. Rev.* 1997;29:413–580.
- Ducharme MP, Warbasse LH, Edwards DJ. Disposition of intravenous and oral cyclosporine after administration with grapefruit juice. *Clin Pharmacol Ther* 1995;57:485–91.
- Yee GC, Stanley DL, Pessa LJ, Dalla CT, Beltz SE, Ruiz J. Effect of grapefruit juice on blood cyclosporin concentration. *Lancet* 1995;345: 955–56.
- Kupferschmidt HH, Ha HR, Ziegler WH, Meier PJ, Krahenbuhl S. Interaction between grapefruit juice and midazolam in humans. *Clin Pharmacol Ther* 1995;58:20–8.
- Bailey DG, Arnold JM, Spence JD. Grapefruit juice and drugs. How significant is the interaction? *Clin Pharmacokinet* 1994;26: 91–8.
- Hukkinen SK, Varhe A, Olkkola KT, Neuvonen PJ. Plasma concentrations of triazolam are increased by concomitant ingestion of grapefruit juice. *Clin Pharmacol Ther* 1995;58:127–31.
- Ohta T, Nagahashi M, Hosoi S, Tsukamoto S. Dihydroxybergamottin caproate as a potent and stable CYP3A4 inhibitor. *Bioorg Med Chem* 2002;10:969–73.
- Ohta T, Maruyama T, Nagahashi K, Miyamoto Y, Hosoi S, Kiuchi F, Yamazoe Y, Tsukamoto S. Paradisin C: a new CYP3A4 inhibitor from grapefruit Juice. *Tetrahedron* 2002;58:6631–35.
- Ohta T, Miyamoto Y, Maruyama T, Kiuchi F, Tsukamoto S. Localization and contents of paradisins, the most potent CYP3A4 inhibitors, in a

grapefruit Citrus paradisii and grapefruit Juice. Nat Med 2002;56: 264-7.

- Tsukamoto S, Cha B-C, Ohta T. Dipiperamides A, B, and C: bisalkaloids from the white pepper *Piper nigrum* inhibiting CYP3A4 activity. *Tetrahedron* 2002;58:1667–71.
- Tsukamoto S, Tomise K, Miyakawa K, Cha B-C, Abe T, Hamada Y, Hirota H, Ohta T. CYP3A4 inhibitory activity of new bisalkaloids dipiperamides D and E and cognate alkaloids from the white pepper. *Bioorg Med Chem* 2002;10:2981–85.
- Tsukamoto S, Tomise K, Aburatani, M, Onuki H, Hirota H, Ishiharajima E, Ohta T. Isolation of cytochrome P450 inhibitors from strawberry fruit, *Fragaria ananassa* Duch. cv. Tochiotome. J Nat Prod 2004;67(11): 1839–41.
- Shao Y, Harris A, Wang M, Zhang H, Cordell GA, Bowman M, Lemmo E. Triterpene glucosides form *Cimicifuga racemosa*. J Nat Prod 2000;63: 905–10.
- Koeda M, Aoki Y, Sakurai N, Nagai M. Studies on the Chinese crude drug 'Shoma.' IX. Three novel cycloanostanol xylosides, cimicifugosides H-1, H-2 and H-5, form Cimicifuga Rhizome. *Chem Pharm Bull* 1995; 43:771–6.
- Chen S-N, Li W, Fabricant DS, Santarsiero BD, Mesecar A, Fitzloff JF, Fong HHS, Farnsworth NR. Isolation, structure elucidation, and absolute configuration of 26-deoxyactein from *Cimicifuga racemosa* and clarification of nomenclature associated with 27-deoxyactein. J Nat Prod 2002;65:601–5.

- 23. Kusano A, Shibano M, Kitagawa S, Kusano G, Nozoe S, Fushiya S. Studies on the constituents of *Cimicifuga* species. XV. Two new diglycosides from the aerial parts of *Cimicifuga simplex* WORMSK. *Chem. Pharm. Bull.* 1994;42:1940–43.
- Wende K, Mugge C, Thurow K, Schopke T, Lindequist U, Actaeaepoxide 3-O-β-D-xylopyranoside, a new cycloartane glycoside from the rhizomes of Actaea racemosa (*Cimicifuga racemosa*). J Nat Prod 2001;64: 986–9.
- Ye W, Zhang J, Che C-T, Ye T, Zhao S, New cycloartane glycosides from Cimicifuga dahurica. Planta Med 1999;65:770–1.
- Kadota S, Li JX, Tanaka K, Namba T. Constituents of Cimicifugae rhizoma II. Isolation and Strucures of new cycloartenol triterpenoids and related compounds from *Cimicifuga foetida L. Tetrahedron* 1995;51:1143–66.
- Takemoto T, Kusano G, Kawahara M. Studies on the constituents of *Cimicifuga* spp. VI. Structures of 25-O-acetylcimigenoside and 25-O-methylcimigenoside. *Yakugaku Zasshi* 1970;90:64–7.
- 'The Pharmacopoeia of Chinese People's Republic' vol. I, The People's Health Publishing House & The Chemical Industry Publishing House, Beijing, 1990, p. 59.
- Iwata H, Tezuka Y, Kadota S, Hiratsuka A, Watabe T. Identification and characterization of potent CYP3A4 inhibitors in Schisandra fruit extract. *Drug Metab Dispos* 2004;32:1351–8.

Received September 30, 2004; revised January 18, 2005; accepted March 28, 2005



The Scientific **World Journal**



Gastroenterology Research and Practice





Journal of Diabetes Research



Disease Markers



Immunology Research





Submit your manuscripts at http://www.hindawi.com





BioMed **Research International**



Journal of Ophthalmology

Computational and Mathematical Methods in Medicine





Behavioural Neurology









Research and Treatment





Oxidative Medicine and Cellular Longevity



Stem Cells International

