

Supplement 1. Differences in the body characteristics and serum lipid parameters among the pattern identification groups of Korean stroke patients

Characteristics	QD	D&P	YD	F&H	p
<i>body characteristics</i>					
weight (kg)	57.16 ±9.3*	63.62 ±10.46	56.84 ±10.07	64.42 ±10.33	<0.001
BMI (kg/m ²)	22.87 ±2.90	24.73 ±3.22	22.65 ±3.15	24.08 ±2.92	<0.001
waist circumference (cm)	85.06 ±8.60	89.0 ±9.46	83.05 ±9.25	88.04 ±8.90	<0.001
WHR	0.927 ±0.100	0.943 ±0.077	0.936 ±0.142	0.948 ±0.092	<0.001
<i>serum lipid parameters</i>					
total cholesterol (mg/dL)	185.29 ±49.18	190.06 ±46.82	186.27 ±40.65	179.46 ±44.930	0.149
triglyceride (mg/dL)	156.06 ±148.78	164.33 ±115.24	145.47 ±95.23	157.99 ±95.43	0.032
HDL-cholesterol					
(mg/dL)	44.85 ±13.31	42.40 ±11.92	44.09 ±13.60	42.37 ±11.90	0.303
Atherogenic index	0.478 ±0.298	0.527 ±0.287	0.472 ±0.304	0.525 ±0.277	0.056

* indicates mean ± standard deviation

The p values among the PI groups were calculated by a general linear model adjusted for sex, age, smoking and drinking