

## Erratum Erratum to "Chemistry and Biology of Essential Oils of Genus Boswellia"

## Hidayat Hussain,<sup>1,2</sup> Ahmed Al-Harrasi,<sup>1</sup> Ahmed Al-Rawahi,<sup>1</sup> and Javid Hussain<sup>1</sup>

<sup>1</sup> Department of Biological Sciences and Chemistry, College of Arts and Sciences, University of Nizwa, Birkat Al-Mouz, 616 Nizwa, Oman

<sup>2</sup> Department of Chemistry, University of Paderborn, Warburger Straße 100, 33098 Paderborn, Germany

Correspondence should be addressed to Ahmed Al-Harrasi; aharrasi@unizwa.edu.om

Received 29 September 2013; Accepted 8 October 2013; Published 12 January 2014

Copyright © 2014 Hidayat Hussain et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the original paper the following two errors were present. Correspondence should be addressed as shown above. In the original paper, the table describing major compounds in the oils of *Boswellia* species including studies on page 7 (Table 2) includes some errors. The value "29.3" should be "29.7" and "61.36" should be "61.4." The corrected table is presented as follows.

Boswellia species	Method of obtaining resin	Predominant compound(s)	Percentage (%)	Literature
B. serrata	Obtained from Willy Benecke GmbH (Hamburg, Germany)	Myrcene	38	[16]
B. serrata	NA	α-Thujene	22.7-47.4	[22]
B. serrata	NA	α-Thujene	29.7	[24]
B. serrata	NA	α-Thujene	61.4	[23]
B. carteri	Purchased from the local market of herbs and spices in Egypt	Duva-3,9,13-triene-1a- ol-5,8-oxide-1-acetate	21.4	[20]
B. sacra	Botanically certified oleogum resin	$E$ - $\beta$ -Ocimene	32.3	[25]
B. carteri/sacra	NM	Octanol acetate	45.2	[26]
B. carteri	Authentic sample from Ethiopia certified for its authenticity from the Agricultural Department of the Ethiopian government	Octyl acetate	39.3	[16]
B. rivae	NA	Limonene	28.0	[24]
B. rivae	Authentic sample from Ethiopia	α-Pinene	16.7	[16]
B. rivae	NA	α-Pinene	13.3	[24]
B. rivae	NA	Octanol	17.8	[24]
B. neglecta	NA	α-Pinene	16.7	[27]
B. neglecta	Authentic sample from Ethiopia	α-Pinene	21.3	[16]
B. papyrifera	NA	Octyl acetate	63.5	[24]
B. papyrifera	NA	Octyl acetate	56.0	[36]
B. pirottae	NA	trans-Verbenol	15.5	[27]
B. pirottae	NA	Terpinen-4-ol	14.6	[27]
B. frereana	Willy Benecke GmbH (Hamburg, Germany)	α-Pinene	38.0	[16]

TABLE 2: Percentages of major compounds in the essential oils of reported Boswellia species.

NA: not available.



**The Scientific** World Journal



Gastroenterology Research and Practice





Journal of Diabetes Research



**Disease Markers** 



Immunology Research









BioMed **Research International** 





Computational and Mathematical Methods in Medicine





Behavioural Neurology



Complementary and Alternative Medicine











Oxidative Medicine and Cellular Longevity