

Research Article

Short-Term Effects of Kinesio Taping and Cross Taping Application in the Treatment of Latent Upper Trapezius Trigger Points: A Prospective, Single-Blind, Randomized, Sham-Controlled Trial

Tomasz Halski,¹ Kuba Ptaszkowski,² Lucyna Słupska,³ Małgorzata Paprocka-Borowicz,³ Robert Dymarek,⁴ Jakub Taradaj,⁵ Gabriela Bidzińska,¹ Daniel Marczyński,¹ Aleksandra Cynarska,¹ and Joanna Rosińczuk⁴

¹Department of Physiotherapy, Opole Medical School, Katowicka 68, 45-060 Opole, Poland

²Department of Obstetrics, Faculty of Health Science, Wrocław Medical University, K. Bartla 5, 51-618 Wrocław, Poland

³Department of Clinical Biomechanics and Physiotherapy in Motor System Disorders, Faculty of Health Science, Wrocław Medical University, Grunwaldzka 2, 50-355 Wrocław, Poland

⁴Department of Nervous System Diseases, Faculty of Health Science, Wrocław Medical University, K. Bartla 5, 51-618 Wrocław, Poland

⁵Department of Physiotherapy Basics, Academy of Physical Education in Katowice, Mikolowska Street 72, Building B, 40-065 Katowice, Poland

Correspondence should be addressed to Kuba Ptaszkowski; kptaszowski@gmail.com

Received 19 May 2015; Revised 14 August 2015; Accepted 8 September 2015

Academic Editor: Ilaria Lampronti

Copyright © 2015 Tomasz Halski et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Kinesio taping (KT) may be a new treatment in patients with myofascial trigger points (MTrPs). A new method available for taping practitioners is cross taping (CT). The main objective was to determine how CT, KT, and medical adhesive tape (sham group) affect the subjective assessment of resting bioelectrical activity and pain of the upper trapezius muscle (UT) in patients with MTrPs. 105 volunteers were recruited to participate. The primary outcome was resting bioelectrical activity of UT muscle as assessed by surface electromyography (sEMG) in each group and pain intensity on a visual analog scale (VAS). Assessments were collected before and after intervention and after the 24-hours follow-up. No significant differences were observed in bioelectrical activity of UT between pre-, post-, and follow-up results. In three groups patients had significantly lower pain VAS score after the intervention (CT— $p < 0.001$, KT— $p < 0.001$, and sham— $p < 0.01$). The Kruskal-Wallis ANOVA showed no significant differences in almost all measurements between groups. The application of all three types of tapes does not influence the resting bioelectrical activity of UT muscle and may not lead to a reduction in muscle tone in the case of MTrPs.

1. Introduction

Myofascial pain syndrome (MPS) is one of the most frequent causes of musculoskeletal problems [1–3]. Myofascial pain is defined as pain which comes from myofascial trigger points (MTrPs) in muscle that are considered as hyperirritable spot located within a taut band of skeletal muscle. MTrPs are characterized by local tenderness without referred pain and local tenderness with referred pain and also by muscle

dysfunction (weakness, fatigue, stiffness, and poor blood flow), restricted range of motion (ROM), change of motor pattern, poor posture, and limited physical, professional, and social activity [3–9]. Two types of MTrPs are distinguished: latent and active. Latent MTrPs cause local and referred pain with palpation and active MTrPs cause pain at rest and on palpation (“spontaneous pain”) [1, 5–10].

Publications about the MTrPs diagnosis indicate that the upper trapezius (UT) is the muscle in which MTrPs occur

very often [7, 9–14]. MTrPs within the UT may cause neck pain, chronic upper-quarter pain, headache, migraine, or shoulder pain [12–15].

There are many ways to treat myofascial pain. The treatment algorithms can contain noninvasive methods like educational programs, behavioral cognitive therapy, medication, and physical therapy or physiotherapy (spray and stretch, general exercises, myofascial release, massage, Jacobson's muscle relaxation, autogenic training, manual therapy, neuromuscular techniques, electrotherapy—ultrasound, interferential therapy, transcutaneous electrical nerve stimulation, pulsed shortwave therapy, and laser therapy) [14–23]. The invasive treatments for MTrPs include injections with dry needling, local anesthetics, corticosteroids, and botulin toxin [21, 22, 24–27].

More and more publications suggest that kinesio taping (KT) may be a new treatment option and indicate the possibility of the use of KT in patients with musculoskeletal problems [28–32], including MTrPs [32–36]. KT is a therapeutic taping technique developed by Dr. Kenzo Kase (Japan, 1979). This technique uses an elastic tape that is thin and more elastic than conventional bandages. The tape can be stretched to 140% of its original length and applied to the skin [28, 31, 34]. KT is used as an alternative to athletic taping to support the fascia, muscles, and joints. In addition, in the literature it can be found that KT can increase the ROM, reduce swelling, inflammation, and bruising, enhance blood circulation, enhance strength and muscle tone, or be used in muscle spasms and cramping prevention and to speed recovery of overused muscles [32–34, 37–42]. Most of the research is related to the use of KT in relieving pain, specifically reducing pain and disability in patients with chronic, nonspecific back pain [29]. It seems that KT can also be used to combat pain in patients with MTrPs [38].

A new form of tapes available for taping practitioners is cross tapes. Cross tapes are small, polyester tapes with an adhesive acrylic coating. The water-resistant cross tapes are free of medication and active ingredients and mostly can be used for local points of pain, trigger or acupuncture points, tense muscles, painful joints, headaches, or painful scars. The tapes are applied directly over points of pain. Depending on the wearing conditions, they can adhere to the skin for a period up to several days. The tapes are available in different sizes (M, L, and XL) and, in contrast to KT, they are not elastic and are unable to follow the skin when it is stretched. Most information about cross tapes is on manufacturer's website; however, to our knowledge, there are no good quality research studies that evaluate the effectiveness of using cross taping (CT) [43]. Therefore, the influence of CT on body structures and functions needs to be confirmed by objective research.

The lack of the strongest types of research (meta-analysis, systematic reviews, or randomized controlled trials) on MTrPs treatment using CT and the lack of clear methodology of cross tapes application prompted the authors to perform an evaluation of effectiveness of CT. The main objective was to determine how CT, KT, and medical adhesive tape (sham group) affect the subjective assessment of pain and resting bioelectrical activity of the UT muscle in patients with MTrPs. A secondary objective was to evaluate cervical ROM before

and after the intervention. Additionally, a comparison of the results was conducted between CT, KT, and sham group.

2. Methods

2.1. Design. We designed a prospective, single-blind, randomized, sham-controlled study in which the CT and KT effects were compared.

2.2. Approval. The study was approved by the Bioethics Committee of Opole Medical School (Poland: no. KB/01/08/2013) and all subjects provided written informed consent.

2.3. Setting, Participants, and Random Allocation of Patients. 105 volunteers were recruited from Opole Medical School population to participate. The inclusion criteria were being between 18 and 26 years, being asymptomatic, latent MTrPs in the upper part of the trapezius muscle (pain during examination), the absence of skin allergies, and the consent to physical examination and taping application. The exclusion criteria included any history of upper limb, back or neck severe injury in the last 12 months, surgical intervention, upper limb fractures, neurological diseases or musculoskeletal disorders, pharmacological treatment at present, infection, open wound, rash, decreased blood circulation in the treatment area, a pacemaker, or epilepsy.

Randomization was conducted a priori using the website <https://www.random.org/>. Participants were randomized into three study groups: cross taping group (CT group), kinesio taping group (KT group), and sham group.

2.4. Outcomes and Assessment Procedures. Assessments were collected before and after intervention and after the 24-hour follow-up. The primary outcome was resting bioelectrical activity of UT muscle as assessed by surface electromyography (sEMG) in each group and pain intensity on a visual analog scale (VAS). The secondary outcome was cervical mobility evaluated by tape measurement.

In all patients, a proper tape was applied on the MTrPs of the upper part of the trapezius muscle for three days (72 hours). The evaluation of the MTrPs was conducted while the patient was in relaxed prone position on an examination table, and the upper body was exposed. An experienced physiotherapist was assessing the trapezius muscle bilaterally by palpation with thumb with the same pressure. Four diagnostic criteria for the MTrPs were assumed: a hypersensitive spot in a taut band, pain on spot palpation, restricted ROM, and a referred pain distant to the spot.

MTrPs application with a cross tape (Kumbrink Cross-Tape, biviax GmbH & Co. KG, 1.5 cm × 2.5 cm, German) was used in CT group (Figure 1). This polyester tape was placed on the upper part of the trapezius, on MTrP spot. In KT group, the Kinesio Tape (Nitto Denko K-Active Tape, 5 m/2.5 cm, Japan) was placed on the same muscle using four "I" strips' application in star shape to create more space directly above an area of pain (space correction) (Figure 2). Each strip was stretched to 50% of available tension [28]. In sham group an adhesive, nonelastic medical tape with no therapeutic influence (Polovis Plus, 5 m/2.5 cm, Poland) was used over



FIGURE 1: A CT application on MTrPs on the upper part of the trapezius.



FIGURE 2: A KT application on MTrPs on the upper part of the trapezius.

the same muscle (four strips in star shape without tension) (Figure 3). Before applications, the skin was shaved, cleaned with alcohol, and dried. All applications were performed by the same researcher (certified KT physiotherapist).

The electromyographic signal was registered by a dual-channel sEMG NeuroTrac ETS device integrated with computer software for digital analysis and report creation (Verity Medical Ltd., United Kingdom). This device is characterized by an amplitude range of 0.2–2000 μV RMS continuous in the frequency band of 2–100 Hz and pulse width from 50 to 450 μs for recording signals generated by muscles. Device sensitivity is established at a level 0.1 μV (4% accuracy; readings \pm 0.3 mV at 200 Hz), with selectable bandpass filter (3 db bandwidth) and 50 Hz notch filter (33 db; 0.1% accuracy). The analogue signal recorded by the sEMG electrodes was amplified, filtered, and subsequently transformed into a digital signal. Such signal facilitated statistical analysis of acquired results and allowed for data representation in a



FIGURE 3: A sham application on MTrPs on the upper part of the trapezius.

graphical form. Mean values of muscle resting bioelectrical activity were given according to root mean square algorithm (RMS). The monopolar, self-adhesive reference electrode was placed on the seventh cervical vertebra.

The electrodes were attached parallel to the muscle fibre orientation over the following muscles: at the UT muscle halfway between the seventh cervical vertebra and the acromion [<http://www.seniam.org/>].

Pain was recorded by the participant using a 10 cm VAS, where 0 represented no pain and 10 represented unbearable pain. This scale was used to assess the pain during palpation assessment. This assessment was conducted by the same experienced researcher (certified physiotherapist). And during this examination all patients had to evaluate the sensations of pain.

Cervical ROM was measured with a tape measure in centimeters (cm). The range of flexion movement was assessed as the distance from sternal notch to the chin while patients were instructed to bend the head forward; the extension movement was the distance from sternal notch to chin while patients were asked to bend the head backward. Lateral flexion movement was the distance from acromion process to the lowest point of the ear lobe when patients were told to tilt the head to the side opposite the involved UT muscle [44–48].

2.5. Statistical Analysis. Data were analysed with the Statistica version 10 for Windows (StatSoft Inc., USA), and the results are presented as the mean \pm SD. In order to analyze the changes in bioelectrical activity, VAS score, and ROM between pre-, post-, and follow-up results, the analysis of variance (ANOVA) of Friedman and the Wilcoxon matched-pairs test were used to examine the changes within each group. An independent Kruskal-Wallis ANOVA test and nonparametric multiple comparison test were used for comparison among the three groups. A value of $p < 0.05$ was considered statistically significant.




FIGURE 4: Flow diagram.

3. Results

A total of 105 people were recruited for this study. 32 were excluded: 12 because they had a previous history of upper limb or back or neck severe injury, 10 because they had not consented to participation in the study, 3 because they had musculoskeletal disorders of the neck or shoulder, 5 because they were allergic to tape, and 2 because they took antidepressants (Figure 4).

The remaining 73 participants were randomized to three groups (CT, KT, and Sham) and all evaluated before, after, and 24 hours after the intervention (follow-up). The demographic characteristics of the participants are presented in Table 1.

In each group, no significant differences were observed in bioelectrical activity between pre-, post-, and follow-up results. There was a significant interaction between results in each group for the VAS score. In fact, in three groups patients had significantly lower pain VAS score after the intervention (main effect in each group: CT— $p < 0.001$, KT— $p < 0.001$, and Sham— $p < 0.01$) (Table 2). Significant differences for the remaining secondary outcomes were detected.

The interaction between pre-, post-, and follow-up results in each group for the range of flexion movement was significant. The range of flexion movement significantly improved in patients after the intervention and after 24 hours (main effect in each group: CT— $p < 0.0001$, KT— $p < 0.0001$, and Sham— $p < 0.001$). Only in KT group, there was a significant interaction between results for the range of lateral flexion movement ($p < 0.01$) (Table 2).

The Kruskal-Wallis ANOVA showed no significant differences in almost all measurements between groups (Table 3). In the KT group, greater decrease of VAS score was found comparatively to sham group ($p = 0.0018$).

4. Discussion

This study was conducted to identify CT and KT effect on bioelectrical activity, the level of myofascial pain, and the cervical ROM when taping was applied to subjects with latent MTrPs in their upper part of trapezius muscle. There is scarce number of published studies which show the influence of CT on muscle functions, and there are only a few publications which show the effect of KT in patients with latent MTrPs [8, 11, 36, 38]. The study failed to identify significant effect of KT and CT on muscle bioelectrical activity. Gómez-Soriano et al. [37] found a short-term increase of gastrocnemius muscle activity after KT but it was not maintained up to 24 h. Additionally, the study was performed in healthy subjects, what could diminish the therapeutic effect of taping. Nonetheless, research findings in the literature, concerning the influence of KT on muscle bioelectrical activity, specifically in patients with musculoskeletal disorders [49, 50] report about decreasing effect of Kinesio Tapes on muscle electrical activity. Takasaki et al. [41] investigated whether the activity of the UT muscle varied with the application of tensioned and nontensioned taping. In both cases, applications reduced the UT activity comparing to control group. The inhibitory effect

TABLE 1: Groups characteristic.

	CT group	KT group	Sham group	<i>p</i> value
Number of patients	<i>n</i> = 24	<i>n</i> = 25	<i>n</i> = 24	—
Age [year]				
Range	19.0–23.0	19.0–24.0	19.0–22.0	
Mean	20.2	20.6	19.9	<i>p</i> = 0.2631*
SD	1.1	1.5	0.8	
Weight [kg]				
Range	43.0–83.0	51.0–87.0	50.0–85.0	
Mean	62.4	66.4	60.6	<i>p</i> = 0.1320*
SD	10.0	11.6	8.77	
Height [m]				
Range	1.51–1.78	1.57–1.82	1.60–1.75	
Mean	1.68	1.70	1.67	<i>p</i> = 0.2016*
SD	0.07	0.06	0.04	
BMI [kg/m ²]				
Range	17.9–28.0	18.6–29.4	18.1–31.2	
Mean	22.0	22.9	21.9	<i>p</i> = 0.4997*
SD	2.47	3.1	3.2	
Sex				
Female	23	21	24	
Male	1	4	0	<i>p</i> = 0.0701**
Dominant lower limb				
Left	3	1	5	
Right	21	24	19	<i>p</i> = 0.2008**

*Kruskal-Wallis test; **chi² test.

TABLE 2: Comparison between preintervention, postintervention, and follow-up results in each group.

Outcomes	Group	Measurement			<i>p</i> value (main effect of Friedman ANOVA)
		Preintervention	Postintervention	Follow-up	
Resting bioelectrical activity (μ V) Mean \pm SD	CT	6.8 \pm 4.8	4.5 \pm 2.7	4.2 \pm 2.6	<i>p</i> = 0.1152
	KT	6.2 \pm 4.3	5.0 \pm 3.3	4.6 \pm 2.8	<i>p</i> = 0.3260
	Sham	7.4 \pm 7.6	4.7 \pm 2.5	3.9 \pm 1.9	<i>p</i> = 0.0542
Visual analogue scale (VAS) Mean \pm SD	CT	7.2 \pm 1.2**	5.8 \pm 1.6*	5.1 \pm 1.8***	<i>p</i> = 0.0001
	KT	6.8 \pm 1.8**	4.0 \pm 2.0*	5.2 \pm 2.4***	<i>p</i> = 0.0001
	Sham	6.4 \pm 1.6**	5.7 \pm 2.0	4.9 \pm 2.2***	<i>p</i> = 0.0011
The range of flexion movement (cm) Mean \pm SD	CT	2.6 \pm 0.7**	2.0 \pm 0.5*	1.4 \pm 0.9***	<i>p</i> = 0.0000
	KT	3.1 \pm 1.0**	2.2 \pm 1.0*	1.7 \pm 1.1***	<i>p</i> = 0.0000
	Sham	3.1 \pm 0.8**	2.7 \pm 0.9*	2.4 \pm 0.7***	<i>p</i> = 0.0004
The range of extension movement (cm) Mean \pm SD	CT	8.0 \pm 1.2	8.0 \pm 1.6	8.0 \pm 1.1	<i>p</i> = 0.8140
	KT	7.8 \pm 1.6	7.9 \pm 1.5	8.6 \pm 1.6	<i>p</i> = 0.3068
	Sham	8.1 \pm 1.3	7.6 \pm 1.0	8.6 \pm 1.2	<i>p</i> = 0.3068
The range of left lateral flexion movement (cm) Mean \pm SD	CT	5.3 \pm 1.3	5.2 \pm 1.0	5.9 \pm 0.9	<i>p</i> = 0.0056
	KT	5.4 \pm 1.2**	6.0 \pm 0.9*	6.0 \pm 0.8	<i>p</i> = 0.0010
	Sham	5.3 \pm 1.0	5.5 \pm 0.7	5.9 \pm 0.7	<i>p</i> = 0.0057
The range of right lateral flexion movement (cm) Mean \pm SD	CT	5.2 \pm 1.1	5.4 \pm 1.2	5.7 \pm 1.0	<i>p</i> = 0.0705
	KT	5.3 \pm 1.1	5.8 \pm 1.0	5.8 \pm 0.9	<i>p</i> = 0.0314
	Sham	5.4 \pm 0.9	5.4 \pm 0.7	6.0 \pm 0.6	<i>p</i> = 0.0051

Post hoc analysis:

*Statistically significant comparison between pre- and postresults (*p* < 0.05).

**Statistically significant comparison between pre- and follow-up results (*p* < 0.05).

***Statistically significant comparison between post- and follow-up results (*p* < 0.05).

TABLE 3: Comparison of the results between CT, KT, and sham groups.

Outcomes	Results*	Differences between groups (post hoc analysis)			<i>p</i> value (main effect of Kruskal-Wallis ANOVA)
		CT/KT	CT/sham	KT/sham	
Resting bioelectrical activity (μ V) Mean \pm SD	I	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.5892$
	II	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.7014$
	III	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.3939$
Visual analogue scale (VAS) Mean \pm SD	I	$p > 0.05$	$p > 0.05$	$p = 0.0018$	$p = 0.0017$
	II	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.6259$
	III	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.1897$
The range of flexion movement (cm) Mean \pm SD	I	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.3859$
	II	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.1440$
	III	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.4227$
The range of extension movement (cm) Mean \pm SD	I	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.4890$
	II	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.2646$
	III	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.0529$
The range of left lateral flexion movement (cm) Mean \pm SD	I	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.1427$
	II	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.9505$
	III	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.0541$
The range of right lateral flexion movement (cm) Mean \pm SD	I	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.2622$
	II	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.8262$
	III	$p > 0.05$	$p > 0.05$	$p > 0.05$	$p = 0.0862$

*Results:

Result I—post- minus pre-intervention results.

Result II—follow-up minus pre-intervention results.

Result III—follow-up minus post-intervention results.

of taping is supported by the findings of Huang et al. [50], who likewise demonstrated that taping over the UT decreased its activity. Lowered electromyographic activity was also noticed by Paoloni et al. [49]. In their research on chronic low back pain patients, KT rapidly reduced abnormal EMG activity of lumbar paraspinal muscles.

According to the study results, only after KT application on UT muscle, the myofascial pain was relieved. However, there was no improvement in range of motion in any of the groups. Bae et al. [8] evaluated the changes in the myofascial pain and ROM of temporomandibular joint when KT was applied to patients with latent MTrPs of the sternocleidomastoid muscle. In this study, they found significant decrease in the VAS score and increase of ROM of temporomandibular joint. Precise mechanism which explains the effect of KT on musculoskeletal pain is not yet fully understood. There are a number of hypotheses indicating a probable analgesic action of KT. The gate control theory seems to be the most fundamental approach, in which the cutaneous stretch stimulation, activated by KT, can interfere nociceptive stimuli reaching the central nervous system and inhibit the pain [11, 49, 51, 52]. Although this theory does not imply to CT, those tapes are unable to provide the stretch to the tissue. The positive effect of KT application on myofascial pain was also observed by García-Muro et al. [38]. In a case report of a patient with shoulder dysfunction caused by the MTrPs, the author showed an objective improvement in

the VAS score and the algometry, what was reported as possible consequence of an inactivation of MTrPs in deltoid muscle. Analgesic effects of KT in patients with musculoskeletal dysfunctions are reported most frequently [8, 11, 38, 49, 53].

It is also stated that KT increases blood and lymphatic fluid circulation under the taped area in a consequence of a lifting effect, which creates a wider space between the skin and the muscle [37, 53–55], what may affect muscle functions and result in pain and ROM improvement. With respect to ROM, great importance gains the theory of the influence of KT on fascial tissue [36]. The last decade abounds in scientific exploration concerning the role of fascial system [36, 52]. The direct contact between the fascia and muscular structures suggests that it can take part in transmitting the relative tensioning (evoked by stretched KT tape) to proper receptors and thus elicit the muscle response.

Only in the intragroup comparison we observed significant increase of range of flexion movement in all three groups. However, only in KT group the relevant increase of lateral flexion movement was noted. Those results are compliant with most of the other researches which show the positive effect of KT on ROM [36, 53]. Yoshida and Kahanov [53] reported that KT applied over the lower trunk may increase the ROM of lower trunk flexion as they did not find significant differences for extension and lateral flexion. Both cases, flexion of the lower trunk and lateral flexion of the

cervical spine, seem to be the movements where muscle fibers of respective muscle are being stretched the most. Hence, the improvement may be most noticeable in those particular movements. That could explain KT stretch activation of cutaneous and fascial mechanoreceptors resulting in the improvement of muscle excitability. This phenomenon has been also reported in several other studies [38, 53–57].

5. Conclusion

The application of all three types of tapes does not influence the resting bioelectrical activity of UT muscle and may not lead to a reduction in muscle tone in the case of MTrPs. However, in comparison to CT and sham, the KT application reduces the subjective pain sensation, what confirms the scientific reports about its analgesic influence. Authors suggest further verification of CT and KT application methods to compare their therapeutic effect and also to compare them with different methods used in the therapy of MTrPs. Therefore, it is appropriate to continue measurements of KT and CT influence on bioelectrical activity of muscles with MTrPs, pain, and cervical ROM. Further experimental research should include a larger number of participants and more objective assessment tools.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] A. L. Rodríguez-Fernández, V. Garrido-Santofimia, J. Güeita-Rodríguez, and C. Fernández-de-Las-Peñas, “Effects of burst-type transcutaneous electrical nerve stimulation on cervical range of motion and latent myofascial trigger point pain sensitivity,” *Archives of Physical Medicine and Rehabilitation*, vol. 92, no. 9, pp. 1353–1358, 2011.
- [2] D. M. Kietrys, K. M. Palombaro, E. Azzaretto et al., “Effectiveness of dry needling for upper-quarter myofascial pain: a systematic review and meta-analysis,” *Journal of Orthopaedic and Sports Physical Therapy*, vol. 43, no. 9, pp. 620–634, 2013.
- [3] D. Celik and P. Yeldan, “The relationship between latent trigger point and muscle strength in healthy subjects: a double-blind study,” *Journal of Back and Musculoskeletal Rehabilitation*, vol. 24, no. 4, pp. 251–256, 2011.
- [4] D. Celik and E. K. Mutlu, “Clinical implication of latent myofascial trigger point,” *Current Pain and Headache Reports*, vol. 17, no. 8, article 353, 2013.
- [5] J. Buchmann, B. Neustadt, K. Buchmann-Barthel et al., “Objective measurement of tissue tension in myofascial trigger point areas before and during the administration of anesthesia with complete blocking of neuromuscular transmission,” *Clinical Journal of Pain*, vol. 30, no. 3, pp. 191–198, 2014.
- [6] K. P. Botwin and B. C. Patel, “Electromyographically guided trigger point injections in the cervicothoracic musculature of obese patients: a new and unreported technique,” *Pain Physician*, vol. 10, no. 6, pp. 753–756, 2007.
- [7] Y. Bae, “Change the myofascial pain and range of motion of the temporomandibular joint following kinesio taping of latent myofascial trigger points in the sternocleidomastoid muscle,” *Journal of Physical Therapy Science*, vol. 26, no. 9, pp. 1321–1324, 2014.
- [8] S. H. Bae, J. H. Lee, K. A. Oh, and K. Y. Kim, “The effects of kinesio taping on potential in chronic low back pain patients anticipatory postural control and cerebral cortex,” *Journal of Physical Therapy Science*, vol. 25, no. 11, pp. 1367–1371, 2013.
- [9] E. A. Tough, A. R. White, S. Richards, and J. Campbell, “Variability of criteria used to diagnose myofascial trigger point pain syndrome—evidence from a review of the literature,” *Clinical Journal of Pain*, vol. 23, no. 3, pp. 278–286, 2007.
- [10] Y.-M. Xu, H.-Y. Ge, and L. Arendt-Nielsen, “Sustained nociceptive mechanical stimulation of latent myofascial trigger point induces central sensitization in healthy subjects,” *The Journal of Pain*, vol. 11, no. 12, pp. 1348–1355, 2010.
- [11] A. M. Castro-Sánchez, I. C. Lara-Palomo, G. A. Matarán-Peñarocha, M. Fernández-Sánchez, N. Sánchez-Labraca, and M. Arroyo-Morales, “Kinesio Taping reduces disability and pain slightly in chronic non-specific low back pain: a randomised trial,” *Journal of Physiotherapy*, vol. 58, no. 2, pp. 89–95, 2012.
- [12] G. Bodes-Pardo, D. Pecos-Martín, T. Gallego-Izquierdo, J. Salom-Moreno, C. Fernández-De-Las-Peñas, and R. Ortega-Santiago, “Manual treatment for cervicogenic headache and active trigger point in the sternocleidomastoid muscle: a pilot randomized clinical trial,” *Journal of Manipulative and Physiological Therapeutics*, vol. 36, no. 7, pp. 403–411, 2013.
- [13] M. R. Suh, W. H. Chang, H. S. Choi, and S. C. Lee, “Ultrasound-guided myofascial trigger point injection into brachialis muscle for rotator cuff disease patients with upper arm pain: a pilot study,” *Annals of Rehabilitation Medicine*, vol. 38, no. 5, pp. 673–681, 2014.
- [14] J. Sarrafzadeh, A. Ahmadi, and M. Yassin, “The effects of pressure release, phonophoresis of hydrocortisone, and ultrasound on upper trapezius latent myofascial trigger point,” *Archives of Physical Medicine and Rehabilitation*, vol. 93, no. 1, pp. 72–77, 2012.
- [15] A. F. Moraska, L. Stenerson, N. Butryn, J. P. Krutsch, S. J. Schmiede, and J. D. Mann, “Myofascial trigger point-focused head and neck massage for recurrent tension-type headache: a randomized, placebo-controlled clinical trial,” *The Clinical Journal of Pain*, vol. 31, no. 2, pp. 159–168, 2015.
- [16] J. Macgregor and D. G. von Schweinitz, “Needle electromyographic activity of myofascial trigger points and control sites in equine cleido-brachialis muscle—an observational study,” *Acupuncture in Medicine*, vol. 24, no. 2, pp. 61–70, 2006.
- [17] R. Llamas-Ramos, D. Pecos-Martín, T. Gallego-Izquierdo et al., “Comparison of the short-term outcomes between trigger point dry needling and trigger point manual therapy for the management of chronic mechanical neck pain: a randomized clinical trial,” *The Journal of Orthopaedic and Sports Physical Therapy*, vol. 44, no. 11, pp. 852–861, 2014.
- [18] D. T. Gulick, “Influence of instrument assisted soft tissue treatment techniques on myofascial trigger points,” *Journal of Bodywork and Movement Therapies*, vol. 18, no. 4, pp. 602–607, 2014.
- [19] H. Gemmell and A. Hilland, “Immediate effect of electric point stimulation (TENS) in treating latent upper trapezius trigger points: a double blind randomised placebo-controlled trial,” *Journal of Bodywork and Movement Therapies*, vol. 15, no. 3, pp. 348–354, 2011.
- [20] H.-Y. Ge, S. Monterde, T. Graven-Nielsen, and L. Arendt-Nielsen, “Latent myofascial trigger points are associated with

- an increased intramuscular electromyographic activity during synergistic muscle activation," *The Journal of Pain*, vol. 15, no. 2, pp. 181–187, 2014.
- [21] D. O. Draper, C. Mahaffey, D. Kaiser, D. Eggett, and J. Jarmin, "Thermal ultrasound decreases tissue stiffness of trigger points in upper trapezius muscles," *Physiotherapy Theory and Practice*, vol. 26, no. 3, pp. 167–172, 2010.
- [22] L.-W. Chou, Y.-L. Hsieh, M.-J. Kao, and C.-Z. Hong, "Remote influences of acupuncture on the pain intensity and the amplitude changes of endplate noise in the myofascial triggerpoint of the upper trapezius muscle," *Archives of Physical Medicine and Rehabilitation*, vol. 90, no. 6, pp. 905–912, 2009.
- [23] L. W. Chou, Y. L. Hsieh, H. S. Chen, C. Z. Hong, M. J. Kao, and T. I. Han, "Remote therapeutic effectiveness of acupuncture in treating myofascial trigger point of the upper trapezius muscle," *American Journal of Physical Medicine & Rehabilitation*, vol. 90, no. 12, pp. 1036–1049, 2011.
- [24] M. F. Aranha, C. E. Müller, and M. B. Gavião, "Pain intensity and cervical range of motion in women with myofascial pain treated with acupuncture and electroacupuncture: a double-blinded, randomized clinical trial," *Brazilian Journal of Physical Therapy*, 2014.
- [25] C. Myburgh, H. H. Lauridsen, A. H. Larsen, and J. Hartvigsen, "Standardized manual palpation of myofascial trigger points in relation to neck/shoulder pain; the influence of clinical experience on inter-examiner reproducibility," *Manual Therapy*, vol. 16, no. 2, pp. 136–140, 2011.
- [26] N. M. Oliveira-Campelo, C. A. de Melo, F. Albuquerque-Sendín, and J. P. Machado, "Short- and medium-term effects of manual therapy on cervical active range of motion and pressure pain sensitivity in latent myofascial pain of the upper trapezius muscle: a randomized controlled trial," *Journal of Manipulative and Physiological Therapeutics*, vol. 36, no. 5, pp. 300–309, 2013.
- [27] R. Renan-Ordine, F. Albuquerque-Sendín, D. P. R. de Souza, J. A. Cleland, and C. Fernández-De-Las-Peñas, "Effectiveness of myofascial trigger point manual therapy combined with a self-stretching protocol for the management of plantar heel pain: a randomized controlled trial," *Journal of Orthopaedic and Sports Physical Therapy*, vol. 41, no. 2, pp. 43–50, 2011.
- [28] K. Kase, J. Wallis, and T. Kase, *Clinical Therapeutic Applications of the Kinesio Taping Method*, Kinesio Taping Association, 2003.
- [29] P. D. C. Silva Parreira, L. D. C. Menezes Costa, R. Takahashi et al., "Do convolutions in Kinesio Taping matter? Comparison of two Kinesio Taping approaches in patients with chronic nonspecific low back pain: protocol of a randomised trial," *Journal of Physiotherapy*, vol. 59, no. 1, p. 52, 2013.
- [30] C. P. Pdo, C. C. Lda, R. Takahashi et al., "Kinesio taping to generate skin convolutions is not better than sham taping for people with chronic non-specific low back pain: a randomised trial," *Journal of Physiotherapy*, vol. 60, no. 2, pp. 90–96, 2014.
- [31] P. D. C. S. Parreira, L. D. C. M. Costa, L. C. Hespanhol Jr., A. D. Lopes, and L. O. P. Costa, "Current evidence does not support the use of Kinesio Taping in clinical practice: a systematic review," *Journal of Physiotherapy*, vol. 60, no. 1, pp. 31–39, 2014.
- [32] A. M. Montalvo, E. L. Cara, and G. D. Myer, "Effect of kinesiology taping on pain in individuals with musculoskeletal injuries: systematic review and meta-analysis," *The Physician and Sportsmedicine*, vol. 42, no. 2, pp. 48–57, 2014.
- [33] R. Merino-Marban, E. Fernandez-Rodriguez, and D. Mayorga-Vega, "The effect of kinesiotaping on calf pain and extensibility immediately after its application and after a duathlon competition," *Research in Sports Medicine*, vol. 22, no. 1, pp. 1–11, 2014.
- [34] A. Kalron and S. Bar-Sela, "A systematic review of the effectiveness of Kinesio Taping—Fact or fashion?" *European Journal of Physical and Rehabilitation Medicine*, vol. 49, no. 5, pp. 699–709, 2013.
- [35] S. J. Kachanathu, A. M. Alenazi, H. E. Seif, A. R. Hafez, and A. M. Alroumim, "Comparison between Kinesio taping and a traditional physical therapy program in treatment of nonspecific low back pain," *Journal of Physical Therapy Science*, vol. 26, no. 8, pp. 1185–1188, 2014.
- [36] A. Gusella, M. Bettuolo, F. Contiero, and G. Volpe, "Kinesio taping and muscular activity: a myofascial hypothesis and a randomised, blinded trial on healthy individuals," *Journal of Bodywork and Movement Therapies*, vol. 18, no. 3, pp. 405–411, 2014.
- [37] J. Gómez-Soriano, J. Abián-Vicén, C. Aparicio-García et al., "The effects of Kinesio taping on muscle tone in healthy subjects: a double-blind, placebo-controlled crossover trial," *Manual Therapy*, vol. 19, no. 2, pp. 131–136, 2014.
- [38] F. García-Muro, Á. L. Rodríguez-Fernández, and Á. Herrero-de-Lucas, "Treatment of myofascial pain in the shoulder with Kinesio Taping. A case report," *Manual Therapy*, vol. 15, no. 3, pp. 292–295, 2010.
- [39] S. Álvarez-Álvarez, "Effects of Kinesio Tape in low back muscle fatigue: randomized, controlled, doubled-blinded clinical trial on healthy subjects," *Journal of Back and Musculoskeletal Rehabilitation*, vol. 27, no. 2, pp. 203–212, 2014.
- [40] M. A. Added, L. O. Costa, T. Y. Fukuda et al., "Efficacy of adding the kinesio taping method to guideline-endorsed conventional physiotherapy in patients with chronic nonspecific low back pain: a randomised controlled trial," *BMC Musculoskeletal Disorders*, vol. 14, article 301, 2013.
- [41] H. Takasaki, B. M. Delbridge, and V. Johnston, "Taping across the upper trapezius muscle reduces activity during a standardized typing task—an assessor-blinded randomized cross-over study," *Journal of Electromyography and Kinesiology*, vol. 25, no. 1, pp. 115–120, 2015.
- [42] B. G. Beutel and D. A. Cardone, "Kinesiology taping and the world wide web: a quality and content analysis of internet-based information," *International Journal of Sports Physical Therapy*, vol. 9, no. 5, pp. 665–673, 2014.
- [43] B. Kumbrink, *K Taping: An Illustrated Guide—Basics—Techniques—Indications*, Springer, 2nd edition, 2014.
- [44] C. Y. Hsieh and B. W. Yeung, "Active neck motion measurements with a tape measure," *Journal of Orthopaedic & Sports Physical Therapy*, vol. 8, no. 2, pp. 88–92, 1986.
- [45] Y. Severinsson, L. Elisson, and O. Bunketorp, "Reliability of measuring the cervical sagittal translation mobility with a simple method in a clinical setting," *Rehabilitation Research and Practice*, vol. 2012, Article ID 629104, 9 pages, 2012.
- [46] C. Ma, S. Wu, G. Li, X. Xiao, M. Mai, and T. Yan, "Comparison of miniscalpel-needle release, acupuncture needling, and stretching exercise to trigger point in myofascial pain syndrome," *Clinical Journal of Pain*, vol. 26, no. 3, pp. 251–257, 2010.
- [47] C. H. P. de Koning, S. P. van den Heuvel, J. B. Staal, B. C. M. Smits-Engelsman, and E. J. M. Hendriks, "Clinimetric evaluation of active range of motion measures in patients with non-specific neck pain: a systematic review," *European Spine Journal*, vol. 17, no. 7, pp. 905–921, 2008.
- [48] J. V. Viitanen, M.-L. Kokko, S. Heikkilä, and H. Kautiainen, "Neck mobility assessment in ankylosing spondylitis: a clinical study of nine measurements including new tape methods

- for cervical rotation and lateral flexion,” *British Journal of Rheumatology*, vol. 37, no. 4, pp. 377–381, 1998.
- [49] M. Paoloni, A. Bernetti, G. Fratocchi et al., “Kinesio Taping applied to lumbar muscles influences clinical and electromyographic characteristics in chronic low back pain patients,” *European Journal of Physical and Rehabilitation Medicine*, vol. 47, no. 2, pp. 237–244, 2011.
- [50] T. Huang, W. Cheng, and J. Lin, “Relationship between trapezius muscle activity and typing speed: taping effect,” *Ergonomics*, vol. 55, no. 11, pp. 1404–1411, 2012.
- [51] J. A. DeLeo, “Basic science of pain,” *The Journal of Bone & Joint Surgery—American Volume*, vol. 88, supplement 2, pp. 58–62, 2006.
- [52] G. Hedley, “Fascia science and clinical applications: editorial,” *Journal of Bodywork and Movement Therapies*, vol. 16, no. 4, pp. 494–495, 2012.
- [53] A. Yoshida and L. Kahanov, “The effect of kinesio taping on lower trunk range of motions,” *Research in Sports Medicine*, vol. 15, no. 2, pp. 103–112, 2007.
- [54] T. Halseth, J. W. McChesney, M. DeBeliso, R. Vaughn, and J. Lien, “The effects of Kinesio taping on proprioception at the ankle,” *Journal of Sports Science and Medicine*, vol. 3, no. 1, pp. 1–7, 2004.
- [55] E. Akbaş, A. O. Atay, and I. Yüksel, “The effects of additional kinesio taping over exercise in the treatment of patellofemoral pain syndrome,” *Acta Orthopaedica et Traumatologica Turcica*, vol. 45, no. 5, pp. 335–341, 2011, Erratum in: *Acta Orthopaedica et Traumatologica Turcica*, vol. 45, no. 6, p. 471, 2011.
- [56] Y. Konishi, “Tactile stimulation with Kinesiology tape alleviates muscle weakness attributable to attenuation of Ia afferents,” *Journal of Science and Medicine in Sport*, vol. 16, no. 1, pp. 45–48, 2013.
- [57] G. M. Bove, “Weaving a mat of fascia research,” *Journal of Bodywork and Movement Therapies*, vol. 16, no. 2, pp. 132–133, 2012.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

