

Letter to the Editor

Comment on “Acupuncture Point “Hegu” (LI4) is Close to the Vascular Branch from the Superficial Branch of the Radial Nerve”

Kanae Umemoto ¹, Munekazu Naito,¹ Kaori Tano,² Hayato Terayama ³, Taro Koike,⁴ Mika Ohmichi,⁵ Yusuke Ohmichi ⁵, Kou Sakabe,³ and Takashi Nakano¹

¹Department of Anatomy, Aichi Medical University School of Medicine, 1-1 Yazakokarimata, Nagakute, Aichi, Japan

²Department of Acupuncture and Moxibustion, Faculty of Health Science, Suzuka University of Medical Science, 1001-1 Kishioka, Suzuka, Mie, Japan

³Department of Anatomy, Division of Basic Medical Science, Tokai University School of Medicine, 143 Shimokasuya, Isehara, Kanagawa, Japan

⁴Department of Anatomy, Kansai Medical University, 2-5-1 Shinmachi, Hirakata, Osaka, Japan

⁵Department of Anatomy II, Kanazawa Medical University, 1-1 Daigaku, Uchinada, Kahoku, Ishikawa, Japan

Correspondence should be addressed to Kanae Umemoto; kanaeteru13@yahoo.co.jp

Received 13 February 2021; Accepted 29 March 2021; Published 21 December 2021

Academic Editor: Mark Moss

Copyright © 2021 Kanae Umemoto et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Acupuncture Point ‘Hegu’ (LI4) is Close to the Vascular Branch from the Superficial Branch of the Radial Nerve” [1], the authors describe LI4 as being located at the base of the first and second metacarpal bones; however, the World Health Organization (WHO) describes LI4 as being on the dorsum of the hand radial to the midpoint of the second metacarpal bone [2] and the authors wish to make the following clarifications.

The contents and notations in textbooks on the topic of acupoints have been written separately in each country, and the standardization of meridian point locations was agreed in the Formal Consultation Meeting leading to the publication of the WHO Standard Acupuncture Point Locations in the Western Pacific Region by the WHO/WPRO in May 2008. The location of LI4 adopted during the meeting was different from that stated in textbooks in South Korea and Japan (i.e., at the base of the first and second metacarpal bones) [3]. In many historical books including “Huang Di Nei Jing Ling Shu (黄帝内经灵枢)”, which is one of the earliest written works on Traditional Chinese Medicine, the location of LI4 was stated to be “between the thumb and index finger.” It has been reported that the location of LI4 on “tai ping sheng hui fang (太平圣惠方)” (992) is considered

to be the lower edge at the base of the first and second metacarpal bones [3, 4].

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] K. Umemoto, M. Naito, K. Tano et al., “Acupuncture Point “Hegu” (LI4) is Close to the Vascular Branch from the Superficial Branch of the Radial Nerve,” *Evidence-Based Complementary and Alternative Medicine*, vol. 2019, Article ID 6879076, 6 pages, 2019.
- [2] World Health Organization, *WHO Standard Acupuncture Point Locations in the Western Pacific Region*, World Health Organization, 2008.
- [3] The Second Japan Meridian Point Committee, *Syokai Keiketsubui Kanzen Gaido (详解経穴部位完全ガイド, The Complete Detailed Guide to Meridian Points)*, Ishiyaku Publishers, Tokyo, Japan, 2009.
- [4] Japanese Acupoint Committee, *Hyojun Keiketsugaku (標準経穴学, Standard Meridian Points)*, Ishiyaku Publishers, Tokyo, Japan, 1989.