

Retraction

Retracted: Computational Analysis of the Related Factors of Deep Vein Thrombosis (DVT) Formation in Patients Undergoing Hip Fracture Surgery

Evidence-Based Complementary and Alternative Medicine

Received 20 June 2023; Accepted 20 June 2023; Published 21 June 2023

Copyright © 2023 Evidence-Based Complementary and Alternative Medicine. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This article has been retracted by Hindawi following an investigation undertaken by the publisher [1]. This investigation has uncovered evidence of one or more of the following indicators of systematic manipulation of the publication process:

- (1) Discrepancies in scope
- (2) Discrepancies in the description of the research reported
- (3) Discrepancies between the availability of data and the research described
- (4) Inappropriate citations
- (5) Incoherent, meaningless and/or irrelevant content included in the article
- (6) Peer-review manipulation

The presence of these indicators undermines our confidence in the integrity of the article's content and we cannot, therefore, vouch for its reliability. Please note that this notice is intended solely to alert readers that the content of this article is unreliable. We have not investigated whether authors were aware of or involved in the systematic manipulation of the publication process.

In addition, our investigation has also shown that one or more of the following human-subject reporting requirements has not been met in this article: ethical approval by an Institutional Review Board (IRB) committee or equivalent, patient/participant consent to participate, and/or agreement to publish patient/participant details (where relevant).

Wiley and Hindawi regrets that the usual quality checks did not identify these issues before publication and have since put additional measures in place to safeguard research integrity.

We wish to credit our own Research Integrity and Research Publishing teams and anonymous and named external researchers and research integrity experts for contributing to this investigation.

The corresponding author, as the representative of all authors, has been given the opportunity to register their agreement or disagreement to this retraction. We have kept a record of any response received.

References

- [1] Y. Zhou, H. Chen, Y. Zhang et al., "Computational Analysis of the Related Factors of Deep Vein Thrombosis (DVT) Formation in Patients Undergoing Hip Fracture Surgery," *Evidence-Based Complementary and Alternative Medicine*, vol. 2022, Article ID 1127095, 6 pages, 2022.

Research Article

Computational Analysis of the Related Factors of Deep Vein Thrombosis (DVT) Formation in Patients Undergoing Hip Fracture Surgery

Yi Zhou,¹ Huali Chen ,² Yan Zhang ,^{3,4} Chao Yang,¹ Xiaohui Yi,¹ Shanshan Liu,¹ and Yao Zeng¹

¹Department of Orthopedics (Department of Osteoarticular and Sports Medicine), Hunan Provincial People's Hospital (The First-Affiliated Hospital of Hunan Normal University), Changsha 410005, China

²Quality Control Office of Nursing Department, Hunan Provincial People's Hospital (The First-Affiliated Hospital of Hunan Normal University), Changsha 410005, China

³Department of Cardiology, Hunan Provincial People's Hospital (The First-Affiliated Hospital of Hunan Normal University), Changsha 410005, China

⁴Clinical Medicine Research Center of Heart Failure of Hunan Province, Hunan Provincial People's Hospital (The First-Affiliated Hospital of Hunan Normal University), Changsha 410000, China

Correspondence should be addressed to Yan Zhang; zhangyan.01@outlook.com

Received 7 May 2022; Accepted 14 May 2022; Published 27 May 2022

Academic Editor: Fenglin Liu

Copyright © 2022 Yi Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A retrospective study was conducted on 51 patients undergoing hip fracture surgery to investigate the factors associated with the formation of deep venous thrombosis (DVT). The independent sample *t*-test and correlation analysis were used to sort out and analyze the data. The findings are as follows. (1) Different gender samples showed significant differences in the Caprini score and thrombus location. Most DVTs in females are located in the posterior tibial vein and intermuscular veins. The Caprini score of females was significantly higher than that of males. (2) Age displays a positive correlation with DVT, coronary heart disease, hypertension, and different surgical types, respectively. (3) There is a correlation between age and operation duration. (4) Hyperlipidemia and cerebrovascular disease show a positive correlation with DVT. (5) There was a significant negative correlation between the Caprini score and the quantification of D-dimer. This indicates that in this sample, the higher the patients' Caprini score is, the lower the quantification of D-dimer will be. (6) Hyperlipidemia and cardiac insufficiency show a positive correlation with cerebrovascular disease. Patients with hyperlipidemia and cardiac insufficiency may also suffer from cerebrovascular diseases.

1. Introduction

Patients undergoing hip fracture surgery are likely to lead to deep venous thrombosis (DVT) due to decreased activity (for one, intraoperative supine position and anesthesia will make the peripheral veins dilate. For another, postoperative bed rest will cause deep venous blood flow in the lower limbs slow), dehydration (diet is prohibited for a long time before and after the operation, and blood viscosity will increase), senile diseases (a variety of chronic diseases), surgical

trauma, and anesthesia factors. Following a hip fracture, vascular damage is serious, and platelet aggregation cannot heal the wound, so it needs to form a blood clot [1]. Thrombin will activate more platelets to promote blood coagulation. It helps soluble fibrin in plasma to be converted into insoluble protein for rapid hemostasis [2–4]. As shown in Figure 1, the interaction among exposed lipid-rich atherosclerotic plaques and platelet receptors and coagulation factors can lead to platelet activation and aggregation. This may lead to DVT, increase the risk of surgery, affect the




FIGURE 1: Causes of thrombosis.

smooth surgery, and even endanger life. It can also lead to heart disease, high blood pressure, high blood fat, and many other diseases [5, 6].

SPSS has been widely used for data analysis in the analysis of risk factors related to DVT, and case analysis is used more frequently [7–10]. Treatment options for DVT have been analyzed and developed in some studies [11–13]. The INARI FlowTriever device can remove DVT and significantly reduce the risk of bleeding [14]. Some studies analyzed and studied the causes of DVT and the locations where DVT frequently occurs [15, 16]. Other studies have found that DVT can trigger other diseases [2, 17–19], such as multiple organ embolism, stroke, and acute myocardial infarction.

In summary, DVT is one of the common and serious complications of hip fracture. Whether the operation is performed or not, the incidence of DVT is high and the harm is great. Once it forms, it will affect the operation effect and even endanger life. At present, there are few studies analyzing the related risk factors of DVT formation in patients undergoing hip fracture surgery. Therefore, in this study, the locations and the related factors of DVT formation were analyzed among 51 patients with hip fractures, including 18 males and 33 females. The independent sample *t*-test and correlation analysis were used to sort out the data. This study will supplement the data on DVT and its related factors in hip fracture patients, so as to provide a theoretical basis for future research in this field.

2. Methods

2.1. General Description. 51 hip fracture patients (18 males and 33 females) were randomly selected as research subjects according to their hospital stay. These patients were admitted to the orthopedics department for surgery from June 1, 2021, to April 7, 2022. The femoral head replacement was performed in 19 patients, intramedullary nailing in 21 patients, and total hip replacement in 11 patients. SPSS analysis was conducted on the related factors of deep vein thrombosis (DVT) formation in these patients. The *t*-test analysis, Pearson correlation, and curve regression of SPSS were used to study the relationship between variables.

2.2. Inclusion and Exclusion Criteria. Inclusion criteria were as follows:

- (1) Patients admitted to our hospital and diagnosed with no DVT using color Doppler ultrasonography of blood vessels of both lower limbs
- (2) Patients with hip fractures and undergoing surgical treatment

Exclusion criteria were as follows:

- (1) Coagulation dysfunction
- (2) Pathological fractures
- (3) Data cannot be collected

3. Results

As given in Table 1, the *t*-test (independent sample *t*-test) was used to analyze the differences between gender for the Caprini score as well as thrombus sites. As given in Table 1, samples of different genders showed the significance for the Caprini score and DVT site. For females, most DVTs are located in the posterior tibial vein and intermuscular veins. This is different from males. Females were significantly higher than males in the Caprini score.

As given in Table 2, correlation analysis was performed to study the correlation between age and DVT, coronary heart disease, hypertension, D-dimer quantification, and operation type changes, respectively. The Pearson correlation coefficient was used to indicate the strength of the correlation. According to the correlation coefficient in Table 2, age displayed a positive correlation with DVT, coronary heart disease, hypertension, and surgical-type changes, respectively. There was a negative correlation between age and the D-dimer quantitative value. This indicates that, in this sample, as the age increases, people will be more likely to suffer from DVT, coronary heart disease, hypertension, and other diseases, but the quantitative value of D-dimer will be lower. In addition, as the age increases, patients will choose different types of surgery.

As given in Table 3, the regression coefficient ($p = 0.006$, less than 0.01) of the curve between age and surgical duration indicates that the age (independent variable) has a correlation with the surgical duration (dependent variable). But the selected effective sample size was too small, and this conclusion is only for describing the situation of this sample. Curvilinear regression is a nonlinear relationship in the relation form, but it can be changed into a linear relationship through various conversions. The relationship between age and surgery duration is clearly shown in Figure 2.

As given in Table 4, correlation analysis was performed to study the correlation between DVT and hyperlipidemia and cerebrovascular disease, respectively. The Pearson correlation coefficient was used to indicate the strength of the correlation. From the correlation coefficient given in Table 4, it can be seen that DVT shows a positive correlation with hyperlipidemia and cerebrovascular disease. It indicates that hyperlipidemia and cerebrovascular disease increase the risk of DVT in patients with hip fractures in this sample.

As given in Table 5, correlation analysis was performed to analyze the correlation between the Caprini score and

TABLE 1: The *t*-test analysis of gender and the Caprini score and thrombus site.

	Gender (mean and standard deviation)		<i>t</i>	<i>P</i>
	Female (<i>n</i> = 33)	Male (<i>n</i> = 18)		
Caprini score	11.21 ± 1.95	9.33 ± 2.66	2.888	0.006**
Thrombus site	2.33 ± 0.85	1.78 ± 0.81	2.261	0.028*

P* < 0.05. *P* < 0.01.

TABLE 2: Pearson correlation of age with DVT, coronary heart disease, hypertension, D-dimer quantification, and surgical-type changes.

		Age (years)
DVT	Correlation coefficient	0.421**
	<i>P</i> value	0.002
Coronary heart disease	Correlation coefficient	0.294*
	<i>P</i> value	0.036
Hypertension	Correlation coefficient	0.301*
	<i>P</i> value	0.032
D-dimer quantification	Correlation coefficient	-0.288*
	<i>P</i> value	0.041
Surgical-type changes	Correlation coefficient	0.307*
	<i>P</i> value	0.028

P* < 0.05. *P* < 0.01.

TABLE 3: Curvilinear regression coefficient of age and operation duration.

	Nonstandardized coefficient		Normalization coefficient	<i>t</i>	<i>P</i>
	<i>B</i>	Standard error	Beta		
Constant	243.905	80.435	—	3.032	0.006**
Age (years)	-3.040	2.890	-1.143	-1.052	0.303
Age (years)**2	0.018	0.024	0.785	0.723	0.477

***P* < 0.01. Dependent variable, operation duration (minutes).




FIGURE 2: Quadratic curve fitting.

TABLE 4: Pearson correlation of DVT with hyperlipidemia and cerebrovascular disease.

		DVT
Hyperlipidemia	Correlation coefficient	0.305*
	<i>P</i> value	0.030
Cerebrovascular disease	Correlation coefficient	0.368**
	<i>P</i> value	0.008

P* < 0.05. *P* < 0.01.

D-dimer quantification. The Pearson correlation coefficient was used to indicate the strength of the correlation. From the correlation coefficient given in Table 5, it can be seen that there is a significant negative correlation between the Caprini score and D-dimer quantification. This indicates that a higher Caprini score is associated with a lower D-dimer quantitative value in this sample. D-dimer is an indicator of risk for DVT. Notably, the higher the Caprini score and D-dimer quantification value are, the higher the risk of DVT will be. Theoretically, the Caprini score and the D-dimer quantification value have a positive correlation with thrombosis. However, in this sample, the Caprini score displayed a negative correlation with D-dimer quantification. The cause of this result remains unclear and deserves attention and discussion in future studies.

As given in Table 6, correlation analysis was performed to study the correlation between cerebrovascular disease and cardiac insufficiency and hyperlipidemia, respectively. The Pearson correlation coefficient was used to indicate the strength of the correlation. From the correlation coefficient given in Table 6, it can be seen that cerebrovascular disease displays a positive correlation with cardiac insufficiency and hyperlipidemia. It indicates that in this sample, patients with hyperlipidemia and cardiac insufficiency may be accompanied by the occurrence of cerebrovascular diseases.

TABLE 5: Pearson correlation of Caprini scores with D-dimer quantification.

	Correlation coefficient	Caprini score
D-dimer quantification	P value	-0.376** 0.006

** $P < 0.01$.

TABLE 6: Pearson correlation of cerebrovascular diseases with cardiac insufficiency and hyperlipidemia.

		Cerebrovascular disease
Cardiac insufficiency	Correlation coefficient	0.327*
	P value	0.020
Hyperlipidemia	Correlation coefficient	0.328*
	P value	0.019

* $P < 0.05$.

4. Conclusion

The following conclusions were derived from this sample:

- (1) Samples of different genders showed significant differences in the Caprini score and thrombus location. The location of thrombosis in females is different from that in males, and most DVTs in females are located in the posterior tibial vein and intermuscular veins. The Caprini score of females was significantly higher than that of males.
- (2) Age displays a positive correlation with DVT, coronary heart disease, hypertension, and different surgical types, respectively. There was a negative correlation between age and D-dimer. It indicates that in this sample, as the age increases, people will be more likely to suffer from DVT, coronary heart disease, hypertension, and other diseases, but the quantitative value of D-dimer will be lower. As the age increases, patients will choose different types of surgery.
- (3) There is a correlation between age and operation duration
- (4) Hyperlipidemia and cerebrovascular disease show a positive correlation with DVT. This indicates that in this sample, the patients with hyperlipidemia and cerebrovascular disease had an increased risk of DVT.
- (5) There was a significant negative correlation between the Caprini score and the quantification of D-dimer. This indicates that in this sample, the higher the patients' Caprini score is, the lower the quantitation of D-dimer will be.
- (6) Hyperlipidemia and cardiac insufficiency show a positive correlation with cerebrovascular diseases. Patients with hyperlipidemia and cardiac insufficiency may suffer from cerebrovascular diseases.

5. Discussion

In this sample, females differ from males in the location of the thrombus, and most DVTs in females are located in the posterior tibial vein and intermuscular veins. However, there is no fixed location in males. This may be due to the fact that the majority of patients with deep venous thrombosis of the lower extremities suffer from varicose veins of the lower extremities [20]. Females are more likely to have varicose veins in the lower extremities [21] and have a higher incidence of hip fractures than in males [22]. There were twice as many females as males in this sample, so there is such a significant difference.

It is also important to note that the Caprini score is significantly higher for females than for males. The Caprini score is related to age, gender, and thrombosis. This is meaningful because previous studies have found that VTE frequently occurs in females who use estrogen, during pregnancy, or in people with thrombophilia [23–26]. Our study was in agreement with other researchers. At present, these studies do not address specific reasons for gender differences. The causes of this situation can be further explored in future studies.

It was concluded that there was a negative correlation between age and D-dimer quantification. In previous studies, D-dimer levels were positively correlated with age [27] and significantly decreased in males [28]. Since the number of young and middle-aged people in the samples of this study is small and the D-dimer level of the young and middle-aged people is already high at the time of going to the hospital, it can be concluded that the D-dimer value decreases with age in statistical significance. The development of age-related changes in microcirculation and blood coagulation was excluded, resulting in results that were inconsistent with the increasing concentrations of D-dimer with age.

The study found that there was a relationship between age and operation duration. The older the patient was, the longer the operation duration would be. This finding is common in surgery [29]. The older the patient is, the more comorbidities he has [30] and the higher the risk of surgery would be.

A higher Caprini score was associated with a lower D-dimer quantitative value. Notably, the D-dimer quantitative value and Caprini scale were used as indicators and assessment tools to determine the risk of DVT, respectively [31]. Theoretically, both have a positive correlation with thrombosis. It indicates that the higher the Caprini score is, the greater the risk of DVT will occur, and the higher the D-dimer quantitative value is, the more likely the patients will suffer from thrombosis [32]. However, in this sample, the Caprini score showed a negative correlation with the

D-dimer quantification value. Our speculation about the cause of this result was that the sample cases were incidental. There was accidental variation in the extracted samples, and the variation of medical data may lead to wrong conclusions [33]. This situation deserves attention and discussion in future studies.

Data Availability

The data used to support this study are included within the article and the supplementary file.

Ethical Approval

This study was approved by the Ethical Review Committee of Hunan Provincial People's Hospital (the first-affiliated hospital of Hunan Normal University).

Disclosure

Yi Zhou and Huali Chen are the co-first authors.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Yi Zhou and Huali Chen conceptualized and designed the study and wrote the first draft of the manuscript. Yan Zhang revised the article and involved in project management. Chao Yang, Xiaohui Yi, Shanshan Liu, and Yao Zeng contributed to data collection and analysis. All the authors approved the submitted version. Yi Zhou and Huali Chen contributed equally to this work.

Acknowledgments

The work was supported by grants from the Key Research and Development Program of Hunan Province (2019SK2021) and the Health Commission of Hunan Province of China (202104072141 and 202214013647).

Supplementary Materials

Raw data for the analysis. (*Supplementary Materials*)

References

- [1] A. L. Fogelson and K. B. Neeves, "Fluid mechanics of blood clot formation," *Annual Review of Fluid Mechanics*, vol. 47, no. 1, pp. 377–403, 2015.
- [2] C. A. Bravo, J. A. Fried, J. Z. Willey et al., "Presence of intracardiac thrombus at the time of left ventricular assist device implantation is associated with an increased risk of stroke and death," *Journal of Cardiac Failure*, vol. 27, no. 12, pp. 1367–1373, 2021.
- [3] L. Herfs, F. Swieringa, N. Jooss et al., "Multiparameter microfluidics assay of thrombus formation reveals increased sensitivity to contraction and antiplatelet agents at physiological temperature," *Thrombosis Research*, vol. 203, pp. 46–56, 2021.
- [4] J. W. Weisel, "Fibrinogen and fibrin," *Advances in Protein Chemistry*, vol. 70, pp. 247–299, 2005.
- [5] M. Rezaeimoghaddam and F. N. van de Vosse, "Continuum modeling of thrombus formation and growth under different shear rates," *Journal of Biomechanics*, vol. 132, Article ID 110915, 2022.
- [6] S. Yesudasan and R. D. Averett, "Recent advances in computational modeling of fibrin clot formation: a review," *Computational Biology and Chemistry*, vol. 83, 2019.
- [7] M. Abbasi, J. Arturo Larco, M. O. Mereuta et al., "Diverse thrombus composition in thrombectomy stroke patients with longer time to recanalization," *Thrombosis Research*, vol. 209, pp. 99–104, 2022.
- [8] C. Cohen, K. Gaillot, H. Ifergan et al., "Quantitative analysis of thrombus migration before mechanical thrombectomy: determinants and relationship with procedural and clinical outcomes," *Journal of Neuroradiology*, 2021.
- [9] T. Kanagami, N. Hayakawa, and J. Kanda, "TCTAP C-144 novel bailout technique using myocardial biopsy forceps with intravascular ultrasound guided in patient with distal embolism of organized thrombus," *Journal of the American College of Cardiology*, vol. 79, no. 15, pp. S341–S343, 2022.
- [10] V. Lau, M. Blaszkak, J. Lam, M. German, and F. Myslik, "Point-of-Care resuscitative echocardiography diagnosis of intracardiac thrombus during cardiac arrest (predict study): a retrospective, observational cohort study," *Resuscitation*, vol. 10, 2022.
- [11] K. A. Honan, A. Jogimahanti, and T. Khair, "An updated review of the efficacy and safety of direct oral anticoagulants in treatment of left ventricular thrombus," *The American Journal of Medicine*, vol. 135, no. 1, pp. 17–23, 2022.
- [12] K. Konstantinou, J. R. Davies, O. Alsanjari et al., "Clinical effectiveness of thrombus aspiration during percutaneous coronary intervention for stent thrombosis in a contemporary setting," *Hellenic Journal of Cardiology*, 2022.
- [13] H. Neki, T. Katano, T. Maeda, A. Shibata, H. Komine, and Y. Kikkawa, "Familiarization with contact aspiration using non-penetrating of the thrombus (CANP) technique as the initial procedure for acute ischemic stroke," *Journal of Stroke and Cerebrovascular Diseases*, vol. 30, no. 11, 2021.
- [14] N. Nezami, I. Latich, N. Murali et al., "Right atrial and massive pulmonary artery mechanical thrombectomy under echocardiography guidance using the FlowTriever system," *EJVES Short Reports*, vol. 45, pp. 22–25, 2019.
- [15] M. Ono, T. Kido, M. Burri et al., "Risk factors for thrombus formation at stage 2 palliation and its effect on long-term outcome in patients with univentricular heart," *Seminars in Thoracic and Cardiovascular Surgery*, vol. 21, 2021.
- [16] G. M. Vasilakis, D. A. Lakhani, A. Adelanwa, J. P. Hogg, and C. Kim, "Atypical imaging presentation of a massive intracavitary cardiac thrombus: a case report and brief review of the literature," *Radiology Case Reports*, vol. 16, no. 10, pp. 2847–2852, 2021.
- [17] Y. Imamura, R. Kowatari, K. Daitoku, S. Goto, and I. Fukuda, "Multiorgan emboli due to an intraluminal thrombus from frozen elephant trunk," *Cardiovascular Pathology*, vol. 52, 2021.
- [18] T. Saito, H. Yamamoto, S. Oishi et al., "Left main trunk occlusion due to impella-related thrombus in a patient with extracorporeal cardiopulmonary resuscitation," *JACC: Cardiovascular Interventions*, vol. 14, no. 22, pp. e313–e316, 2021.
- [19] J. Zhou, S. Yu, P. Zhou et al., "Impact of residual thrombus burden on ventricular deformation after acute myocardial infarction: a sub-analysis from an intravascular optical

- coherence tomography study," *EclinicalMedicine*, vol. 39, 2021.
- [20] T. Alsaigh and E. Fukaya, "Varicose veins and chronic venous disease," *Cardiology Clinics*, vol. 39, no. 4, pp. 567–581, 2021.
- [21] H. Nemoto, M. Mo, T. Ito et al., "Venous thromboembolism complications after endovenous laser ablation for varicose veins and role of duplex ultrasound scan," *Journal of Vascular Surgery: Venous and Lymphatic Disorders*, vol. 7, no. 6, pp. 817–823, 2019.
- [22] T. T. Fung, H. E. Meyer, W. C. Willett, and D. Feskanich, "Association between diet quality scores and risk of hip fracture in postmenopausal women and men aged 50 years and older," *Journal of the Academy of Nutrition and Dietetics*, vol. 118, no. 12, pp. 2269–2279, 2018.
- [23] C. A. Hennessey, V. K. Patel, E. A. Tefera, and V. Gomez-Lobo, "Venous thromboembolism in female adolescents: patient characteristics," *Journal of Pediatric and Adolescent Gynecology*, vol. 31, no. 5, pp. 503–508, 2018.
- [24] M. A. Kohorst, D. M. Warad, A. A. Nageswara Rao, and V. Rodriguez, "Obesity, sedentary lifestyle, and video games: the new thrombophilia cocktail in adolescents," *Pediatric Blood and Cancer*, vol. 65, no. 7, 2018.
- [25] B. Lacruz, G. Tiberio, A. Latorre et al., "Venous thromboembolism in young adults: findings from the RIETE registry," *European Journal of Internal Medicine*, vol. 63, pp. 27–33, 2019.
- [26] K. A. Meier, E. Clark, C. Tarango, R. S. Chima, and E. Shaughnessy, "Venous thromboembolism in hospitalized adolescents: an approach to risk assessment and prophylaxis," *Hospital Pediatrics*, vol. 5, no. 1, pp. 44–51, 2015.
- [27] A. Rumley, J. R. Emberson, S. G. Wannamethee, L. Lennon, P. H. Whincup, and G. D. O. Lowe, "Effects of older age on fibrin D-dimer, C-reactive protein, and other hemostatic and inflammatory variables in men aged 60–79 years," *Journal of Thrombosis and Haemostasis*, vol. 4, no. 5, pp. 982–987, 2006.
- [28] C. Legnani, M. Cini, B. Cosmi et al., "Age and gender specific cut-off values to improve the performance of d-dimer assays to predict the risk of venous thromboembolism recurrence," *Internal and Emergency Medicine*, vol. 8, no. 3, pp. 229–236, 2013.
- [29] S. Moriguchi, Y. Maehara, D. Korenaga, K. Sugimachi, and Y. Nose, "Relationship between age and the time of surgery and prognosis after gastrectomy for gastric cancer," *Journal of Surgical Oncology*, vol. 52, no. 2, pp. 119–123, 1993.
- [30] J. K. Prümmer, J. Howard, L. M. Grandt, R. Aguilar, F. Meneses, and L. M. Peters, "Hyperlipasemia in critically ill dogs with and without acute pancreatitis: prevalence, underlying diseases, predictors, and outcome," *Journal of Veterinary Internal Medicine*, vol. 34, no. 6, pp. 2319–2329, 2020.
- [31] I. Golemi, J. P. Salazar Adum, A. Tafur, and J. Caprini, "Venous thromboembolism prophylaxis using the caprini score," *Disease-a-Month*, vol. 65, no. 8, pp. 249–298, 2019.
- [32] K. Lobastov, G. Dementieva, N. Soshitova et al., "Utilization of the caprini score in conjunction with thrombodyamic testing reduces the number of unpredicted postoperative venous thromboembolism events in patients with colorectal cancer," *Journal of Vascular Surgery: Venous and Lymphatic Disorders*, vol. 8, no. 1, pp. 31–41, 2020.
- [33] A. P. Dempster, "Purposes and limitations of data analysis," in *Scientific Inference, Data Analysis, and Robustness*, G. E. P. Box, T. Leonard, and C.-F. Wu, Eds., pp. 117–133, Academic Press, Cambridge, MA, USA, 1983.