

Review Article

Cardiovascular Effects of *Cydonia oblonga* Miller (Quince)

Atefeh Amerizadeh ^{1,2}, Golnaz Vaseghi ², Nazgol Esmailian,² and Sedigheh Asgary ¹

¹Isfahan Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

²Applied Physiology Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

Correspondence should be addressed to Sedigheh Asgary; sedighehasgary@gmail.com

Received 21 May 2022; Revised 28 July 2022; Accepted 21 August 2022; Published 10 October 2022

Academic Editor: Fadia S. Youssef

Copyright © 2022 Atefeh Amerizadeh et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cydonia oblonga Miller (quince) is a monotypic genus in the Rosaceae family which used to treat or prevent many medical conditions such as diabetes, cancer, infections, and ulcer. This review will summarize the current state of knowledge available on botany, phytochemistry, and pharmacology of this plant with a focus on its effect on some cardiovascular risk factors such as diabetes, blood pressure, lipid profile, and body weight. Databases of Google Scholar, Web of Science, PubMed, Scopus, and SID were searched systematically for English published articles with no date limitation. There were no human studies found, and all of the studies were conducted on animals or *in vitro* models. Reviewing of all 12 included articles showed that different types of quince extract have positive effects on cardiovascular-related factors such as blood pressure, diabetes, glucose metabolism, lipid profile, ROS, body weight, liver dysfunction, and thrombosis. An antihypertensive effect of quince showed to be a magnitude that is comparable to captopril and the lipid-lowering effect of quince showed to be a magnitude that is comparable to simvastatin. However, two studies evaluated the effect of quince fruit extract on insulin levels; one of them reported no positive effect, and the other one reported a significant positive effect. It can be concluded that different parts of quince including leaf, seed, and fruit could be used for improving cardiovascular-related factors including blood pressure, glucose metabolism and diabetes, obesity, and lipid-adjusting purposes. Quince was also found to have strong anti-inflammatory and antioxidant capacity. This study paves the way for further studies on the cardiovascular effect of quince consumption as a beneficial nutraceutical in humans.

1. Introduction

Dr. Stephen DeFelice coined the term “Nutraceutical” from “Nutrition” and “Pharmaceutical” in 1989 [1]. Nutraceuticals, in contrast to pharmaceuticals, are substances, which usually have no patent protection. Both pharmaceutical and nutraceutical compounds might be used to cure or prevent diseases, but only pharmaceutical compounds have the governmental sanction. Consumers are turning increasingly toward food supplements to improve well-being when pharmaceuticals fail [2, 3]. Nowadays, nutraceuticals have received considerable interest due to their potential nutritional, safety, and therapeutic effects [4, 5]. According to previous studies, many medicinal plants such as fenugreek, sumac, dill, ginger, grape, and green tea improve the lipid profile significantly [6–10].

Quince is one of the herbal nutraceuticals that has been used throughout olden times [11]. *Cydonia oblonga* Miller (COM) leaves are used in traditional medicines for cardiovascular disease (CVD) treatment or prevention [12]. Its fruit is employed in the food sector as a source of pectin, which defends the colon from damage in IBS and peptic ulcers [13–15]. The plant’s seeds have long been used to cure cough, diarrhea, dysentery, constipation, sore throat, and bronchitis [16–18]. Quince extract has been proven to have cancer-fighting, antihemolytic, and antidiabetic properties [19–22]. It is also found to have an anti-inflammatory and antioxidant capacity [23], antiallergic [24], antidepressant, and antistress effect [25, 26]. In addition, the plant contains phenol peroxidase, an enzyme that decolorizes hazardous aromatic colors in industrial wastewater [27, 28]. Quince’s widespread therapeutic use and valuable phytochemical

composition prompted us to write a review on it, with a focus on its cardiovascular effects.

1.1. General Characteristics. The quince (*Cydonia oblonga*), is a nondrought-tolerant small deciduous tree with fruits that look like pears and a superficial and fasciculate root [11]. When exposed to air, the white-yellow pulp becomes hard, or sour, bitter, and acidic, making it unfit for consumption as a fresh product (just a few cultivars are), but when mature, the quince has a pleasant, durable, and powerful flavor. The existence of essential oil in quince fruit, whose odor-active components are considered to be monoterpene lactones and oxides, is responsible for the fruit's strong and distinctive odor [29].

1.2. Distribution. The quince fruit is a native of Western Asia and Southwest Europe and has been originated from Armenia, Iran, Azerbaijan, Turkmenistan, and southwest Russia [29]. Turkey produces the most quinces worldwide, ahead of China, Uzbekistan, Morocco, Iran, Azerbaijan, Argentina, and Spain (FAO, 2013).

1.3. Taxonomy. The quince is the only member of the genus *Cydonia*, which belongs to the Rosaceae family, subfamily Maloideae, and tribe Pireas. [11, 30].

Domain: Eukaryota
 Kingdom: Plantae
 Phylum: Spermatophyta
 Subphylum: Angiospermae
 Class: Dicotyledonae
 Subclass: Caryophyllidae
 Order: Rosales
 Tribe: Pireas
 Subfamily: Maloideae
 Family: Rosaceae
 Genus: *Cydonia*.
 Species: *Cydonia oblonga*

1.4. Phytochemistry. Different portions of the quince contain phenolic compounds, essential oils, organic acids, ionone glycosides, and tetracyclic sesterterpenes [31–35]. In the pulp, leaves, peel, seeds, and complete fruits of quinces, 27 polyphenolic substances have been discovered including citric, ascorbic, malic, oxalic, quinic, fumaric, and shikimic acids. The sugar profile of the quince fruit includes glucose, fructose, maltose, and sucrose with substantial levels of quinic and malic acids [11, 26, 29]. The fruit contains vitamin C and various minerals such as sodium, calcium, phosphorus, potassium, and nitrogen [36]. COM seeds contain triterpenes, sterols, and tannins that are reported to be responsible for their antidiarrheal activity [37–39].

2. Methodology

Databases of Google Scholar, Web of Science, PubMed, Scopus, and SID were searched systematically with no date limitation for English published articles.

The utilized phrases were “phytochemical,” “biological investigations,” “traditional uses,” “pharmacological activity,” “anticancer activity,” “antifungal activity,” “antibacterial activity,” “anti-inflammatory activity,” “medicinal plants,” “antioxidants,” “toxicity,” “commercial products,” “ethnobotany,” and “immunological activity,” with “*Cydonia oblonga*,” or “quince” that reflect subjects of interest up to Feb 2022.

3. Results

Figure 1 shows the selection process. Of 58 articles, 45 were excluded due to duplications and unrelated content/abstract/topic. In among 13 continuous papers, 2 were excluded because they were on Japanese quince [40] and Chinese quince [41]. Two studies were found through manual searching, and finally, 12 were included in our study [12, 22, 42–52]. No human experimental or clinical trial was found, and all studies were on animal or in *in vitro* models. Table 1 listed the details of the included paper.

4. Discussion

In 2016, an evaluation of quince's possible bioactive components and nutritional applications was published. We discuss and update the cardiovascular effects of quince, as well as its botany, phytochemistry, and pharmacology, in this review. Reviewing of all 13 included articles on this topic showed that different types of quince extract have positive and beneficial effects on cardiovascular-related factors such as blood pressure [50, 52, 53], diabetes, and glucose metabolism but not on the insulin level [22, 47, 48], lipid profile [12, 43–46, 49], serum biomarkers of liver function [12, 43, 44, 46], body weight [45], and thrombosis [51]. The quince leaf extract's effect on lipid profile, liver enzymes, and atherosclerotic plaque formation in the coronary artery was not significantly different from effects reported in the atorvastatin group [43]. The antihypertensive effect of 320 mg/kg per day of quince leaf extract reported to be not very different from using captopril [52]. None of the studies reported a negative or no positive effect of quince consumption on cardiac-related factors.

4.1. Lipid Profile and Body Weight. Of six studies that reported outcomes related to the effect of quince on the lipid profile, all reported significant positive improvement [12, 43–46, 49]. Umar et al. showed that total flavonoids of quince fruit and leaves could adjust the lipid profile in hyperlipidemic rats significantly [49]. In their study high dose of COM total leave flavonoids (160 mg/kg), had lipid-lowering effect almost same as 5 mg/day of simvastatin [49]. According to Khademi et al., the quince group had significantly decreased TC, TG, and LDL_C levels compared to the high-cholesterol diet group ($P < 0.05$), and it was more similar to the normal diet group. Their results showed that quince leaf extract, such as atorvastatin, can successfully adjust the lipid profile; however, both cannot improve atherosclerosis caused on by a high-fat diet significantly [43]. Another study also reported that the aqueous extract of




FIGURE 1: Flowchart of study selection.

quince fruit significantly reduced serum TG, TC, and LDL-C levels while increasing HDL-C levels in diabetic rats [46]. Abliz et al. found that in hyperlipidemia rats, COM lowered TC, TG, and LDL-C while increasing HDL-C, which was significant at high and medium doses [12]. Lee et al. reported that COM has lipid-adjusting and antiobesity effects, and it can reduce fat mass and adipose tissue weight [45].

4.2. Diabetes, Glucose Metabolism, and the Insulin Level.

Of five studies that reported outcomes related to the effect of quince on glucose metabolism and diabetes, all reported significant positive improvement [22, 40, 41, 47, 48]. Tang et al. using the *in vitro* model showed that by stimulating the PI3K/AKT signaling pathway, COM prompted glucose metabolism [48]. However, in terms of the insulin level, results were inconsistent, and procyanidin-rich extract of Chinese quince fruit showed no positive effect on the insulin level, while elevation of adiponectin in obese mice resulted in insulin level elevation [45].

4.3. Blood Pressure.

Of two studies that investigated the effect of COM on blood pressure, both reported significant positive results [50, 52]. It has been shown that extracts of COM leaves (80, 160, and 320 mg/kg) at the highest doses have an antihypertensive effect comparable to captopril [52, 53]. Another study using aqueous-methanolic extract from seeds of quince (200, 400, and 600 mg/kg doses)

showed that extract in 600 mg/kg dose produced the most effect and avoided a rise in blood pressure [50].

4.4. Oxidative Stress. Oxidative stress is defined as increased intracellular levels of reactive oxygen species (ROS). ROS plays a role in the onset and development of coronary artery disease [54]. ROS is involved in the development of oxidized LDL, which is the first stage in the pathogenesis of atherosclerosis. Furthermore, ROS may activate matrix metalloproteinases, leading to plaque rupture. However, ROS appears to have a significant function in the setting of acute MI and after reperfusion treatment [55]. ROS may oxidize myofibrillar proteins, causing contractile dysfunction in heart failure patients [56]. An *in vitro* study on a polyphenolic extract of Japanese quince fruit showed that it can reduce ROS intracellular [40]. HPLC analysis of the phenolic profile of COM showed that chlorogenic acid was the major phenolic compound in quince leaf extract [19, 45]. Chlorogenic acid is a bioactive polyphenol with potent antioxidant properties which have a heart-protective capacity [57]. Quince fruit had a lot of quinic acid and shikimic derivatives which can protect against cardiovascular disease and cancer [58], as well as procyanidins and flavonoids in it. The presence of 4-caffeoyl shik, which has antioxidative activity, in quince is significantly higher than that of an apple [57].

Quince leaves had a much stronger lowering power than green tea in a comparison research study [19]. In a similar

TABLE 1: Included studies on *Cydonia oblonga* Miller.

Authors, Year	Model	Intervention	Duration	Studied factors	Results
Zhou et al. [52]	Renal hypertensive rat	Captopril administration 25 mg/kg/day Various doses of COM leaf extracts	8 weeks	SBP&DBP	Blood pressure was considerably and dose-dependently lowered by COM leaf extracts. The greatest. COM dose produced the same results as captopril.
Abliz et al. [12]	Sprague–Dawley rats received high-fat diet	Simvastatin administration Various doses of COM leaf extracts	56 days	Lipid profile Liver enzymes MDA Liver steatosis	COM reduced TC, TG, LDL-C and increased HDL-C dose-dependently. COM inhibited the ALT, AST enzymes activity, and reduced liver steatosis.
Zhou et al. [51]	Mice	Aspirin 5 mg/kg/day COM aqueous extracts 20, 40, and 80 mg/kg/day	2 weeks	Bleeding and clotting times	In comparison to aspirin, COM extracts prolonged bleeding and clotting times in a dose-dependent manner.
Khademi et al. [43]	Rabbits	Atorvastatin (0.5 mg/kg) COM leaf extract (50 mg/kg)	12 weeks	Biopsy of coronary artery Lipid profile, AST, ALT, AP, BUN, Cr	Treatment with the COM extract at 50 mg/kg and 0.5 mg/kg atorvastatin lowered plasma lipid profile (P.0.05), liver enzymes, BUN, and Cr levels while also increasing HDL.
Khademi et al. [44]	Hypercholesterolemic rabbits	Atorvastatin (0.5 mg/kg) COM leaf extract (50 mg/kg)	8 weeks	Lipid profile, Liver enzymes	Treatment with the COM extract at 50 mg/kg and 0.5 mg/kg atorvastatin lowered plasma lipid profile (P.0.05), liver enzymes while also increasing HDL-cholesterol levels.
Mirmohammad et al. [46]	Diabetic rats	Various doses of aqueous extract of COM fruit		Lipid profile, AST, ALT, AP, BUN, Cr	LDL, TC, TG were significantly decreased. HDL was significantly increased.
Aslan et al. [22]	Healthy and diabetic rats	Various doses of COM leaf extract	5 days	Fasting blood glucose	ALT, AST and ALP were significantly reduced. Blood glucose levels dropped by 33.8 percent
Tang et al. [48]	L6 skeletal muscle cells	Various doses of COM seed extract	1 h	Glucose consumption and glycogen synthesis	At 12.5 g/ml, glucose intake and glycogen production was boosted
Umar et al. [49]	Hyperlipidaemic rats	Various doses of total leaves COM flavonoids Various doses of total fruit COM flavonoids	4 weeks	Lipid profile, Liver enzymes	Total flavonoids from COM fruit and leaves dramatically lowered TC, TG, LDL, ALT, AST, and MDA while considerably increasing HDL.
Mohebbi et al. [47]	Diabetic rats	Simvastatin (5 mg/kg) Various doses of COM fruit aqueous extract	28 days	Fasting blood glucose	A significant collapse in FBG in a dose-dependent manner SBP decreased significantly in a dose-dependent manner.
Ur rahman et al. [50]	Normotensive, high cholesterol, and glucose fed hypertensive rats	COM extract in 200, 400, and 600 mg/kg dose	21 days	SBP, DBP	COM extract in 600 mg/kg dose produced maximum effect and prevented a rise in blood pressure.
Lee et al. [45]	High-fat diet (HFD)-induced obese C57BL/6 mice	HFD + 50 mg/kg BW/day COM (COM 50) HFD + 100 mg/kg BW/day COM (COM 100) HFD + 200 mg/kg BW/day COM (COM 200)	8 weeks	Body weight, fat mass, adipose tissue weight, lipid profile, and leptin	COM showed an anti-obesity effect. COM improved blood HDL, TG, adiponectin levels, body weight, fat mass, adipose tissue weight, insulin resistance, and leptin levels.

COM: *Cydonia oblonga* Miller, ALT: alanine transaminase, AST: aspartate transaminase, ALP: alkaline phosphatase, FBG: fasting blood glucose, SBP: systolic blood pressure, DBP: diastolic blood pressure, TC: total cholesterol, LDL: low-density lipoprotein, HDL: high-density lipoprotein, TG: triglycerides.

way to green tea, the methanolic extract of quince leaves strongly protected the erythrocyte membrane against hemolysis [19]. The phenolic fraction of quince has been found to have stronger antioxidant activity than the entire methanolic extract [59]. According to Silva et al., organic acid extracts consistently had the lowest antiradical activity, showing that the phenolic component of quince fruit contributes significantly to its antioxidant ability [59]; however, Khademi et al. mentioned that the antioxidant properties of the quince leaf are most likely to be responsible for its antiatherosclerotic effects [43]. In study by Silva et al., methanolic extracts were tested for the antioxidant capacity, and peel extract was shown to have the highest antioxidant activity [59].

A study by Magalhães et al. reported that the DPPH free radical scavenging capabilities of pulp and peel extracts were comparable, whereas seed extract had a substantially lower antioxidant capacity. Pulp and peel extracts significantly protected the erythrocyte membrane from hemolysis under the oxidative action of AAPH in a time-dependent and concentration-dependent manner. Seed extracts alone caused significant hemolysis [60]. These findings imply that the quince leaf or fruit could be employed as a prophylactic or therapeutic agent in free radical-related diseases such as CVD.

Doxorubicin is one of the most regularly used medications to treat a wide range of cancers, and its most common adverse effects include anemia and cardiovascular toxicity. Doxorubicin increases malondialdehyde (MDA) while decreasing glutathione (GSH) and catalase activity in rat heart tissue. Two weeks of treatment with *Cydonia* hydroalcoholic extract significantly reduced MDA levels while raising GSH, demonstrating that *Cydonia* can relieve oxidative stress and minimize cardiac toxicity [61].

4.5. Inflammation. Zhou et al. in their study besides showing the significant antihypertensive effect of quince leaf extract also reported a considerable decrease in the contents of IL-6, IL-1 β , TNF- α , and CRP levels in total flavonoids of COM leaves [53]. It has been documented that chronic inflammation is a hallmark of atherosclerosis [23]. According to a study on a polyphenolic extract from Tunisian COM, it can prevent high levels of the proinflammatory cytokine TNF-, chemokine IL-8, and LPS-mediated activation of three main cellular proinflammatory effectors; p38MAPK, nuclear factor-kappa B, and Akt [23]. Reduced anti-inflammatory cytokine levels, such as CRP, together with lower LDL-C and TC levels, inhibited the development of atherosclerosis [62].

4.6. Other Factors. In a study on male rabbits, Khademi et al. discovered that high-cholesterol diet significantly elevated alkaline phosphatase (ALP), ALT, AST, Cr, and BUN. Biochemical markers; AST, ALT, and Cr levels were dramatically reduced when quince leaf extract (50 mg/kg) and atorvastatin 0.5 mg/kg were used in comparison to the control group; however, the difference was not significant [43]. Other studies also proved the positive effect of *Cydonia*

on liver function biomarkers (ALP, AST, and ALT) and kidney function biomarkers (Cr and BUN) [44, 46, 49]. However, histological analysis in their study revealed that atorvastatin and COM leaf extract were unable to prevent plaque accumulation in the coronary artery following plaque formation for 12 weeks.

Zhou et al. found that in comparison to aspirin (2.58), extract from *Cydonia* (20, 40, 80 mg/kg/day) prolonged bleeding times by 2.17, 2.78, and 3.63 and the clotting time by 1.44, 2.47, and 2.48 times, respectively [51]. *Cydonia* extract also reduced pulmonary emboli mortality, increased thrombolysis, decreased euglobulin lysis time, and increased the length of venous occlusion in rats compared to the aspirin group [51].

5. Conclusion

It can be concluded that *Cydonia oblonga* Miller (quince) has cardiovascular preventive properties. In animal studies, it regulates blood lipids, lowers blood pressure and glucose levels, and protects the liver and kidneys. It also helps with thrombosis and body weight and has anti-inflammatory and antioxidant properties.

Data Availability

No data were used to support the findings of the study.

Ethical Approval

This article does not feature any of the authors' human subject research.

Conflicts of Interest

All the authors declare that they have no conflicts of interest.

References

- [1] E. K. Kalra, "Nutraceutical--definition and introduction," *AAPS PharmSciTech*, vol. 5, no. 3, pp. 27-28, 2003.
- [2] V. Brower, "A nutraceutical a day may keep the doctor away," *EMBO Reports*, vol. 6, no. 8, pp. 708-711, 2005.
- [3] S. Basak and J. Gokhale, "Immunity boosting nutraceuticals: current trends and challenges," *Journal of Food Biochemistry*, vol. 46, no. 3, 2022.
- [4] S. Ansari, B. Chauhan, N. Kalam, and G. Kumar, "Current concepts and prospects of herbal nutraceutical: a review," *Journal of Advanced Pharmaceutical Technology & Research*, vol. 4, no. 1, pp. 4-8, 2013.
- [5] D. C. Mora, G. Overvag, M. C. Jong et al., "Complementary and alternative medicine modalities used to treat adverse effects of anti-cancer treatment among children and young adults: a systematic review and meta-analysis of randomized controlled trials," *BMC Complementary Medicine and Therapies*, vol. 22, no. 1, 2022.
- [6] G. K. Heshmat, N. Mashayekhiasi, A. Amerizadeh, Z. Teimouri Jervekani, and M. Sadeghi, "Effect of fenugreek consumption on serum lipid profile: a systematic review and meta analysis," *Phytotherapy Research*, vol. 34, no. 9, pp. 2230-2245, 2020.

- [7] I. Onakpoya, E. Spencer, C. Heneghan, and M. Thompson, "The effect of green tea on blood pressure and lipid profile: a systematic review and meta-analysis of randomized clinical trials," *Nutrition, Metabolism, and Cardiovascular Diseases*, vol. 24, no. 8, pp. 823–836, 2014.
- [8] S. Ehsani, H. Zolfaghari, S. Kazemi, and F. Shidfar, "Effects of sumac (*Rhus coriaria*) on lipid profile, leptin and steatosis in patients with non-alcoholic fatty liver disease: a randomized double-blind placebo-controlled trial," *Journal of Herbal Medicine*, vol. 31, Article ID 100525, 2022.
- [9] M. Sadeghi, S. Kabiri, A. Amerizadeh et al., "Anethum graveolens L.(Dill) effect on human lipid profile: an updated systematic Review," *Current Problems in Cardiology*, vol. 47, no. 11, Article ID 101072, 2022.
- [10] M. Akbari-Fakhrabadi, J. Heshmati, M. Sepidarkish, and F. Shidfar, "Effect of sumac (*Rhus Coriaria*) on blood lipids: a systematic review and meta-analysis," *Complementary Therapies in Medicine*, vol. 40, pp. 8–12, 2018.
- [11] M. Zubair, S. Mahmood Sajid, M. Waqas, M. Nawaz, and Z. Ahmad, "A review on quince (*Cydonia oblonga*)," *A Useful Medicinal Plant*, vol. 14, pp. 517–524, 2015.
- [12] A. Abliz, Q. Aji, E. Abdusalam et al., "Effect of *Cydonia oblonga* Mill. leaf extract on serum lipids and liver function in a rat model of hyperlipidaemia," *Journal of Ethnopharmacology*, vol. 151, no. 2, pp. 970–974, 2014.
- [13] K. Usmanghani, A. Saeed, and M. T. Alam, *Indusyunic Medicine: Traditional Medicine of Herbal Animal and Mineral Origin in Pakistan*, Department of Pharmacognosy, Faculty of Pharmacy, University of Karachi, Karachi, Sindh, Pakistan, 1997.
- [14] Y. Hamauzu, M. Irie, M. Kondo, and T. Fujita, "Antiulcerative properties of crude polyphenols and juice of apple, and chinese quince extracts," *Food Chemistry*, vol. 108, no. 2, pp. 488–495, 2008.
- [15] M. Minaiyan, A. Ghannadi, M. Etemad, and P. Mahzouni, "A study of the effects of *Cydonia oblonga* Miller (Quince) on TNBS-induced ulcerative colitis in rats," *Research in pharmaceutical sciences*, vol. 7, no. 2, pp. 103–110, 2012.
- [16] K. Nadkarni, *Indian Materia Medica: With Ayurvedic, Unani-Tibbi, Siddha, Allopathic, Homeopathic, Naturopathic And Home Remedies, Appendices And Indexes*, Ramdas Bhatkal, Popular Prakashan Private Ltd, Cumballa Hill, Mumbai, Maharashtra, 1976.
- [17] J. A. Duke, *Handbook of Medicinal Herbs*, CRC Press, FL, USA, 2002.
- [18] N. D. Prajapati, S. Purohit, A. K. Sharma, and T. Kumar, *A Handbook of Medicinal Plants: A Complete Source Book*, Agrobios, Sardarpura, Jodhpur, Rajasthan, 2003.
- [19] R. M. Costa, A. S. Magalhães, J. A. Pereira et al., "Evaluation of free radical-scavenging and antihemolytic activities of quince (*Cydonia oblonga*) leaf: a comparative study with green tea (*Camellia sinensis*)," *Food and Chemical Toxicology*, vol. 47, no. 4, pp. 860–865, 2009.
- [20] M. Carvalho, B. M. Silva, R. Silva, P. Valentao, P. B. Andrade, and M. L. Bastos, "First report on *Cydonia oblonga* Miller anticancer potential: differential antiproliferative effect against human kidney and colon cancer cells," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 6, pp. 3366–3370, 2010.
- [21] A. Jouyban, M. M. Shoja, M. R. Ardalan et al., "The effect of quince leaf decoction on renal injury induced by hypercholesterolemia in rabbits: a pilot study," *Journal of Medicinal Plants Research*, vol. 5, no. 21, pp. 5291–5295, 2011.
- [22] M. Aslan, N. Orhan, D. D. Orhan, and F. Ergun, "Hypoglycemic activity and antioxidant potential of some medicinal plants traditionally used in Turkey for diabetes," *Journal of Ethnopharmacology*, vol. 128, no. 2, pp. 384–389, 2010.
- [23] B. K. Essafi, A. Refai, I. Riahi, S. Fattouch, H. Karoui, and M. Essafi, "Quince (*Cydonia oblonga* Miller) peel polyphenols modulate LPS-induced inflammation in human THP-1-derived macrophages through NF- κ B, p38MAPK and Akt inhibition," *Biochemical and Biophysical Research Communications*, vol. 418, no. 1, pp. 180–185, 2012.
- [24] F. Shinomiya, Y. Hamauzu, and T. Kawahara, "Anti-allergic effect of a hot-water extract of quince (*Cydonia oblonga*)," *Bioscience, Biotechnology, and Biochemistry*, vol. 73, no. 8, pp. 1773–1775, 2009.
- [25] F. Z. Sakhri, N. Adachi, S. Zerizer et al., "Behavioral and neurological improvement by *Cydonia oblonga* fruit extract in chronic immobilization stress rats," *Phytotherapy Research*, vol. 35, no. 4, pp. 2074–2084, 2021.
- [26] M. U. Ashraf, G. Muhammad, M. A. Hussain, and S. N. A. Bukhari, "*Cydonia oblonga* M., A medicinal plant rich in phytonutrients for pharmaceuticals," *Frontiers in Pharmacology*, vol. 7, 2016.
- [27] B. K. Nandi, A. Goswami, and M. K. Purkait, "Adsorption characteristics of brilliant green dye on kaolin," *Journal of Hazardous Materials*, vol. 161, no. 1, pp. 387–395, 2009.
- [28] G. Arabaci and A. Usluoglu, "The enzymatic decolorization of textile dyes by the immobilized polyphenol oxidase from quince leaves," *The Scientific World Journal*, vol. 2014, pp. 1–5, 2014.
- [29] F. Hernández-García, *Quinces*, 2020.
- [30] S. Z. Hussain, B. Naseer, T. Qadri et al., "Quince (*Cydonia oblonga*) morphology, taxonomy, composition and health benefits," in *Fruits Grown in Highland Regions of the Himalayas: Nutritional and Health Benefits*, pp. 49–62, Springer International Publishing, Cham, Switzerland, 2021.
- [31] T. Erdoğan, T. Gönenç, Z. Hortoğlu, B. Demirci, K. Başer, and B. Kivçak, "Chemical composition of the essential oil of quince (*Cydonia Oblonga miller*) leaves," *Medicinal & Aromatic Plants*, vol. 01, no. 08, p. 134, 2012.
- [32] A. P. Oliveira, J. A. Pereira, P. B. Andrade, P. Valentão, R. M. Seabra, and B. M. Silva, "Phenolic profile of *Cydonia oblonga* Miller leaves," *Journal of Agricultural and Food Chemistry*, vol. 55, no. 19, pp. 7926–7930, 2007.
- [33] A. G. Osman, M. Koutb, and A. E. D. H. Sayed, "Use of hematological parameters to assess the efficiency of quince (*Cydonia oblonga miller*) leaf extract in alleviation of the effect of ultraviolet-A radiation on African catfish *Clarias gariepinus* (Burchell, 1822)," *Journal of Photochemistry and Photobiology B: Biology*, vol. 99, no. 1, pp. 1–8, 2010.
- [34] N. De Tommasi, F. De Simone, C. Pizza, and N. Mahmood, "New tetracyclic sesterterpenes from *Cydonia vulgaris*," *Journal of Natural Products*, vol. 59, no. 3, pp. 267–270, 1996.
- [35] A. Lutz-Röder, M. Schneider, and P. Winterhalter, "Isolation of two new ionone glucosides from quince (*Cydonia oblonga* Mill.) leaves," *Natural Product Letters*, vol. 16, no. 2, pp. 119–122, 2002.
- [36] O. Rop, J. Balik, V. Řezníček et al., *Chemical Characteristics of Fruits of Some Selected Quince (*Cydonia Oblonga* Mill.) Cultivars*, Czech Journal of Food Sciences, Prague, Czech, 2011.
- [37] N. Kirimer, Z. Tunalier, K. Can Başer, and I. Cingi, "Anti-spasmodic and spasmogenic effects of *Scolymus hispanicus* and taraxasteryl acetate on isolated ileum preparation," *Planta Medica*, vol. 63, no. 06, pp. 556–558, 1997.

- [38] S. Ammar, H. Edziri, M. A. Mahjoub, R. Chatter, A. Bouraoui, and Z. Mighri, "Spasmolytic and anti-inflammatory effects of constituents from *hertia cheirifolia*," *Phytomedicine*, vol. 16, no. 12, pp. 1156–1161, 2009.
- [39] R. Budriesi, P. Ioan, M. Micucci, E. Micucci, V. Limongelli, and A. Chiarini, "Stop Fitán: antispasmodic effect of natural extract of chestnut wood in Guinea pig ileum and proximal colon smooth muscle," *Journal of Medicinal Food*, vol. 13, no. 5, pp. 1104–1110, 2010.
- [40] M. Zakłós-Szyda and N. Pawlik, "Japanese quince (*Chaenomeles japonica* L.) fruit polyphenolic extract modulates carbohydrate metabolism in HepG2 cells via AMP-activated protein kinase," *Acta Biochimica Polonica*, vol. 65, no. 1, pp. 67–78, 2018.
- [41] N. Nagahora, Y. Ito, and T. Nagasawa, "Dietary Chinese quince polyphenols suppress generation of α -dicarbonyl compounds in diabetic KK-*A*' mice," *Journal of Agricultural and Food Chemistry*, vol. 61, no. 27, pp. 6629–6635, 2013.
- [42] S. Miraghaee, G. Bahrami, B. Izadi et al., "Antidiabetic potential of the isolated fractions from the plants of Rosaceae family in streptozotocin-induced diabetic rats," *Research in Pharmaceutical Sciences*, vol. 16, no. 5, pp. 505–515, 2021.
- [43] F. Khademi, B. Danesh, A. Delazar, D. Mohammad Nejad, M. Ghorbani, and J. Soleimani Rad, "Effects of quince leaf extract on biochemical markers and coronary histopathological changes in rabbits," *ARYA Atherosclerosis*, vol. 9, no. 4, pp. 223–231, 2013.
- [44] F. Khademi, B. Danesh, D. Mohammad Nejad, and J. Soleimani Rad, "The comparative effects of atorvastatin and quince leaf extract on atherosclerosis," *Iranian Red Crescent Medical Journal*, vol. 15, no. 8, pp. 639–643, 2013.
- [45] H. S. Lee, Y. E. Jeon, J. I. Jung et al., "Anti-obesity effect of *Cydonia oblonga* miller extract in high-fat diet-induced obese C57BL/6 mice," *Journal of Functional Foods*, vol. 89, Article ID 104945, 2022.
- [46] M. Mirmohammadlu, S. H. Hosseini, M. Kamalinejad, M. Esmaeili Gavvani, M. Noubarani, and M. R. Eskandari, "Hypolipidemic, hepatoprotective and renoprotective effects of *Cydonia oblonga* mill. Fruit in streptozotocin-induced diabetic rats," *Iranian Journal of Pharmaceutical Research*, vol. 14, no. 4, pp. 1207–1214, 2015.
- [47] S. Mohebbi, M. Naserkheil, M. Kamalinejad et al., "Anti-hyperglycemic activity of quince (*Cydonia oblonga* Mill.) fruit extract and its fractions in the rat model of diabetes," *International Pharmacy Acta*, vol. 2, no. 1, pp. 1–8, 2019.
- [48] D. Tang, L. Xie, X. Xin, and H. Aisa, "Anti-diabetic action of *Cydonia oblonga* seed extract: improvement of glucose metabolism via activation of PI3K/AKT signaling pathway," *Journal of Pharmacognosy and Phytochemistry*, vol. 4, no. 2, pp. 7–13, 2016.
- [49] A. Umar, G. Iskandar, A. Aikemu et al., "Effects of *Cydonia oblonga* miller leaf and fruit flavonoids on blood lipids and anti-oxidant potential in hyperlipidemia rats," *Journal of Ethnopharmacology*, vol. 169, pp. 239–243, 2015.
- [50] M. S. Ur Rahman, M. Saleem, M. Alamgeer et al., "Antihypertensive and safety studies of *Cydonia oblonga* Miller," *Pakistan Journal of Pharmaceutical Sciences*, vol. 34, pp. 687–691, 2021.
- [51] W. Zhou, A. Abdurahman, A. Umar et al., "Effects of *Cydonia oblonga* miller extracts on blood hemostasis, coagulation and fibrinolysis in mice, and experimental thrombosis in rats," *Journal of Ethnopharmacology*, vol. 154, no. 1, pp. 163–9, 2014.
- [52] W. T. Zhou, A. Abdurahman, E. Abdusalam et al., "Effect of *Cydonia oblonga* miller. leaf extracts or captopril on blood pressure and related biomarkers in renal hypertensive rats," *Journal of Ethnopharmacology*, vol. 153, no. 3, pp. 635–640, 2014.
- [53] W. T. Zhou, W. L Y Yiming, H. Ma, G. Mamat, and A. Umar, "Anti-hypertensive effect of total flavonoids of *Cydonia oblonga* leaves and its mechanism based on anti-inflammatory function," *Zhong Yao Cai*, vol. 38, no. 10, pp. 2134–2138, 2015.
- [54] M. Schieber and N. S. Chandel, "ROS function in redox signaling and oxidative stress," *Current Biology*, vol. 24, no. 10, pp. 453–462, 2014.
- [55] D. Moris, M. Spartalis, E. Spartalis et al., "The role of reactive oxygen species in the pathophysiology of cardiovascular diseases and the clinical significance of myocardial redox," *Annals of Translational Medicine*, vol. 5, no. 16, 2017.
- [56] P. C. Schenkel, A. M. V. Tavares, R. O. Fernandes et al., "Redox-sensitive prosurvival and proapoptotic protein expression in the myocardial remodeling post-infarction in rats," *Molecular and Cellular Biochemistry*, vol. 341, no. 1–2, pp. 1–8, 2010.
- [57] S. Sut, S. Dall'Acqua, G. Poloniato, F. Maggi, and M. Malagoli, "Preliminary evaluation of quince (*Cydonia oblonga* Mill.) fruit as extraction source of antioxidant phytoconstituents for nutraceutical and functional food applications," *Journal of the Science of Food and Agriculture*, vol. 99, no. 3, pp. 1046–1054, 2019.
- [58] M. N. Cha, H. J. Kim, B. G. Kim, and J. H. Ahn, "Synthesis of chlorogenic acid and p-coumaroyl shikimates from glucose using engineered *Escherichia coli*," *Journal of Microbiology and Biotechnology*, vol. 24, no. 8, pp. 1109–1117, 2014.
- [59] B. M. Silva, P. B. Andrade, P. Valentão, F. Ferreres, R. M. Seabra, and M. A. Ferreira, "Quince (*Cydonia oblonga* Miller) fruit (pulp, peel, and seed) and Jam: antioxidant activity," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 15, pp. 4705–4712, 2004.
- [60] A. S. Magalhães, B. M. Silva, J. A. Pereira, P. B. Andrade, P. Valentão, and M. Carvalho, "Protective effect of quince (*Cydonia oblonga* Miller) fruit against oxidative hemolysis of human erythrocytes," *Food and Chemical Toxicology*, vol. 47, no. 6, pp. 1372–1377, 2009.
- [61] F. Fallahi, M. Roghani, and I. Ansari, "The effect of hydroalcoholic extract of *Cydonia oblonga* Miller leaf on doxorubicin-induced cardiac injury in rat," *Journal of Basic and Clinical Pathophysiology*, vol. 6, no. 2, pp. 7–12, 2018.
- [62] M. Setorki, S. Asgary, A. Eidi, A. H. Rohani, and N. Esmaeil, "Effects of apple juice on risk factors of lipid profile, inflammation and coagulation, endothelial markers and atherosclerotic lesions in high cholesterolemic rabbits," *Lipids in Health and Disease*, vol. 8, no. 1, p. 39, 2009.