Review Article

Research Progress on the Pharmacological Action of Schisantherin A

Zehao Xiao,¹ Wen Xiao,² and Guilin Li³

¹Queen Mary School, Medical College of Nanchang University, Nanchang 330006, China

²Department of Pathology, Medical College of Nanchang University, Nanchang 330006, China

³Department of Physiology, Medical College of Nanchang University, Nanchang 330006, China

Correspondence should be addressed to Guilin Li; liguilin@ncu.edu.cn

Received 16 August 2021; Revised 23 October 2021; Accepted 21 January 2022; Published 12 February 2022

Academic Editor: Md. Areeful Haque

Copyright © 2022 Zehao Xiao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Schisantherin A (Sch A) is a dibenzocyclooctadiene lignan monomer isolated from the fruit of *Schisandra chinensis* (Turcz.) Baill. (*S. chinensis*). At present, many studies have shown that Sch A has a wide range of pharmacological effects, including its anti-Parkinson and anti-inflammatory effects and ability to protect the liver, protect against ischemia-reperfusion (I/R) injury, suppress osteoclast formation, and improve learning and memory. Its mechanism may be related to the antioxidant, anti-inflammatory, and antiapoptotic properties of Sch A through the MAPK, NF- κ B, AKT/GSK3 β , and PI3K/AKT pathways. This is the first review of the recent studies on the pharmacological mechanism of Sch A.

1. Introduction

Natural products play a very important role in drug discovery and development. Many kinds of small-molecule drugs have been isolated from plant sources. *Schisandra chinensis* (Turcz.) Baill. (*S. chinensis*) is a traditional Chinese medicine (TCM) in the Angiosperm Phylogeny Group (APG)-II system, in which Schisandraceae and Magnoliaceae are placed in the same category as Austrobaileyales. *S. chinensis* has hepatoprotective, neuroprotective, anti-inflammatory, antiviral, antioxidative, anticancer, detoxification, antiapoptosis, and immunostimulant activities [1–4]. *S. chinensis* has been used in the treatment of many diseases, such as gastrointestinal diseases, cardiovascular diseases, bodily fatigue and weakness, respiratory failure, excessive sweating, and insomnia [2, 5]. Clinical studies over the last ten years have indicated that *S. chinensis* can act on the nervous system, reduce fatigue, and treat palpitation and hot flashes [3, 6], and *S. chinensis* was suggested to be used to improve and treat cognitive impairment in models of neurodegenerative diseases and Alzheimer's disease [7]. The ability to resist oxidative stress in the liver is also evident. *S. Chinensis* can significantly

improve the antioxidant capacity of the body, reduce oxidative active substances, and effectively improve liver function in patients [8–10]. *S. chinensis* can also be used in combination with tacrolimus to protect the liver and reduce tacrolimus requirements in the treatment of transplant patients [11, 12]. Additionally, clinical studies have shown that the extract of *S. chinensis* can be used as a health supplement, has beneficial effects on muscle strength and lactic acid levels [13], and can also be used to regulate the composition of the intestinal flora [14].

The biological activity and pharmacological use of *S. chinensis* are related to dibenzocyclooctadiene lignans. Schisantherin A (Sch A), a dibenzocyclooctadiene lignan monomer and primary biologically active lignan also known as gomisin C, schizandrin A, and wuweizi ester A, is isolated from the fruit of *S. chinensis*. Its molecular weight is 536.57 g/mol, its molecular formula is C₃₀H₃₂O₉, and its chemical structure is shown in Figure 1. Sch A is highly soluble in benzene, chloroform, and acetone; soluble in methanol and ethanol; and insoluble in petroleum ether and water. The structure of Sch A contains a methylenedioxy group, methyl groups, methoxy groups, a benzoyloxy group, and a hydroxyl group. Therefore, the loss of C₂H₂O₂, benzoic

FIGURE 1: Chemical structural formula and pharmacological effects of schisantherin A.

acid, H₂O, methyl, and methoxy groups are the main and natural metabolic pathways.

Sch A has the highest content and the most effective advantage in the lignan extract of *S. chinensis* [15, 16]. In a 2019 study, the metabolic profiles of Sch A *in vitro* (rat liver microsomes) and *in vivo* (rat plasma, urine, and bile) were assessed by ultra-high-performance liquid chromatography-quadrupole time-of-flight tandem mass spectrometry (UPLC-Q-TOF-MS/MS) [17]. Sch A was found to be more complex *in vivo* than *in vitro* and undergo both hepatic and extrahepatic metabolic pathways in its biotransformation. The metabolic process mainly involves the loss of C₇H₄O and C₇H₄O₂, as well as further oxidative transformation, carboxylic acid reaction, and the loss of CH₂ and CH₂O groups. Moreover, the coupling of Sch A with glucuronic acid, sulfate, taurine, glutathione, and glucose may also account for its multiple effects. This study showed that the methoxy group and biphenyl cyclooctene are the main metabolic sites in Sch A. This was also demonstrated in a trial in which the efficacy of Sch A and other major lignans in the treatment of Parkinson's disease (PD) was compared [16]. Sch A has a greater neuroprotective function than schizandrin A and schizandrol A, which have no methylenedioxy groups, and this activity may be enhanced by the addition of a benzoyloxy substituent on the cyclooctadiene ring. This may be because the benzoyloxy and methoxy groups in Sch A inhibit the production of 1-methyl-4-phenylpyridinium ion (MPP⁺), similar structures to which are also present in the antipsychotic drugs sulpiride and amishupiride [18].

1.1. Extraction Technology. Lignans are traditionally separated by similar procedures. Ground seeds are extracted with ethanol/water or by using heat reflux [19, 20] and

Soxhlet [19]. At present, most Sch A is extracted by ethanol extraction and heating. However, the extraction efficiency is low due to the long heating time and the instability of Sch A under heat [19]. Additionally, these methods cannot avoid the problem of the excessive consumption of medicinal materials and are labor-intensive, time-consuming, and harmful to human health and the environment. Therefore, new extraction methods, such as microwave-assisted extraction (MAE) [21] and supercritical CO₂ fluid extraction (SFE) [22], have been developed. However, due to the high cost of these technologies, which limits their large-scale application, the industrial applications of this technology are still limited. Currently, an extraction method consisting of an aqueous two-phase system coupled with ultrasound is also being used [15]. Three steps are needed: solid-liquid extraction, liquid-liquid micro-extraction, and ultrasonic extraction. The main principle of the application of ultrasonic extraction technology is the use of the ultrasonic cavitation effect to accelerate the dissolution of the active ingredients of plant cell membranes, allowing the solvent to enter the cells, releasing the medicinal compounds, and the secondary effect of the ultrasonic wave, such as that upon mechanical vibration, emulsification, diffusion, crushing, chemistry, and so on, can also accelerate the diffusion of the extract released with the solvent mixture. Compared with the traditional solvent extraction method, this method has the advantages of a shorter extraction time, no need for heating, good suitability for extracting substances, and a higher target components extraction rate, and it can reduce the required amount of solvent and is conducive to further refinement and purification. Under optimal conditions with a solvent-to-solid ratio of 20:1, ultrasonic power of 800 W, and extraction time of 61.1 min, 13.10 mg/g schizandrin, 1.87 mg/g Sch A, and 1.84 mg/g schizandrin could be

extracted [15]. As a new extraction technology for TCM, ultrasonic extraction has good application prospects.

Recently, increasing amounts of data have shown that Sch A has broad pharmacological functions, including its antioxidant, sedative [23], anti-Parkinson, and anti-inflammatory effects and ability to protect the liver and protect against ischemia-reperfusion (I/R) injury, suppress osteoclast formation, and improve learning and memory. This summary aims to present the pharmacological effects and mechanisms of Sch A and provides a theoretical reference for future research.

2. The Role of Sch A in Diseases

2.1. Sch A against PD. PD, one of the most common neurodegenerative diseases, is characterized by degeneration of dopaminergic neurons in the substantia nigra with the formation of Lewy bodies. Previous studies have suggested that PD is a movement disorder as it mainly manifests by movement symptoms such as rigidity, trembling, and/or postural gait instability [24].

The pathological processes of PD include oxidative stress, impairment of the antioxidant response, mitochondrial dysfunction, and the accumulation of α -synuclein aggregates. These changes are known to aggravate damage to dopaminergic neurons [25].

In recent years, modern medical research has shown that the pathogenesis of PD is complex and involves multiple factors. Past experiments have shown that the pathological changes in PD involve not only dopaminergic neurons in the substantia nigra and striatum system but also involve other parts of the dopaminergic system and the 5-hydroxytryptamergic, norepinephrinergic, and cholinergic systems, as well as the dynorphin, enkephalin, and other peptide transmitter systems [26–28].

Sch A has the potential to become an oral drug for the treatment of PD and to provide a new way to improve the bioavailability of PD drugs. Sch A was reported to act against PD through *in vitro* and *in vivo* experiments. In a PD mouse model, Sch A has been widely used to protect against the loss of TH-positive dopaminergic neurons induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP). Among the five dibenzocyclooctadiene lignans of *Schisandra* (schizandrin A, schizandrin B, schizandrin C, schizandrol A, and Sch A), Sch A presented the strongest neuroprotective activity. This suggests that Sch A is a key component of dibenzocyclooctadiene lignans with neuroprotective activity. The results also indicate that unique positions and constituents on different chemical arms (including methylenedioxy, methoxy, and benzoyloxy groups) in the dibenzocyclooctadiene lignan may play an important role in the neuroprotective activity of Sch A. *In vitro* experiments showed that Sch A protected SH-SY5Y cells and significantly decreased the cytotoxicity induced by MPP⁺. The possible molecular mechanisms are related to two pathways: the phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K)/protein kinase B (Akt) signaling and cAMP-response-element-binding protein-mediated B-cell lymphoma 2 (Bcl2). It is well known that PI3K/Akt signaling promotes cell

survival. Sch A could increase the phosphorylation level of Akt and rescue the decrease in Akt phosphorylation induced by MPP⁺. Furthermore, Sch A could improve cell viability, which was inhibited by MPP⁺, while Akt inhibitor IV, a PI3K/Akt inhibitor, could abolish the effect of Sch A. Thus, it can be inferred that the neuroprotective effects of Sch A are exerted through the PI3K/Akt pathway. The ratio of Bcl2-associated X (Bax)/Bcl2 is one of the major biomarkers used to evaluate MPP⁺-induced apoptosis. Sch A decreased the Bax/Bcl2 ratio by upregulating Bcl2 expression, which was reduced by MPP⁺. The inhibitory effect of Sch A on apoptosis was closely related to the phosphorylation of CREB upstream of Bcl2. Therefore, the neuroprotective effect of Sch A is also related to CREB-mediated Bcl2 [16].

To improve the poor water solubility of Sch A, Chen et al. formulated it as nanocrystals. They found that Sch A concentration was considerably increased in the brain after oral administration of the nanocrystals. Sch A nanocrystals remarkably improved swimming behavior defects induced by MPTP and significantly decreased the death of dopaminergic neurons and locomotion deficiency induced by MPTP in zebrafish. The authors also found that activation of the Akt/glycogen synthase kinase-3 β (Gsk3 β) pathway may be involved in the strong neuroprotective effects of Sch A [29].

In addition, many studies have provided evidence that the formulation of Sch A in nanoemulsion can significantly increase the absolute bioavailability of Sch A from 4.3% to 47.3% [30], and its formulation in small methoxy poly (ethylene glycol) (mPEG)-poly (lactic-co-glycolic acid) (PLGA) nanoparticles improved the bioactivity, oral bioavailability, cross-barrier transportation, and brain uptake of Sch A [31].

Other studies showed that Sch A protected neurons from toxicity induced by glutamate [32] and dopaminergic neuron damage and cytotoxicity induced by 6-hydroxydopamine (6-OHDA) [33] in zebrafish models and human SH-SY5Y neuroblastoma cells. In zebrafish, Sch A prevented 6-OHDA-induced dopaminergic neuron loss. In 6-OHDA-treated SH-SY5Y cells, Sch A improved intracellular reactive oxygen species (ROS) accumulation and reduced nitric oxide (NO) overproduction by inhibiting the expression of inducible nitric oxide synthase (iNOS). Sch A also repressed the activation of GSK3 β , mitogen-activated protein kinases (MAPKs), and PI3K/Akt signaling pathway members [33].

2.2. Sch A Improves Learning and Memory. Over the past century, the treatment of disease has significantly increased life expectancy. However, age-related cognitive decline has become one of the greatest threats to the health of elderly individuals [34]. In the aging process, a decline in learning and memory is a prominent manifestation of aging of the central nervous system. Therefore, the development of drugs with auxiliary memory improvement functions will play an important role in improving memory and the quality of life of elderly individuals.

Sch A has been suggested to protect against oxidative stress, neurodegeneration, and cognitive impairment, and

Sch A is a potential drug for the treatment of Alzheimer's disease [35]. Studies have shown that successive intracerebroventricular administration of Sch A (0.01 and 0.1 mg/kg) for 5 days notably improved A β 1–42-induced cognitive deficits. Sch A increased spontaneous alternation behavior (examined by the Y-maze) and the number of avoidances (tested by the shuttle-box) and significantly reduced the escape latency (measured by the Morris water maze test). Additionally, Sch A restored the activities of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) and the levels of A β 1–42, glutathione (GSH), and malondialdehyde (MDA) in the hippocampus and cerebral cortex. These effects of Sch A might be related to its antioxidant capacity [36].

Sch A also drastically improved D-galactose-induced learning and memory impairment in mice. Furthermore, Sch A significantly improved learning and memory behavior. For example, Sch A increased the latency to enter a dark compartment, reduced the number of errors (examined by the step-through test) and shortened the time required to find the platform, and reduced the number of platform passes (as measured by the Morris water maze test). Sch A also increased SOD activity, decreased MDA content in both the hippocampus and peripheral blood of mice, and decreased the 8 hydroxy deoxyguanosine (8-OHdG) content in the hippocampus of the mice. Moreover, Sch A reduced the expression levels of p19, p53, and p21 and elevated the expression levels of cyclin D1 and CDK4 and retinoblastoma protein (Rb) phosphorylation in the hippocampus [37].

2.3. Anti-Inflammatory Effects of Sch A in Alcoholic Liver Disease, Acute Lung Injury, and Osteoarthritis. Long-term heavy drinking has been suggested to cause alcoholic liver disease. Many influencing factors, such as sex (women are more susceptible) [38], genetics [39], malnutrition, infection with hepatitis c virus (HCV) [40], obesity, and smoking [41], affect an alcoholic liver disease.

At present, Western medicine provides nutritional support, especially branched-chain amino acids, to treat alcoholic liver disease [42]. Other compounds have also been proven to be useful, such as glutathione, cysteine, tiopronin, metadoxine [43], S-adenosylmethionine, polyene phosphatidyl choline [5, 44], compound glycyrrhizin, silymarin [45], taurine [46, 47], and glucocorticoid, and if all else fails, liver transplantation is performed [48, 49]. However, their curative effect is not obvious, and many drugs have serious adverse reactions and complications.

In an alcohol-induced liver injury mouse model treated with Sch A, aspartate aminotransferase (AST), alcohol dehydrogenase, and alanine transaminase (ALT) activities and vesicular steatosis were significantly lower than those in the control group, suggesting that Sch A had alleviated alcohol-induced pathologic changes in the liver and improved liver function. Sch A decreased the activity of alcohol dehydrogenase (ADH), increased the activity of acetaldehyde dehydrogenase (ALDH), and reversed the metabolic imbalance induced by alcohol. The elevated MDA level and myeloperoxidase (MPO) activity and reduced activities of GSH

activity induced by alcohol were strongly abrogated by the administration of Sch A. Sch A also suppressed the microsomal ethanol oxidizing system and alcohol-induced activation of the nuclear factor-kappa B (NF- κ B) pathway. The hepatoprotective effects of Sch A may be related to its regulation of alcohol metabolism and inhibition of the NF- κ B pathway [50].

The pathogenesis of acute lung injury (ALI) is attributed to an excessive inflammatory response leading to the destruction of lung tissue, which is manifested as tumor necrosis factor-necrosis alpha (TNF- α) induction by various cytokines, the increased expression of interleukin-1 (IL-1), and an inflammatory cascade mediated by cytokines, which aggravates lung tissue injury. Moreover, a more severe form of ALI called acute respiratory distress syndrome (ARDS) can occur. Therefore, inhibiting inflammatory overreaction is an effective way to prevent and treat ALI/ARDS [31, 51].

Many studies have shown the anti-inflammatory functions of Sch A. When mice with ALI and ARDS were treated with Sch A, the levels of TNF- α , IL-6, and NF- κ B decreased. Sch A may also play a role in treating ALI/ARDS by blocking the NF- κ B pathway [52, 53].

Osteoarthritis (OA), a chronic degenerative disease characterized by articular cartilage destruction, synovial membrane inflammation, joint space narrowing, osteophytosis, and subchondral bone remodeling with joint pain, swelling, deformity, and movement disorder, is often found in weight-bearing joints such as the knees and hips [54–56].

The onset of OA is closely related to imbalance between the synthesis and degradation of the extracellular matrix in joints caused by the accumulation of senescent chondrocytes, the degradation of articular cartilage, and persistent pathological inflammation [57]. Therefore, inhibiting the inflammatory response in OA and protecting the integrity of articular cartilage may improve the pathological changes that occur in OA and relieve the pain of OA patients. Sch A exerted a dose-dependent anti-inflammatory effect by inhibiting the production of NO, prostaglandin E2 (PGE2), and TNF- α induced by IL-1 β in human OA chondrocytes [58]. In a lipopolysaccharide (LPS)-induced ARDS mouse model, pretreatment with Sch A significantly improved LPS-induced histopathological changes and reduced the levels of IL-1 β , IL-6, and TNF- α in bronchoalveolar lavage fluid [53]. The release of proinflammatory cytokines affects the expression of various molecules and eventually leads to neutrophil aggregation and lymphocyte activation [59–61]. The reduction in neutrophil infiltration in an MPO immunofluorescence assay further confirmed the anti-inflammatory activity of Sch A.

2.4. Sch A against Cholestasis. Cholestasis is a deficiency of bile acid formation and flow, leading to residual toxic bile acids in the liver. Intrahepatic cholestasis can cause hepatobiliary damage due to the accumulation of excess bile acids throughout the body and liver. Without proper treatment, long-term cholestasis can cause symptoms such as itching, osteoporosis, fatigue, jaundice, and hypercholesterolemia and lead to liver fibrosis and cirrhosis [62–64].

Thus, the treatment of cholestasis urgently needs to be improved.

Ursodeoxycholic acid (UDCA), an anti-inflammatory agent that reduces bile salt toxicity, is the only Food and Drug Administration (FDA)-approved treatment for cholestasis, but UDCA is not used for all patients, as it can cause severe abnormal alkaline phosphatase levels in some cases, which may lead to death [65–67].

Moreover, obeticholic acid has the potential to treat cholestasis. On the one hand, obeticholic acid improved cholestasis by decreasing the alkaline phosphatase content and total bilirubin level, but on the other hand, it was more prone to pruritus compared with placebo [68]. Therefore, it is necessary to develop new drugs with fewer adverse events as soon as possible.

An interesting study has shown that *Schisandra sphenanthera*, including Sch A, can activate pregnane X receptor (PXR) in mice, which increases the expression of UDP glucuronosyltransferase family 1 member A1 (UGT1A1) and the cytochrome P450 (CYP) family members CYP2B10 and CYP3A11 [69]. As a bile acid homeostasis regulator, PXR can modulate metabolism (Cyp3a11, Ugt1a1), transport (Oatp2, Mrp3), biosynthesis (Cyp7a1), and the excretion of xenobiotics and bile acids *in vivo* [70–72]. In addition, a dual-luciferase reporter gene assay was used to study the transactivation effect of lignans on the human pregnane X receptor (hPXR), and the results showed a similar effect [73]. These analyses revealed that Sch A can protect against lithocholic acid (LCA)-induced cholestatic liver injury.

2.5. Sch A against I/R Injury. I/R injury, which consists of two stages, ischemia and reperfusion injury, occurs after the restoration of oxygen delivery to a hypoxic organ [74, 75]. The liver is one of the organs most injured by I/R [76]. To illustrate, in the early stage of hepatic I/R, due to increased blood viscosity, a large number of white blood cells activate and block blood vessels, accompanied by vasoconstriction, and endothelial cell swelling, microcirculation disorders, and cell ischemia and hypoxia are further aggravated [74, 77, 78]. Eventually, this can lead to liver failure and multiple organ dysfunction syndromes (MODS) [79].

The protective effect of Sch A against hepatic I/R injury in mice was studied. After hepatic reperfusion, Sch A pretreatment remarkably restored liver function, as exhibited by limited histological damage; decreased levels of serum ALT, AST, and lactate dehydrogenase (LDH); and reduced neutrophil infiltration. Pretreatment with Sch A attenuated the expression of proinflammatory cytokines, such as IL-1 β , IL-6, and TNF- α , in an I/R-induced inflammatory state. The results showed that the protective effect of Sch A was related to a decrease in oxidative/nitrosative stress, as characterized by a decrease in MDA production and total NO content and a high GSH/glutathione disulfide (GSSG) ratio. Moreover, ischemic mice pretreated with Sch A showed a similar increase in adenosine triphosphate (ATP) content in the liver. These findings suggest that Sch A improves mitochondrial function in hepatocytes by alleviating oxidative/nitrosative stress. Sch A pretreatment not only reduced I/R-induced

oxidative stress and inflammation but also remarkably decreased hepatocyte apoptosis, which was evident in the diminished number of terminal deoxynucleotidyl-transferase-mediated dUTP Nick end-labeling (TUNEL)-positive hepatocytes and reductions in the apoptosis-related proteins Bax, Bcl2, and cleaved caspase-3. Sch A pretreatment reversed the upregulated phosphorylation of c-Jun N-terminal kinase (JNK), p38, and extracellular regulated protein kinases (ERKs) induced by hepatic I/R injury, suggesting that Sch A plays a role in protecting against I/R-induced liver injury by hindering activation of the MAPK pathway. Therefore, Sch A pretreatment is a promising method for hepatic protection in I/R-induced pathological environments, such as blood occlusion during liver resection and liver transplantation [80]. Ischemic heart disease is one of the most common cardiovascular diseases in clinical practice. Reperfusion of ischemic coronary arteries can effectively reduce total mortality [81]; however, myocardial ischemia-reperfusion injury (MIRI) occurs after coronary reperfusion. MIRI includes microvascular injury, arrhythmia, reversible cardiac dysfunction, and myocyte apoptosis. Various mechanisms that include reperfusion injury, intracellular Ca²⁺ overload, oxidative stress, and inflammatory factors are involved in changes in myocardial structure and function [82].

For the treatment of I/R injury, compared with the Western medicine reperfusion therapy, which exhibits limitations, the advantages of TCM are particularly prominent; for instance, TCM supports the use of multiple approaches and multiple targets. TCM compounds and related ingredients can alleviate MIRI injury through a variety of mechanisms, and remote ischemic treatment may have a higher clinical value than ischemic preconditioning (IPC) and ischemic postconditioning (IPOC) [83, 84]. Chang et al. studied the protective effect of Sch A against MIRI. The results showed that Sch A has a strong protective effect on cardiac function by reducing the left ventricular systolic pressure (LVSP), left ventricular end-diastolic pressure (LVEDP), and occurrence of arrhythmia. Furthermore, Sch A significantly reduced the infarction area and release of MDA. However, SOD activity increased, and the changes in myocardial histopathology were significantly alleviated, showing the strong protective effect of Sch A against myocardial injury. In addition, Sch A could significantly reduce the apoptosis of myocardial cells and caspase-3 activity, showing a strong protective effect. These results illustrate that Sch A has a protective effect against MIRI [85].

Inflammation and oxidative stress are key to acute and long-term nerve tissue injury after cerebral ischemic stroke [86]. After cerebral ischemic stroke, blood flow in the local brain tissue is reduced, and disorder in the tissue blood supply leads to injury and the death of neurons at the damaged site and in the cerebral cortex, as well as the impairment of synaptic structure and conduction function [87]. Timely repair of the damaged neuronal structure and prevention of the loss of neuronal function is important for the treatment of this disease. Studies have shown that both autophagy and apoptosis signals in neurons can be activated after cerebral I/R. Excessive cellular autophagy will cause

TABLE 1: Summary table for disease, study model, route of administration, dose, related signaling pathways, genes, and therapeutics outcome of Sch A.

Disease	Study model	Route of administration	Dose	Pathways and related genes	Therapeutics outcome	References
Parkinson's disease	Human neuroblastoma SH-SY5Y cells	Medium	3–25 μ M	PI3K/Akt	Protect against MPP+-induced cell damage in a concentration-dependent manner	[16]
	Male C57BL/6 mice	Intragastric gavage	30, 100, 300 mg/kg/d, 14 d			
	Male SD mice	Intragastric gavage	SA-NC 4 mg/kg		Increase SA oral absorption	[29]
	Zebrafish embryo	Embryo medium	SA-NC 1, 3, 10 μ M		Attenuated MPTP-induced neurotoxicity on zebrafish	
	Human neuroblastoma SH-SY5Y	Medium	SA-NC 30 μ M	Akt/Gsk3 β	Blocked the MPP+-induced repression of p-Gsk3 β	
	Cortical cells of fetal SD mice	Medium	0.1, 1.0, 5.0 μ M		Possess therapeutic potential against oxidative neuronal damage induced by excitotoxins	[32]
	Human neuroblastoma SH-SY5Y	Medium	3, 6, 12, 25, 50, 100 μ g/ml	ROS/NO AKT/GSK3 β		[33]
Alzheimer's disease	A β 1–42-induced learning and memory impairment in male 12-week-old KM mice	Intragastric gavage	0.01, 0.1 mg/kg for 5 days		Protect against cognitive deficits, oxidative stress, and neurodegeneration	[24, 36]
	D-galactose-induced learning and memory impairment in male ICR mice	Orally	1.25, 2.50, and 5.00 mg/kg for 42 days.	p19/p53/p21/cyclinD1/CDK4 genes	Antioxidation	
Alcohol-induced liver injury	Alcohol-containing diet-induced liver injury in eight-week-old male C57BL/6 mice	Oral gavage	100 mg/kg and 200 mg/kg	NF- κ B	Anti-inflammation and antioxidation	[39, 50]
Acute lung injury	Lipopolysaccharide-induced acute lung injury in male BALB/c mice	Intragastric gavage	25, 50 mg/kg, and 40 mg/kg	NF- κ B and MAPK	Anti-inflammation	[41, 42, 52, 53]
Osteoarthritis	Primary chondrocytes from articular cartilage samples	Medium	9.5, 19, 38 μ M	NF- κ B and MAPK	Anti-inflammation	
Cholestasis	Lithocholic acid-induced cholestasis in adult male C57BL/6J mice	Oral gavage	100 mg/kg,	hPXR-regulate genes, CYP3A4, UGT1A1 and OATP2.	Hepatoprotection and detoxification and efflux of toxic bile acids	[58, 69]
Liver ischemia-reperfusion injury	70% hepatic warm ischemia was used in male C57BL/6 mice	Gavage	200 mg/kg/day gavage for 5 days,	MAPK	Anti-inflammation, antioxidation against apoptosis	[69, 80]
Myocardial ischemia-reperfusion injury (MIRI)	MIRI in male Wistar rats by occluding the left coronary artery for 45 min followed by 2 h of reperfusion	Microemulsion injection	40 mmol/kg	Bcl2/Bax/Caspase-3	Inhibiting cardiomyocyte apoptosis	[74, 85]

TABLE 1: Continued.

Disease	Study model	Route of administration	Dose	Pathways and related genes	Therapeutics outcome	References
Cerebral ischemic stroke	I/R-induced neuronal injury in SD rat embryos in the hypoxic chamber	Medium	1.25, 2.5, and 5 μ g/ml	TLR4 and C5aR1	Alleviated neurological deficits, reduced infarct volume, and attenuated oxidation stress, inflammation, and apoptosis in the ischemic parietal cortex of rats after MCAO/R injury	[79, 90]
Acute kidney injury	Hypoxia/reoxygenation injury in renal tubular epithelial cells by incubating in the hypoxic chamber	Medium	5, 10, 20 μ M	PI3K/Akt	Attenuated oxidation stress, cell apoptosis, and inflammation	[84, 95]
Ulcerative colitis	HCT116 cells	Medium	1 μ g/ml		Inhibits H ₂ O ₂ -induced apoptotic cell death in intestinal epithelial cells	[97]

cells to undergo autophagic death [88], while moderate autophagy activation is conducive to the survival of neurons and induces neuroprotection [89].

Sch A may be an effective neuroprotective agent with antioxidant, anti-inflammatory, and antiapoptotic effects on ischemia and related diseases. Shi et al. assessed the protective effect of SCA on cerebral I/R injury and explored the mechanisms related to Toll-like receptor 4 (TLR4) and complement C5a receptor (C5aR1, CD88) signaling pathways *in vitro* and *in vivo*. The results showed the potent neuroprotective effects of Sch A, which reduced oxygen and glucose deprivation and the reperfusion injury-induced apoptosis of primary cultured cortical neurons. It restored homeostasis through different features, such as inflammation, oxidative stress, and apoptosis, by inhibiting the TLR4 and C5aR1 signaling pathways [90]. Furthermore, the activation of C5aR1, a G-protein-coupled receptor highly expressed in neutrophils, can aggravate acute and chronic inflammation [91]. The NF- κ B signaling pathway can be activated by TLR4 and TNFR1, which induces high cytokine expression levels [92, 93].

With the increasing incidence of kidney transplantation and partial nephrectomy, renal I/R injury is difficult to avoid. Blood reperfusion, the main factor leading to kidney injury, can cause acute tubular necrosis, induce renal failure, affect the recovery of renal function after surgery, and in severe cases endanger patient survival [94]. Gong and Wang examined the effect of Sch A on renal I/R injury *in vitro*. Sch A decreased the levels of MDA and ROS and reduced the production of TNF- α , IL-1 β , and IL-6 in a hypoxia/reoxygenation-stimulated human renal tubular epithelial cell line (HK-2). Moreover, Sch A suppressed I/R-induced HK-2 cell apoptosis through the PI3K/Akt signaling pathway [95], which is part of the casein kinase signaling pathway and involved in cell proliferation, angiogenesis, and other activities through its effects on a variety of intracellular second messengers and participation in several important steps in the process of I/R injury

[96]. These experiments showed the potent protective effects of Sch A against renal I/R injury.

3. Summary and Prospects

Sch A has primary anti-inflammatory, antioxidant, and antiapoptotic properties (Figure 1). It is mainly involved in the MAPK, NF- κ B, AKT/GSK3 β , and PI3K/AKT pathways, reducing oxidative damage by reducing the production of MDA and ROS, and reducing inflammation by reducing IL-1 β and TNF- α (Table 1). This may be due to the methoxy group and the biphenyl cyclooctene in its structure, which effectively improve the AKT pathway and reduce MPP+ and protect nerves. In addition, metabolic derivatives of Sch A can be coupled with glucuronic acid, sulfate, taurine, glutathione, and glucose to produce possible anti-inflammatory and antioxidant effects.

Our understanding of Sch A is still limited, and the specific mechanism by which Sch A affects signaling pathways and genes is still unknown, even though some progress has been made. Many derivatives of Sch A have been obtained by chemical synthesis, but their pharmacological activities also need further study. To obtain the maximum advantages of TCM, understanding the interaction between different components of *S. chinensis* is also indispensable. With further research, it is hoped that Sch A and its derivatives can become new clinical drugs due to their characteristics, which include their high efficiency, low adverse reactions, and easy access.

I/R: ischemia/reperfusion; MCAO/R: middle cerebral artery occlusion and reperfusion.

Abbreviation

ROS:	Reactive oxygen species
NO:	Nitric oxide
MPP+:	1-methyl-4-phenylpyridinium ion
Bax:	Bcl2-associated X

Bcl2:	B-cell lymphoma 2
PI3K:	Phosphatidylinositol-4,5-bisphosphate 3-kinase
Akt:	Protein kinase B
Gsk3 β :	Glycogen synthase kinase-3 β
MDA:	Malondialdehyde
SOD:	Superoxide dismutase
GSH:	Glutathione
GSH-Px:	Glutathione peroxidase
TLR4:	Toll-like receptor 4
C5aR1:	Complement C5a receptor
IRI:	Ischemia-reperfusion injury
TNF- α :	Tumor necrosis factor-alpha
IL-6:	Interleukin-6
IL-1 β :	Interleukin-1 beta
NF- κ B:	Nuclear factor-kappa B
MAPK:	Mitogen-activated protein kinases
AST:	Aspartate aminotransferase
ALT:	Alanine transaminase
ADH:	Alcohol dehydrogenase
MPO:	Myeloperoxidase
ALDH:	Acetaldehyde dehydrogenase
GSSG:	Glutathione disulfide
ATP:	Adenosine triphosphate
PXR:	Pregnane X receptor
PGE2:	Prostaglandin E2.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Z. Xiao wrote the initial draft. W. Xiao and G. Li edited the manuscript. All authors edited and approved the manuscript.

Acknowledgments

This study was supported by grants from the National Natural Science Foundation of China (82160163, 81860217, 81560219, and 81200853) and a grant from the Key Research and Development Programs of Jiangxi Province (20192BBH80017).

References

- [1] A. Nowak, M. Zakłós-Szyda, J. Błasiak, A. Nowak, Z. Zhang, and B. Zhang, "Potential of *Schisandra chinensis* (Turcz.) Baill. In human health and nutrition: a review of current knowledge and therapeutic perspectives," *Nutrients*, vol. 11, no. 2, Article ID 333, 2019.
- [2] D. M. Kopustinskiene and J. Bernatoniene, "Antioxidant effects of *Schisandra chinensis* fruits and their active constituents," *Antioxidants*, vol. 10, no. 4, Article ID 620, 2021.
- [3] J. Fu, H. Zhang, S. Liu et al., "An integrated strategy using LC-MS/MS combined with in vivo microdialysis for the simultaneous determination of lignans of *Schisandra chinensis* (Turcz.) Baill. Fructus and endogenous neurotransmitters: application in pharmacokinetic and pharmacodynamic studies," *Food & Function*, vol. 12, no. 19, pp. 8932–8945, 2021.
- [4] Y. Zhou, L. Men, Y. Sun, M. Wei, and X. Fan, "Pharmacodynamic effects and molecular mechanisms of lignans from *Schisandra chinensis* Turcz. (Baill.), a current review," *European Journal of Pharmacology*, vol. 892, Article ID 173796, 2021.
- [5] Z. Li, X. He, F. Liu, J. Wang, and J. Feng, "A review of polysaccharides from *Schisandra chinensis* and *Schisandra sphenanthera*: properties, functions and applications," *Carbohydrate Polymers*, vol. 184, pp. 178–190, 2018.
- [6] J. Y. Park and K. H. Kim, "A randomized, double-blind, placebo-controlled trial of *Schisandra chinensis* for menopausal symptoms," *Climacteric*, vol. 19, no. 6, pp. 574–580, 2016.
- [7] G. Aslanyan, E. Amroyan, E. Gabrielyan, M. Nylander, G. Wikman, and A. Panossian, "Double-blind, placebo-controlled, randomised study of single dose effects of ADAPT-232 on cognitive functions," *Phytomedicine*, vol. 17, no. 7, pp. 494–499, 2010.
- [8] H.-F. Chiu, T.-Y. Chen, Y.-T. Tzeng, and C.-K. Wang, "Improvement of liver function in humans using a mixture of *Schisandra* fruit extract and sesamin," *Phytotherapy Research*, vol. 27, no. 3, pp. 368–373, 2013.
- [9] T. Zhao, L. Mao, Z. Yu et al., "Therapeutic potential of bicyclol in liver diseases: lessons from a synthetic drug based on herbal derivative in traditional Chinese medicine," *International Immunopharmacology*, vol. 91, Article ID 107308, 2021.
- [10] G. Xu, X. Lv, Y. Feng et al., "Study on the effect of active components of *Schisandra chinensis* on liver injury and its mechanisms in mice based on network pharmacology," *European Journal of Pharmacology*, vol. 910, Article ID 174442, 2021.
- [11] J. Li, S. Chen, X. Qin et al., "Wuzhi tablet (*Schisandra sphenanthera* extract) is a promising tacrolimus-sparing agent for renal transplant recipients who are CYP3A5 expressers: a two-phase prospective study," *Drug Metabolism and Disposition*, vol. 45, no. 11, pp. 1114–1119, 2017.
- [12] H.-W. Xin, Q. Li, X.-C. Wu et al., "Effects of *Schisandra sphenanthera* extract on the blood concentration of tacrolimus in renal transplant recipients," *European Journal of Clinical Pharmacology*, vol. 67, no. 12, pp. 1309–1311, 2011.
- [13] J. Park, S. Han, and H. Park, "Effect of *Schisandra chinensis* extract supplementation on quadriceps muscle strength and fatigue in adult women: a randomized, double-blind, placebo-controlled trial," *International Journal of Environmental Research and Public Health*, vol. 17, no. 7, Article ID 2475, 2020.
- [14] M.-Y. Song, J.-H. Wang, T. Eom, and H. Kim, "*Schisandra chinensis* fruit modulates the gut microbiota composition in association with metabolic markers in obese women: a randomized, double-blind placebo-controlled study," *Nutrition Research*, vol. 35, no. 8, pp. 655–663, 2015.
- [15] Y. X. Guo, J. Han, D. Y. Zhang, L. H. Wang, and L. L. Zhou, "Aqueous two-phase system coupled with ultrasound for the extraction of lignans from seeds of *Schisandra chinensis* (Turcz.) Baill.," *Ultrasonics Sonochemistry*, vol. 20, no. 1, pp. 125–132, 2013.
- [16] F. Sa, L. Q. Zhang, C. M. Chong et al., "Discovery of novel anti-parkinsonian effect of schisantherin A in in vitro and in vivo," *Neuroscience Letters*, vol. 593, pp. 7–12, 2015.
- [17] W. Feng, L.-Y. Zhou, R.-F. Mu et al., "Screening and identification of the main metabolites of schisantherin A in vivo and in vitro by using UHPLC-Q-TOF-MS/MS," *Molecules*, vol. 25, no. 2, Article ID 258, 2020.

- [18] S. Becker, M. Ceko, M. Louis-Foster et al., "Dopamine and pain sensitivity: neither sulpiride nor acute phenylalanine and tyrosine depletion have effects on thermal pain sensations in healthy volunteers," *PLoS One*, vol. 8, no. 11, Article ID e80766, 2013.
- [19] C. W. Halstead, S. Lee, C. S. Khoo, J. R. Hennell, and A. Bensoussan, "Validation of a method for the simultaneous determination of four *Schisandra* lignans in the raw herb and commercial dried aqueous extracts of *Schisandra chinensis* (Wu Wei Zi) by RP-LC with DAD," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 45, no. 1, pp. 30–37, 2007.
- [20] Y. Lu and D.-F. Chen, "Analysis of *Schisandra chinensis* and *Schisandra sphenanthera*," *Journal of Chromatography A*, vol. 1216, no. 11, pp. 1980–1990, 2009.
- [21] Z. Cheng, H. Song, Y. Yang et al., "Optimization of microwave-assisted enzymatic extraction of polysaccharides from the fruit of *Schisandra chinensis* Baill.," *International Journal of Biological Macromolecules*, vol. 76, pp. 161–168, 2015.
- [22] M. Razgonova, A. Zakharenko, K. Pikula et al., "Rapid mass spectrometric study of a supercritical CO₂-extract from woody liana *Schisandra chinensis* by HPLC-SPD-ESI-MS/MS," *Molecules*, vol. 25, no. 11, Article ID 2689, 2020.
- [23] E. J. Jeong, H. K. Lee, K. Y. Lee et al., "The effects of lignan-riched extract of *Schisandra chinensis* on amyloid- β -induced cognitive impairment and neurotoxicity in the cortex and hippocampus of mouse," *Journal of Ethnopharmacology*, vol. 146, no. 1, pp. 347–354, 2013.
- [24] D. Grimes, M. Fitzpatrick, J. Gordon et al., "Canadian guideline for Parkinson disease," *Canadian Medical Association Journal*, vol. 191, no. 36, pp. E989–E1004, 2019.
- [25] M. Golpich, B. Rahmani, N. Mohamed Ibrahim et al., "Preconditioning as a potential strategy for the prevention of Parkinson's disease," *Molecular Neurobiology*, vol. 51, no. 1, pp. 313–330, 2015.
- [26] C. Kwan, I. Frouni, S. G. Nuara et al., "Combined 5-HT_{2A} and mGlu₂ modulation for the treatment of dyskinesia and psychosis in Parkinson's disease," *Neuropharmacology*, vol. 186, Article ID 108465, 2021.
- [27] J. Marinus, K. Zhu, C. Marras, D. Aarsland, and J. J. van Hilten, "Risk factors for non-motor symptoms in Parkinson's disease," *The Lancet Neurology*, vol. 17, no. 6, pp. 559–568, 2018.
- [28] Y. Vermeiren and P. P. De Deyn, "Targeting the nor-epinephrinergic system in Parkinson's disease and related disorders: the locus coeruleus story," *Neurochemistry International*, vol. 102, pp. 22–32, 2017.
- [29] T. Chen, C. Li, Y. Li, X. Yi, S. M.-Y. Lee, and Y. Zheng, "Oral delivery of a nanocrystal formulation of schisantherin A with improved bioavailability and brain delivery for the treatment of Parkinson's disease," *Molecular Pharmaceutics*, vol. 13, no. 11, pp. 3864–3875, 2016.
- [30] F. Sa, B. J. Guo, S. Li et al., "Pharmacokinetic study and optimal formulation of new anti-Parkinson natural compound schisantherin A," *Parkinson's Disease*, vol. 2015, Article ID 951361, 7 pages, 2015.
- [31] T. Chen, C. Li, Y. Li et al., "Small-sized mPEG-PLGA nanoparticles of schisantherin A with sustained release for enhanced brain uptake and anti-parkinsonian activity," *ACS Applied Materials & Interfaces*, vol. 9, no. 11, pp. 9516–9527, 2017.
- [32] S. R. Kim, M. K. Lee, K. A. Koo et al., "Dibenzocyclooctadiene lignans from *Schisandra chinensis* protect primary cultures of rat cortical cells from glutamate-induced toxicity," *Journal of Neuroscience Research*, vol. 76, no. 3, pp. 397–405, 2004.
- [33] L. Q. Zhang, F. Sa, C. M. Chong et al., "Schisantherin A protects against 6-OHDA-induced dopaminergic neuron damage in zebrafish and cytotoxicity in SH-SY5Y cells through the ROS/NO and AKT/GSK3 β pathways," *Journal of Ethnopharmacology*, vol. 170, pp. 8–15, 2015.
- [34] N. A. Bishop, T. Lu, and B. A. Yankner, "Neural mechanisms of ageing and cognitive decline," *Nature*, vol. 464, no. 7288, pp. 529–535, 2010.
- [35] K. Sowndhararajan, P. Deepa, M. Kim, S. J. Park, and S. Kim, "An overview of neuroprotective and cognitive enhancement properties of lignans from *Schisandra chinensis*," *Biomedicine & Pharmacotherapy*, vol. 97, pp. 958–968, 2018.
- [36] X. Li, X. Zhao, X. Xu et al., "Schisantherin A recovers A β -induced neurodegeneration with cognitive decline in mice," *Physiology & Behavior*, vol. 132, pp. 10–16, 2014.
- [37] C. Liu, W. Sun, N. Li et al., "Schisantherin A improves learning and memory of mice with D-galactose-induced learning and memory impairment through its antioxidation and regulation of p19/p53/p21/Cyclin D1/CDK4/RB gene expressions," *Journal of Medicinal Food*, vol. 21, no. 7, pp. 678–688, 2018.
- [38] V. Medici and C. H. Halsted, "Folate, alcohol, and liver disease," *Molecular Nutrition & Food Research*, vol. 57, no. 4, pp. 596–606, 2013.
- [39] Y. Liu, L. Y. Zhou, and X. W. Meng, "Genetic polymorphism of two enzymes with alcoholic liver disease in Northeast China," *Hepato-Gastroenterology*, vol. 59, no. 113, pp. 204–207, 2012.
- [40] A. Mota, F. Guedes, J. Areias, L. Pinho, and M. F. Cardoso, "Alcohol consumption among patients with hepatitis B infection in northern Portugal considering gender and hepatitis B virus genotype differences," *Alcohol*, vol. 44, no. 2, pp. 149–156, 2010.
- [41] H. Hagström, "Alcohol, smoking and the liver disease patient," *Best Practice & Research Clinical Gastroenterology*, vol. 31, no. 5, pp. 537–543, 2017.
- [42] M. Charlton, "Branched-chain amino acid enriched supplements as therapy for liver disease," *Journal of Nutrition*, vol. 136, no. 1, pp. 295S–298S, 2006.
- [43] M. C. Gutiérrez-Ruiz, L. Bucio, A. Correa et al., "Metadoxine prevents damage produced by ethanol and acetaldehyde in hepatocyte and hepatic stellate cells in culture," *Pharmacological Research*, vol. 44, no. 5, pp. 431–436, 2001.
- [44] M. Cao, X. Li, B. Zhang et al., "The effect of polyene phosphatidyl choline intervention on nonalcoholic steatohepatitis and related mechanism," *American Journal of Tourism Research*, vol. 8, no. 5, pp. 2325–2330, 2016.
- [45] Z. Yu, F. Wu, J. Tian, X. Guo, and R. An, "Protective effects of compound ammonium glycylrhizin, L-arginine, silymarin and glucuro-lactone against liver damage induced by ochratoxin A in primary chicken hepatocytes," *Molecular Medicine Reports*, vol. 18, no. 3, pp. 2551–2560, 2018.
- [46] S. Li, H. Guan, Z. Qian et al., "Taurine inhibits 2,5-hexanedione-induced oxidative stress and mitochondria-dependent apoptosis in PC12 cells," *Industrial Health*, vol. 55, no. 2, pp. 108–118, 2017.
- [47] C. Başaran-Küçükgergin, İ. Bingöl, M. S. Tekkeşin, V. Olgaç, S. Doğru-Abbassoğlu, and M. Uysal, "Effects of carnitine, taurine, and betaine pretreatments on diethylnitrosamine-induced oxidative stress and tissue injury in rat liver," *Toxicology and Industrial Health*, vol. 32, no. 8, pp. 1405–1413, 2016.
- [48] J.-H. Bockmann, M. Dandri, S. Lüth, N. Pannicke, and A. W. Lohse, "Combined glucocorticoid and antiviral therapy

- of hepatitis B virus-related liver failure,” *World Journal of Gastroenterology*, vol. 21, no. 7, pp. 2214–2219, 2015.
- [49] L. Gao, J.-F. Wang, M. Xiang, Y.-C. Fan, Z.-G. Zhang, and K. Wang, “Expression of human glucocorticoid receptor in T lymphocytes in acute-on-chronic hepatitis B liver failure,” *Digestive Diseases and Sciences*, vol. 56, no. 9, pp. 2605–2612, 2011.
 - [50] B. Li, D. Li, Y. Wang et al., “Schisantherin A alleviated alcohol-induced liver injury by the regulation of alcohol metabolism and NF- κ B pathway,” *Experimental Animals*, vol. 67, no. 4, pp. 451–461, 2018.
 - [51] W. Liu, H. Zhu, and H. Fang, “Propofol potentiates sevoflurane-induced inhibition of nuclear factor- κ B-mediated inflammatory responses and regulation of mitogen-activated protein kinases pathways via toll-like receptor 4 signaling in lipopolysaccharide-induced acute lung injury in mice,” *The American Journal of the Medical Sciences*, vol. 354, no. 5, pp. 493–505, 2017.
 - [52] D. Li, X. Ci, Y. Li et al., “Alleviation of severe inflammatory responses in LPS-exposed mice by Schisantherin A,” *Respiratory Physiology & Neurobiology*, vol. 202, pp. 24–31, 2014.
 - [53] E. Zhou, Y. Li, Z. Wei et al., “Schisantherin A protects lipopolysaccharide-induced acute respiratory distress syndrome in mice through inhibiting NF- κ B and MAPKs signaling pathways,” *International Immunopharmacology*, vol. 22, no. 1, pp. 133–140, 2014.
 - [54] S. J. Nho, S. M. Kymes, J. J. Callaghan, and D. T. Felson, “The burden of hip osteoarthritis in the United States: epidemiologic and economic considerations,” *Journal of the American Academy of Orthopaedic Surgeons*, vol. 21, pp. S1–S6, 2013.
 - [55] M. Kapoor, J. Martel-Pelletier, D. Lajeunesse, J.-P. Pelletier, and H. Fahmi, “Role of proinflammatory cytokines in the pathophysiology of osteoarthritis,” *Nature Reviews Rheumatology*, vol. 7, no. 1, pp. 33–42, 2011.
 - [56] L. Kalichman, Y. Zhang, J. Niu et al., “The association between patellar alignment on magnetic resonance imaging and radiographic manifestations of knee osteoarthritis,” *Arthritis Research and Therapy*, vol. 9, no. 2, Article ID R26, 2007.
 - [57] D. R. Eyre, “Collagens and cartilage matrix homeostasis,” *Clinical Orthopaedics and Related Research*, vol. 427, pp. S118–S122, 2004.
 - [58] S. Liao, K. Zhou, D. Li, X. Xie, F. Jun, and J. Wang, “Schisantherin A suppresses interleukin-1 β -induced inflammation in human chondrocytes via inhibition of NF- κ B and MAPKs activation,” *European Journal of Pharmacology*, vol. 780, pp. 65–70, 2016.
 - [59] C. Peralta, L. Fernandez, J. Panes et al., “Preconditioning protects against systemic disorders associated with hepatic ischemia-reperfusion through blockade of tumor necrosis factor-induced P-selectin up-regulation in the rat,” *Hepatology*, vol. 33, no. 1, pp. 100–113, 2001.
 - [60] R. F. Schwabe and D. A. Brenner, “Mechanisms of Liver Injury. I. TNF- α -induced liver injury: role of IKK, JNK, and ROS pathways,” *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 290, no. 4, pp. G583–G589, 2006.
 - [61] M. Rajesh, H. Pan, P. Mukhopadhyay et al., “Pivotal Advance: cannabinoid-2 receptor agonist HU-308 protects against hepatic ischemia/reperfusion injury by attenuating oxidative stress, inflammatory response, and apoptosis,” *Journal of Leukocyte Biology*, vol. 82, no. 6, pp. 1382–1389, 2007.
 - [62] J. Yan, G. Xie, C. Liang et al., “Herbal medicine Yinchenhaotang protects against α -naphthylisothiocyanate-induced cholestasis in rats,” *Scientific Reports*, vol. 7, no. 1, Article ID 4211, 2017.
 - [63] S. Al-Dury, A. Wahlström, S. Wahlin et al., “Pilot study with IBAT inhibitor A4250 for the treatment of cholestatic pruritus in primary biliary cholangitis,” *Scientific Reports*, vol. 8, no. 1, Article ID 6658, 2018.
 - [64] N. Guañabens and A. Parés, “Osteoporosis in chronic liver disease,” *Liver International*, vol. 38, no. 5, pp. 776–785, 2018.
 - [65] M. K. Bothe, C. Meyer, U. Mueller et al., “Characterization of a rat model of moderate liver dysfunction based on alpha-naphthylisothiocyanate-induced cholestasis,” *Journal of Toxicological Sciences*, vol. 42, no. 6, pp. 715–721, 2017.
 - [66] T. Khamphaya, P. Chansela, P. Piyachaturawat, A. Suksamrarn, M. H. Nathanson, and J. Weerachayaphorn, “Effects of andrographolide on intrahepatic cholestasis induced by alpha-naphthylisothiocyanate in rats,” *European Journal of Pharmacology*, vol. 789, pp. 254–264, 2016.
 - [67] E. J. Carey, A. H. Ali, and K. D. Lindor, “Primary biliary cirrhosis,” *The Lancet*, vol. 386, no. 10003, pp. 1565–1575, 2015.
 - [68] F. Nevens, P. Andreone, G. Mazzella et al., “A placebo-controlled trial of obeticholic acid in primary biliary cholangitis,” *New England Journal of Medicine*, vol. 375, no. 7, pp. 631–643, 2016.
 - [69] S. Fan, C. Liu, Y. Jiang et al., “Lignans from *Schisandra sphenanthera* protect against lithocholic acid-induced cholestasis by pregnane X receptor activation in mice,” *Journal of Ethnopharmacology*, vol. 245, Article ID 112103, 2019.
 - [70] J. Y. L. Chiang, “Bile acid metabolism and signaling,” *Comprehensive Physiology*, vol. 3, no. 3, pp. 1191–1212, 2013.
 - [71] U. Beuers, M. Trauner, P. Jansen, and R. Poupon, “New paradigms in the treatment of hepatic cholestasis: from UDCA to FXR, PXR and beyond,” *Journal of Hepatology*, vol. 62, no. 1, pp. S25–S37, 2015.
 - [72] C. Lv and L. Huang, “Xenobiotic receptors in mediating the effect of sepsis on drug metabolism,” *Acta Pharmaceutica Sinica B*, vol. 10, no. 1, pp. 33–41, 2020.
 - [73] H. Zeng, D. Li, X. Qin et al., “Hepatoprotective effects of *Schisandra sphenanthera* extract against lithocholic acid-induced cholestasis in male mice are associated with activation of the pregnane X receptor pathway and promotion of liver regeneration,” *Drug Metabolism and Disposition*, vol. 44, no. 3, pp. 337–342, 2016.
 - [74] Y. Zhai, H. Petrowsky, J. C. Hong, R. W. Busuttill, and J. W. Kupiec-Weglinski, “Ischaemia-reperfusion injury in liver transplantation—from bench to bedside,” *Nature Reviews Gastroenterology & Hepatology*, vol. 10, no. 2, pp. 79–89, 2013.
 - [75] X.-J. Zhang, X. Cheng, Z.-Z. Yan et al., “An ALOX12-12-HETE-GPR31 signaling axis is a key mediator of hepatic ischemia-reperfusion injury,” *Nature Medicine*, vol. 24, no. 1, pp. 73–83, 2018.
 - [76] Y. Li, Y. Yang, Y. Feng et al., “A review of melatonin in hepatic ischemia/reperfusion injury and clinical liver disease,” *Annals of Medicine*, vol. 46, no. 7, pp. 503–511, 2014.
 - [77] C. Peralta, M. B. Jiménez-Castro, and J. Gracia-Sancho, “Hepatic ischemia and reperfusion injury: effects on the liver sinusoidal milieu,” *Journal of Hepatology*, vol. 59, no. 5, pp. 1094–1106, 2013.
 - [78] H. Huang, S. Tohme, A. B. Al-Khafaji et al., “Damage-associated molecular pattern-activated neutrophil extracellular trap exacerbates sterile inflammatory liver injury,” *Hepatology*, vol. 62, no. 2, pp. 600–614, 2015.
 - [79] J.-Y. Han, Q. Li, Z.-Z. Ma, and J.-Y. Fan, “Effects and mechanisms of compound Chinese medicine and major

- ingredients on microcirculatory dysfunction and organ injury induced by ischemia/reperfusion,” *Pharmacology & Therapeutics*, vol. 177, pp. 146–173, 2017.
- [80] N. Zheng, F. Liu, H. Lu et al., “Schisantherin A protects against liver ischemia-reperfusion injury via inhibition of mitogen-activated protein kinase pathway,” *International Immunopharmacology*, vol. 47, pp. 28–37, 2017.
- [81] J. Han, D. Wang, B. Yu et al., “Cardioprotection against ischemia/reperfusion by licochalcone B in isolated rat hearts,” *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 134862, 11 pages, 2014.
- [82] F. Jiang, J. Yang, L. Zhang et al., “Rosuvastatin reduces ischemia-reperfusion injury in patients with acute coronary syndrome treated with percutaneous coronary intervention,” *Clinical Cardiology*, vol. 37, no. 9, 2014.
- [83] J. Q. Zhang, Q. Wang, F. S. Xue et al., “Ischemic preconditioning produces more powerful anti-inflammatory and cardioprotective effects than limb remote ischemic postconditioning in rats with myocardial ischemia-reperfusion injury,” *Chinese Medical Journal*, vol. 126, no. 20, pp. 3949–3955, 2013.
- [84] G. Tarantini, E. Favaretto, M. P. Marra et al., “Post-conditioning during coronary angioplasty in acute myocardial infarction: the POST-AMI trial,” *International Journal of Cardiology*, vol. 162, no. 1, pp. 33–38, 2012.
- [85] R. Chang, Y. Li, X. Yang et al., “Protective role of deoxyschizandrin and schisantherin A against myocardial ischemia-reperfusion injury in rats,” *PLoS One*, vol. 8, no. 4, Article ID e61590, 2013.
- [86] S. Lorenzano, N. S. Rost, M. Khan et al., “Oxidative stress biomarkers of brain damage,” *Stroke*, vol. 49, no. 3, pp. 630–637, 2018.
- [87] L. Zhu, L. Wang, F. Ju, A. Khan, X. Cheng, and S. Zhang, “Reversible recovery of neuronal structures depends on the degree of neuronal damage after global cerebral ischemia in mice,” *Experimental Neurology*, vol. 289, pp. 1–8, 2017.
- [88] Q. Yang, Q. Huang, Z. Hu, and X. Tang, “Potential neuro-protective treatment of stroke: targeting excitotoxicity, oxidative stress, and inflammation,” *Frontiers in Neuroscience*, vol. 13, Article ID 1036, 2019.
- [89] C. Thornton, B. Leaw, C. Mallard, S. Nair, M. Jinnai, and H. Hagberg, “Cell death in the developing brain after hypoxia-ischemia,” *Frontiers in Cellular Neuroscience*, vol. 11, Article ID 248, 2017.
- [90] Y. W. Shi, X. C. Zhang, C. Chen et al., “Schisantherin A attenuates ischemia/reperfusion-induced neuronal injury in rats via regulation of TLR4 and C5aR1 signaling pathways,” *Brain, Behavior, and Immunity*, vol. 66, pp. 244–256, 2017.
- [91] P. A. Ward, “The harmful role of c5a on innate immunity in sepsis,” *Journal of Innate Immunity*, vol. 2, no. 5, pp. 439–445, 2010.
- [92] L. Xu, R. Sang, Y. Yu, J. Li, B. Ge, and X. Zhang, “The polysaccharide from *Inonotus obliquus* protects mice from *Toxoplasma gondii*-induced liver injury,” *International Journal of Biological Macromolecules*, vol. 125, pp. 1–8, 2019.
- [93] Y. Lu, X. Xu, T. Jiang et al., “Sertraline ameliorates inflammation in CUMS mice and inhibits TNF- α -induced inflammation in microglia cells,” *International Immunopharmacology*, vol. 67, pp. 119–128, 2019.
- [94] H. Nishikawa, Y. Taniguchi, T. Matsumoto et al., “Knockout of the interleukin-36 receptor protects against renal ischemia-reperfusion injury by reduction of proinflammatory cytokines,” *Kidney International*, vol. 93, no. 3, pp. 599–614, 2018.
- [95] J. Gong and X. Wang, “Schisantherin A protects renal tubular epithelial cells from hypoxia/reoxygenation injury through the activation of PI3K/Akt signaling pathway,” *Journal of Biochemical and Molecular Toxicology*, vol. 32, no. 7, Article ID e22160, 2018.
- [96] H. H. Arab, S. A. Salama, and I. A. Maghrabi, “Camel milk attenuates methotrexate-induced kidney injury via activation of PI3K/Akt/eNOS signaling and intervention with oxidative aberrations,” *Food & Function*, vol. 9, no. 5, pp. 2661–2672, 2018.
- [97] B. H. Gu, N. V. Minh, S. H. Lee et al., “Deoxyschisandrin inhibits H₂O₂-induced apoptotic cell death in intestinal epithelial cells through nuclear factor-kappaB,” *International Journal of Molecular Medicine*, vol. 26, no. 3, pp. 401–406, 2010.