



## Research Article

# ***In Vitro* Assessment of the Cercaricidal Activity of *Sida acuta* Burm. F. and *Sida rhombifolia* Linn. (Malvaceae) Hydroethanolic Extracts, Cytotoxicity, and Phytochemical Studies**

**Ulrich Membe Femoe** <sup>1,2</sup>, **Joseph Bertin Kadji Fassi**,<sup>1,2</sup> **Hermine Boukeng Jatsa** <sup>1,2</sup>,  
**Yannick Leonel Tchoffo**,<sup>1,2</sup> **David Carel Amvame Nna**,<sup>1,2</sup> **Blaise Cédric Kamdoum**,<sup>3</sup>  
**Steven Collins Njonte Wouamba**,<sup>4</sup> **Billy Toussie Tchegnitegni**,<sup>5</sup>  
**Bonaventure Tchaleu Ngadjui**,<sup>3</sup> **Norbert Sewald**,<sup>5</sup> **Bruno Ndjakou Lenta**,<sup>4</sup>  
**Louis-Albert Tchuem Tchuente**,<sup>1,2</sup> and **Theophile Dimo** <sup>1</sup>

<sup>1</sup>Department of Animal Biology and Physiology, Faculty of Science, University of Yaounde I, P.O. Box 812, Yaounde, Cameroon

<sup>2</sup>Centre for Schistosomiasis and Parasitology, P.O. Box 7244, Yaounde, Cameroon

<sup>3</sup>Department of Organic Chemistry, Faculty of Science, University of Yaounde I, P.O. Box 812, Yaounde, Cameroon

<sup>4</sup>Department of Chemistry, Higher Teacher Training College, University of Yaounde I, P.O. Box 47, Yaounde, Cameroon

<sup>5</sup>Department of Organic and Bioorganic Chemistry, Faculty of Chemistry, Bielefeld University, P.O. BOX 100131, D-33501, Bielefeld, Germany

Correspondence should be addressed to Theophile Dimo; [dimo59@yahoo.com](mailto:dimo59@yahoo.com)

Received 24 November 2021; Accepted 17 December 2021; Published 10 January 2022

Academic Editor: Daniel Dias Rufino Arcanjo

Copyright © 2022 Ulrich Membe Femoe et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Despite the global efforts, schistosomiasis remains a public health problem in several tropical and subtropical countries. One of the major challenges in the fight against schistosomiasis is the interruption of the parasite life cycle. Here, we evaluated the anticercarial, cytotoxicity, and phytochemical profiles of *Sida acuta* (HESa) and *Sida rhombifolia* (HESr) hydroethanolic extracts (Malvaceae). *Schistosoma mansoni* cercaria was collected from fifteen *Biomphalaria pfeifferi*-infected snails. Twenty-five cercariae were incubated in duplicate with different concentrations (31.25–1,000 µg/mL) of HESa or HESr. The cercaria viability was monitored at 30 min time intervals for 150 min, and the concentration-response curve of each plant extract was used to determine their respective lethal concentration 50 (LC<sub>50</sub>). Additionally, the cytotoxicity profile of each plant extract was evaluated on the Hepa 1–6 cell line at a concentration range of 15.625–1,000 µg/mL using the WST-8 assay method and its inhibitory concentration 50 (IC<sub>50</sub>) was calculated. Moreover, phytochemical characterization of each plant extract was carried out by HPLC-MS. Both extracts exhibited cercaricidal activity in a time- and concentration-dependent manner. At 30 min time point, HESa (LC<sub>50</sub> = 28.41 ± 3.5 µg/mL) was more effective than HESr (LC<sub>50</sub> = 172.42 ± 26.16 µg/mL) in killing *S. mansoni* cercariae. Regarding the cytotoxicity effect of both extracts, the IC<sub>50</sub> of HESa (IC<sub>50</sub> = 109.67 µg/mL) was lower than that of HESr (IC<sub>50</sub> = 888.79 µg/mL). The selectivity index was 3.86 and 5.15 for HESa and HESr, respectively. Fifteen compounds were identified from HESa and HESr after HPLC-MS analysis. N-Feruloyltyramine, a polyphenol, and thamnimonin, a coumarin, were identified in both extracts. HESa and HESr displayed cercaricidal activity and were not toxic on Hepa 1–6 cell line. Based on the selectivity index of these extracts, *S. rhombifolia* extract could be more effective on *S. mansoni* cercariae than *S. acuta* extract. This study could provide baseline information for further investigations aiming to develop plant-based alternative drugs against *S. mansoni*.

## 1. Introduction

Schistosomiasis, also known as bilharzia, is an infectious disease caused by trematodes flatworms of the genus *Schistosoma*. It is known as one of the most prevalent tropical diseases worldwide. It is estimated that 229 million people required preventive treatment in 2018 and close to 800 million are at risk of infection [1]. After an infected person releases *Schistosoma* eggs into the water by defecation or urination, the ripe miracidia hatch out and invade the intermediate host freshwater snail where they form sporocysts. They then matured to give cercariae that emerges from snails and swims to penetrate the skin of humans and/or animals that are definitive hosts [2]. Therefore, to control schistosomiasis, the life cycle of the parasite can be interrupted by killing cercariae and miracidia [3–5]. The use of chemical compounds to control the aquatic snails, cercariae, or miracidia is not recommended because of their adverse effects on the environment [6]. A large number of plant families with potential schistosomicidal activity have been identified through plant screening, which represents a continuous effort to find new bioactive molecules. The anti-cercarial activities of several plants such as *Rauwolfia vomitoria* [7], *Nigella sativa* [8] *Balanites aegyptica*, *Azadirachta indica*, *Nauclea latifolia*, *Morinda lucida*, *Phyllanthus amarus*, *Vernonia amygdalina* [9], and *Ozoroa pulcherrima* [10] have been reported. In several countries, *Sida L.* (Malvaceae) has been used for centuries in traditional medicines for the prevention and treatment of different diseases such as diarrhea, dysentery, gastrointestinal and urinary infections, cancer, malaria, and helminth infections [11–18]. Moreover, during the last decade, Jatsa et al. [19–21] have demonstrated the *in vitro* and *in vivo* anti-schistosomal potency of *Sida pilosa*. In addition, previous investigations on two of the plants of the genus *Sida*, namely *Sida acuta* and *Sida rhombifolia*, reported their anthelmintic and larvicidal activities [22–24]. Considering the above, this study was therefore carried out to evaluate the *in vitro* cercaricidal activity of *S. acuta* and *S. rhombifolia* hydroethanolic extracts against *Schistosoma mansoni* cercaria, their cytotoxic profile, and determine their phytoconstituents.

## 2. Materials and Methods

**2.1. Collection, Preparation, and Extraction Process of Plant Specimens.** The whole plant of *S. acuta* (Figure 1(a)) was collected in February 2017 at Nkoemvone near Ebolowa in the South Region while the aerial parts of *S. rhombifolia* (Figure 1(b)) were harvested in March 2017 at Bazou, a village of Ndé division in the West region of Cameroon. The plant specimens were identified and authenticated at Cameroon's National Herbarium under a voucher specimen HNC n°46188 (*S. acuta*) and HNC n°20113 (*S. rhombifolia*). Each plant species was air-dried for 2 weeks, and the dried plant materials were ground into powder. Exactly 400 g of each pulverized plant material was macerated with 800 mL of ethanol/water mixture (70/30) at room temperature for 48 h. The macerates were filtered, and the filtrate was concentrated using a rotary evaporator (Buchi Rotavapor, R

200) under reduced pressure at 40°C. The crude extract was finally air-dried, and the yield of obtained hydroethanolic extracts of *S. acuta* (HESa) and *S. rhombifolia* (HESr) was calculated.

### 2.2. In Vitro Cercaricidal Activity of the Plant Extracts

**2.2.1. Snails Infection and Preparation of the Cercarial Suspension.** Schistosome cercariae were obtained from experimentally infected juvenile *Biomphalaria pfeifferi* snails that were maintained in the snail laboratory at the Centre Schistosomiasis and Parasitology of Yaoundé, Cameroon, under standard laboratory conditions of 18–26°C. Briefly, each snail was exposed for 12 hours to 4–5 miracidia obtained after hatching of eggs isolated from *S. mansoni*-infected mice liver. Four weeks later, 14 infected snails, known to be shedding cercariae, were pooled into a glass Beaker with 20 mL of distilled water and allowed to shed cercariae by exposing them to artificial light for 60 min. After the exposition time, snails were removed from the beaker and 100 µL of the cercariae suspension was transferred on three microscope slides. A volume of 100 µL of Lugol was added on each slide to stain and fix cercariae, and the slides were transferred onto a light microscope to estimate the number of cercariae per 100 µL, and the volume of the cercariae suspension containing approximately 25 viable cercariae was determined [7, 9].

**2.2.2. In Vitro Cercaricidal Assay.** The concentrations ranging from 1,000 to 31.25 µg/mL are widely used to evaluate the cercaricidal activity of medicinal plant crude extracts [7, 9]. According to this protocol, serial dilutions (1,000, 500, 250, 125, 62.5, and 31.25 µg/mL) of each of the test plant extract were prepared using distilled water as diluent. A volume of 1 mL of each dilution was added into a well of a 24-well culture plate. Niclosamide-olamide 5% (1 µg/mL; Jiangsu Aijin Agrochemical Co. Ltd., China) was used as a reference control. Approximately 25 cercariae were pipetted into each of the wells. Cercaria mortality was observed for 150 min at 30 min intervals since the infectivity of cercariae is known to be rapidly lost after 12 h [25]. The bottom of each well was observed at 4X magnification using an inverted microscope and cercariae survival and mortality were recorded at specific 30 min time intervals (30, 60, 90, 120, and 150 min). Cercariae were presumed dead when they stopped movement and sank or their tail was detached [26]. All experiments were carried out in duplicate and repeated twice. The viability percentage and the lethal concentration 50 (LC<sub>50</sub>) values of plant extracts on schistosome cercariae were calculated at each time point.

**2.3. Cytotoxicity Assay of the Extracts against Hepatocyte Cell Lines.** Cytotoxicity of *S. acuta* and *S. rhombifolia* extracts was investigated on C57/L mouse melanoma liver cells line (*Hepa 1-6*, ATCC CRL-1830) using WST-8-based assay as previously described by Murata et al. [27]. The cells were cultured in high-glucose Dulbecco's Modified Eagle




FIGURE 1: Aerial parts of (a) *Sida acuta* and (b) *Sida rhombifolia*.

Medium (DMEM) with pyruvate and L-glutamine (Gibco, Life Technologies, USA), supplemented with 10% (v/v) fetal bovine serum (FBS) heat-inactivated (Serana, Australia) and 1% (v/v) penicillin/streptomycin (Gibco, Life Technologies, USA). The cells were grown in a growth medium at 37°C in a 95% air, 5% CO<sub>2</sub>-humidified incubator. Monolayer cultures reaching a confluence between 80 and 90% were detached using trypsin solution (Sigma Aldrich, Germany) and calibrated using a cell counter chamber (Fast Read 102). The calibrated cell suspension was seeded into 96-well tissue culture microtiter plates at a density of  $1 \times 10^4$  cells per well and incubated overnight at 37°C in a 5% CO<sub>2</sub> incubator for cell adhesion. Following incubation, the medium was removed from the cells and replaced with a fresh one followed by the addition of extracts at different concentrations (1,000, 500, 250, 125, 62.5, 31.25, and 15.625 µg/mL). Control wells with cells only were added, and the plates were incubated for 24 h in the same culture conditions. After incubation, cell viability was measured by mitochondrial activity in reducing 2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium monosodium salt to formazan using the cell counting kit-8 (WST-8, Abcam, ab228554, UK) according to the manufacturer instructions. After 4 hours of incubation, optical density was measured at 450 nm using a Dynex MRX TC II microplate reader (Dynex Technologies, USA). The results were expressed as a percentage growth of the control cells and the inhibitory concentration 50 (IC<sub>50</sub>) values were calculated as the concentration of the resulting in 50% reduction of absorbance compared to untreated cells. Tests were carried out in duplicate, and each experiment was repeated three times.

**2.4. Selectivity Index.** In the present study, the degree of selectivity of each hydroethanolic plant extract is expressed as the ratio of the IC<sub>50</sub> obtained for the cell line to the LC<sub>50</sub> for *S. mansoni* cercariae.

$$SI = \frac{\text{IC}_{50} \text{ of extract in cell line}}{\text{LC}_{50} \text{ of the same extract in } S. \text{ mansoni cercariae}} \quad (1)$$

## 2.5. Phytochemical Analysis

**2.5.1. Phytochemical Screening.** The hydroethanolic extracts of *S. acuta* and *S. rhombifolia* were subjected to qualitative chemical tests to identify phytochemical constituents. The screening of alkaloids, anthraquinones, and cardiac glycosides was performed by Mayer's test, Bontrager's test, and the Keller–Killiani's test, respectively. Ferric chloride (FeCl<sub>3</sub>) test was used for the identification of phenols and tannins, Fehling's test for reducing sugars, foam test for saponins, and Liebermann–Burchard's test for steroids. The presence of flavonoids, lipids, and terpenoids was performed using the ammonia test, the grease spot test, and Salkowski's test, respectively [28].

## 2.5.2. Qualitative Determination of Compounds Using LC-DAD-(HR) ESI-MS

**Sample Preparation.** HES<sub>a</sub> and HES<sub>r</sub> were dissolved in HPLC grade methanol at a concentration of 5 mg/mL and then filtrated through a syringe-filter-membrane. Aliquots of 5 µL were injected into the UPLC-DAD/MS Dionex Ultimate 3000 HPLC (Germany).

**HPLC-MS Conditions.** High-resolution mass spectra were obtained with a quadrupole-time-of-flight (QTOF) spectrometer (Bruker, Germany) equipped with a HESI source. The spectrometer was operated in positive mode (mass range: 100–1,500, with a scan rate of 1.00 Hz) with automatic gain control to provide high-accuracy mass measurements within 0.40 ppm deviation using Na formate as calibrant. The following parameters were used for experiments: spray voltage of 4.5 kV and the capillary temperature of 200°C.

Nitrogen was used as sheath gas (10 L/min). The spectrometer was attached to an Ultimate 3000 UHPLC system (Thermo Fisher, USA) consisting of an LC-pump, Diode Array Detector (DAD;  $\lambda$ : 190–600 nm), autosampler (injection volume: 10  $\mu$ L), and a column oven (40°C). The separations were performed using a Synergi MAX-RP 100A (50  $\times$  2 mm, 2.5  $\mu$  particle size) with a H<sub>2</sub>O (+0.1% HCOOH) (A)/acetonitrile (+0.1% HCOOH) (B) gradient (flow rate: 500  $\mu$ L/min and injection volume: 5  $\mu$ L). Samples were analyzed using a gradient program as follows: A linear gradient starting with 95% eluent A isocratic over 1.5 min to linear gradient with 100% eluent B over 6 min; 2 min after, the system returned to its initial condition (90% of eluent A) within 1 min and was equilibrated for 1 min.

**Identification of Peaks.** Identification of all constituents was performed by UPLC-DAD-MS analysis and by comparing the UV and MS spectra and MS/MS fragmentation of the peaks in the samples with those of data reported in the literature of the SciFinder, NIST/EPA/NIH Mass Spectral Library (NIST 14), and Mass Bank of North America (MoNA) databases.

**2.6. Statistical Analysis.** Graph drawing and statistical analysis were performed using GraphPad Software version 8.01 (GraphPad Software, San Diego, USA). The results were expressed as means  $\pm$  SEM, and Student's *t*-test and two-way ANOVA followed by Tukey multiple comparison post-test were used to determine the significance of differences between mean values. A *p*-value less than 0.05 was considered statistically significant.

### 3. Results

**3.1. Cercaricidal Activity of Plant Extracts.** The mortality of *S. mansoni* cercariae following *in vitro* exposure to different concentrations of *S. acuta* and *S. rhombifolia* extracts was both time- and concentration-dependent increased. In the absence of the plant extract, cercariae showed normal viability without any morphological changes (tail loss) for up to 2 h (Figures 2(a) and 3(a)).

**3.1.1. Cercaricidal Activity of *Sida acuta* Hydroethanolic Extract.** As shown in Figure 2(b), following 30 min incubation of cercariae with *S. acuta* hydroethanolic extract with 1,000, 500, or 250  $\mu$ g/mL, 100% of the cercariae died. At the same time, we recorded mortality rates of 83%, 88%, and 91% of cercariae incubated with 125, 62.5, and 31.25  $\mu$ g/mL of the hydroethanolic extract of *S. acuta*, respectively. However, from 60 min to 150 min, we obtained a mortality rate of 100% with all concentrations of the extract. At 30 minutes time point, the cercaricidal activity of HESa at all concentrations was significantly higher (*p* < 0.01) than the niclosamide-olamide one (Figure 2(b)).

**3.1.2. Cercaricidal Activity of *Sida rhombifolia* Hydroethanolic Extract.** The incubation of cercariae with *S. rhombifolia* hydroethanolic extract at 1,000, 500, and

250  $\mu$ g/mL induced after 30 min, mortality rates of 99%, 97%, and 66%, respectively. At the 60 min time point, we observed a mortality rate of 100% from the cercariae incubated with niclosamide-olamide or HESr at 1,000 and 500  $\mu$ g/mL, while the same rate (100%) was obtained after 90 min and 120 min of incubation respectively with 250 and 125  $\mu$ g/mL of HESr. The concentrations of 62.5 and 31.25  $\mu$ g/mL showed a 100% cercariae mortality rate at the 150 min time point. In addition, at 30 minutes time-point, HESr at 500 and 1,000  $\mu$ g/mL showed a significantly high cercaricidal activity (*p* < 0.001) in comparison to niclosamide one (Figure 3(b)).

**3.1.3. Median Lethal Concentration of *Sida acuta* and *Sida rhombifolia* Hydroethanolic Extracts.** As shown in Table 1, the cercaricidal median lethal concentration of *S. acuta* extract was almost constant from 60 to 150 mins of incubation. Conversely, the LC<sub>50</sub> profile was gradually decreased after cercariae incubation with *S. rhombifolia* extract. The LC<sub>50</sub> of HESa was then 28.41  $\pm$  3.45  $\mu$ g/mL and 18.63  $\pm$  0.28  $\mu$ g/mL after 30 min and 60 min of incubation respectively, while that of HESr were 172.42  $\pm$  26.16  $\mu$ g/mL and 56.60  $\pm$  4.07  $\mu$ g/mL at the same time points. After 150 min of incubation, both extracts disclosed the same LC<sub>50</sub> (18.15  $\pm$  0.00  $\mu$ g/mL).

**3.2. Effect of *Sida acuta* and *Sida rhombifolia* Hydroethanolic Extracts on Mouse Hepatic (Hepa 1–6) Cell Growth.** As indicated in Figure 4, the results of the effect of HESa and HESr extracts on the growth of the Hepa 1–6 cells at different concentrations (15.625–1,000  $\mu$ g/mL) showed a decrease in the viability of cells in a concentration-dependent manner after 24 h of incubation.

The inhibitory rates of *S. acuta* and *S. rhombifolia* extracts are shown in Table 2. At 31.25  $\mu$ g/mL, the inhibitory rate was 28.69% and 14.78% for HESa and HESr, respectively. The same tendency was observed at 125  $\mu$ g/mL where, after 24 h of incubation, inhibitory rates of 51.66% and 29.91% were recorded for HESa and HESr, respectively. Furthermore, the incubation of cells with the highest concentration (1,000  $\mu$ g/mL) shows inhibition rates of 85.69% for HESa and 51.49% for HESr (Table 2). Based on their inhibitory activity on the Hepa 1–6 cells, the IC<sub>50</sub> of HESa was 109.67  $\pm$  6.99  $\mu$ g/mL, and that of HESr was 888.79  $\pm$  29.94  $\mu$ g/mL.

Based on their cercaricidal activity and their cytotoxicity, the selectivity index of *S. acuta* and *S. rhombifolia* hydroethanolic extracts was calculated. At 30 min post-incubation, this index was 3.86 for HESa and 5.15 for HESr.

**3.3. Phytochemical Constituents of *Sida acuta* and *Sida rhombifolia* Hydroethanolic Extracts.** Qualitative phytochemical analysis of hydroethanolic extracts of *S. acuta* (HESa) and *S. rhombifolia* (HESr) revealed the presence of some secondary metabolites such as flavonoids, polyphenols, saponins, alkaloids, tannins, anthocyanins, triterpenes, sterols, anthraquinones, coumarins, and cardiac glycosides (Table 3).




FIGURE 2: Effect of *Sida acuta* hydroethanolic extract on *Schistosoma mansoni* cercariae viability (a) and mortality rate (b). All bars are expressed as mean  $\pm$  SEM. \*  $p < 0.05$  and \*\*\*  $p < 0.001$ , significantly different from controls (distilled water). #  $p < 0.05$  and ###  $p < 0.001$ , significantly different from reference control (niclosamide).

The HPLC-MS analysis of *S. rhombifolia* and *S. acuta* hydroethanolic extracts confirmed the presence of alkaloids, polyphenols, and flavonoids. The HPLC-MS chromatogram recorded for HESr was composed of peaks of chemical compounds, with retention times below 10 min (Figure 5) while those of HESa were below 15 min (Figure 6). The combination of the MS spectral data and the information from the literature allows tentative identification of 23 compounds (numbered 1–23 on the chromatograms).

The peak eluted at 0.34 min (compound 1), corresponding to the molecular ion  $[M + Na]^+$  detected at  $m/z$  203 was credited to glucose with a molecular mass of 203.05 (Figure 5(a)). Compound 2 appears at RT 3.06 min and produced the molecular ion  $[M + H]^+$  detected at  $m/z$  219. According to the obtained data, it was possible to infer the molecular mass of 219.09 and identify it as quindoline, an alkaloid (Figure 5(b)). Compound 3 was observed at RT

3.13 min with a molecular ion  $[M + Na]^+$  detected at  $m/z$  503 and was identified as 20-hydroxyecdysone, an ecdysteroid with the molecular mass of 503.29 (Figure 5(c)). Concerning compound 4 identification, it appeared at RT 3.50 min with a molecular ion  $[M + H]^+$  detected at  $m/z$  314 and was identified as the polyphenol, N-feruloyltyramine with 314.13 as molecular mass (Figure 5(d)). The peak eluted at 4.39 min was allotted to compound 8 with a molecular ion  $[M + Na]^+$  at  $m/z$  617 and was identified as tiliroside, a flavonoid with 617.12 as molecular mass (Figure 5(h)). The analysis of the HeSr spectrum also allowed to identify two coumarins, thamnsonin (compound 9, RT 4.77 min with  $[M + H]^+$  detected at  $m/z$  277) and scoparone (compound 11, RT 5.01 min with  $[M + H]^+$  detected at  $m/z$  207) with molecular masses of 277.14 and 207.09, respectively (Figures 5(i) and 5(k)). Moreover, at RT 6.30 min, compound 16 corresponding to the molecular ion  $[M + Na]^+$  detected at  $m/z$  413 was




FIGURE 3: Effect of *Sida rhombifolia* hydroethanolic extract on *Schistosoma mansoni* cercariae viability (a) and mortality rate (b). All bars are expressed as mean  $\pm$  SEM. \*  $p < 0.05$  and \*\*\*  $p < 0.001$ , significantly different from controls (distilled water). ###  $p < 0.001$ , significantly different from reference control (niclosamide).

TABLE 1: LC<sub>50</sub> values of *Sida acuta* and *Sida rhombifolia* hydroethanolic extracts on *Schistosoma mansoni* cercariae at different time points.

Time (min)	LC <sub>50</sub> (µg/mL)	
	HESa	HESr
30	28.41 $\pm$ 3.45	172.42 $\pm$ 26.16
60	18.63 $\pm$ 0.28	56.60 $\pm$ 4.07
90	18.74 $\pm$ 0.59	32.29 $\pm$ 4.05
120	18.15 $\pm$ 0.00	21.95 $\pm$ 1.54
150	18.15 $\pm$ 0.00	18.15 $\pm$ 0.00

identified as di-(2-ethylhexyl) phthalate with the molecular mass of 413.26 (Figure 5(p)). In addition to these identified compounds, compounds 10, 12, 13, 14, and 15 showed at 4.94, 5.07, 5.25, 5.38, and 5.57 min, respectively, were unidentified alkaloids. Also, compounds 5, 6, and 7, with RT 3.91 min, 3.98 min, and 4.20 min, respectively, showed a

molecular ion [M + Na]<sup>+</sup> at m/z 353, 239, and 259 (Figures 5(e)–5(g)) and were not identified (Table 4).

Regarding the HESa chromatogram (Figure 6), the peak eluted at 4.80 min (compound 17), corresponding to the molecular ion [M + Na]<sup>+</sup> detected at m/z 277 was credited to chrysin with a molecular mass of 277.09 (Figure 6(a)). Compound 18 appears at RT 5.19 min and produced the molecular ion [M + H]<sup>+</sup> detected at m/z 233 and according to the obtained data, it was possible to infer the molecular mass of 219.10 and identify it as cryptolepine, an alkaloid (Figure 6(b)). Compound 19 was observed at RT 5.61 min with a molecular ion [M + H]<sup>+</sup> detected at m/z 287 and was identified as kaempferol, a flavonoid with the molecular mass of 287.24 (Figure 5(c)). Concerning compound 20 identification, it appeared at RT 6.05 min with a molecular ion [M + H]<sup>+</sup> detected at m/z 229 and was identified as the coumarin, xanthyletin with 229.24 as molecular mass




FIGURE 4: Concentration-response curve of the effect of *Sida acuta* and *Sida rhombifolia* hydroethanolic extracts on Hepa 1–6 cells.

TABLE 2: Inhibition rates of the *Sida acuta* and *Sida rhombifolia* hydroethanolic extracts on the growth of the Hepa 1–6 cells.

Concentrations	Percentage of inhibition	
	HESa	HESr
15.625 µg/mL	12.42 ± 1.54	12.42 ± 0.54
31.25 µg/mL	28.69 ± 1.33	14.78 ± 0.73
62.5 µg/mL	44.20 ± 2.33	23.11 ± 1.33
125 µg/mL	51.66 ± 1.12	29.91 ± 0.91
250 µg/mL	64.53 ± 1.21	36.12 ± 1.41
500 µg/mL	75.62 ± 1.80	41.11 ± 1.44
1000 µg/mL	85.69 ± 0.41	51.49 ± 0.45

TABLE 3: Secondary metabolites of *Sida acuta* and *Sida rhombifolia* hydroethanolic extracts.

Phytochemicals	HESa	HESr
Anthocyanins	+	–
Triterpenes	+	+
Sterols	+	–
Flavonoids	+	+
Polyphenols	+	+
Saponins	+	+
Essential oils	–	–
Anthraquinones	+	–
Alkaloids	+	+
Tannins	–	+
Coumarins	+	+
Reducing sugar	–	+
Cardiac glycosides	+	+



(+): presence; (–): absence.

(Figure 6(d)). The peaks eluted at 6.48 min and 7.48 min were N-feruloyltyramine and thamnimonin corresponding to compounds 4 and 9, respectively. These compounds were previously identified in HESr (Figures 6(e) and 6(f)). The analysis of the HESr spectrum also allowed to identify two flavonoids, luteolin (compound 22, RT 8.26 min with [M + Na]<sup>+</sup> detected at m/z 309) and acetin (compound 23,


RT 12.30 min with [M + Na]<sup>+</sup> detected at m/z 284) with molecular masses of 309.24 and 284.26, respectively (Figures 6(h) and 6(i)). Moreover, at RT 7.75 min, compound 21, an alkaloid identified as cryptolepinone was obtained with the molecular ion [M + Na]<sup>+</sup> detected at m/z 249 and the molecular mass of 249.28 (Figure 6(g); Table 5). The chemical structures of all identified compounds from *S. acuta* and *S. rhombifolia* hydroethanolic extracts are shown in Figure 7.

#### 4. Discussion


The present investigation showed that hydroethanolic (30:70) extracts of *S. acuta* whole plant and *S. rhombifolia* aerial parts disclosed cercaricidal activity against *S. mansoni* cercariae in both time- and concentration-dependent manner. This confirms the larvicidal activity of isolated compounds from *S. rhombifolia* previously highlighted by Islam et al. [23]. Previous studies conducted by some authors have also reported the cercaricidal activity of *Glinus lotoides* fruits aqueous extract [29], *R. vomitoria* stem bark, roots ethanolic extract [7], and *O. pulcherrima* roots methanolic extract and fractions [10]. Other studies have also reported a time- and concentration-dependent molluscicidal, cercaricidal, and/or schistosomicidal activity of various plant




(a)




(b)



(c)




(d)




(e)

FIGURE 5: Continued.







(f)




(g)




(h)



(i)



(j)



(k)

FIGURE 5: Continued.







FIGURE 5: UPLC-DAD-MS UV profile of *Sida rhombifolia* hydroethanolic extract, identified peaks (1–16), and each identified compound spectrum (a–p).

extracts such as *A. indica* and *V. amygdalina* [9], *Milletia thonningii* [30, 31], and *Jatropha elliptica* [32]. Based on the median lethal concentration  $LC_{50}$  and at any defined period, the cercaricidal activity of *S. acuta* hydroethanolic extract was more important than that of *S. rhombifolia*. As compared with results from previous studies, *S. acuta* and *S. rhombifolia* hydroethanolic 30:70 extracts were more potent, regarding the  $LC_{50}$  and the effective time, in killing *S. mansoni* cercariae than *G. lotoides* fruits aqueous extract [29], *R. vomitoria* stem bark and roots ethanolic extracts [7],


*A. indica* and *V. amygdalina* methanolic extracts [9] but less effective than *O. pulcherrima* roots methanolic extract [10]. The cercaricidal activity of *S. acuta* and *S. rhombifolia* extracts might be due to their secondary metabolites such as flavonoids, alkaloids, and saponins. Abo-Zeid and Shohayeb [8] have shown the anti-miracidial and anti-cercarial activities of total alkaloids, saponins, and volatile oil extracted from *N. sativa* seeds hydroethanolic extract. It has also been demonstrated that alpinumisoflavone and robustic acid, two isoflavonoids isolated from *M. thonningii* seeds exhibit in



(a)




(b)




(c)


FIGURE 6: Continued.




(d)



(e)



(f)



(g)

FIGURE 6: Continued.




FIGURE 6: UPLC-DAD-MS UV profile of *Sida acuta* hydroethanolic extract, identified peaks, and each identified compound spectrum (a–i).

TABLE 4: Main signals exhibited in the LC-DAD-MS spectra of compounds detected in the hydroethanolic extract from *Sida rhombifolia* aerial parts and proposed attribution.

RT (min)	Exp. mass	Cald. mass	Molecular formula	Identified compounds	Structure
0.34	203.0527; [M + Na] <sup>+</sup>	203.0526	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub> Na	Glucose (sugar)	<b>1</b>
3.06	219.0986; [M + H] <sup>+</sup>	219.0922	C <sub>15</sub> H <sub>11</sub> N <sub>2</sub>	Quindoline (alkaloid)	<b>2</b>
3.13	503.2959; [M + Na] <sup>+</sup>	503.2979	C <sub>27</sub> H <sub>44</sub> O <sub>7</sub> Na	20-Hydroxycycdysone (steroid)	<b>3</b>
3.50	314.1371; [M + H] <sup>+</sup>	314.1387	C <sub>18</sub> H <sub>20</sub> NO <sub>4</sub>	N-Feruloyltyramine (polyphenol)	<b>4</b>
3.91	353.2281; [M + Na] <sup>+</sup>	353.2298	C <sub>20</sub> H <sub>34</sub> O <sub>5</sub> Na	NI	<b>5</b>
3.98	239.1246; [M + Na] <sup>+</sup>	239.1254	C <sub>11</sub> H <sub>20</sub> O <sub>4</sub> Na	NI	<b>6</b>
4.20	259.1693; [M + Na] <sup>+</sup>	259.1669	C <sub>15</sub> H <sub>24</sub> O <sub>2</sub> Na	NI	<b>7</b>
4.39	617.1259; [M + Na] <sup>+</sup>	617.1271	C <sub>30</sub> H <sub>26</sub> O <sub>13</sub> Na	Tiliroside (flavonoid)	<b>8</b>
4.77	277.1409; [M + H] <sup>+</sup>	277.0895	C <sub>15</sub> H <sub>17</sub> O <sub>5</sub>	Thamnosmonin (coumarin)	<b>9</b>
4.94	319.2227; [M + H] <sup>+</sup>	319.2227	C <sub>15</sub> H <sub>31</sub> N <sub>2</sub> O <sub>5</sub>	NI (alkaloids)	<b>10</b>
5.01	207.0982; [M + H] <sup>+</sup>	207.0657	C <sub>11</sub> H <sub>11</sub> O <sub>4</sub>	Scoparone (coumarin)	<b>11</b>
5.07	463.2637; [M + Na] <sup>+</sup>	463.2626	C <sub>19</sub> H <sub>40</sub> N <sub>2</sub> O <sub>9</sub>	NI (alkaloid)	<b>12</b>
5.25	363.2484; [M + H] <sup>+</sup>	363.2490	C <sub>17</sub> H <sub>35</sub> N <sub>2</sub> O <sub>6</sub>	NI (alkaloid)	<b>13</b>
5.38	635.4454; [M + Na] <sup>+</sup>	635.4453	C <sub>30</sub> H <sub>64</sub> N <sub>2</sub> O <sub>10</sub> Na	NI (alkaloid)	<b>14</b>
5.57	377.2641; [M + H] <sup>+</sup>	377.2646	C <sub>18</sub> H <sub>37</sub> N <sub>2</sub> O <sub>6</sub>	NI (alkaloid)	<b>15</b>
6.30	413.2643; [M + Na] <sup>+</sup>	413.2663	C <sub>24</sub> H <sub>38</sub> O <sub>4</sub> Na	Di-(2-ethylhexyl) phthalate	<b>16</b>

NI: not identified.

*in vitro* cercaricidal activity [30, 31]. Moreover, Dos Santos et al. [32] reported a strong cercaricidal activity of diethyl 4-phenyl-2,6-dimethyl-3,5 pyridine dicarboxylate, a penta-substituted pyridine alkaloid from the rhizome of *J. elliptica* with an LC<sub>100</sub> of 2 µg/mL. The cercaricidal potential of these secondary metabolites could strongly support the efficacy of *S. acuta* and *S. rhombifolia* extracts in killing *S. mansoni* cercariae. Indeed, phytochemicals belonging to flavonoids, polyphenols, coumarins, alkaloids, and steroids were

isolated from *S. rhombifolia* hydroethanolic 30:70 extract in this study. The cercaricidal activity of *S. acuta* and *S. rhombifolia* may be due to their ability to damage the cercariae tegument and disturb its motor activity. Xiao et al. [33] have previously demonstrated that exposure of *S. japonicum* cercariae to praziquantel results in intensive disturbance in motor activity and lysis of cercarial tissues, followed by an extensive release of gland contents and separation of the tail from the body. These observations were

TABLE 5: Main signals exhibited in the LC-DAD-MS spectra of compounds detected in the hydroethanolic extract from *Sida acuta* the whole plant and proposed attribution.

RT (min)	Exp. mass	Cald. mass	Molecular formula	Identified compounds	Structure
4.80	277.0977; [M + Na] <sup>+</sup>	277.24	C <sub>15</sub> H <sub>10</sub> O <sub>4</sub> Na	Chrysin (flavonoid)	<b>17</b>
5.19	233.1090; [M + H] <sup>+</sup>	233.28	C <sub>16</sub> H <sub>12</sub> N <sub>2</sub>	Cryptolepine (alkaloid)	<b>18</b>
5.61	287.0545; [M + H] <sup>+</sup>	287.24	C <sub>15</sub> H <sub>10</sub> O <sub>6</sub>	Kaempferol (flavonoid)	<b>19</b>
6.05	229.0828; [M + H] <sup>+</sup>	229.24	C <sub>14</sub> H <sub>13</sub> O <sub>3</sub>	Xanthyletin (coumarin)	<b>20</b>
6.48	314.1371; [M + H] <sup>+</sup>	314.13	C <sub>18</sub> H <sub>21</sub> NO <sub>4</sub>	N-Feruloyltyramine (polyphenol)	<b>4</b>
7.48	277.2147; [M + H] <sup>+</sup>	277.14	C <sub>11</sub> H <sub>20</sub> O <sub>4</sub>	Thamnosmonin (coumarin)	<b>9</b>
7.75	249.1035; [M + H] <sup>+</sup>	249.28	C <sub>16</sub> H <sub>13</sub> N <sub>2</sub> O	Cryptolepinone (alkaloid)	<b>21</b>
8.26	309.2067; [M + Na] <sup>+</sup>	309.24	C <sub>15</sub> H <sub>10</sub> O <sub>6</sub> Na	Luteolin (flavonoid)	<b>22</b>
12.30	284.2629; [M + Na] <sup>+</sup>	284.26	C <sub>16</sub> H <sub>12</sub> O <sub>5</sub> Na	Acacetin (flavonoid)	<b>23</b>




FIGURE 7: Chemical structures of identified compounds from *Sida acuta* and *Sida rhombifolia* hydroethanolic extracts.

followed by the cercariae surface damages that are characterized by the decrease of the membranous glycocalyx, the swelling, and the degeneration of mitochondria distributed in the muscle and parenchymal cells as well as the lysis of the tegumental muscular layer. In fact, in our study, the incubation of *S. mansoni* cercaria with *S. acuta* and *S. rhombifolia* hydroethanolic extracts resulted in the separation of their tail from their body.

The genus *Sida* L. is one of the most diverse in the Malvaceae family, with about 200 species scattered in every part of tropical and subtropical regions in the world [18, 34, 35]. In addition, 142 chemical constituents belonging to various classes have been reported for *Sida* sp. Alkaloids, flavonoids, and ecdysteroids were predominant and reported mostly from *S. acuta*, *S. cordifolia*, *S. rhombifolia*, *S. glutinosa*, and *S. spinosa*. [36]. In our study, alkaloids were the major bioactive principles of *S. acuta* and *S. rhombifolia*

hydroethanolic 30:70 extracts. Moreover, among the 15 compounds identified in those extracts, only thamnosmonin and xanthyletin (coumarins), and tiliroside (flavonoid) were recently identified in *S. rhombifolia* hydroethanolic extract [37]. Some of these compounds have been previously identified and/or isolated from the *Sida* L. genus. In fact, quindoline has been isolated from *S. cordifolia* and *S. rhombifolia* [38, 39], while 20-hydroxyecdysone has been isolated from *S. spinosa* [40] *S. cordifolia* [41], *S. tuberculata* [42–44], and *S. acuta* [37]. Furthermore, N-feruloyltyramine and di-(2-ethylhexyl) phthalate have been isolated from *S. acuta* [45] and scoparone from *S. rhombifolia* [46]. Moreover, Das et al. [47] identified chrysin in *S. glutinosa* and Silva et al. [48] isolated luteolin from *S. galheirensis*. Acacetin and kaempferol were isolated from *S. rhombifolia* [38, 39], while cryptolepine and cryptolepinone were characterized from *S. acuta* [38, 39]. These

phytoconstituents exhibited various pharmacological activities such as anti-microbial, anti-plasmodial, vasorelaxant, anti-oxidant, and anti-inflammatory [38, 42, 44, 49, 50].

Regarding the cytotoxic activity of both extracts on the growth of the Hepa 1–6 cells, *S. acuta* extract disclosed the highest anti-proliferative activity than *S. rhombifolia*. Anti-proliferative activity of a plant extract may be linked to its phytochemical constituents. It has been shown that quindoline and cryptolepinone, as well as N-trans-feruloyltyramine identified in *S. acuta* and *S. rhombifolia* hydroethanolic extracts have a significant anti-proliferative activity on mouse hepatoma cells (Hepa 1c1c7) by inhibiting the quinone reductase activity [46].

## 5. Conclusions

*S. acuta* and *S. rhombifolia* hydroethanolic extracts disclosed cercaricidal activity and were not toxic on Hepa 1–6 cell line. However, Based on the LC<sub>50</sub>, the cercaricidal activity was more pronounced with *S. acuta* extract. The cercaricidal activity of these plant extracts may be linked to the presence of several secondary metabolite such as alkaloids, flavonoids, and coumarins. Based on their activity and their selectivity index, *S. rhombifolia* extract could be more effective on *S. mansoni* cercariae than *S. acuta* extract. This study could provide baseline information for further investigations aiming to develop plant-based alternative drugs against *S. mansoni*.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this study.

## Acknowledgments

The authors are grateful to the “Deutscher Akademischer Austauschdienst” (DAAD) for financial support through the Yaounde-Bielefeld Graduate School for Natural Products with Antiparasitic and Antibacterial Activities (YaBINaPA, grant no. 57316173).

## References

- [1] WHO, *Schistosomiasis*, WHO, Geneva, Switzerland, 2020, <https://www.who.int/news-room/fact-sheets/detail/schistosomiasis>.
- [2] A. P. Castro, A. C. Alves de Mattos, A. C. Alves de Mattos, R. L. Martins Souza, M. José Marques, and M. Henrique Dos Santos, “Medicinal plants and their bioactive constituents: a review of bioactivity against *Schistosoma mansoni*,” *Journal of Medicinal Plants Research*, vol. 7, no. 21, pp. 1515–1522, 2013.
- [3] C. Collins, J. Xu, and S. Tang, “Schistosomiasis control and the health system,” *Infectious Diseases of Poverty*, vol. 1, 2012.
- [4] K. H. Abo Zaid, H. El-Wakil, A. El-Hussein, S. Jomaa, and M. Shohayeb, “Evaluation of the molluscicidal activity of *Punica granatum*, *Calotropis procera*, *Solanum incanum* and *Citrullus colocynthis* against *Biomphalaria arabica*,” *World Applied Sciences Journal*, vol. 26, no. 7, pp. 873–879, 2013.
- [5] WHO, *Report of an Informal Consultation on Schistosomiasis Control*, WHO, Geneva, Switzerland, 2013.
- [6] I. Takougang, J. Meli, J. Wabo Poné, and F. Angwafo, “Community acceptability of the use of low-dose niclosamide (Bayluscide), as a molluscicide in the control of human schistosomiasis in Sahelian Cameroon,” *Annals of Tropical Medicine and Parasitology*, vol. 101, no. 6, pp. 479–486, 2007.
- [7] E. M. Tekwu, K. M. Bosompem, W. K. Anyan et al., “In vitro assessment of anthelmintic activities of *Rauwolfia vomitoria* (Apocynaceae) stem bark and roots against parasitic stages of *Schistosoma mansoni* and cytotoxic study,” *Journal of parasitology research*, vol. 2017, Article ID 2583969, 11 pages, 2017.
- [8] K. Abo-zeid and M. Shohayeb, “Evaluation of the biocidal activity of alkaloids, saponins and volatile oil extracted from *Nigella sativa* seeds against miracidia and cercariae of *Schistosoma mansoni*,” *International Journal of Pharmaceutical Science Invention*, vol. 4, no. 1, pp. 47–54, 2015.
- [9] D. O. Acheampong, N. Owusu-Adzorah, F. A. Armah et al., “Ethnopharmacological evaluation of schistosomicidal and cercaricidal activities of some selected medicinal plants from Ghana,” *Tropical Medicine and Health*, vol. 48, no. 1, p. 19, 2020.
- [10] N. G. Feussom, H. B. Jatsa, M. C. Kenfack et al., “In vitro activity of *Ozoroa pulcherrima* Schweinf. extracts and fractions on *Schistosoma mansoni* cercariae and adult worms,” *European Journal of Medicinal Plants*, vol. 31, pp. 17–30, 2020.
- [11] R. S. Pawar, A. Jain, P. Sharma, P. K. Chaurasiya, and P. K. Singour, “In vitro studies on *Sida cordifolia* Linn for anthelmintic and antioxidant properties,” *Chinese Medicine*, vol. 2, no. 2, pp. 47–52, 2011.
- [12] R. Wake, N. Patil, and U. K. Halde, “Genus *Sida*—the plants with ethno medicinal and therapeutic potential,” *Golden Research Thoughts*, vol. 1, no. 2231, 2011.
- [13] F. G. Coe and G. J. Anderson, “Screening of medicinal plants used by the Garífuna of Eastern Nicaragua for bioactive compounds,” *Journal of Ethnopharmacology*, vol. 53, no. 1, pp. 29–50, 1996.
- [14] D. Karou, M. H. Dicko, S. Sanon, J. Simporé, and A. S. Traore, “Antimalarial activity of *Sida acuta* Burm. F. (Malvaceae) and *Pterocarpus erinaceus* Poir. (Fabaceae),” *Journal of Ethnopharmacology*, vol. 89, no. 2–3, pp. 291–294, 2003.
- [15] O. Tene Tcheghebe, F. Ngouafong Tatong, A. Seukep, J. Kamga, and J. Nenwa, “Ethnobotanic survey of medicinal plants used for malaria therapy in western Cameroon,” *Journal of medicinal plants studies*, vol. 4, no. 3, pp. 248–258, 2016.
- [16] G. Mallikarjuna, Prabhakaran, and B. Saratkumar Reddy, “Anticancer activity of *Sida acuta* Burm.F against nitrosodiethylamine and CCl<sub>4</sub> induced hepatocellular carcinoma,” *Indo American Journal of Pharmaceutical Research*, vol. 3, no. 54, 2013.
- [17] J. Adjanohoun, N. Aboubakar, and K. Dramane, *Traditional Medicine and Pharmacopeia- Contribution to Ethnobotanical and Floristic Studies in Cameroon*, Scientific, Technical, and Research Commission of the Organization of African Unity, Lagos, Nigeria, 1996.
- [18] F. C. Rodrigues and A. F. M. de Oliveira, “The genus *Sida* L. (Malvaceae): an update of its ethnomedicinal use, pharmacology and phytochemistry,” *South African Journal of Botany*, vol. 132, pp. 432–462, 2020.

- [19] H. Boukeng Jatsa, C. A. de Jesus Pereira, A. B. Dias Pereira et al., "In vitro evaluation of *Sida pilosa* Retz (Malvaceae) aqueous extract and derived fractions on *Schistosoma mansoni*," *Pharmacology & Pharmacy*, vol. 6, no. 8, pp. 380–390, 2015.
- [20] H. B. Jatsa, R. C. Russo, C. A. D. J. Pereira et al., "Improvement of the liver pathology by the aqueous extract and the n-butanol fraction of *Sida pilosa* Retz in *Schistosoma mansoni*-infected mice," *Journal of Ethnopharmacology*, vol. 180, pp. 114–123, 2016.
- [21] H. B. Jatsa, U. M. Femoe, J. Njiza et al., "Efficacy of *Sida pilosa* Retz aqueous extract against *Schistosoma mansoni*-induced granulomatous inflammation in the liver and the intestine of mice: histomorphometry and gastrointestinal motility evaluation," *BMC Complementary and Alternative Medicine*, vol. 18, no. 1, p. 247, 2018.
- [22] S. Akilandeswari, R. Senthamarai, S. Prema, and R. Valarmathi, "Antimicrobial activity of leaf extracts of *Sida acuta* Burm.," *International Journal of Pharma Sciences and Research*, vol. 1, no. 5, pp. 248–250, 2010.
- [23] M. E. Islam and N. A. Khatune, "Larvicidal activity of a new glycoside, phenyl ethyl  $\beta$ -D glucopyranoside from the stem of the plant *Sida rhombifolia*," *Pakistan Journal of Biological Sciences*, vol. 6, no. 1, pp. 73–75, 2002.
- [24] M. Govindarajan, "Larvicidal and repellent activities of *Sida acuta* Burm. F. (Family: Malvaceae) against three important vector mosquitoes," *Asian Pacific Journal of Tropical Medicine*, vol. 3, no. 9, pp. 691–695, 2010.
- [25] M. C. Holtfreter, M. Lobermann, S. Klammt et al., "*Schistosoma mansoni*: schistosomicidal effect of mefloquine and primaquine in vitro," *Experimental Parasitology*, vol. 127, no. 1, pp. 270–276, 2011.
- [26] M. M. Eissa, S. El Bardicy, and M. Tadros, "Bioactivity of miltefosine against aquatic stages of *Schistosoma mansoni*, *Schistosoma haematobium* and their snail hosts, supported by scanning electron microscopy," *Parasites & Vectors*, vol. 4, no. 1, pp. 73–11, 2011.
- [27] S. Murata, R. Shiragami, C. Kosugi et al., "Antitumor effect of 1, 8-cineole against colon cancer," *Oncology Reports*, vol. 30, no. 6, pp. 2647–2652, 2013.
- [28] G. E. Trease and W. C. Evans, *Pharmacognosy*, London, UK, 1989.
- [29] G. Kiros, B. Erko, M. Giday, and Y. Mekonnen, "Laboratory assessment of molluscicidal and cercariacidal effects of *Glinus lotoides* fruits," *BMC Research Notes*, vol. 7, no. 1, pp. 220–227, 2014.
- [30] S. Perrett, P. J. Whitfield, A. Bartlett, and L. Sanderson, "Attenuation of *Schistosoma mansoni* cercariae with a molluscicide derived from *Milletia thonningii*," *Parasitology*, vol. 109, no. 5, pp. 559–563, 1994.
- [31] J. R. A. Lyddiard, P. J. Whitfield, and A. Bartlett, "Antischistosomal bioactivity of isoflavonoids from *Milletia thonningii* (leguminosae)," *The Journal of Parasitology*, vol. 88, no. 1, pp. 163–170, 2002.
- [32] A. F. Dos Santos, S. A. Fonseca, F. A. César, M. C. P. de Azevedo Albuquerque, J. V. Santana, and A. E. G. Santana, "A penta-substituted pyridine alkaloid from the rhizome of *Jatropha elliptica* (Pohl) Muell. Arg. is active against *Schistosoma mansoni* and *Biomphalaria glabrata*," *Parasitology Research*, vol. 113, no. 3, pp. 1077–1084, 2014.
- [33] S.-H. Xiao, J. Keiser, M.-G. Chen, M. Tanner, and J. Utzinger, "Research and development of antischistosomal drugs in the people's Republic of China," *Important Helminth Infections in Southeast Asia: Diversity and Potential for Control and Elimination, Part B*, vol. 73, pp. 231–295, 2010.
- [34] J. L. Brandão, G. S. Baracho, M. F. de Sales, and M. P. V. Filho, "Synopsis of *Sida* (Malvaceae, malvoideae, malveae) in the state of Pernambuco, Brazil," *Phytotaxa*, vol. 307, no. 3, pp. 205–227, 2017.
- [35] N. S. Aminah, E. R. Laili, M. Rafi, A. Rochman, M. Insanu, and K. N. W. Tun, "Secondary metabolite compounds from *Sida* genus and their bioactivity," *Heliyon*, vol. 7, no. 4, p. e06682, 2021.
- [36] B. Dinda, N. Das, S. Dinda, M. Dinda, and I. Silsarma, "The genus *Sida* L.-a traditional medicine: its ethnopharmacological, phytochemical and pharmacological data for commercial exploitation in herbal drugs industry," *Journal of Ethnopharmacology*, vol. 176, pp. 135–176, 2015.
- [37] B. C. Kamdoum, I. Simo, and S. C. N. Woumba, "Chemical constituents of two Cameroonian medicinal plants: *Sida rhombifolia* L. and *Sida acuta* Burm. f. (Malvaceae) and their antiplasmodial activity," *Natural Product Research*, vol. 14, 2021.
- [38] O. S. Chaves, Y. C. Teles, M. M. Monteiro et al., "Alkaloids and phenolic compounds from *Sida rhombifolia* L. (Malvaceae) and vasorelaxant activity of two indoquinoline alkaloids," *Molecules*, vol. 22, no. 1, 2017.
- [39] O. Chaves, R. Gomes, A. Tomaz et al., "Secondary metabolites from *Sida rhombifolia* L. (Malvaceae) and the vasorelaxant activity of cryptolepinone," *Molecules*, vol. 18, no. 3, pp. 2769–2777, 2013.
- [40] F. M. M. Darwish and M. G. Reinecke, "Ecdysteroids and other constituents from *Sida spinosa* L.," *Phytochemistry*, vol. 62, no. 8, pp. 1179–1184, 2003.
- [41] A. N. Jadhav, C. S. Rumalla, B. Avula, and I. A. Khan, "HPTLC method for determination of 20-hydroxyecdysone in *Sida rhombifolia* L. and dietary supplements," *Chromatographia*, vol. 66, no. 9–10, pp. 797–800, 2007.
- [42] H. S. Da Rosa, M. Koetz, M. C. Santos et al., "Extraction optimization and UHPLC method development for determination of the 20-hydroxyecdysone in *Sida tuberculata* leaves," *Steroids*, vol. 132, pp. 33–39, 2018.
- [43] H. S. Rosa, I. S. Coelho, M. D. Silva et al., "*Sida tuberculata* extract reduces the nociceptive response by chemical noxious stimuli in mice: implications for mechanism of action, relation to chemical composition and molecular docking," *Phytotherapy Research*, vol. 33, no. 1, pp. 224–233, 2019.
- [44] S. Kumar, P. K. Lakshmi, C. Sahi, and R. S. Pawar, "*Sida cordifolia* accelerates wound healing process delayed by dexamethasone in rats: effect on ROS and probable mechanism of action," *Journal of Ethnopharmacology*, vol. 235, pp. 279–292, 2019.
- [45] H. S. Da Rosa, V. B. De Camargo, G. Camargo, C. V. Garcia, A. M. Fuentefria, and A. S. L. Mendez, "Ecdysteroids in *Sida tuberculata* R.E. Fries (Malvaceae): chemical composition by LC-ESI-MS and selective anti-*Candida krusei* activity," *Food Chemistry*, vol. 182, pp. 193–199, 2015.
- [46] D. S. Jang, E. J. Park, Y.-H. Kang et al., "Compounds obtained from *Sida acuta* with the potential to induce quinone reductase and to inhibit 7,12-dimethylbenz-[a]anthracene-induced preneoplastic lesions in a mouse mammary organ culture model," *Archives of Pharmacal Research*, vol. 26, no. 8, pp. 585–590, 2003.
- [47] N. Das, J. Nath, and B. Dinda, "Antioxidant phytochemicals from *Sida glutinosa*," *Journal of Pharmacy Research*, vol. 5, no. 9, pp. 4845–4848, 2012.



- [48] D. A. E. Silva, T. M. S. D. Silva, A. C. D. S. Lins et al., "Constituintes químicos e atividade antioxidante de *Sida galheirensis* Ulbr. (Malvaceae)," *Química Nova*, vol. 29, no. 6, pp. 1250–1254, 2006.
- [49] D. S. Preethidan, G. Arun, and M. P. Surendran, "Lip-oxygenase inhibitory activity of some *Sida* species due to di (2-ethylhexyl) phthalate," *Current Science*, vol. 105, pp. 232–234, 2013.
- [50] D. Karou, A. Savadogo, and A. Canini, "Antibacterial activity of alkaloids from *Sida acuta*," *African Journal of Biotechnology*, vol. 4, no. 12, pp. 1452–1457, 2005.