

Supplementary Table 1: Risk factors for bacteremia (n = 34)

	No. of patients (n)	No. of bacteremia patients (n, %)	Adjusted OR (95% CI)	p-value
ISS				*0.003
≤ 16	416	7 (1.9%)		
17 >	443	27 (6.1%)		
Head AIS				0.705
0	572	22 (3.8%)		
1-3	210	11 (5.2%)		
4-6	77	1 (1.3%)		
Face AIS				0.125
0	630	23 (3.7%)		
1-3	213	8 (3.8%)		
4-6	16	3 (19.7%)		
Chest AIS				0.378
0	392	15 (3.8%)		
1-3	406	14 (3.4%)		
4-6	61	5 (8.2%)		
Abdominal AIS				*0.001
0	444	8 (1.8%)		
1-3	322	17 (5.3%)		
4-6	93	9 (9.7%)		
Extremity AIS				*0.028
0	403	11 (2.7%)		
1-3	382	17 (4.5%)		
4-6	74	6 (8.1%)		
External AIS				0.086
0	718	24 (3.3%)		
1-3	131	10 (7.6%)		
4-6	10	0		
Rib fracture	335	17 (5.1%)		0.245
Hemothorax	111	8 (7.2%)		0.105
Liver injury	92	8 (8.7%)	2.709 (1.027-6.415)	*0.022
Pancreas injury	19	3 (15.8%)	4.874 (0.869-18.362)	*0.036
Bowel perforation	57	9 (15.8%)	5.804 (2.256-13.759)	*0.001
Pelvic injury	217	14 (6.5%)	2.143 (0.982-4.554)	*0.047
Degloving injury	22	1 (4.5%)		0.593
Urogenital injury	27	2 (7.4%)		0.290
Shock	161	13 (8.1%)	2.827 (1.349-5.764)	*0.006
Massive Transfusion	70	10 (14.3%)	5.337 (2.323-11.452)	*0.001

406ISS, Injury severity score; AIS, abbreviated injury scale.

407* Significance set as $p = 0.05$