

Research Article

Multivariate Analysis of Fruit Antioxidant Activities of Blackberry Treated with 1-Methylcyclopropene or Vacuum Precooling

Jian Li, Guowei Ma, Lin Ma, Xiaolin Bao, Liping Li, Qian Zhao, and Yousheng Wang 

Beijing Advanced Innovation Center for Food Nutrition and Human Health, Beijing Engineering and Technology Research Center of Food Additives, Beijing Technology and Business University, Beijing 100048, China

Correspondence should be addressed to Yousheng Wang; wangys@btbu.edu.cn

Received 29 September 2017; Accepted 31 December 2017; Published 1 February 2018

Academic Editor: Ying Yang

Copyright © 2018 Jian Li et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Effects of 1-methylcyclopropene (1-MCP) and vacuum precooling on quality and antioxidant properties of blackberries (*Rubus* spp.) were evaluated using one-way analysis of variance, principal component analysis (PCA), partial least squares (PLS), and path analysis. Results showed that the activities of antioxidant enzymes were enhanced by both 1-MCP treatment and vacuum precooling. PCA could discriminate 1-MCP treated fruit and the vacuum precooled fruit and showed that the radical-scavenging activities in vacuum precooled fruit were higher than those in 1-MCP treated fruit. The scores of PCA showed that H_2O_2 content was the most important variables of blackberry fruit. PLSR results showed that peroxidase (POD) activity negatively correlated with H_2O_2 content. The results of path coefficient analysis indicated that glutathione (GSH) also had an indirect effect on H_2O_2 content.

1. Introduction

Blackberries (*Rubus* spp.) are notable for their antioxidant activities, particularly due to their high contents of polyphenolic compounds, such as ellagic acid, quercetin, gallic acid, anthocyanins, and cyanidins, plus excellent contents of the antioxidant vitamins A and C [1–3]. However, blackberries have a short market life and the fruit quality is rapidly reduced after harvesting [4, 5].

Reactive oxygen species (ROS), like hydrogen peroxide (H_2O_2) and superoxide $O^{\bullet-}$, could promote the oxidation of proteins and lipids and thus lead to fruit senescence and a decrease of fruit quality [6, 7]. The antioxidant enzymes, including catalase (CAT), ascorbate peroxidase (APX), peroxidase (POD), and polyphenol oxidase (PPO), can prevent the accumulation of ROS and repair oxidative damage [8]. The antioxidants, such as polyphenolic compounds, also play an important role in scavenging excess ROS [9]. There are many correlated factors affecting the fruit quality. Hence, application of a multivariate technique to characterize the relationship among the antioxidant enzymes' activities, ROS levels, and antioxidant activities seems to be appropriate.

Principal component analysis (PCA) is a multivariate technique used to extract the important information from multivariate data [10]. Partial least squares (PLS) and path analysis can be especially useful to find the correlations between intercorrelated quantitative dependent variables. PLS has been used to predict the changes of quality of pasteurized pineapple juice during storage [11].

In this study, the quality and antioxidant parameters were collected from 1-MCP treated, vacuum precooled, and control fruit. Principal component analysis (PCA) was applied to evaluate the effect of 1-MCP and vacuum precooling on physiological properties of fruit. Correlations between quality and antioxidant parameters were studied through partial least squares (PLS) and path analysis.

2. Materials and Methods

2.1. Plant Material and Treatment. Blackberry fruit (*Rubus* spp. cv. Triple Crown) at the commercially mature stage was harvested from an orchard near Beijing. Fruit was sorted to eliminate damaged and diseased fruit and selected for uniformity in size and color. Fruit was randomized and

divided into 3 lots for the following treatments: (1) untreated and referred to as the control and (2) treated with 5 $\mu\text{g/L}$ 1-methylcyclopropene (1-MCP). Fruits were placed in sealed 250 L plastic chambers with 1-MCP powdered formulation (1250 μg 1-MCP release) at room temperature for 24 h and (3) precooled in vacuum cooler at 10 °C for 1 h.

After the treatments, all the fruits were placed in 60 cm \times 37 cm \times 20 cm plastic containers and then stored at 0 °C with 85–95% RH.

2.2. Fruit Quality Measurement. Flesh firmness of the fruit was measured using a texture analyzer (LFRA, Brookfield Ltd., USA). Fifteen fruits were measured at each sampling time.

Fruit of each treatment was distributed into three groups (30 fruit per group), and each group represented one replicate. The number of decayed fruits in each plastic container was counted and decay rate was calculated as percentage of decayed fruit versus total fruit.

2.3. Enzymatic Activity Analysis. For the ascorbate peroxidase (APX), peroxidase (POD), and polyphenol oxidase (PPO) activities assay, 10.0 g samples were thoroughly homogenized with 20 mL extracting buffer (pH 7.8 100 mM phosphate buffer containing 0.2 g polyvinylpyrrolidone) and centrifuged at 10,000 $\times g$ at 4 °C for 20 min, and the supernatant was collected and stored at –80 °C for further analysis.

APX activity was determined spectrophotometrically by monitoring the decline in absorbance at 290 nm as ascorbate was oxidized [12]. APX activity was expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$.

POD activity was measured as the oxidation of guaiacol in the presence of H_2O_2 by measuring the absorbance at 460 nm [13]. The POD activity was expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$.

PPO activity was measured according to the method of Jang et al. and expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$ [13].

2.4. GSH Assays. Glutathione (GSH) was extracted from 10.0 g of the flesh tissue with 20 mL of ice-cold 5% trichloroacetic acid containing 5 mM Ethylenediaminetetraacetic Acid (EDTA) and then centrifuged at 4 °C for 10 min at 10,000 $\times g$. The supernatant was assayed for GSH according to the method of Guri, and the GSH content was expressed as $\text{mg}\cdot 100\text{g}^{-1}\text{FW}$ [14].

2.5. H_2O_2 Assays. H_2O_2 was extracted by homogenizing 10.0 g of fruit tissue in 20 ml of cold acetone and was measured according to the method of Brennan and Frenkel [15]. The H_2O_2 content was expressed as $\text{mg}\cdot 100\text{g}^{-1}\text{FW}$.

2.6. Antioxidant Activities Assay. For antioxidant activities assay, 10.0 g samples were thoroughly homogenized with 20 mL methyl alcohol and centrifuged at 10,000 $\times g$ at 4 °C for 20 min. The supernatant was used for antioxidant activities, total phenol, and total flavonoid concentration assays.

Total antioxidant activity was measured using ferric reducing antioxidant potential assay (FARP) [16]. The values were expressed as the concentration of antioxidants having a ferric reducing ability equivalent to that of 1 mmol/L FeSO_4 .

Trolox equivalent antioxidant capacity (TEAC) was determined according to the method of Arts et al. [17]. 50% of the 2,2'-Azinobis-(3-ethylbenzothiazoline-6-sulphonate) (ABTS) radical-scavenging activity is defined as one activity unit. TEAC activity was expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$.

The 2,2-diphenylpicrylhydrazyl (DPPH) radical-scavenging activity was assayed by the method of Shon et al. [18]. 50% of the DPPH radical-scavenging activity is defined as an activity unit and the DPPH radical-scavenging activity was expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$.

Superoxide anion scavenging activity was measured by Nitrotetrazolium Blue chloride (NBT) reduction method [19]. 50% of the superoxide anion scavenging activity is defined as an activity unit. Superoxide anion scavenging activity was expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$.

Hydroxyl radical-scavenging activity is determined by the method of Shon et al. [18]. 50% of the hydroxyl radical-scavenging activity is defined as an activity unit. Hydroxyl radical-scavenging activity was expressed as $\text{U}\cdot\text{g}^{-1}\text{FW}$.

The total phenolic concentration of flesh extracts was measured using a modified Folin–Ciocalteu colorimetric method [20]. Absorbance was measured at 760 nm after 60 min at room temperature. The results were expressed as micrograms of gallic acid equivalents per gram of fresh weight.

The total flavonoid concentration of flesh extracts was determined using a colorimetric assay [20]. The absorbance of the solution versus a blank at 510 nm was measured after 60 min. The results were expressed as micrograms of catechin equivalents per gram of fresh weight.

2.7. Statistical Analysis. All data were analyzed by one-way analysis of variance (ANOVA) with SPSS 11.0 statistical software. Significant differences were performed by a least significant difference method (LSD test, $P \leq 0.05$) for all treatments at different sampling times.

For multivariate analysis, data were centered and weighted by the inverse of the standard deviation of each variable in order to avoid dependence on measured units. Principal component analysis (PCA) and partial least squares (PLS) in this study were performed using Unscrambler 9.7 statistical software. Path analysis model was developed by DPS (v.8.01) software.

3. Results and Discussion

3.1. One-Way Analysis of Variance. 1-Methylcyclopropene (1-MCP) has been proved to slow down the ripening of some fruits [21, 22]. In this work, the decay rate of blackberry fruit was reduced significantly ($P < 0.05$) by 1-MCP treatment (Table 1). The antioxidant enzymes, polyphenol oxidase (PPO), in 1-MCP treated fruit were also higher than control. On the other hand, a higher firmness was observed with vacuum precooling compared to control after 38 days of storage. The PPO activities were also enhanced by vacuum precooling after storage at 0 °C for 38 days. Our finding suggested that both 1-MCP and vacuum precooling treatment had a potential value in delaying the senescence of blackberries.

TABLE 1: Changes of quality and reactive oxygen metabolism indexes in blackberries during postharvest storage with different treatments.

	Harvest	Days of storage					
		Control		1-MCP		Vacuum precooling	
		21 d	38 d	21 d	38 d	21 d	38 d
Firmness	21.38 ^d	12.33 ^b	6.09 ^a	15.00 ^{bc}	5.84 ^a	17.76 ^{bcd}	18.31 ^{cd}
Decay rate (%)	0.00 ^a	0.00 ^a	21.67 ^c	0.00 ^a	11.67 ^b	0.00 ^a	18.34 ^c
H ₂ O ₂ (mg/100 g fw)	6.28 ^a	16.76 ^b	23.52 ^d	18.92 ^c	31.76 ^f	16.99 ^b	26.22 ^e
GSH (mg/100 g fw)	61.55 ^d	65.89 ^e	26.62 ^a	80.80 ^f	39.43 ^c	97.02 ^g	35.22 ^b
APX (U/g fw)	0.02 ^a	0.03 ^a	0.09 ^c	0.05 ^{ab}	0.08 ^{bc}	0.02 ^a	0.04 ^a
POD (U/g fw)	105.71 ^b	5.58 ^a	1.32 ^a	5.41 ^a	4.17 ^a	5.09 ^a	4.96 ^a
PPO (U/g fw)	1.54 ^f	0.75 ^b	0.59 ^a	0.92 ^c	0.91 ^c	1.2 ^e	1.1 ^d
FRAP (U/g fw)	45.43 ^c	30.04 ^b	36.30 ^c	28.03 ^a	42.50 ^d	36.86 ^c	48.96 ^f
TEAC (U/g fw)	0.2232 ^b	0.1980 ^a	0.2172 ^b	0.2609 ^c	0.3192 ^f	0.2937 ^e	0.2824 ^d
DPPH (U/g fw)	31.74 ^c	30.51 ^a	31.24 ^b	31.73 ^c	38.30 ^e	31.22 ^b	36.61 ^d
NBT (U/g fw)	7.80 ^a	33.18 ^e	26.18 ^c	35.69 ^f	24.25 ^b	36.96 ^g	30.77 ^d
Hydroxyl radical-scavenging activity (U/g fw)	42.51 ^f	38.17 ^a	40.75 ^e	38.90 ^b	39.33 ^c	39.97 ^d	38.05 ^a
Total phenol (μ g/g)	6.15 ^a	8.43 ^c	7.68 ^b	9.02 ^d	9.43 ^f	9.18 ^e	9.37 ^f
Total flavonoid (μ g/g)	9.93 ^f	8.15 ^b	7.28 ^a	9.18 ^d	9.37 ^e	8.57 ^c	9.01 ^d

The different superscript letters in the same row indicated significant difference ($P < 0.05$).

The antioxidant activities can be characterized by TEAC, DPPH, FRAP, and NBT radical-scavenging activities. Compared with vacuum precooling, at the end of storage, 1-MCP treated fruit had higher TEAC and DPPH radical-scavenging activities but lower FRAP and NBT radical-scavenging activities. So, it was difficult to compare the effect between 1-MCP and vacuum precooling. For this reason, the PCA model was performed.

3.2. Principal Component Analysis. The parameters in Table 1 were used to develop the PCA model. The first three PCs explained 89% of the variance in the data, which was high enough to represent all the variables. The score plot for PC1 versus PC2 (Figure 1(a)) clearly distinguished three groups defined by length of storage, indicating that storage time had a major influence on the quality and reactive oxygen metabolism parameters of blackberry fruit.

Following PC3, the 1-MCP treated fruit and the vacuum precooled fruit were discriminated (Figure 1(b)). The loading plot of the variables showed that the TEAC and DPPH radical-scavenging activities had a heavy load on the positive coordinate of PC3. So, PC3 could be defined by antioxidant activities. The vacuum precooled fruit had higher positive scores for PC3 than 1-MCP treated fruit. These results suggested that precooled fruit might have values larger than the mean of the antioxidant activities, while 1-MCP treated fruit had relatively lower values.


3.3. Partial Least Squares. H₂O₂ as signal molecule plays an important role inside plant bodies [23]. From the PCA study (Figure 1(a)), H₂O₂ content had a heavy load on the negative coordinate of PC1, suggesting that H₂O₂ content was the most important variable of blackberry fruit. We chose H₂O₂ content as X variable and the other parameters as Y variable

TABLE 2: The result of path analysis taking H₂O₂ as dependent variable.


	GSH	APX	POD	DPPH	Total flavonoid
Direct	0.0441	0.2559	-0.4243	0.6499	-0.1618
→GSH		-0.0307	0.0037	-0.0231	0.0101
→APX	-0.178		-0.116	0.0959	-0.1044
→POD	-0.0359	0.1923		0.0792	-0.2541
→DPPH	-0.3397	0.2436	-0.1213		0.2654
→Total flavonoid	-0.0371	0.066	-0.0969	-0.0661	

to develop the PLS model to obtain a closer understanding of the relation between them. 64% of X variables explained 88% of the variability of Y variables. Many studies have showed that the production of H₂O₂ increased when plants were exposed to various biotic and abiotic stresses [23]. Peroxidase (POD) can decompose H₂O₂ by oxidation of cosubstrates, such as phenolic compounds and antioxidants [24], which could explain why a strongly negative correlation between POD activity and H₂O₂ content was found in the present study (Figure 2). It also showed that TEAC, FRAP, and NBT radical-scavenging activities slightly correlated with H₂O₂ (Figure 2), which indicated that they had little influence on H₂O₂ content.

3.4. Path Analysis. The direct effect of physiological parameters on H₂O₂ content was analyzed by PLS model. To find the indirect factors, the path analysis model was developed. As shown in Table 2, the indirect path coefficient of GSH based on DPPH was -0.3397, which suggested that GSH was also a factor affecting H₂O₂ content.



(a)



(b)

FIGURE 1: Loadings and scores from PCA of blackberries. “c”: the control group; “t1”: treatment of 1-MCP; “t2”: treatment of vacuum precooling; “dr”: decay rate; “cv”: hydroxyl radical-scavenging activity; “tp”: total phenol; “tf”: total flavonoids.




FIGURE 2: Correlation loading plot from a PLSR model. The inner and outer ellipses represent 50% and 100% of explained variance, respectively. “dr”: decay rate; “cv”: hydroxyl radical-scavenging activity; “tp”: total phenol; “tf”: total flavonoids.

4. Conclusion

Based on the results of PCA, vacuum precooling treatment could play a stronger role in keeping the antioxidant activities of blackberry fruit than did the 1-MCP treatment. The score of PCA also revealed that H_2O_2 was the most important variable of blackberry fruit. Results from partial least squares regression and path analysis showed that POD activity had a

direct effect and GSH content had an indirect effect on H_2O_2 content, while TEAC, FRAP, and NBT radical-scavenging activities had little effect on H_2O_2 content.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this article.

Acknowledgments

This research was supported by the National Natural Science Foundation of China (31471626), the National Science & Technology Support Program of China (2015BAD16B06), and the Importation and Development of High-Caliber Talents Project of Beijing Municipal Institutions (CIT&TCD201504008).

References

- [1] J. Gudej and M. Tomczyk, “Determination of flavonoids, tannins and ellagic acid in leaves from *Rubus* L. species,” *Archives of Pharmacological Research*, vol. 27, no. 11, pp. 1114–1119, 2004.
- [2] S. Y. Wang and H. Lin, “Antioxidant activity in fruits and leaves of blackberry, raspberry, and strawberry varies with cultivar and developmental stage,” *Journal of Agricultural and Food Chemistry*, vol. 48, no. 2, pp. 140–146, 2000.
- [3] I. Koca and B. Karadeniz, “Antioxidant properties of blackberry and blueberry fruits grown in the black sea region of Turkey,” *Scientia Horticulturae*, vol. 121, no. 4, pp. 447–450, 2009.
- [4] M. Joo, N. Lewandowski, R. Auras, J. Harte, and E. Almenar, “Comparative shelf life study of blackberry fruit in bio-based and petroleum-based containers under retail storage conditions,” *Food Chemistry*, vol. 126, no. 4, pp. 1734–1740, 2011.
- [5] G. A. Manganaris, I. F. Ilias, M. Vasilakakis, and I. Mignani, “The effect of hydrocooling on ripening related quality attributes and cell wall physicochemical properties of sweet cherry fruit (*Prunus avium* L.),” *International Journal of Refrigeration*, vol. 30, no. 8, pp. 1386–1392, 2007.
- [6] G. Cheng, X. Duan, J. Shi et al., “Effects of reactive oxygen species on cellular wall disassembly of banana fruit during ripening,” *Food Chemistry*, vol. 109, no. 2, pp. 319–324, 2008.
- [7] Y. Imahori, M. Takemura, and J. Bai, “Chilling-induced oxidative stress and antioxidant responses in mume (*Prunus mume*) fruit during low temperature storage,” *Postharvest Biology and Technology*, vol. 49, no. 1, pp. 54–60, 2008.
- [8] D. Inzé and M. V. Montagu, “Oxidative stress in plants,” *Current Opinion in Biotechnology*, vol. 6, no. 2, pp. 153–158, 1995.
- [9] U. Imeh and S. Khokhar, “Distribution of conjugated and free phenols in fruits: antioxidant activity and cultivar variations,” *Journal of Agricultural and Food Chemistry*, vol. 50, no. 22, pp. 6301–6306, 2002.
- [10] Y. Y. Voon, N. Sheikh Abdul Hamid, G. Rusul, A. Osman, and S. Y. Quek, “Volatile flavour compounds and sensory properties of minimally processed durian (*Durio zibethinus* cv. D24) fruit during storage at 4°C,” *Postharvest Biology and Technology*, vol. 46, no. 1, pp. 76–85, 2007.
- [11] H. Zheng and H. Lu, “Use of kinetic, Weibull and PLSR models to predict the retention of ascorbic acid, total phenols and antioxidant activity during storage of pasteurized pineapple juice,” *LWT- Food Science and Technology*, vol. 44, no. 5, pp. 1273–1281, 2011.

- [12] A. Jiménez, J. A. Hernández, L. A. del Río, and F. Sevilla, "Evidence for the presence of the ascorbate-glutathione cycle in mitochondria and peroxisomes of pea leaves," *Plant Physiology*, vol. 114, no. 1, pp. 275–284, 1997.
- [13] A. Jiang, S. P. Tian, and Y. Xu, "Effects of controlled atmospheres with high-O₂ or high CO₂ concentrations on postharvest physiology and storability of "Napoleon" sweet cherry," *Acta Botanica Sinica*, vol. 44, pp. 925–930, 2002.
- [14] A. Guri, "Variation in glutathione and ascorbic acid content among selected cultivars of *Phaseolus vulgaris* prior to and after exposure to ozone," *Canadian Journal of Plant Science*, vol. 63, no. 3, pp. 733–737, 1983.
- [15] T. Brennan and C. Frenkel, "Involvement of Hydrogen Peroxide in the Regulation of Senescence in Pear," *Plant Physiology*, vol. 59, no. 3, pp. 411–416, 1977.
- [16] I. F. F. Benzie and J. J. Strain, "The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay," *Analytical Biochemistry*, vol. 239, no. 1, pp. 70–76, 1996.
- [17] M. J. T. J. Arts, J. S. Dallinga, H.-P. Voss, G. R. M. M. Haenen, and A. Bast, "A critical appraisal of the use of the antioxidant capacity (TEAC) assay in defining optimal antioxidant structures," *Food Chemistry*, vol. 80, no. 3, pp. 409–414, 2003.
- [18] M.-Y. Shon, T.-H. Kim, and N.-J. Sung, "Antioxidants and free radical scavenging activity of *Phellinus baumii* (Phellinus of Hymenochaetaceae) extracts," *Food Chemistry*, vol. 82, no. 4, pp. 593–597, 2003.
- [19] C. Larrigaudière, A. P. Candan, D. Ubach, and J. Graell, "Physiological response of "Larry Ann" plums to cold storage and 1-MCP treatment," *Postharvest Biology and Technology*, vol. 51, no. 1, pp. 56–61, 2009.
- [20] F. Bamdad, M. Kadivar, and J. Keramat, "Evaluation of phenolic content and antioxidant activity of Iranian caraway in comparison with clove and BHT using model systems and vegetable oil," *International Journal of Food Science & Technology*, vol. 41, no. 1, pp. 20–27, 2006.
- [21] R. A. Spotts, P. L. Sholberg, P. Randall, M. Serdani, and P. M. Chen, "Effects of 1-MCP and hexanal on decay of d'Anjou pear fruit in long-term cold storage," *Postharvest Biology and Technology*, vol. 44, no. 2, pp. 101–106, 2007.
- [22] R. Zhou, Y. Li, L. Yan, and J. Xie, "Effect of edible coatings on enzymes, cell-membrane integrity, and cell-wall constituents in relation to brittleness and firmness of Huanghua pears (*Pyrus pyrifolia* Nakai, cv. Huanghua) during storage," *Food Chemistry*, vol. 124, no. 2, pp. 569–575, 2011.
- [23] S. Navabpour, K. Morris, R. Allen, E. Harrison, S. A-H-Mackerness, and V. Buchanan-Wollaston, "Expression of senescence-enhanced genes in response to oxidative stress," *Journal of Experimental Botany*, vol. 54, no. 391, pp. 2285–2292, 2003.
- [24] Y. F. Xue and Z. P. Liu, "Antioxidant enzymes and physiological characteristics in two Jerusalem artichoke cultivars under salt stress," *Russian Journal of Plant Physiology*, vol. 55, no. 6, pp. 776–781, 2008.



Hindawi

Submit your manuscripts at
www.hindawi.com

