




Supplementary Figure 1 Dose and exposure relationship over the single-ascending dose range of 10 to 50 mg in healthy Chinese subjects. Filled circles and lines in orange indicate the theoretical exposure according to a proportional increase along with the dose levels, and those in blue indicate the actual exposure after single-ascending doses.



Supplementary Figure 2 Dose and exposure relationship over the single- and multiple-ascending dose range of 15 to 180 (90) mg in Japanese healthy subjects from study D5611C00005 (NCT02176252). Filled circles and lines in orange indicate the theoretical exposure according to a proportional increase along with the dose levels, and those in blue indicate the actual exposure under single-ascending doses.





Supplementary Table 1 Comparison of PK parameters of tenapanor-M1 in healthy Chinese and Caucasian subjects receiving 50 mg single and repeated dose of tenapanor.

	Geometric mean and ratios			90% CI of Chinese/Caucasian
	Chinese	Caucasian	Chinese/Caucasian	
50 mg-Single dose				
C _{max} (ng/mL)	5.52 (n=12)	5.25 (n=11)	1.05	0.78-1.41
AUC _{0-t} (hour*ng/mL)	169.70 (n=12)	129.42 (n=11)	1.31	0.92-1.87
AUC _{0-∞} (hour*ng/mL)	195.09 (n=12)	172.30 (n=9)	1.13	0.84-1.53
50 mg-Repeated dose				
C _{ss} (ng/mL)	13.46 (n=12)	12.16 (n=11)	1.11	0.87-1.41
C _{max_ss} (ng/mL)	19.18 (n=12)	17.60 (n=11)	1.09	0.88-1.34
C _{min_ss} (ng/mL)	13.34 (n=12)	12.16 (n=11)	1.10	0.86-1.39
C _{av_ss} (ng/mL)	16.81 (n=12)	15.21 (n=11)	1.10	0.89-1.37
AUC _{0-τ} (hour*ng/mL)	201.71 (n=12)	182.57 (n=11)	1.10	0.89-1.37

AUC area under curve, CI confidence interval